

RW-CLOSED MAPS AND RW-OPEN MAPS IN TOPOLOGICAL SPACES

M.Karpagadevi ,
Karpagam College of Engineering,
Coimbatore, India

A.Pushpalatha
Government Arts College,
Udumalpet, India

Abstract:In this paper we introduce rw-closed map from a topological space X to a topological space Y as the image of every closed set is rw-closed and also we prove that the composition of two rw-closed maps need not be rw-closed map. We also obtain some properties of rw-closed maps.

Mathematics Subject Classification: 54C10

Keywords: rw-closed maps, rw-open maps.

1. INTRODUCTION

Generalized closed mappings were introduced and studied by Malghan[5].wg-closed maps and rwg-closed maps were introduced and studied by Nagaveni[6].Regular closed maps,gpr-closed maps and rg-closed maps have been introduced and studied by Long[4], Gnanambal[3] and Arockiarani[1] respectively.

In this paper, a new class of maps called regular weakly closed maps (briefly, rw-closed) maps have been introduced and studied their relations with various generalized closed maps. We prove that the composition of two rw-closed maps need not be rw-closed map. We also obtain some properties of rw-closed maps.

S.S. Benchalli and R.S Wali [2] introduced new class of sets called regular weakly - closed (briefly rw - closed) sets in topological spaces which lies between the class of all w - closed sets and the class of all regular g - closed sets.

Throughout this paper (X, τ) and (Y, σ) (or simply X and Y) represents the non-empty topological spaces on which no separation axiom are assumed, unless otherwise mentioned. For a subset A of X , $cl(A)$ and $int(A)$ represents the closure of A and interior of A respectively.

2. PRELIMINARIES

In this section we recollect the following basic definitions which are used in this paper.

Definition 2.1 [2]: A subset A of a topological space (X, τ) is called rw-closed (briefly rw-closed) if $cl(A) \subseteq U$, whenever $A \subseteq U$ and U is regular semiopen in X .

Definition 2.2 [7]: A subset A of a topological space (X, τ) is called regular generalized closed (briefly rg-closed) if $cl(A) \subseteq U$ whenever $A \subseteq U$ and U is regular open in X .

Definition 2.3 [9]: A subset A of a topological space (X, τ) is called weakly closed (briefly w-closed) if $cl(A) \subseteq U$ whenever $A \subseteq U$ and U is semi open in X .

Definition 2.4 [7] :A map $f: (X, \tau) \rightarrow (Y, \sigma)$ from a topological space X into a topological space Y is called rg continuous if the inverse image of every closed set in Y is rg-closed in X .

Definition 2.5 [9] :A map $f: (X, \tau) \rightarrow (Y, \sigma)$ from a topological space X into a topological space Y is called w-continuous if the inverse image of every closed set in Y is w-closed in X .

Definition 2.6 [5]: A map $f: (X, \tau) \rightarrow (Y, \sigma)$ is called g-closed if $f(F)$ is g-closed in (Y, σ) for every closed set F of (X, τ) .

Definition 2.7 [8]: A map $f: (X, \tau) \rightarrow (Y, \sigma)$ is called w-closed if $f(F)$ is w-closed in (Y, σ) for every closed set F of (X, τ) .

Definition 2.8 [1]:A map $f: (X, \tau) \rightarrow (Y, \sigma)$ is called rg-closed if $f(F)$ is rg-closed in (Y, σ) for every closed set F of (X, τ) .

Definition 2.9 [10]:A map $f: (X, \tau) \rightarrow (Y, \sigma)$ is called g-open if $f(U)$ is g-open in (Y, σ) for every open set U of (X, τ) .

Definition 2.10 [8]:A map $f: (X, \tau) \rightarrow (Y, \sigma)$ is called w-open if $f(U)$ w-open in (Y, σ) for every open set U of (X, τ) .

Definition 2.11[1]:A map $f: (X, \tau) \rightarrow (Y, \sigma)$ is called rg-open if $f(U)$ rg-open in (Y, σ) for every open set U of (X, τ) .

3. Rw-closed maps

We introduce the following definition

Definition : 3.1 A map $f : (X, \tau) \rightarrow (Y, \sigma)$ is said to be regular weakly (briefly rw-closed) if the image of every closed set in (X, τ) is rw-closed in (Y, σ)

Theorem: 3.2 Every closed map is rw-closed map but not conversely.

Proof: The proof follows from the definitions and fact that every closed set is rw-closed.

Remark: 3.3 The converse of the above theorem need not be true as seen from the following example.

Example : 3.4 Consider $X=Y=\{a,b,c\}$ with topologies $\tau = \{X, \phi, \{c\}\}$ and $\sigma = \{Y, \phi, \{a\}, \{b\}, \{a,b\}\}$. Let $f : (X, \tau) \rightarrow (Y, \sigma)$ be the identity map. Then this function is rw-closed but not closed as the image of closed set $\{a,b\}$ in X is $\{a,b\}$ which is not closed set in Y .

Theorem: 3.5 Every rw-closed map is rg-closed map but not conversely.

Proof: The proof follows from the definitions and fact that every rw-closed set is rg-closed.

Remark: 3.6 The converse of the above theorem need not be true as seen from the following example.

Example : 3.7 Consider $X=Y=\{a,b,c,d\}$ with topologies $\tau = \{X, \phi, \{b\}, \{a,b,d\}\}$ and $\sigma = \{Y, \phi, \{a\}, \{b\}, \{a,b\}, \{a,b,c\}\}$. Let $f : (X, \tau) \rightarrow (Y, \sigma)$ be the identity map. Then this function is rg-closed but not rw-closed as the image of closed set $\{c\}$ in X is $\{c\}$ which is not rw-closed set in Y .

Theorem: 3.8 Every w-closed map is rw-closed map but not conversely.

Proof: The proof follows from the definitions and fact that every w-closed set is rw-closed.

Remark: 3.9 The converse of the above theorem need not be true as seen from the following example.

Example: 3.10 Consider $X=Y=\{a,b,c\}$ with topologies $\tau = \{X, \phi, \{c\}\}$ and $\sigma = \{Y, \phi, \{a\}, \{b\}, \{a,b\}\}$. Let $f : (X, \tau) \rightarrow (Y, \sigma)$ be the identity map. Then this function is rw-closed but not w-closed as the image of closed set $\{a,b\}$ in X is $\{a,b\}$ which is not closed set in Y .

Theorem: 3.11 A map $f : (X, \tau) \rightarrow (Y, \sigma)$ is rw-closed if and only if for each subset S of (Y, σ) and each open set U containing $f^{-1}(S) \subset U$, there is a rw-open set of (Y, σ) such that $S \subset V$ and $f^{-1}(V) \subset U$.

Proof: Suppose f is rw-closed. Let $S \subset Y$ and U be an open set of (X, τ) such that $f^{-1}(S) \subset U$. Now $X-U$ is closed set in (X, τ) . Since f is rw-closed, $f(X-U)$ is rw-closed set in (Y, σ) . Then $V=Y-f(X-U)$ is a rw-open set in (Y, σ) . Note that $f^{-1}(S) \subset U$ implies $S \subset V$ and $f^{-1}(V) = X-f^{-1}(f(X-U)) \subset X-(X-U) = U$. That is $f^{-1}(V) \subset U$.

For the converse, let F be a closed set of (X, τ) . Then $f^{-1}(f(F)^c) \subset F^c$ and F^c is an open set in (X, τ) . By hypothesis, there exists a rw-open set V in

(Y, σ) such that $f(F)^c \subset V$ and $f^{-1}(V) \subset F^c$ and so $F \subset (f^{-1}(V))^c$. Hence $V^c \subset f(F) \subset f((f^{-1}(V))^c) \subset V^c$ which implies $f(V) \subset V^c$. Since V^c is rw-closed, $f(F)$ is rw-closed. That is $f(F)$ rw-closed in (Y, σ) and therefore f is rw-closed.

Remark: 3.12 The composition of two rw-closed maps need not be rw-closed map in general and this is shown by the following example.

Example: 3.13 Consider $X=Y=\{a,b,c\}$ with topologies $\tau = \{X, \phi, \{b\}, \{a,b\}\}$, $\sigma = \{Y, \phi, \{a\}, \{b\}, \{a,b\}\}$, $\eta = \{Z, \phi, \{a\}, \{c\}, \{a,c\}\}$. Define $f : (X, \tau) \rightarrow (Y, \sigma)$ by $f(a) = a, f(b) = b$ and $f(c) = c$ and $g : (Y, \sigma) \rightarrow (Z, \eta)$ be the identity map. Then f and g are rw-closed maps but their composition $g \circ f : (X, \tau) \rightarrow (Z, \eta)$ is not rw-closed map because $F = \{c\}$ is closed in (X, τ) but $g \circ f(\{a\}) = g(f(\{c\})) = g(\{c\}) = \{c\}$ which is not rw-closed in (Z, η) .

Theorem: 3.14 If $f : (X, \tau) \rightarrow (Y, \sigma)$ is closed map and $g : (Y, \sigma) \rightarrow (Z, \eta)$ is rw-closed map, then the composition $g \circ f : (X, \tau) \rightarrow (Z, \eta)$ is rw-closed map.

Proof: Let F be any closed set in (X, τ) . Since f is closed map, $f(F)$ is closed set in (Y, σ) . Since g is rw-closed map, $g(f(F))$ is rw-closed set in (Z, η) . That is $g \circ f(F) = g(f(F))$ is rw-closed and hence $g \circ f$ is rw-closed map.

Remark: 3.15 If $f : (X, \tau) \rightarrow (Y, \sigma)$ is rw-closed map and $g : (Y, \sigma) \rightarrow (Z, \eta)$ is closed map, then the composition need not be rw-closed map as seen from the following example.

Example: 3.16 Consider $X=Y=Z=\{a,b,c\}$ with topologies $\tau = \{X, \phi, \{b\}, \{a,b\}\}$, $\sigma = \{Y, \phi, \{a\}, \{b\}, \{a,b\}\}$, $\eta = \{Z, \phi, \{a\}, \{c\}, \{a,c\}\}$. Define $f : (X, \tau) \rightarrow (Y, \sigma)$ by $f(a) = a, f(b) = b$ and $f(c) = c$ and $g : (Y, \sigma) \rightarrow (Z, \eta)$ be the identity map. Then f is rw-closed map and g is a closed map but their composition $g \circ f : (X, \tau) \rightarrow (Z, \eta)$ is not rw-closed map since for the closed set $\{c\}$ in (X, τ) but $g \circ f(\{c\}) = g(f(\{c\})) = g(\{c\}) = \{c\}$ which is not rw-closed in (Z, η) .

Theorem: 3.17 Let $(X, \tau), (Z, \eta)$ be topological spaces and (Y, σ) be topological space where every rw-closed subset is closed. Then the composition $g \circ f : (X, \tau) \rightarrow (Z, \eta)$ of the rw-closed maps $f : (X, \tau) \rightarrow (Y, \sigma)$ and $g : (Y, \sigma) \rightarrow (Z, \eta)$ is rw-closed.

Proof: Let A be a closed set of (X, τ) . Since f is rw-closed, $f(A)$ is rw-closed in (Y, σ) . Then by hypothesis $f(A)$ is closed. Since g is rw-closed, $g(f(A))$ is rw-closed in (Z, η) and $g(f(A)) = g \circ f(A)$. Therefore $g \circ f$ is rw-closed.

Theorem: 3.18 If $f : (X, \tau) \rightarrow (Y, \sigma)$ is g-closed, $g : (Y, \sigma) \rightarrow (Z, \eta)$ be rw-closed and (Y, σ) is $T_{1/2}$ -space then their composition $g \circ f : (X, \tau) \rightarrow (Z, \eta)$ is rw-closed map.

Proof: Let A be a closed set of (X, τ) . Since f is g-closed, $f(A)$ is g-closed in (Y, σ) . Since g is rw-closed, $g(f(A))$ is rw-closed in (Z, η) and $g(f(A)) = g \circ f(A)$. Therefore $g \circ f$ is rw-closed.

Theorem: 3.19 Let $f : (X, \tau) \rightarrow (Y, \sigma)$ and $g : (Y, \sigma) \rightarrow (Z, \eta)$ be two mappings such that their

composition $g \circ f: (X, \tau) \rightarrow (Z, \eta)$ be rw-closed mapping. Then the following statements are true.

- i) If f is continuous and surjective, then g is rw-closed
- ii) If g is rw-irresolute and injective, then f is rw-closed.
- iii) If f is g -continuous, surjective and (X, τ) is a $T_{1/2}$ - space, then g is rw-closed.

Proof: i) Let A be a closed set of (Y, σ) . Since f is continuous, $f^{-1}(A)$ is closed in (X, τ) . $g \circ f (f^{-1}(A))$ is rw-closed in (Z, η) . That is $g (A)$ is rw-closed in (Z, η) , since f is surjective. Therefore g is rw-closed.

ii) Let B be a closed set of (X, τ) . Since $g \circ f$ is rw-closed, $g \circ f (B)$ is rw-closed in (Z, η) .

Since g is rw-irresolute, $g^{-1}(g \circ f (B))$ is rw-closed set in (Y, σ) . That is $f (B)$ is rw-closed in (Y, σ) , since f is injective. Therefore f is rw-closed.

iii) Let c be a closed set of (Y, σ) . Since f is g -continuous, $f^{-1}(c)$ is g -closed set in (X, τ) . Since (X, τ) is a $T_{1/2}$ -space, $f^{-1}(c)$ is closed set in (X, τ) . Since $g \circ f$ is rw-closed $(g \circ f) (f^{-1}(c))$ is rw-closed in (Z, η) . That is $g(c)$ is rw-closed in (Z, η) , since f is surjective. Therefore g is rw-closed.

4. Rw-open maps

Definition: 4.1 A map $f: (X, \tau) \rightarrow (Y, \sigma)$ is called a rw-open map if the image $f(A)$ is rw-open in (Y, σ) for each open set A in (X, τ)

Theorem: 4.2 For any bijection map $f: (X, \tau) \rightarrow (Y, \sigma)$ the following statements are equivalent.

- i) $f^{-1}: (Y, \sigma) \rightarrow (X, \tau)$ is rw-continuous
- ii) f is rw-open map and
- iii) f is rw-closed map.

Proof: (i) \Rightarrow (ii) Let U be an open set of (X, τ) . By assumption, $(f^{-1})^{-1}(U) = f(U)$ is rw-open in (Y, σ) and so f is rw-open.

(ii) \Rightarrow (iii) Let F be a closed set of (X, τ) . Then F^c is open set in (X, τ) . By assumption $f(F^c)$ is rw-open in (Y, σ) . That is $f(F^c) = f(F)^c$ is rw-open in (Y, σ) and therefore $f(F)$ is rw-closed in (Y, σ) . Hence F is rw-closed.

(iii) \Rightarrow (i) Let F be a closed set of (X, τ) . By assumption, $f(F)$ is rw-closed in (Y, σ) . But $f(F) = (f^{-1})^{-1}(F)$ and therefore f^{-1} is continuous.

Theorem: 4.3 A map $f: (X, \tau) \rightarrow (Y, \sigma)$ is rw-open if and only if for any subset S of (Y, σ) and any closed set of (X, τ) containing $f^{-1}(S)$, there exists a rw-closed set K of (Y, σ) containing S such that $f^{-1}(K) \subset F$.

Proof: Suppose f is rw-open map. Let $S \subset Y$ and F be a closed set of (X, τ) such that $f^{-1}(S) \subset F$. Now $X-F$ is an open set in (X, τ) . Since f is rw-open map, $f(X-F)$ is rw-open set in (Y, σ) . Then $K=Y- f(X-F)$ is a rw-closed set in (Y, σ) . Note that $f^{-1}(S) \subset F$ implies $S \subset K$ and $f^{-1}(K) = X- f^{-1}(X-F) \subset X-(X-F) = F$. That is $f^{-1}(K) \subset F$.

For the converse let U be an open set of (X, τ) . Then $f^{-1}((f(U))^c) \subset U^c$ and U^c is a closed set in

(X, τ) . By hypothesis, there exists a rw-closed set K of (Y, σ) such that $(f(U))^c \subset K$ and $f^{-1}(K) \subset U^c$ and so $U \subset (f^{-1}(K))^c$. Hence $K^c \subset f(U) \subset f((f^{-1}(K))^c)$ which implies $f(U) = K^c$. Since K^c is a rw-open, $f(U)$ is rw-open in (Y, σ) and therefore f is rw-open map.

4. REFERENCES

- [1] Arockiarani. I, Studies on generalizations of generalized closed sets and maps in topological spaces, Ph.D., Thesis, Bharathiar Univ., Coimbatore, 1997.
- [2] Benchalli.S.S and Wali.R.S., On $R\omega$ -closed sets in topological spaces, Bull.Malays.math.Sci.Soc(2) 30(2) (2007), 99-110.
- [3] Gnanambal Y, On Generalized Pre-regular Closed sets in Topological Spaces, Indian J.Pure Appl.Math.,28(1997), 351-360.
- [4] Long P.E, and Herington L.L, Basic properties of Regular Closed Functions, Rend.Cir.Mat.Palermo, 27(1978) , 20-28.
- [5] Malghan S.R, Generalized Closed maps, J.Karnatk Univ.Sci.,27(1982), 82-88
- [6] Nagaveni N, Studies on Generalizations of Homeomorphisms in Topological spaces, Ph.D., Thesis, Bharathiar University, Coimbatore(1999).
- [7] Palaniappan N and Rao K C, Regular generalized closed sets, Kyungpook Math. J. 33(1993), 211-219
- [8] Sheik John M, A Study on Generalizations of Closed Sets on Continuous maps in Topological and Bitopological Spaces, Ph.D, Thesis Bharathiar University, Coimbatore,(2002)
- [9] Sundaram P and Sheik John M, On w-closed sets in topology, Acta Ciencia Indica 4(2000), 389-392
- [10] Sundaram P, Studies on Generalizations of Continuous Maps in Topological Spaces, Ph.D, Thesis, Bharathiar University, Coimbatore,(1991).
- [11] Vadivel A and Vairamanickam K, $rg\alpha$ -Closed Sets and $rg\alpha$ -Open Sets in Topological spaces, Int.Journal of Math.Analysis, Vol.3, 2009, no.37, 1803-1819.
- [12] Vadivel A and Vairamanickam K, $rg\alpha$ -Closed and $rg\alpha$ -Open Maps in Topological spaces, Int.Journal of Math.Analysis, Vol.4, 2010, no.10, 453-468