

Dartmouth College

Dartmouth Digital Commons

Dartmouth Scholarship

Faculty Work

7-2006

Saccharomyces Cerevisiae-Based Molecular Tool Kit for Manipulation of Genes from Gram-Negative Bacteria

Robert M. Q. Shanks
Dartmouth College

Nicky C. Caiazza
Dartmouth College

Shannon M. Hinsa
Dartmouth College

Christine M. Toutain
Dartmouth College

George A. O'Toole
Dartmouth College

Follow this and additional works at: <https://digitalcommons.dartmouth.edu/facoa>



Part of the [Biotechnology Commons](#), [Environmental Microbiology and Microbial Ecology Commons](#), and the [Genetics Commons](#)

Dartmouth Digital Commons Citation

Shanks, Robert M. Q.; Caiazza, Nicky C.; Hinsa, Shannon M.; Toutain, Christine M.; and O'Toole, George A., "Saccharomyces Cerevisiae-Based Molecular Tool Kit for Manipulation of Genes from Gram-Negative Bacteria" (2006). *Dartmouth Scholarship*. 490.
<https://digitalcommons.dartmouth.edu/facoa/490>

This Article is brought to you for free and open access by the Faculty Work at Dartmouth Digital Commons. It has been accepted for inclusion in Dartmouth Scholarship by an authorized administrator of Dartmouth Digital Commons. For more information, please contact dartmouthdigitalcommons@groups.dartmouth.edu.

Saccharomyces cerevisiae-Based Molecular Tool Kit for Manipulation of Genes from Gram-Negative Bacteria†

Robert M. Q. Shanks, Nicky C. Caiazza,‡ Shannon M. Hinsla,§
Christine M. Toutain, and George A. O'Toole*

Department of Microbiology and Immunology, Dartmouth Medical School, Hanover, New Hampshire 03755

Received 24 March 2006/Accepted 3 May 2006

A tool kit of vectors was designed to manipulate and express genes from a wide range of gram-negative species by using in vivo recombination. *Saccharomyces cerevisiae* can use its native recombination proteins to combine several amplicons in a single transformation step with high efficiency. We show that this technology is particularly useful for vector design. Shuttle, suicide, and expression vectors useful in a diverse group of bacteria are described and utilized. This report describes the use of these vectors to mutate *clpX* and *clpP* of the opportunistic pathogen *Pseudomonas aeruginosa* and to explore their roles in biofilm formation and surface motility. Complementation of the rhamnolipid biosynthetic gene *rhlB* is also described. Expression vectors are used for controlled expression of genes in two pseudomonad species. To demonstrate the facility of building complicated constructs with this technique, the recombination of four PCR-generated amplicons in a single step at >80% efficiency into one of these vectors is shown. These tools can be used for genetic studies of pseudomonads and many other gram-negative bacteria.

Cloning using *Saccharomyces cerevisiae* homologous recombination is a powerful technique in which PCR primers with stretches of homologous DNA are used to target recombination of an amplicon with a vector (9, 21, 25, 27). Cloning of an amplicon with yeast recombination into an intact vector can be done without restriction enzymes, but this requires the use of a selectable or counterselectable marker. The formation of a gap or double-stranded break in the vector by restriction enzyme digestion allows selective cloning of unmarked amplicons (29). An amplicon can seamlessly replace any part of the DNA in a vector with no need for enzyme sites at the junctions of the recombined pieces of DNA, as long as the yeast replication machinery is not replaced and the ends of the amplicons (or other DNA with homologous ends) to be added to a vector are on either side of the digest site (gap) in the vector. Therefore, multiple unmarked pieces of DNA can efficiently be assembled together in a single step using this method. In addition to fusions, directed mutations, restriction sites, and other changes can be designed into the primers. This technique is particularly useful for building vectors and complex constructs.

The basic concept of cloning with gap repair is as follows. A vector able to replicate in yeast with a selective marker, such as *URA3*, is digested (gapped) with a restriction enzyme. Primers that amplify the region of DNA that one wants to insert into the vector are designed, containing sequences of homology to the yeast-replicating plasmid (30 to 40 bp in length). One or

more targeted amplicons and the gapped vector are simultaneously introduced into yeast cells. A subset of these cells recombines the amplicon(s) and vector via *Saccharomyces cerevisiae* native recombination enzymes, generating stable circularized plasmids. Linearized or gapped vectors are not stable in yeast, while vectors that circularize through homologous recombination are stable. Cells containing circularized plasmids can be selected for by being plated on medium selective for the vector backbone (e.g., deficient in uracil).

We present a tool kit of vectors designed to manipulate and express genes from a wide range of gram-negative species by using *S. cerevisiae* in vivo recombination.

MATERIALS AND METHODS

Strains and growth conditions. Strains and plasmids used in this study are listed in Table 1. *Escherichia coli* strain DH5 α was used as a host strain for plasmid construction and propagation, while S17-1 was used for conjugations with *Pseudomonas* species (37). *Pseudomonas aeruginosa* strain PA14 and *Pseudomonas fluorescens* strain WCS365 were used for phenotypic experiments (13, 31). *S. cerevisiae* strains DC49-7.1c and InvScl (Invitrogen Corporation) were used interchangeably for the in vivo homologous recombination used to modify plasmids (1). *E. coli* and *Pseudomonas* strains were grown with LB medium, *S. cerevisiae* was grown with yeast extract-peptone-dextrose (1% Bacto yeast extract, 2% Bacto peptone, and 2% dextrose), and selections were performed with synthetic defined agar-uracil (Obiogene 4813-065). *E. coli* was transformed by electroporation, whereas *Pseudomonas* was transformed by conjugation or electroporation (5). Ampicillin at 75 μ g/ml and 150 μ g/ml was used for plasmid selection and propagation, respectively, for *E. coli* or 500 to 1,000 μ g/ml for *P. aeruginosa*. Gentamicin was used at 10 μ g/ml for *E. coli* and 50 to 100 μ g/ml for *Pseudomonas* strains. Kanamycin was used at 50 μ g/ml for *E. coli* and 150 to 250 μ g/ml for *P. fluorescens*. Tetracycline was used at 15 μ g/ml for *E. coli* and 75 to 150 μ g/ml for *Pseudomonas* species. Nalidixic acid was used at 20 μ g/ml for selection against *E. coli* following *E. coli*-*Pseudomonas* sp. conjugations.

Plasmid construction. All vectors, with the exception of pMQ87, pMQ89, pMQ57, pMQ65, and pMQ90, were built using yeast gap repair as follows. A yeast-replicating vector to be modified was cut with a restriction enzyme(s). Primers were made to amplify DNA to be joined with the cut vector, including ~40-bp tails that are homologous to the cut vector on each side of the restriction enzyme cut site (up to 2,000 bp from the cut site). A small sample of the amplicon was run out on a gel to ensure that the desired band was in molar excess to any other

* Corresponding author. Mailing address: Department of Microbiology and Immunology, Dartmouth Medical School, Hanover, NH 03755. Phone: (603) 650-1248. Fax: (603) 650-1245. E-mail: georgeo@Dartmouth.edu.

† Supplemental material for this article may be found at <http://aem.asm.org/>.

‡ Present address: Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena, CA 91125.

§ Present address: Center for Microbial Ecology, Michigan State University, East Lansing, MI 48824.

TABLE 1. Strains and plasmids used in this study

Strain or plasmid	Genotype or description	Source or reference
Strains		
<i>Escherichia coli</i>		
DH5 α	<i>supE44 ΔlacU169(ϕ80dlacZΔM15) hsdR17 thi-1 relA1 recA1</i>	Life Technologies
S17-1	<i>thi pro hsdR hsdM⁺ ΔrecA RP4-2::TcMu-Km::Tn</i>	37
<i>P. aeruginosa</i>		
PA14	Wild type, clinical isolate	31
RSS255	PA14 <i>clpX</i> ::pRMQS84	This study
EP1	PA14 <i>clpP</i> :: <i>GFP_{uvr}</i>	This study
SMC2758	PA14 <i>rhlB</i> :: <i>kanMX4</i>	4
<i>P. fluorescens</i>		
WCS365	Wild type, environmental isolate	13
SMC2836	WCS365 Δ <i>lapD</i>	17
<i>S. cerevisiae</i>		
DC49-7.1c	<i>MATa leu2-3,112 trp1-289 ura3-52 arg4-Δ57,RV</i>	1
InvSc1	<i>MATa/MATα leu2/leu2 trp1-289/trp1-289 ura3-52/ura3-52 his3-Δ1/his3-Δ1</i>	Invitrogen
Plasmids		
pBAD18	<i>E. coli</i> expression vector with <i>P_{BAD}</i> and <i>araC</i>	15
Mini-CTX	CTX phage neutral site integration vector	19
pEPSA5	p15a containing <i>P_{T5X}</i> expression vector	11
pEX18-Gm	Gram-negative allelic replacement vector	18
pGFPmut3	<i>E. coli</i> vector with mutated <i>GFP</i> allele	6
pJB785TTKml	RK2 broad-host-range reporter construct	34
pRS415	<i>S. cerevisiae-E. coli</i> shuttle vector	36
pRS416	<i>S. cerevisiae-E. coli</i> shuttle vector	36
pRS426	<i>S. cerevisiae-E. coli</i> shuttle vector	36
pUCP20	<i>Pseudomonas-E. coli</i> shuttle vector	39
pUG6	<i>loxP-kanMX-loxP</i> construct	14
pYC2-CT	<i>S. cerevisiae-E. coli</i> shuttle vector	Invitrogen

bands, and if it was not, the desired band was gel purified. Overnight cultures of *S. cerevisiae* were cotransformed by the addition of 20 to 200 ng of the cut vector (directly from the restriction enzyme mix) and 50 to 500 ng of the amplicon. These modified vectors all contained a *URA3* gene capable of complementing the *ura3-52* mutation of the strains used. Transformed cells were either tested individually for the desired construct with colony PCR or pooled, and a preparation of plasmid DNA was made and moved into *E. coli* cells for screening.

Construction of pMQ30. Plasmid pMQ30 is derived from pEX18-Gm, a vector used for two-step gene replacement experiments with *Pseudomonas* species (18), onto which were added elements needed for replication and selection in *S. cerevisiae*. pMQ30 was made by homologous recombination in yeast (see below for protocol); a 1,633-bp PCR amplicon containing the selectable and counter-selectable yeast gene *URA3* and the *CEN6* and *ARSH4* elements was amplified from pYC2-CT (Invitrogen) and recombined with pEX18-Gm that had been linearized with AatII. Both primers had regions homologous to pYC2-CT and ~40-bp tails, identical to regions of DNA flanking the AatII site of pEX18-Gm. See Table S1 in the supplemental material for primer sequences (primers Ex18-Up and Ex18-Dn).

S. cerevisiae strain DC49-7.1c (*ura3-52*) was cotransformed with both linear pieces of DNA, and recombinants were selected on uracil dropout medium. Because the *CEN6-URA3* amplicon is linear and without telomeres, it is not stably maintained in the yeast cell, but cells in which the linear *CEN6-URA3* amplicon and the digested pEX18-Gm had recombined to form a circular stable replicon were able to grow on selective medium. Plasmids were liberated from uracil-prototrophic yeast and electroporated into *E. coli*, and gentamicin-resistant transformants were selected. Candidates were isolated and screened by restriction analysis and phenotype (uracil prototroph in yeast, gentamicin resistance, and sucrose sensitivity [data not shown]). All of the candidates were verified as correct by restriction analysis, PCR, and phenotypic assays.


Construction of pMQ87 and pMQ89. pMQ30 was chosen as a backbone for suicide vectors. The *sacB* gene was removed to enhance stability of single-crossover constructs as follows. pMQ30 was gapped via digestion with MslI and

MscI, whose restriction sites are both in the *sacB* open reading frame (ORF). We then amplified an internal fragment from the *clpX* gene of *P. aeruginosa* strain PA14 by using primers that target the amplicon to recombine with pMQ30 on either side of *sacB* and replacing *sacB* and some dispensable vector sequence flanking *sacB*. The primers each contained an XhoI site so that the *clpX* sequence could be removed completely following digestion with XhoI and subsequent unimolecular ligation. We also included a StuI site proximal to the yeast replication machinery so that it could be removed by digestion with StuI to generate a very small suicide vector not capable of replication in yeast. Thus, we made pRMQS84, which is pMQ30 with *sacB* replaced by a *P. aeruginosa clpX* internal sequence; pMQ87, where the entire *clpX* sequence had been removed through digestion with XhoI; and pMQ89, which had been digested with StuI to remove the yeast replicon and selectable marker (Fig. 1).

Construction of shuttle vectors pMQ51, pMQ52, and pMQ61. Homologous recombination was used to add the pRO1600 replicon to pRS416 and pRS426, resulting in plasmids pV204 and pV204-2 μ m, respectively. We were unable to transform *Pseudomonas aeruginosa* using these plasmids, although we were able to do so with a positive-control pUCP20 with the same replication machinery (39). The *URA3* gene was proximal to the *Pseudomonas* origin, and we hypothesized that transcriptional read-through from this gene in *Pseudomonas* could inhibit replication. Therefore, we added a kanamycin resistance cassette between the *URA3* gene and the pRO1600 origin by using yeast homologous recombination. The kanamycin cassette from pUG6 was chosen rather than a transcriptional terminator because the addition of a selectable marker broadened the usefulness of these plasmids as well as aided in recovery of the correct construct (14). The resulting plasmids, pMQ51 (2 μ m) and pMQ52 (*CEN6*), were able to replicate in both *P. aeruginosa* and *P. fluorescens* (Fig. 1). The kanamycin resistance cassette from pUG6 is useful for conferring antibiotic resistance in *E. coli* and *S. cerevisiae*, where it confers resistance to kanamycin and G418 sulfate, respectively, and *P. fluorescens*, but not in *P. aeruginosa*, because of *P. aeruginosa*'s intrinsic high level of kanamycin resistance.

To generate pMQ61, pMQ51 was digested in the ampicillin resistance deter-


Allelic Replacement Vectors



Suicide Vector



Shuttle Vectors



Expression Vectors




FIG. 1. Tool kit for use with genes from gram-negative bacteria. Shown are maps of vectors for manipulating bacterial genes by using *S. cerevisiae*-based homologous recombination. Vector maps are shown as linearized at the first base pair and are not shown to scale. Constructs are clustered based upon suggested utility. *sacB*, *Bacillus subtilis* levansucrase gene for counterselection; *oriT*, origin of conjugal transfer; T1T2, *E. coli* *rmB* transcriptional terminators; *lacZα*, *lacZα* with multicloning site driven by the lactose promoter; *ColE1*, high-copy-number variants of the narrow-host-range *ColE1* origin of replication; *p15a*, narrow-host-range, low-copy origin of replication; *aacC1*, gentamicin resistance determinant from Tn1696; *URA3*, orotidine-5'-phosphate decarboxylase gene from *S. cerevisiae*; *CEN/ARSH*, low-copy yeast replication and segregation machinery; *kanMX*, kanamycin/G418 resistance cassette from pUG6; *P_{oriV}-rep*, pRO1600 broad-host-range replicon; *bla*, ampicillin and carbenicillin resistance determinant; *2μm*, yeast episomal plasmid replicon; *P_{BAD}-araC*, arabinose-inducible promoter system from *E. coli*; *RK2-trfA*, minimal replicon from the extremely-broad-host-range RK2 plasmid; *GFP*, bright and stable GFPmut3 variant. Plasmid request information can be found at <http://www.dartmouth.edu/~gotoole/vectors.html>.

minant with BcgI and a fragment 644 bp upstream and 234 bp downstream of the *bla* gene was replaced with a gentamicin resistance-conferring gene (*aacC1*) from pEX18-Gm (Fig. 1).

Construction of conjugal shuttle vectors pMQ56, pMQ57, pMQ64, and pMQ65. pMQ51 and pMQ61 were digested with MluI and modified with an *oriT*-containing amplicon with 40-bp tails that target replacement of the *kanMX* cassette by using yeast homologous recombination. The primers used to make the *oriT* amplicon also contained a PmlI site that was subsequently used to remove the yeast replication machinery. From these recombination events, we made pMQ56 (*CEN*, ampicillin resistance, *oriT*) and pMQ64 (*2μm*, gentamicin resistance, *oriT*). The yeast machinery of pMQ56 was removed by digestion with PmlI; one site was engineered between *oriT* and *URA3*, and the other is within the *CEN6* region. The resulting fragment was recircularized with T4 DNA ligase, generating pMQ57. A

similar strategy was used with pMQ64, except that we used XmnI, which cuts in the *2μm* origin, and PmlI, both of which leave blunt ends. The ligated product lacks the ability to replicate in yeast and was dubbed pMQ65 (Fig. 1).

Primer design. DNA sequences to be recombined were placed together in silico by using Gene Construction Kit software (Textco, Inc.). Additional features, such as synthetic ribosome binding sites, were added, and the junctions were selected and incorporated as part of the primer sequence.

In vivo cloning procedures (gap repair) and yeast minipreps. Yeast was grown overnight in yeast extract-peptone-dextrose at 30°C and transformed using a modified "lazy bones" transformation protocol (3, 10). Plasmids were recovered from yeast using the "smash and grab" method (3, 20).

Phenotypic assays. Rhamnolipid preparation and analysis, surface motility assays, and immunoblotting were performed using published protocols (4, 17,

38). Antihemagglutinin (anti-HA) antibodies were from QED Bioscience, Inc. (18850-01), and anti-green fluorescent protein (GFP) antibodies were from U.S. Biological, Inc. (G8965-10C).

Phase-contrast and epifluorescent microscopy were performed with a model DM IRBE inverted microscope (Leica Microsystems, Wetzlar, Germany), and photomicrographs were taken with a charge-coupled-device camera (model Orca C4742-5; Hamamatsu, Hamamatsu City, Japan) and analyzed with Open Lab 4.0.2 software (Improvision, Coventry, England). Flow cell experiments were performed as previously described (8).

Plasmid maps, sequences, and request information can be found at <http://www.dartmouth.edu/~gotoole/vectors.html>.

Nucleotide sequence accession numbers. Nucleotide sequences for vectors have been deposited in GenBank under accession numbers DQ230317 to DQ230324 and DQ642033 to DQ642043.

RESULTS

Modeling of gram-negative bacterial vectors using *S. cerevisiae*. (i) **Vectors for allelic replacement in gram-negative organisms unable to support replication of the ColE1 origin.** An existing gene replacement vector (pEX18-Gm) (18) was modified such that it was able to replicate in *S. cerevisiae*, resulting in plasmid pMQ30 (Fig. 1). pEX18-Gm has many fine features, including the counterselectable marker *sacB*, a high-copy ColE1 variant origin of replication, an origin for conjugal transfer, and a multicloning site in *lacZ* α for blue-white screening (18). Sequences required for stable replication in *S. cerevisiae* and a selectable/counterselectable marker, *URA3*, were added to pEX18-Gm by using homologous recombination in yeast. For more details on the construction of pMQ30 and all subsequent vectors, see Materials and Methods; also see Table S1 in the supplemental material for primers, Table S2 in the supplemental material for a key to vector construction, and Table S3 in the supplemental material for host ranges of the replicons used in this study.

pMQ30 was subsequently modified using yeast homologous recombination. The low-copy p15a origin of replication (also referred to as the pACYC origin) for *E. coli* was used to replace the high-copy ColE1-based origin to generate pMQ75 (Fig. 1). We hypothesized that the lower copy number afforded by the p15a origin would circumvent many toxicity problems associated with excess gene copy number.

pMQ30 was used to mutate the *clpP* gene of *P. aeruginosa* (Fig. 2). In this case, we replaced the *clpP* open reading frame with the *GFP* open reading frame. We also made clean deletion constructs by amplifying 500- to 1,000-bp regions upstream and downstream of a gene with primers that have homology on one side with the vector and the other side with each other. The two amplicons and the gapped vector were used to cotransform yeast, and recombinant constructs were selected for on medium deficient in uracil (not shown).

(ii) ***E. coli*-*S. cerevisiae* shuttle vectors usable as suicide vectors in gram-negative organisms unable to support replication of the ColE1 origin.** Vectors were designed specifically to disrupt genes via single-crossover events or to complement genes in their native context. The use of gene disruption by single-crossover events resulting in internal duplication is very quick and easy compared to allelic replacement and is also useful to rapidly test the importance of a gene to a biological phenomenon. This approach is especially useful when genomic or proteomic data implicate multiple genes in a biological




FIG. 2. *P. aeruginosa* *clpP* deletion and *clpX* disruption mutants have altered surface-associated phenotypes. (A) Overnight cultures of *P. aeruginosa* strain PA14 (wild type [wt]) and isogenic $\Delta clpP$ and *clpX*::pMQ84 strains were spotted on swarming agar plates and incubated at 37°C for 24 h (4, 38). Swarming in the wt strain is indicated by the tendrill-like structures originating from the point of inoculation and spreading across the agar surface. The absence of ClpP and ClpX correlated with a reduction in surface motility. (B) $\Delta clpP$ mutants exhibit increased biofilm formation under flow conditions in M63 medium supplemented with arginine. Shown here are photomicrographs of wt, $\Delta clpP$, and *clpX*::pMQ84 biofilms 24 and 48 h after inoculation.

process that would otherwise require considerable time to mutate through traditional allelic replacement.

We chose pMQ30 as a backbone for a suicide vector. The *sacB* gene was removed via homologous recombination to in-

crease the stability of single-crossover constructs in the bacterial chromosome, generating pMQ87 (Fig. 1).

The yeast replication machinery was also removed by using a yeast recombination-introduced *StuI* restriction site, yielding pMQ89 (not shown).

(iii) *E. coli*-*S. cerevisiae*-*Pseudomonas* shuttle vectors. Shuttle vectors that replicate in a range of gram-negative organisms as well as *S. cerevisiae* were made for complementation and other purposes. To make yeast-*E. coli*-*Pseudomonas* shuttle vectors, a *Pseudomonas* stability-conferring sequence from pUCP20 (originally from the cryptic *P. aeruginosa* plasmid pRO1600) was added to two yeast-*E. coli* shuttle vectors of the pRS400 series (35, 36). The pRO1600 replicon is sufficient for replication in many *Pseudomonas* species as well as *Klebsiella* species and has been shown to be stable even in the lack of selection (see Table S3 in the supplemental material) (28, 39). The pRS400 backbones we chose were pRS416 and pRS426, which differ by both copy number in the yeast cell and size. pRS416 is a centromeric (*CEN*) plasmid and therefore is maintained at 1 to 2 copies per cell, while pRS426 is a 2 μ m plasmid, maintained at about 25 copies per cell (36). Both plasmids have the ColE1-derived pBluescript high-copy origin of replication, a *lacZ* α -associated multicloning site, an ampicillin resistance gene (*bla*) for gram-negative bacteria, and a *URA3* marker for yeast. These plasmids also have T7 and *P_{lac}* promoters that can be utilized for gene expression. The *P_{lac}* promoter is useful in pseudomonads as a low-level constitutive promoter.

Homologous recombination was used to add the pRO1600 replicon and a kanamycin resistance cassette to pRS416 and pRS426, resulting in plasmids pMQ51 (2 μ m) and pMQ52 (*CEN6*), which were able to replicate in both *P. aeruginosa* and *P. fluorescens* (Fig. 1 and data not shown).

pMQ51 was further modified with addition of a gentamicin resistance gene by use of yeast homologous recombination to generate pMQ61 (Fig. 1).

(iv) Shuttle vectors containing *oriT*. Homologous recombination was used to add the RP4 *oriT* to pMQ51 and pMQ61 shuttle vectors. The resulting plasmids, pMQ56 (*CEN*, ampicillin resistance, *oriT*) and pMQ64 (2 μ m, gentamicin resistance, *oriT*), were made with primers that contained a *PmlI* site that was subsequently used to remove the yeast replication machinery from these plasmids. The yeast machinery of pMQ56 and pMQ61 was removed, generating vectors pMQ57 and pMQ65, respectively (Fig. 1 and data not shown).

These *oriT*-containing plasmids are capable of replication in and conjugation into *P. aeruginosa* (data not shown). Conjugations were done using *E. coli* strain S17-1, which bears transfer elements from RP4 on its chromosome (37).

(v) Expression vectors. All of the preceding vectors contain the *P_{lac}* promoter driving the *lacZ* α gene. These vectors could be used for expression of genes in *Pseudomonas* simply by recombining or ligating an open reading frame and ribosome binding site of interest downstream from the *P_{lac}* promoter; however, *Pseudomonas* spp. do not have the *lac* repressor, and therefore expression from *P_{lac}* is constitutive. The *P_{BAD}* promoter and the *araC* repressor are known for tight control of gene expression in *E. coli* and other species, so we chose this regulatory system for use with pseudomonads (15, 26). Newman and Fuqua have previously described the utility of the *P_{BAD}* promoter-bearing vectors for pseudomonads and other

gram-negative organisms (26). *P_{BAD}* and *araC* and transcriptional terminators were amplified from pBAD18 and recombined onto several of the shuttle vectors described above by completely replacing the *lacZ* α gene and *P_{lac}* promoter, generating pMQ70 from pMQ56, pMQ71 from pMQ61, and pMQ72 from pMQ64 (Fig. 1). The *PmlI*-*PmlI* yeast replication fragment was removed from pMQ70 to yield pMQ90. The *lacZ* α gene was recombined into pMQ70 under the control of the *P_{BAD}* promoter to enable arabinose-dependent blue-white screening in appropriate *E. coli* strains (pMQ91) (Fig. 1). *E. coli* strain DH5 α bearing pMQ91 forms blue colonies on selective medium containing X-Gal (5-bromo-4-chloro-3-indolyl- β -D-galactopyranoside) plus L-arabinose but not on the same plates without L-arabinose (data not shown).

(vi) Increased host range. The very-broad-host-range replication machinery of the RK2 plasmid was used to extend the usefulness of this tool kit. RK2-based vectors are commonly used in the study of diverse gram-negative genera (see Table S2 in the supplemental material). We replaced the pRO1600 replication machinery with the minimal RK2 replicon by homologous recombination. This replicon has been shown to be remarkably stable without selection in *P. aeruginosa* and *E. coli* but requires selection in *Azotobacter vinelandii* (2). The genetic information swap was performed with pMQ78 and pMQ80 to generate pMQ95 and pMQ97, respectively (Fig. 1). PA14 was transformed with these vectors, and *GFP* expression was found to be similar to that exhibited by PA14 with pMQ78 and pMQ80 (data not shown).

Inactivation of *clpP* but not *clpX* stimulates biofilm formation in *P. aeruginosa*. The ClpP protease has been shown to be conditionally required for *Pseudomonas fluorescens* biofilm formation (30). With *Staphylococcus aureus*, *clpP* mutants have been shown to exhibit elevated levels of biofilm production, while mutants of the ClpP-binding partner, *clpX*, produce diminished biofilms (12). We used some of the tools described in this work to examine the role of these two important proteins in *P. aeruginosa* surface-associated behaviors.

The *clpP* ORF was replaced directly with *GFP_{uvr}* by using a pMQ30-based construct (data not shown). A *clpP*::*GFP_{uvr}* allelic replacement construct was generated in a single recombination step. Briefly, primers were used to amplify *clpP* with flanking regions homologous to pMQ30. A second pair of primers was used to amplify a *GFP_{uvr}* with flanking regions homologous to *clpP*. First, the *clpP* amplicon was digested in the open reading frame with *AatII*. Then, both of these amplicons were added to a yeast transformation reaction along with pMQ30 that had been digested with *SmaI*. Uracil prototrophs were pooled, and plasmid DNA was acquired and transformed into *E. coli*. Gentamicin-resistant colonies that were also white on X-Gal were collected and analyzed by restriction digestion and sequenced. The proximity of the homologous targeting DNA to the restriction site does not seem to be important for the efficacy of these reactions. The *clpP* amplifying primers were also used to add the wild-type locus to pMQ30 for future allelic replacement or complementation scenarios (pRMQS44).

The *clpP* deletion strain grew slowly and exhibited reduced surface swarming motility (Fig. 2A and data not shown). Although *clpP* mutants grew slowly, they formed more robust

biofilms than the wild-type strain in M63 medium supplemented with arginine under flow conditions (Fig. 2B).

We used pRMQS84, described above, to disrupt *clpX*. The plasmid was introduced into *P. aeruginosa* by conjugation, and all gentamicin-resistant candidates were found to have pRMQS84 disrupting *clpX* (data not shown). We found that *clpX* was not essential for *Pseudomonas aeruginosa*, unlike for some other bacteria, such as *Caulobacter crescentus* (22). *clpX*::pRMQS84 mutants exhibited slightly reduced growth rates and altered surface motility (Fig. 2A and data not shown). The effect of *clpX* disruption upon biofilm formation under shear conditions was determined with M63 medium using arginine as a sole carbon source. Under the conditions utilized, *clpX* mutants formed biofilms in a manner similar to that of the wild type (Fig. 2B).

pMQ87 was subsequently used to mutate a number of genes implicated in biofilm formation by simply transforming *P. aeruginosa* with the pMQ87-based construct via conjugation or electroporation followed by selection for gentamicin-resistant colonies (Daniel Kadouri, Daniel MacEachran, and G. A. O'Toole, unpublished observations).

Use of inducible expression vectors in pseudomonads. *GFPmut3* from pGFPmut3 was introduced into vectors pMQ70 to pMQ72 by homologous recombination for use as a reporter. GFPmut3 is a stable and bright variant of GFP (6). The primers used to amplify *GFPmut3* contained an engineered ribosome binding site 8 bp upstream of the start codon of *GFP* as well as 40-bp tails that target recombination downstream of the P_{BAD} promoter on pMQ70 to pMQ72. Uracil-auxotrophic yeast (InvSc1) cells were transformed by the addition of both the *GFP*-containing amplicon and pMQ70 (or pMQ71 or pMQ72 in separate reactions) that had been digested with SmaI. Yeast transformants were pooled, and plasmid DNA was prepared and electroporated into *E. coli* which was subsequently plated on selective medium containing 100 mM L-arabinose. Antibiotic-resistant colonies were screened for *GFP* fluorescence by use of epifluorescent microscopy. Constructs verified to contain GFP were dubbed pMQ78, pMQ79, and pMQ80, respectively (Fig. 1). Fluorescence in *E. coli* was found to be dependent upon the presence of L-arabinose, and there was little leakiness, based on the lack of observable *GFP* fluorescence, even though these plasmids have very high copy numbers in *E. coli* (data not shown). The plasmids were conjugated into *P. aeruginosa* strain PA14 and tested for fluorescence during growth in a dilution series of L-arabinose. We found that at 100 mM L-arabinose the majority of cells fluoresced brightly, while PA14 bearing a no-*GFP* vector control (pMQ72) did not fluoresce (Fig. 3A and data not shown). Cells grown with 1 mM arabinose were clearly less bright, cells grown with 0.1 mM arabinose had detectable but very low levels of fluorescence, and those grown without arabinose exhibited no fluorescence at the exposure used (Fig. 3A).

The efficacy of these expression vectors was also assessed in *P. fluorescens*. The *lapD* gene of *P. fluorescens* is adjacent to the *lapA* gene and is implicated in biofilm formation (16, 17). The open reading frame of *lapD*, including 34 bp upstream and 25 bp downstream, was recombined into pMQ72 via yeast homologous recombination (pMQ72+*lapD*). Additionally, a truncated version of the gene lacking the C-terminal domain was cloned into pMQ72 (pMQ72+*lapD*ΔC-term). These plasmids

were electroporated into *P. fluorescens* and grown in broth culture supplemented with either 0% or 0.2% L-arabinose. Membrane fractions were prepared, run on polyacrylamide gels, transferred to nylon membrane, and immunoblotted with polyclonal anti-LapD antibodies (17). We observed relatively high levels of LapD in cultures grown with 0.2% arabinose and very little LapD from those grown without arabinose (Fig. 3B, lanes 3 and 4). We also observed stable expression of the LapDΔC-term polypeptide (Fig. 3B, lane 5). LapD antibodies showed predicted patterns with control wild-type and *lapD* deletion strains (Fig. 3B, lanes 1 and 2).

Complementation of Δ*rhlB* with *rhlB*-GFP. pMQ30 has previously been utilized to mutate the *rhlB* (rhamnolipid biosynthetic pathway) gene in *P. aeruginosa* (4). Rhamnolipids are biosurfactants and are hypothesized to modulate surface motility and biofilm architecture (4, 8). A biochemical approach was taken to directly determine the formation of rhamnolipids in the wild-type and *rhlB*::*kanMX* strains. Rhamnolipids were extracted from culture medium, separated on thin-layer chromatography plates, and detected by exposure to an anthrone-sulfuric acid mixture (Fig. 3C, lanes 1 and 3). RhlB is a rhamnosyltransferase that acts to add rhamnose moieties to rhamnolipid precursors, generating monorhamnolipids. Monorhamnolipids are subsequently converted to dirhamnolipids by RhlC. In the absence of RhlB, no rhamnolipids should be present, and that is what we observed from rhamnolipid preparations from *rhlB*::*kanMX* cultures, as was previously reported (4) (Fig. 3C, lane 3).

One advantage of using yeast homologous recombination is that seamless translational fusions can be made without multiple, possibly mutagenic, rounds of PCR associated with classical "sewing" techniques. We designed primers to fuse the *rhlB* ORF to the N terminus of GFP in pMQ78 and pMQ80. These primers direct a translational fusion of the last codon of *rhlB* to the first codon of *GFP*. pMQ78 and pMQ80 were digested with SmaI between the P_{BAD} promoter and *GFPmut3* and added to transformation reactions along with the targeted *rhlB* amplicon. The resulting plasmids, pRMQS96 and pRMQS98, were introduced into PA14 and the *rhlB*::*kanMX* strain. Rhamnolipid production was assessed as noted above for strains incubated in the presence of 100 mM L-arabinose. We observed that pRMQS98 was able to restore rhamnolipid production to the *rhlB*::*kanMX* strain (Fig. 3C, compare lanes 3 and 4). The GFP fusion can subsequently be used for microscopic visualization of the protein or as a convenient epitope tag.

This protocol is especially useful in making artificial operons for the study of prokaryotic genes. For example, in a single step we have cloned a two-gene operon while fusing a gene for cyan fluorescent protein or yellow fluorescent protein to the 3' end of the first gene in the operon (C. M. Toutain and G. A. O'Toole, unpublished results). Each of these synthetic operons can then be expressed under the control of its own promoter or under regulation of P_{BAD} or other promoters of choice or moved to pMQ30 (Fig. 1) to replace the chromosomal locus.

High-efficiency recombination of multiple pieces of DNA. The fusion of several pieces of DNA can easily be performed using yeast. This is especially useful for building complicated vectors and making gene fusions and site-directed mutations. As an example, we performed a complex reaction where we




FIG. 3. P_{BAD} -driven expression in *Pseudomonas* species. (A) Aliquots of *P. aeruginosa* cultures bearing pMQ78 grown overnight in LB with ampicillin and various concentrations of L-arabinose were viewed with either phase-contrast microscopy (left) or epifluorescent microscopy (right). Micrographs of GFP fluorescence were taken with identical exposure times. No fluorescence was detected in the absence of L-arabinose or in cells containing an identical vector without GFP (pMQ72 [not shown]). Bar = 10 μ m. (B) LapD immunoblot of membrane fractions from *P. fluorescens*. (Lane 1) Wild-type samples exhibit the predicted 71-kDa LapD protein. (Lane 2) Anti-LapD antibodies do not react with samples from a $\Delta lapD$ strain. (Lane 3) *P. fluorescens* bearing pMQ72 plus *lapD* exhibits relatively high levels of LapD in the presence of 0.2% L-arabinose (grown in M63 plus 0.2% glucose plus 0.5% Casamino Acids) and (lane 4) very little LapD in the absence of L-arabinose. (Lane 5) Fractions with pMQ72+*lapD*ΔC-term exhibited a predicted 45-kDa protein in the presence of 0.2% L-arabinose. (C) Thin-layer chromatographic analysis of *P. aeruginosa*. The wild type and the *rhlB::kanMX* mutant were grown in PPGAS with 100 mM L-arabinose for 24 h at 30°C as reported previously (4). The arrow indicates dirhamnolipids. (Lane 1) Wild type (wt) grown with vector alone (V) (pMQ78). (Lane 2) Wild type grown with a plasmid-borne *rhlB-GFPmut3* fusion construct (B-gfp) under the control of the P_{BAD} promoter (pRMQS96). (Lane 3) *rhlB::kanMX* mutant bearing the vector alone. (Lane 4) *rhlB::kanMX* mutant bearing the P_{BAD} -*rhlB-GFPmut3* plasmid. These data show that the *rhlB::kanMX* mutant is deficient in formation of dirhamnolipids and that this mutation is complemented by an RhlB-GFPmut3 fusion protein.

added four pieces of DNA to a vector in a single step. With pMQ87 (Fig. 4), in one step we (i) added the T5X inducible promoter-repressor system (11), (ii) added *GFP*, (iii) added F64L and S65T mutations to *GFP*, and (iv) fused a C-terminal 3HA epitope tag to *GFP*. The resulting plasmid was named pRMQS106 (Fig. 4A and B). At the same time, we removed the entire *lacZ α* gene and P_{lac} promoter of pMQ87.

The four amplicons are (i) a xylose repressor-promoter with primers to target recombination with pMQ87 and *GFP* including an artificial ribosome binding site, (ii) a portion of the GFP_{uv} ORF, corresponding to the first 63 amino acids, with

primers that target recombination with the promoter sequence including an artificial ribosome binding site on one side and GFP_{uv} targeting sequence with F64L and S65T mutations on the other side, (iii) a portion of the GFP_{uv} ORF, corresponding to the rest of the gene, including targeting sequences to the 5' portion of GFP_{uv} including the F64L and S65T mutations and with HA epitope targeting sequence on the 3' end, and (iv) a 3HA epitope tag with targeting sequences to *GFP* and pMQ87 (Fig. 4A and B). *S. cerevisiae* was cotransformed by the four amplicons and pMQ87 that had been digested with SmaI.

We observed a >10-fold ratio of transformants from the cut




FIG. 4. Yeast recombination facilitates the construction of multiple pieces of DNA. (A and B) Schematics depicting the recombination of four PCR-based amplicons into a gapped vector in a single step. (A) pMQ87 that has been digested with *Sma*I (gapped) and the four PCR-based amplicons: 1, *xylR* repressor-T5X promoter; 2, a 5' region of *GFP_{uvr}* with two mutations added to the primer; 3, a 3' region of *GFP_{uvr}* with two mutations added to the primer; 4, a 3HA tag. (B) Schematic diagram representing the recombinant product which has a modified *GFP-3HA* allele under the control of a xylose-regulated promoter (no. 1 to 4) all upon a yeast-*E. coli* shuttle vector. (C) Phase-contrast and epifluorescent micrographs of *E. coli* strains bearing either pMQ87 (no GFP) or pRMQS106 (*GFP-3HA*). Images were taken with identical exposure times. Magnification, $\times 400$. Bar = 10 μ m. (D) Immunoblots of lysates of *E. coli* cells bearing pMQ80 (*P_{BAD}-GFP*), pMQ87 (vector alone), or pRMQS106 (*P_{T5X}-GFP-3HA*). Blots were probed with either anti-

vector plus all four amplicons compared to the number of transformants from the cut vector alone. Hundreds of transformants were pooled, and their plasmids were obtained and used to transform *E. coli*. Colonies grown on selective medium plus 0.4% D-xylose were observed with fluorescent microscopy, and 11 of the 12 colonies tested exhibited bright green fluorescence (Fig. 4C). Additional candidate colonies were transferred to individual wells of a microtiter dish, each containing LB plus gentamicin plus 0.4% D-xylose, and grown overnight at 37°C. We examined the wells for *GFP* fluorescence with an inverted microscope and found that 52 out of 60 (86.7%) wells exhibited bright fluorescence. Cells in the microtiter plate were subjected to several rounds of freezing and thawing, and detergent was added to lyse cells. Lysates were then spotted onto a nitrocellulose membrane and immunoblotted using anti-HA antibodies. In control experiments, these commercially available antibodies detected no background bands in *E. coli* lysates on Western blots (Fig. 4D and data not shown). We found that while a control lysate bearing an empty vector exhibited no cross-reactivity to the HA antibody, 53 out of 60 (88%) candidate lysates reacted with anti-HA antibodies. Together, these results are consistent with >80% of candidate colonies containing constructs that had undergone five-way recombination events producing functional fusion proteins. This frequency is likely an underrepresentation of the actual frequency of recombinants for two reasons. (i) Original yeast transformants contained a mixture of uncut vectors that had not been digested in the original gap-generating digestion, gapped vectors that had closed through nonhomologous end joining, and gapped vectors that had closed through the targeted recombination events. Rather than screening through yeast transformants, we pooled them and electroporated yeast-plasmid preparations into *E. coli*. (ii) Gap-repaired vectors bearing the targeted recombinant construct may contain PCR-generated or recombination-generated mutations leading to a lack of *GFP-3HA* expression or a lack of production of a functional protein.

Western blot analysis was performed with cells bearing pRMQS106 (*GFP-3HA*) to further verify the construct. Cell lysates of *E. coli* strains bearing pMQ80 (*P_{BAD}-GFP*), pMQ87 (negative control), or pRMQS106 (*P_{T5X}-GFP-3HA*) were immunoblotted with anti-GFP and anti-HA antibodies (Fig. 4D). With anti-GFP antibodies, a single band was identified from pMQ80 lysates, corresponding to approximately 25 kDa (GFP has a predicted mass of 26.9 kDa); no band was observed with the pMQ87 lysate, as expected; and a larger band was observed associated with pRMQS106 lysates (*GFP-3HA* has a predicted mass of ~ 31 kDa). Anti-HA antibodies revealed bands only with pRMQS106 lysates, and these corresponded to the same size as those observed with the anti-GFP antibodies (Fig. 4D).

GFP or anti-HA antiserum. Cells were grown overnight in the presence of gentamicin and 0.4% glucose and then pelleted and diluted at 1:20 in LB supplemented with gentamicin and either 40 mM L-arabinose for the pMQ80 culture (GFP) or 0.4% (vol/vol) D-xylose for pMQ87 (vector) and pRMQS106 (*GFP-3HA*). Diluted cultures were grown for 3 h at 37°C before aliquots were taken for analysis. Lysates of cells bearing two independent pRMQS106 constructs are shown.

The two plasmids used in Fig. 4 were sequenced, and all five recombinant junctions from each construct were as predicted. Likewise, we sequenced the junctions for the majority of recombinant junctions for the vectors described here and all were as predicted, consistent with reports showing the high fidelity of recombination in *S. cerevisiae*.

DISCUSSION

In vivo cloning via gap repair is efficient, easy, and amenable to high-throughput applications. Here we describe a complete set of tools made for modification of bacterial genes by using yeast homologous recombination. While others have previously described the use of gap repair in cloning, a bacterium-centered set of tools has not been made. For example, Raymond and colleagues have previously described using this technique for cloning of *Pseudomonas* DNA and have described the clever use of oligonucleotide linkers that further broaden the flexibility of yeast recombination-mediated cloning (32, 33).

We used these vectors to manipulate genes from *Pseudomonas aeruginosa* and *P. fluorescens* to emphasize the power of a new set of molecular tools that take advantage of the *S. cerevisiae* native recombination proteins. Although these vectors were made with *Pseudomonas* spp. in mind, they can be used with many gram-negative organisms and thus are broadly useful. The bacterial origins of replication placed upon one or more of these vectors include ColE1, p15a, pRO1600, and RK2, allowing the use of these vectors to replicate in more than 30 species of bacteria. These tools will allow researchers who study many types of prokaryotic organisms, including non-model species, to conveniently modify DNA by using recombination.

Other systems for in vivo cloning have been described previously, and some can be used in conjunction with the vectors and system described here (7, 23). The vectors described here can be used with many existing tools designed for use with yeast, such as a set of constructs that allow epitope tagging of a protein of interest with any of a group of several epitopes with a single set of primers via yeast homologous recombination (24).

Cloning with yeast homologous recombination is an additional tool to facilitate molecular studies of a wide range of gram-negative organisms rather than a replacement for traditional in vitro cloning. However, this tool kit can be readily substituted for many applications of traditional cloning, particularly for large-scale cloning projects and for the construction of plasmids requiring multiple inserts, site-directed mutations, or precise junctions. While multiple amplicons can also be fused via splicing by overlap extension PCR, this method requires serial rounds of amplification, increasing the frequency of PCR-generated mutations, and is more limited with regard to the size of the final amplicon. Also, recombination can be targeted hundreds of base pairs away from the gap site on a vector, providing a large advantage over traditional cloning methods where modifications are confined within existing restriction sites. This allows complete exchange of features on a vector, such as replicons or promoter/regulator systems. Another advantage is that, although yeast-based cloning can take longer, as yeast colonies can take 2 to 4 days to appear on

selective plates, these methods often require much less handling time. Amplicons are used directly, and vector digests can be used without any purification or desalination. Additionally, the method of yeast transformation described here requires no lengthy preparation of competent cells. The disadvantages are the increased cost due to longer primers and the increased amount of time for yeast cultures to grow.

ACKNOWLEDGMENTS

We thank Dan MacEachran, Benedict Kemp, and Stu Powers for critical reading of the manuscript and members of the O'Toole lab for trying many of the described constructs. We thank Herbert Schweizer, Dean Dawson, and Janet Ajioka for the kind gifts of plasmids, strains, and glass beads.

This work was supported by NIH training grants T32 AI07363 and F32 GM66658-01A1 to R.M.Q.S. and NIH R21-AI055774 and AI51360 to G.A.O.

REFERENCES

1. Bascom-Slack, C. A., and D. Dawson. 1998. A physical assay for detection of early meiotic recombination intermediates in *Saccharomyces cerevisiae*. *Mol. Gen. Genet.* **258**:512–520.
2. Blatny, J. M., T. Brautaset, H. C. Winther-Larsen, K. Haugan, and S. Valla. 1997. Construction and use of a versatile set of broad-host-range cloning and expression vectors based on the RK2 replicon. *Appl. Environ. Microbiol.* **63**:370–379.
3. Burke, D., D. Dawson, and T. Stearns. 2000. *Methods in yeast genetics: a Cold Spring Harbor Laboratory course manual*. Cold Spring Harbor Laboratory Press, Plainview, N.Y.
4. Caiazza, N. C., R. M. Shanks, and G. A. O'Toole. 2005. Rhamnolipids modulate swarming motility patterns of *Pseudomonas aeruginosa*. *J. Bacteriol.* **187**:7351–7361.
5. Choi, K. H., A. Kumar, and H. P. Schweizer. 2005. A 10-min method for preparation of highly electrocompetent *Pseudomonas aeruginosa* cells: application for DNA fragment transfer between chromosomes and plasmid transformation. *J. Microbiol. Methods* **64**:391–397.
6. Cormack, B. P., R. H. Valdivia, and S. Falkow. 1996. FACS-optimized mutants of the green fluorescent protein (GFP). *Gene* **173**:33–38.
7. Datsenko, K. A., and B. L. Wanner. 2000. One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proc. Natl. Acad. Sci. USA* **97**:6640–6645.
8. Davey, M. E., N. C. Caiazza, and G. A. O'Toole. 2003. Rhamnolipid surfactant production affects biofilm architecture in *Pseudomonas aeruginosa* PAO1. *J. Bacteriol.* **185**:1027–1036.
9. DeMarini, D. J., C. L. Creasy, Q. Lu, J. Mao, S. A. Sheardown, G. M. Sathe, and G. P. Livi. 2001. Oligonucleotide-mediated, PCR-independent cloning by homologous recombination. *BioTechniques* **30**:520–523.
10. Elble, R. 1992. A simple and efficient procedure for transformation of yeasts. *BioTechniques* **13**:18–20.
11. Forsyth, R. A., R. J. Haselbeck, K. L. Ohlsen, R. T. Yamamoto, H. Xu, J. D. Trawick, D. Wall, L. Wang, V. Brown-Driver, J. M. Froelich, G. C. Kedar, P. King, M. McCarthy, C. Malone, B. Misner, D. Robbins, Z. Tan, Z. Zhang, G. Carr, D. A. Mosca, C. Zamudio, J. G. Foulkes, and J. W. Zyskind. 2002. A genome-wide strategy for the identification of essential genes in *Staphylococcus aureus*. *Mol. Microbiol.* **43**:1387–1400.
12. Frees, D., A. Chastanet, S. Qazi, K. Sorensen, P. Hill, T. Msadek, and H. Ingmer. 2004. Clp ATPases are required for stress tolerance, intracellular replication and biofilm formation in *Staphylococcus aureus*. *Mol. Microbiol.* **54**:1445–1462.
13. Geels, F. P., and B. Schippers. 1983. Reduction of yield depressions in high frequency potato cropping soil after seed tuber treatments with antagonistic fluorescent *Pseudomonas* spp. *Phytopathol. Z.* **108**:207–214.
14. Guldener, U., S. Heck, T. Fiedler, J. Behnhauer, and J. H. Hegemann. 1996. A new efficient gene disruption cassette for repeated use in budding yeast. *Nucleic Acids Res.* **24**:2519–2524.
15. Guzman, L.-M., B. Carson, M. J. Carson, and J. Beckwith. 1995. Tight regulation, modulation, and high-level expression by vectors containing the arabinose P_{BAD} promoter. *J. Bacteriol.* **177**:4121–4130.
16. Hinsa, S. M., M. Espinosa-Urgel, J. L. Ramos, and G. A. O'Toole. 2003. Transition from reversible to irreversible attachment during biofilm formation by *Pseudomonas fluorescens* WCS365 requires an ABC transporter and a large secreted protein. *Mol. Microbiol.* **49**:905–918.
17. Hinsa, S. M., and G. A. O'Toole. 2006. Identification and characterization of the LapD protein: LapD modulates the secretion of LapA. *Microbiology* **152**:1375–1383.
18. Hoang, T. T., R. R. Karkhoff-Schweizer, A. J. Kutchma, and H. P. Schweizer. 1998. A broad-host-range Flp-FRT recombination system for site-specific

- excision of chromosomally-located DNA sequences: application for isolation of unmarked *Pseudomonas aeruginosa* mutants. *Gene* **212**:77–86.
19. **Hoang, T. T., A. J. Kutchma, A. Becher, and H. P. Schweizer.** 2000. Integration-proficient plasmids for *Pseudomonas aeruginosa*: site-specific integration and use for engineering of reporter and expression strains. *Plasmid* **43**:59–72.
 20. **Hoffman, C. S., and F. Winston.** 1987. A ten-minute DNA preparation from yeast efficiently releases autonomous plasmids for transformation of *Escherichia coli*. *Gene* **57**:267–272.
 21. **Jansen, G., C. Wu, B. Schade, D. Y. Thomas, and M. Whiteway.** 2005. Drag&Drop cloning in yeast. *Gene* **344**:43–51.
 22. **Jenal, U., and T. Fuchs.** 1998. An essential protease involved in bacterial cell-cycle control. *EMBO J.* **17**:5658–5669.
 23. **Li, M. Z., and S. J. Elledge.** 2005. MAGIC, an in vivo genetic method for the rapid construction of recombinant DNA molecules. *Nat. Genet.* **37**:311–319.
 24. **Longtine, M. S., A. McKenzie III, D. J. Demarini, N. G. Shah, A. Wach, A. Brachat, P. Philippsen, and J. R. Pringle.** 1998. Additional modules for versatile and economical PCR-based gene deletion and modification in *Saccharomyces cerevisiae*. *Yeast* **14**:953–961.
 25. **Mallet, L., and M. Jacquet.** 1996. Intergenic Flip Flop, a method for systematic gene disruption and cloning in yeast. *Yeast* **12**:1351–1357.
 26. **Newman, J. R., and C. Fuqua.** 1999. Broad-host-range expression vectors that carry the L-arabinose-inducible *Escherichia coli* *araBAD* promoter and the *araC* regulator. *Gene* **227**:197–203.
 27. **Oldenburg, K. R., K. T. Vo, S. Michaelis, and C. Paddon.** 1997. Recombination-mediated PCR-directed plasmid construction in vivo in yeast. *Nucleic Acids Res.* **25**:451–452.
 28. **Olsen, R. H., G. DeBusscher, and W. R. McCombie.** 1982. Development of broad-host-range vectors and gene banks: self-cloning of the *Pseudomonas aeruginosa* PAO chromosome. *J. Bacteriol.* **150**:60–69.
 29. **Orr-Weaver, T. L., and J. W. Szostak.** 1983. Yeast recombination: the association between double-strand gap repair and crossing-over. *Proc. Natl. Acad. Sci. USA* **80**:4417–4421.
 30. **O'Toole, G. A., and R. Kolter.** 1998. Flagellar and twitching motility are necessary for *Pseudomonas aeruginosa* biofilm development. *Mol. Microbiol.* **30**:295–304.
 31. **Rahme, L. G., E. J. Stevens, S. F. Wolfort, J. Shao, R. G. Tompkins, and F. M. Ausubel.** 1995. Common virulence factors for bacterial pathogenicity in plants and animals. *Science* **268**:1899–1902.
 32. **Raymond, C. K., E. H. Sims, A. Kas, D. H. Spencer, T. V. Kuttyavin, R. G. Ivey, Y. Zhou, R. Karul, J. B. Clendenning, and M. V. Olson.** 2002. Genetic variation at the O-antigen biosynthetic locus in *Pseudomonas aeruginosa*. *J. Bacteriol.* **184**:3614–3622.
 33. **Raymond, C. K., E. H. Sims, and M. V. Olson.** 2002. Linker-mediated recombinational subcloning of large DNA fragments using yeast. *Genome Res.* **12**:190–197.
 34. **Santos, P. M., I. D. Bartolo, J. M. Blatny, E. Zennaro, and V. Svein.** 2001. New broad-host-range promoter probe vectors based on the plasmid RK2 replicon. *FEMS Microbiol. Lett.* **195**:91–96.
 35. **Sikorski, R. S.** 1993. Extrachromosomal cloning vectors of *Saccharomyces cerevisiae*, p. 225–241. *In* K. G. Hardy (ed.), *Plasmids, a practical approach*, 2nd ed. IRL Press, Oxford, United Kingdom.
 36. **Sikorski, R. S., and P. Hieter.** 1989. A system of shuttle vectors and yeast host strains designed for efficient manipulation of DNA in *Saccharomyces cerevisiae*. *Genetics* **122**:19–27.
 37. **Simon, R., U. Priefer, and A. Puhler.** 1983. A broad host range mobilization system for in vivo genetic engineering: transposon mutagenesis in gram-negative bacteria. *BioTechnology* **1**:784–790.
 38. **Toutain, C. M., M. E. Zegans, and G. A. O'Toole.** 2005. Evidence for two flagellar stators and their role in the motility of *Pseudomonas aeruginosa*. *J. Bacteriol.* **187**:771–777.
 39. **West, S. E., H. P. Schweizer, C. Dall, A. K. Sample, and L. J. Runyen-Janecky.** 1994. Construction of improved *Escherichia-Pseudomonas* shuttle vectors derived from pUC18/19 and sequence of the region required for their replication in *Pseudomonas aeruginosa*. *Gene* **148**:81–86.