



Revista Portuguesa
de

irurgia

II Série • N.º 19 • Dezembro 2011

ISSN 1646-6918

Órgão Oficial da Sociedade Portuguesa de Cirurgia

SAGES guidelines for the clinical application of laparoscopic biliary tract surgery*

D. Wayne Overby¹, Keith N. Apelgren², William Richardson³, Robert Fanelli⁴

¹Department of Surgery, University of North Carolina, Chapel Hill, NC, USA

²Department of Surgery, Michigan State University, Lansing, MI, USA

³Department of Surgery, Ochsner Clinic Foundation, Jefferson, LA, USA

⁴R. Fanelli (&) Pittsfield, MA, USA

Received: 7 April 2010 / Accepted: 27 May 2010 / Published online: 13 August 2010

© Springer Science+Business Media, LLC 2010

PREAMBLE

Laparoscopic cholecystectomy (LC) has become the standard of care for patients requiring removal of the gallbladder. In 1992, a National Institutes of Health (NIH) consensus development conference concluded that “laparoscopic cholecystectomy provides a safe and effective treatment for most patients with symptomatic gallstones, laparoscopic cholecystectomy appears to have become the procedure of choice for many of these patients” [1].

The Society of American Gastrointestinal and Endoscopic Surgeons (SAGES) first offered guidelines for the clinical application of laparoscopic cholecystectomy in May 1990. These guidelines have periodically been updated, and the last guideline in November 2002 expanded the guidelines to include all laparoscopic biliary tract surgery.

This document updates and replaces the previous guideline.

The current recommendations are graded and linked to the evidence utilizing the definitions in Appendices 1 and 2.

DISCLAIMER

Guidelines for clinical practice are intended to indicate preferable approaches to medical problems as established by experts in the field. These recommendations will be based on existing data or a consensus of expert opinion when little or no data are available.

Guidelines are applicable to all physicians who address the clinical problem(s) without regard to specialty training or interests, and are intended to indicate the preferable, but not necessarily the only, acceptable approaches due to the complexity of the healthcare environment. Guidelines are intended to be flexible. Given the wide range of specifics in any health care problem, the surgeon must always choose the course best suited to the individual patient and the variables in existence at the moment of decision.

Guidelines are developed under the auspices of the Society of American Gastrointestinal and Endoscopic Surgeons and its various committees, and approved by the Board of Governors. Each clinical practice guideline has been systematically researched, reviewed, and revised by the guidelines committee, and reviewed by

* Publicado com autorização de SAGES e de Springer (with kind permission of Springer"), publicado em Surgical Endoscopy Volume 24, Number 10, 2368-2386, DOI: 10.1007/s00464-010-1268-77



an appropriate multidisciplinary team. The recommendations are therefore considered valid at the time of its production based on the data available. Each guideline is scheduled for periodic review to allow incorporation of pertinent new developments in medical research knowledge, and practice.

INDICATIONS

The indications for laparoscopic operations on the gallbladder and biliary tree have not changed since the 1992 National Institutes of Health *Consensus Development Conference Statement on Gallstones and Laparoscopic Cholecystectomy* [1]; they remain similar to the indications for open surgery with relative and absolute contraindications as noted below. As stated in the NIH report “most patients with symptomatic gallstones are candidates for laparoscopic cholecystectomy, if they are able to tolerate general anesthesia and have no serious cardiopulmonary diseases or other co-morbid conditions that preclude operation.” The indications include but are not limited to symptomatic cholelithiasis, biliary dyskinesia, acute cholecystitis, and complications related to common bile duct (CBD) stones including pancreatitis (see additional references provided in sections below). Asymptomatic gallstones are generally not an indication for laparoscopic cholecystectomy [2-7].

Indications for laparoscopic operations on the gallbladder and biliary tree

- Include but are not limited to symptomatic cholelithiasis, biliary dyskinesia, acute cholecystitis, and complications related to common bile duct stones including pancreatitis with few relative or absolute contraindications (Level II, Grade A)

RELATIVE CONTRAINDICATIONS AND INDICATIONS FOR PLANNED OPEN PROCEDURES

Relative contraindications for laparoscopic biliary tract surgery include many of the usual contraindi-

cations for laparoscopic surgery in general. These include, but are not limited to, generalized peritonitis, septic shock from cholangitis, severe acute pancreatitis, untreated coagulopathy, lack of equipment, lack of surgeon expertise, previous abdominal operations which prevent safe abdominal access or progression of the procedure, advanced cirrhosis with failure of hepatic function, and suspected gallbladder cancer [1]. Laparoscopic cholecystectomy may be performed safely in patients with cirrhosis and acute cholecystitis (see additional references provided in sections below), but there are cases in which the open approach may be safer. Indications for planned open procedures include a patient’s informed request for an open procedure, known dense adhesions in the upper abdomen, known gallbladder cancer, and surgeon preference.

Relative contraindications for laparoscopic biliary tract surgery

- Untreated coagulopathy, lack of equipment, lack of surgeon expertise, hostile abdomen, advanced cirrhosis/liver failure, and suspected gallbladder cancer (Level II, Grade A)

PREOPERATIVE PREPARATION

Antibiotic prophylaxis

Preoperative antibiotics in elective laparoscopic biliary tract surgery have been discussed with strong opinions on both sides. A recent meta-analysis of randomized controlled trials concluded that prophylactic antibiotics do not prevent infections in low-risk patients undergoing laparoscopic cholecystectomy, while the usefulness of prophylaxis in high-risk patients (age > 60 years, presence of diabetes, acute colic within 30 days of operation, jaundice, acute cholecystitis, or cholangitis) remains uncertain [8]. The most recent randomized, prospective study included in the above-mentioned meta-analysis showed no difference in the postoperative wound infection rate, although the control group had a 1.5% infection rate and the antibiotic group had a 0.7% infection rate; since



there was a total of 277 patients in the study, a type II error might have been committed [9]. Among papers suggesting that antibiotic prophylaxis is helpful is a recent randomized study which found fewer wound infections with ampicillin–sulbactam versus cefuroxime, particularly for infection caused by enterococcus in the setting of high-risk patients undergoing elective cholecystectomy [10]. If antibiotics are used they should be limited to a single preoperative dose given within 1 h of skin incision, and redosed if the procedure is more than 4 h long [11].

Antibiotic prophylaxis

- Antibiotics are not required in low-risk patients undergoing laparoscopic cholecystectomy (Level I, Grade A)
- Antibiotics may reduce the incidence of wound infection in high-risk patients (age \geq 60 years, presence of diabetes, acute colic within 30 days of operation, jaundice, acute cholecystitis, or cholangitis) (Level I, Grade B)
- If given, they should be limited to a single preoperative dose given within 1 h of skin incision (Level II, Grade A)

Deep venous thrombosis prophylaxis

This prophylaxis is necessary for most laparoscopic biliary tract procedures, is addressed in a separate SAGES guideline [12], and should consist of either pneumatic compression stockings or subcutaneous heparin given prior to operation in patients with two or more risk factors. See the above-referenced citation for further information.

Deep venous thrombosis prophylaxis

- Prophylaxis is addressed in a separate SAGES guideline [12]

BASIC OPERATIVE TECHNIQUE

Room setup and patient positioning

There are two basic room setups for performing

laparoscopic biliary tract surgery. The first is the standard supine position with the surgeon standing at the patient's left and monitors at the head of the bed on both sides. The second is with the patient in stirrups and the surgeon standing between the legs. The latter is commonly used in Europe and the former in the Americas. Some surgeons tuck the left arm to improve the working space of the operating surgeon. The patient is generally placed in reverse Trendelenburg position and rotated right side up. The SAGES manual [13] describes room setup, patient positioning, and the remainder of the procedure in further detail.

Room setup and patient positioning

- With no data to guide choices, surgeon preference should dictate room setup (Level III, Grade A)

Equipment needed for laparoscopic cholecystectomy

The equipment needed for laparoscopic cholecystectomy and intraoperative cholangiography is well established, with specific preferences left to the discretion of the operating surgeon. The equipment needed for laparoscopic common bile duct exploration is also at the discretion of the operating surgeon and should be available if that is a possibility when performing cholecystectomy. One potential approach to equipment selection is covered in the SAGES manual [13].

Equipment

- In the absence of data, surgeon preference should dictate choice of equipment (Level III, Grade A)

Abdominal access

There are a variety of techniques for gaining initial abdominal access for laparoscopic surgery; these include: (1) Veress needle, (2) the open Hasson technique, (3) direct trocar placement without prior pneumoperitoneum, and (4) the optical view technique, in which the laparoscope is placed within the trocar so that the layers of the abdominal wall



are visualized as they are being traversed. In general, all of the mentioned approaches to abdominal access are safe. A recent meta-analysis [14] of 17 randomized controlled trials studying a total of 3,040 individuals comparing a variety of open and closed access techniques found no difference in complication rates; potentially lifethreatening injuries to blood vessels occurred in 0.9 per 1,000 procedures and to the bowel in 1.8 per 1,000 procedures. Currently, there are no demonstrable differences in the safety of open versus closed techniques for establishing access and creating the initial pneumoperitoneum, therefore decisions regarding choice of technique are left to the surgeon and should be based on individual training, skill, and case assessment [15].

Abdominal access

- There are no demonstrable differences in the safety of open versus closed techniques for establishing access; decisions regarding choice of technique are left to the surgeon and should be based on individual training, skill, and case assessment (Level I, Grade A)

Safe technique

The safety of laparoscopic cholecystectomy is based largely on determining the anatomy of the cystic duct, common bile duct, cystic artery, and hepatic arteries. Since major bile duct injuries with laparoscopic cholecystectomy are most frequently due to duct misidentification [16, 17], techniques for prevention and/or recognition focus primarily on careful anatomic definition [18] to ensure the “critical view” prior to dividing any structures [19, 20] including dissection (1) to completely expose and delineate the hepatocystic triangle, (2) to identify a single duct and a single artery entering the gallbladder, and (3) to completely dissect the lower part of the gallbladder off the liver bed. Though the protective effect of the practice continues to be debated, routine use of intraoperative cholangiography may decrease the risk or severity of injury and improve injury recognition [17, 21-23]. The general principle of not dividing any structure until

you are certain of its identification applies here; the need for caution and vigilance cannot be overstated given evidence which supports visual misperception as an underlying cause of major bile duct injury [24], coupled with the potential for complacency which may result from the rarity of bile duct injuries.

Safe technique

- The safety of laparoscopic cholecystectomy requires correct identification of relevant anatomy (Level I, Grade A)
- Intraoperative cholangiogram may reduce the rate or severity of injury and improve injury recognition (Level II, Grade B)

Common bile duct assessment

The primary methods for assessing the common bile duct for stones or injury during cholecystectomy are intraoperative cholangiogram and intraoperative ultrasound:

1. Intraoperative cholangiography has been used for many years; fluoroscopy saves time and has improved its usefulness. The issue of routine versus selective cholangiography has been long debated. Studies have suggested that routine use of intraoperative cholangiography may decrease the risk of injury and improve injury recognition, while others have suggested that cholecystectomy may be performed without cholangiogram with low rates of injury [17, 21-23]. In residency programs, a policy of routine cholangiography may be supported by the need to train residents how to do that portion of the procedure [25]. In addition, the skills developed and maintained by routine cholangiography provide a platform for progression to transcystic clearing or stenting of the common bile duct [25]; in many cases, clearing can be accomplished with simple measures such as administration of glucagon and flushing with saline [26]. In terms of detecting bile duct stones, 2-12% of patients will have choledocholithiasis on routine intraoperative cholangiogram, and recent studies suggest that as many as 10% of these are unsuspected prior to operation [27-29]. A meta-analysis performed in 2004 [30] re-



vealed that the incidence of unsuspected retained stones was 4%, with only 15% of these going on to cause clinical problems. The conclusion from that study was that a selective policy should be advocated, though creating a reliable algorithm for predicting the presence of stones and thus the need for selective cholangiogram has been unsuccessful [31, 32].

2. *Laparoscopic ultrasound* This technique has been used increasingly; while it does not by itself offer potentially therapeutic access to the bile ducts, it does help delineate relevant anatomy including bile ducts and vascular structures, and can diagnose choledocholithiasis without opening the biliary system, all without exposure to ionizing radiation. Several recent studies have examined the use of laparoscopic ultrasound during cholecystectomy. Potential advantages and disadvantages of the technique have been summarized by Perry et al.; advantages include high rates of successful studies, the ability to repeat the examination during difficult dissections, less time required for completion, and lower overall cost, while disadvantages include technical difficulties for certain patients, inability to confirm the flow of bile into the duodenum, and the experience required to learn the technique of examination and image interpretation [33]. The authors of the included studies used the technique routinely with no reported bile duct injuries, and minor bile leak secondary to liver bed injury was a rare event (0.2%); high sensitivity and specificity for detection of common bile duct stones were reported [33-36].

Common bile duct assessment

- Intraoperative cholangiography may decrease the risk of bile duct injury when used routinely and allows access to the biliary tree for therapeutic intervention; reliable algorithms to determine the need for selective cholangiography have yet to be developed (Level II, Grade B)
- In experienced hands, intraoperative laparoscopic ultrasound helps delineate relevant anatomy, detect bile duct stones, and decrease the risk of bile duct injury (Level II, Grade B)

Management of choledocholithiasis

Approaches to suspected choledocholithiasis

With increasing laparoscopic expertise, exploration of the common bile duct either via the cystic duct or by primary choledochotomy has become a viable option, but the treatment of symptomatic or suspected common bile duct stones in the era of laparoscopic cholecystectomy remains a complex and controversial issue. Leaving aside open cholecystectomy/bile duct exploration, which is superior to endoscopic retrograde cholangiopancreatography (ERCP) for stone clearance [37], as described by Kharbutli and Velanovich [38] there are two approaches to patients with possible choledocholithiasis who are undergoing laparoscopic cholecystectomy, both for patients who are asymptomatic undergoing elective cholecystectomy and for patients with recent episodes of jaundice or gallstone pancreatitis: (1) laparoscopic cholecystectomy with intraoperative cholangiogram, then address choledocholithiasis if found, or (2) preoperative ERCP to diagnosis and remove choledocholithiasis, followed by laparoscopic cholecystectomy. For choice (1), a number of additional choices are possible for stones found during intraoperative imaging studies: (A) transcystic laparoscopic common bile duct exploration, (B) common bile duct exploration via choledochotomy, (C) placement of an endobiliary stent, (D) intraoperative ERCP and postoperative ERCP. Several recent studies including at least two meta-analyses have attempted to compare the relative merits of the above approaches, and one-stage treatment combining laparoscopic cholecystectomy with laparoscopic common bile duct exploration usually prevails in terms of cost with no discernable difference in morbidity and mortality. With that said, preoperative ERCP should not be used for diagnosis alone; routine preoperative ERCP will likely result in higher than acceptable mortality and morbidity rates, with some unnecessary procedures. The single-stage laparoscopic or combined laparoscopic with intraoperative endoscopic approaches require time, equipment, and a degree of skill and experience which are not univer-



sal among surgeons and facilities performing laparoscopic cholecystectomy. Finally, postoperative ERCP leads to longer hospital stays with increased numbers of procedures required to treat the problem [37-44].

A. *Transcystic common bile duct exploration*

Given the scope of issues detailed above, the choice of technique to treat common duct stones will likely depend largely on local expertise. However, both short and long-term data from a number of studies suggest that transcystic common bile duct exploration, which may be augmented by choledochoscopy, is as safe and efficacious as other minimally invasive approaches [31, 37, 40, 45-49]. The postoperative course after successful transcystic clearance is similar to after laparoscopic cholecystectomy alone [25, 45]. Transcystic stone clearance may be hampered by anomalous anatomy, proximal (hepatic duct) stones, strictures, and large ([6 mm) or numerous ([5) stones [25, 31, 40, 47].

B. *Choledochotomy*

Laparoscopic common bile duct exploration via choledochotomy requires advanced laparoscopic skills and longer operative times; most authors see choledochotomy as an alternative to failed transcystic exploration, though some explore via choledochotomy exclusively, all with generally good results in terms of stone clearance. The open bile duct may be addressed with closure over a T-tube, an exteriorized transcystic drain, or primary closure with or without endoluminal drainage [49-51]. Closure over a T-tube may be required if the common bile duct is inflamed [52] and in any case allows for postoperative radiographic evaluation of the biliary system, the possibility of extraction of retained stones, and the possibility of a controlled biliary fistula, but can be complicated by premature dislodgement, bile leak and peritonitis, localized pain, prolonged fistula, and late biliary stricture [50]. Studies comparing primary closure versus T-tube drainage suggest similar rates of complications with shorter operating times and a trend toward shorter hospital stays with primary closure [51, 53].

C. *Laparoscopic endobiliary stent placement*

This treatment option for choledocholithiasis effectively bridges the gap between laparoscopic common bile duct exploration and ERCP; the technique involves placing a stent through the cystic duct into the common bile duct and across the ampulla of Vater, then closing the cystic duct. The advantages of this approach include decompression of the biliary tree allowing the option of semi-elective postoperative ERCP, which for most patients maintains the minimally invasive approach and ambulatory nature of laparoscopic cholecystectomy; the stent adds little operative time to the procedure, facilitating ERCP and stone clearance while potentially reducing the incidence of post-ERCP pancreatitis, and deployment does not require advanced laparoscopic skills [54-57].

D. *ERCP with stone extraction*

ERCP with stone extraction is another alternative when faced with choledocholithiasis; it may be performed before, during or after cholecystectomy. As discussed by Costi et al. [58], “performing ERCP before surgery raises questions regarding patient selection because systematic preoperative ERCP before LC means an intolerably great number of unnecessary and potentially harmful procedures. Complex scoring systems aimed at identifying asymptomatic patients to undergo ERCP have not been adopted as clinical practice, nor have new examinations such as echoendoscopy and biliary magnetic resonance imaging (MRCP), which are costly and not always available. Performing ERCP contextually to LC implies organizational problems concerning the availability of an endoscopist in the operating theater whenever needed. Finally, performing ERCP after surgery would raise the dilemma of managing CBD stones whenever ERCP fails to retrieve them because a third procedure would then be needed.” With no discernable difference in morbidity and mortality and similar clearance rates when compared with laparoscopic common bile duct exploration, duct clearance with



postoperative ERCP is a viable alternative [37-44]. While, in experienced hands, the two approaches are at least equivalent, there are surgeons for whom the preferred approach is ERCP with stone extraction [41]. However, unless performed intraoperatively, ERCP requires at least one additional procedure, and does have associated complications such as pancreatitis, bleeding, and duodenal perforation, and as noted above, ERCP may fail, leading to multiple procedures for stone clearance. As described by Karaliotas et al., the following entities increase the possibility of failure of endoscopic CBD stone clearance: stone impaction, gastrectomy or Roux-en-Y anatomy, recurrent bile duct stones after prior open exploration of the CBD and biliodigestive anastomosis, periampullary diverticula, and Mirizzi syndrome [52].

Altered anatomy

Rearrangement of the upper gastrointestinal tract can make it difficult, if not impossible, to perform standard ERCP. With the recent increase in the number of Roux-en-Y gastric bypass procedures performed for morbid obesity, it becomes ever more likely that surgeons will encounter patients who have gallstone disease and limited endoscopic access to the biliary system. As described by Ahmed et al., options for treatment include percutaneous transhepatic instrumentation of the common bile duct, percutaneous transgastric ERCP, laparoscopic transgastric ERCP, transenteric ERCP, retrograde endoscopy in which the scope is passed antegrade down to the jejunojejunostomy and then retrograde up the biliopancreatic limb, and open or laparoscopic common bile duct exploration [59].

Management of choledocholithiasis

- There are several approaches, and current data do not suggest clear superiority of any one approach; decisions regarding treatment are most appropriately made based on surgeon preference as well as the availability of equipment and skilled personnel (Level I, Grade A)
- Laparoscopic transcystic common bile duct exploration may employ a number of techniques

from simple to advanced; it is frequently successful, but may be hampered by anomalous anatomy, proximal stones, strictures, and large or numerous stones (Level II, Grade B)

- Laparoscopic choledochotomy requires advanced laparoscopic skills, but has good clearance rates; the open bile duct may be addressed with closure over a T-tube, an exteriorized transcystic drain, or primary closure with or without endoluminal drainage (Level II, Grade B)
- Laparoscopic endobiliary stent placement adds little operative time to the cholecystectomy, and facilitates ERCP and stone clearance (Level II, Grade B)
- ERCP with stone extraction may be performed selectively before, during or after cholecystectomy with little discernable difference in morbidity and mortality and similar clearance rates when compared with laparoscopic common bile duct exploration, though routinely performed pre-operative ERCP will likely result in unnecessary procedures with higher than acceptable mortality and morbidity rates (Level I, Grade A)

Dissection of the gallbladder from the liver bed

The conventional technique for dissection of the gallbladder from the liver bed is to start from the gallbladder infundibulum and work superiorly using electrocautery to remove the gallbladder from the bed. The technique of topdown dissection has also been advocated, particularly in cases with significant inflammation [60-62]. Ultrasonic dissection has been studied for dissection of the gallbladder from the liver bed, as well as division and sealing of the cystic artery and cystic duct without clips; in prospective randomized trials, ultrasonic dissection has been found to be comparable in terms of operative time, gallbladder perforation, bleeding, and bile leak [61, 63]. In addition, hydrodissection with a high-pressure water stream has been used to dissect the gallbladder from the liver bed [64]. The standard technique works well and, with no compelling data to use these alternative techniques, the choice is left to the operating surgeon.



Dissection of the gallbladder from the liver bed

- The more conventional approach starting at the gallbladder infundibulum and working superiorly, or the top-down approach, may be used with electrocautery, ultrasonic dissection or hydrodissection as the surgeon prefers (Level II, Grade B)

Extraction of the gallbladder

The gallbladder is generally extracted from either the epigastric port or the umbilical port. The decision is left up to the operating surgeon. Some surgeons use a 5-mm port in the epigastric position, necessitating removal through the umbilicus. Likewise, most difficult extractions due to large size of the gallbladder should be done through the umbilicus, because it is easier to expand the fascial incision. The use of an endoscopic bag is also at the discretion of the operating surgeon. There are no randomized studies to guide use of these techniques.

Extraction of the gallbladder

- With no data to guide choice of technique, the gallbladder may be extracted as the surgeon prefers (Level III, Grade C)

Use of drains

While use of drains postoperatively after laparoscopic biliary tract surgery is at the discretion of the operating surgeon, recent studies including a randomized controlled trial and meta-analysis of six randomized controlled trials found that drain use after elective laparoscopic cholecystectomy increases postoperative pain and wound infection rates and delays hospital discharge; the authors further stated they could not find evidence to support use of drains after laparoscopic cholecystectomy [65, 66].

Use of drains

- Drains are not needed after elective laparoscopic cholecystectomy, and their use may increase complication rates (Level I, Grade A)
- Drains may be useful in complicated cases particularly if choledochotomy is performed (Level III, Grade C)

Conversion to laparotomy

Conversion from laparoscopic to open cholecystectomy should not be considered a complication, but is rather an attempt to avoid complications and ensure patient safety [67]. Factors which are associated with conversion to open cholecystectomy include: acute cholecystitis with a thickened gallbladder wall, previous upper abdominal surgery, male gender, advanced age, obesity, bleeding, bile duct injury, and choledocholithiasis [67-73]. Ultimately, individual surgeons must base the decision to convert to an open procedure on their own intraoperative assessment, weighing the severity of inflammatory changes, clarity of the anatomy, and their skill/comfort in proceeding [72]. Overall conversion rates have been reported to be between 2 and 15% [67], and in cases of acute cholecystitis from 6 to 35% [71].

Conversion to laparotomy

- Conversion should not be considered a complication, and surgeons should have a low threshold for conversion; the decision to convert to an open procedure must be based on intraoperative assessment weighing the clarity of the anatomy and the surgeon's skill/comfort in proceeding (Level II, Grade A)

INTRAOPERATIVE COMPLICATIONS

Access injuries

Establishing access and creating the initial pneumoperitoneum necessary to perform laparoscopic biliary tract procedures may lead to significant complications. Reviews of data regarding device-related injury and death as reported to the Food and Drug Administration (FDA) [74] as well as thorough reviews of the available literature [15] suggest that vascular and visceral injuries are the major causes of morbidity and mortality related to abdominal access. The true rates of injury are difficult to gauge; injuries are probably underreported both to the FDA and in the literature, and there is a paucity of prospective data, but it is



likely that injuries which occur while establishing pneumoperitoneum account for a significant proportion of complications during laparoscopy [15, 74, 75]. Laparoscopic cholecystectomy is the procedure most frequently associated with both fatal and nonfatal trocar injuries, and almost all fatal injuries were made with shielded or optical trocars [74]. A recent meta-analysis of 17 randomized controlled trials studying a total of 3,040 individuals comparing a variety of open and closed access techniques found no difference in complication rates; potentially life-threatening injuries to blood vessels occurred in 0.9 per 1,000 procedures and to the bowel in 1.8 per 1,000 procedures [14]. Currently, there are no demonstrable differences in the safety of open versus closed techniques for establishing access and creating the initial pneumoperitoneum; therefore, decisions regarding choice of technique are left to the surgeon and should be based on individual training, skill, and case assessment [15]. A high index of suspicion and prompt conversion to laparotomy are required to recognize and treat complications related to access.

Access injuries

- There are no demonstrable differences in the safety of open versus closed techniques for establishing access; decisions regarding choice of technique are left to the surgeon and should be based on individual training, skill, and case assessment (Level I, Grade A)
- A high index of suspicion and prompt conversion to laparotomy are required to recognize and treat complications related to access (Level III, Grade A)

Common bile duct injuries

A great deal continues to be written about bile duct injuries in laparoscopic cholecystectomy, which serves to underscore the seriousness of the complication and the perception that it can and should be avoided. The current rate of major bile duct injury in laparoscopic cholecystectomy has stabilized at 0.1-0.6% [18, 21-23, 76-78], and series with no major bile duct inju-

ries have been reported [20]; while many believe that the rate of major bile duct injury in open cholecystectomy is lower than in laparoscopic cholecystectomy, controversy remains [76, 78]. A host of factors have been associated with bile duct injury, including surgeon experience, patient age, male sex [22], and acute cholecystitis, though the effect that acute cholecystitis has on injury rates remains controversial [23, 79, 80]. Bile duct injuries which occur with laparoscopic cholecystectomy frequently involve complete disruption and excision of ducts, and may be associated with hepatic vascular injuries [81-83]. If major bile duct injuries do occur, whether recognized at the time of the primary operation or in the postoperative period, outcomes are improved by early recognition and by referring patients immediately to experienced specialists for further diagnosis and treatment. Repair should not be attempted by the primary surgeon unless the primary surgeon has significant experience in biliary reconstruction [77, 84-86]. Since major bile duct injuries with laparoscopic cholecystectomy are most frequently due to duct misidentification [16, 17], techniques for prevention and/or recognition focus primarily on careful anatomic definition [18] to ensure the "critical view" prior to dividing any structures [19, 20], and though the protective effect of the practice continues to be debated, use of intraoperative cholangiography may decrease the rate or the severity of common bile duct injury [17, 21-23].

Common bile duct injuries

- Factors which have been associated with bile duct injury include surgeon experience, patient age, male sex, and acute cholecystitis (Level II, Grade C)
- The safety of laparoscopic cholecystectomy requires correct identification of relevant anatomy (Level I, Grade A)
- Intraoperative cholangiogram may reduce the rate or severity of injury and improve injury recognition (Level II, Grade B)
- If major bile duct injuries occur, outcomes are improved by early recognition and immediate re-



ferral to experienced hepatobiliary specialists for further treatment before any repair is attempted by the primary surgeon, unless the primary surgeon has significant experience in biliary reconstruction (Level II, Grade A)

SPECIAL CONSIDERATIONS

Biliary dyskinesia

Patients with symptoms of biliary obstruction without evidence of gallstones but with abnormal gallbladder emptying may benefit from laparoscopic cholecystectomy [87-92]. Symptoms may include episodic, severe, steady pain, frequently with fatty food intolerance, located in the right upper quadrant or epigastrium, with or without radiation to the back or shoulder lasting at least 30 min but less than several hours, and may potentially be associated with nausea and vomiting [89, 90]. Abnormal gallbladder emptying is usually defined as gallbladder ejection fraction of less than 35% on cholescintigraphy after injection of cholecystokinin [88-90]. Severe symptoms, very low gallbladder ejection fraction (<14%), and reproduction of symptoms with cholecystokinin administration may be more predictive of resolution of symptoms after cholecystectomy [88, 90]. In patients who undergo laparoscopic cholecystectomy for biliary dyskinesia, stones are found in specimens 10-12% of the time, indicating a significant false-negative rate for gallbladder ultrasound in this group of patients [88, 90].

Biliary dyskinesia

- Patients with symptoms of biliary obstruction without evidence of gallstones but with abnormal gallbladder emptying may benefit from laparoscopic cholecystectomy (Level II, Grade B)

Acute cholecystitis

About 10-15% of all cholecystectomies performed are for acute cholecystitis [93]. Laparoscopic cholecystectomy has become the preferred approach in patients with acute cholecystitis [93-101] with rates

of conversion to an open procedure of 6-35% [70, 71, 73, 102-108]. For patients who can tolerate the procedure, early cholecystectomy (within 24-72 h of diagnosis) in cases of acute cholecystitis is increasingly advocated; when compared with planned open and/or delayed cholecystectomy, early laparoscopic cholecystectomy: reduces the rate of symptom relapse; may be performed without increased rates of conversion to an open procedure, without an increased risk of complications, including bile duct injury; and may decrease cost and total length of stay [5, 79, 98, 99, 101, 103, 104, 107-114]. In critically ill patients with acute cholecystitis, radiographically guided percutaneous cholecystostomy is an effective temporizing measure until the patient recovers sufficiently to undergo cholecystectomy [99, 115-121]. Laparoscopic cholecystectomy in the elderly (age [65 years) may be associated with higher morbidity and mortality [122, 123].

Acute cholecystitis

- Laparoscopic cholecystectomy has become the preferred approach in patients with acute cholecystitis (Level II, Grade B)
- Early cholecystectomy (within 24-72 h of diagnosis) may be performed without increased rates of conversion to an open procedure, without an increased risk of complications, and may decrease cost and total length of stay (Level I, Grade A)
- In critically ill patients with acute cholecystitis, radiographically guided percutaneous cholecystostomy is an effective temporizing measure until the patient recovers sufficiently to undergo cholecystectomy (Level II, Grade B)

Gallstone pancreatitis

Acute pancreatitis caused by gallstones is an important indication for cholecystectomy. The incidence of acute pancreatitis due to gallstones appears to be increasing [124, 125]. Based on a study of one large state's discharge data, one-third of cases of acute pancreatitis among US adults are caused by gallstones, with an incidence of gallstone pancreatitis of approximately 14.5 per 100,000 [125], which translates into 31,500



cases per year nationally. While laparoscopic cholecystectomy has become the preferred approach for removing the source of stones [126], the timing of the cholecystectomy, as well as the choice and timing of procedures for evaluating and clearing associated common bile duct stones, remain controversial, particularly in cases of mild, self-limited gallstone pancreatitis. There is agreement that severe pancreatitis with ongoing multisystem organ failure requires immediate clearing of any biliary obstruction, usually with ERCP, followed by supportive care until the patient recovers sufficiently to tolerate cholecystectomy [127]. However, when pancreatitis caused by gallstones is mild and self-limited, the issue becomes prevention of recurrent episodes of biliary symptoms, including acute pancreatitis. Currently, the majority of surgeons advocate and perform cholecystectomy urgently, when symptoms have subsided and laboratory values have normalized, usually during the same hospital admission [96, 126-133], while others delay cholecystectomy for weeks; decision-making algorithms regarding approaches to pre-versus intraoperative common bile duct evaluation and clearance are even more provider dependent, though patients with mild pancreatitis generally do not benefit from preoperative ERCP [126, 134]. A recent meta-analysis [39] showed no difference in morbidity and mortality when endoscopic removal of common bile duct stones with cholecystectomy was compared with cholecystectomy with intraoperative removal of common bile duct stones; the authors went on to state that treatment should be determined by local resources and expertise.

Gallstone pancreatitis

- Laparoscopic cholecystectomy has become the preferred approach for removing the source of stones in cases of acute pancreatitis due to gallstones (Level II, Grade B)
- Severe pancreatitis with ongoing multisystem organ failure requires immediate clearing of any biliary obstruction followed by supportive care until the patient recovers sufficiently to tolerate cholecystectomy (Level I, Grade A)

- When pancreatitis caused by gallstones is mild and self-limited, urgent cholecystectomy should be performed after symptoms have subsided and laboratory values have normalized, usually during the same hospital admission (Level II, Grade B)

Laparoscopic cholecystectomy in the setting of pregnancy

Please see the published SAGES guidelines and associated review article regarding diagnosis and laparoscopic treatment of surgical diseases during pregnancy [135].

Laparoscopic cholecystectomy in the setting of pregnancy

- Please see the published SAGES guidelines and associated review article regarding diagnosis and laparoscopic treatment of surgical diseases during pregnancy [135]

Laparoscopic cholecystectomy surgery in the setting of cirrhosis

Cirrhosis places patients at increased risk for gallstone formation [136-138]. Since the NIH consensus conference on gallstones and laparoscopic cholecystectomy in 1992 suggested that patients with cirrhosis were “not usually candidates for laparoscopic cholecystectomy” [1] studies continue to be published supporting the safety of the approach in patients with Child’s A or B cirrhosis (including downgrading from C after appropriate treatment) [39] with almost no data using the Model for EndStage Liver Disease (MELD) score to compare patients [139]; though there is little published data for Child’s C patients, what is available suggests it should be avoided in favor of nonoperative approaches such as percutaneous cholecystostomy [140]. Recent studies generally agree that laparoscopic cholecystectomy in selected cirrhotics has a relatively low conversion rate (0-11%), complication rate (9.5-21%), and risk of dying (0-6.3%), with most showing worsening liver failure, including the presence of ascites and coagulopathy, predicting poorer outcomes [139-144]; a recent prospective ran-



domized trial found that laparoscopic cholecystectomy was safer than open cholecystectomy in cirrhotics [145]. Some authors have suggested laparoscopic subtotal cholecystectomy as an alternative to laparoscopic cholecystectomy [146, 147]. Most authors caution that bleeding is the most frequent and worrisome complication, suggesting that coagulopathy and thrombocytopenia be corrected preoperatively, and that dilated pericholecystic and abdominal wall veins or recanalized umbilical veins be treated with care, with one author noting “conversion to open does not correct coagulopathy” [142, 143].

Laparoscopic cholecystectomy surgery in the setting of cirrhosis

- Laparoscopic cholecystectomy is relatively safe in patients with Child’s A or B cirrhosis (Level I, Grade B)
- Laparoscopic cholecystectomy is not recommended for Child’s C patients (Level III, Grade C)
- Bleeding is the most frequent complication; coagulopathy and thrombocytopenia should be corrected preoperatively, and dilated pericholecystic and abdominal wall veins or recanalized umbilical veins be treated with care (Level II, Grade A)

Laparoscopic cholecystectomy in the setting of systemic anticoagulation

There is little published data regarding laparoscopic cholecystectomy in the setting of systemic anticoagulation, but there are at least two recently published studies of patients taking warfarin for long-term systemic anticoagulation [148, 149]. In both, patients had their warfarin discontinued and were bridged to surgery with low-molecular-weight heparin as inpatients, and laparoscopic cholecystectomy was performed after their international normalized ratio (INR) was 1.5 or less. In one study of 44 anticoagulated patients, postoperative bleeding was significantly more common in the oral anticoagulation group (25%) versus the control group (1.5%), and in the majority of cases, bleeding in the oral anticoagulation group

was serious, requiring blood transfusion or reoperation with concomitantly longer hospital stay with standard laboratory tests not predicting postoperative hemorrhage [148], while the other study with 33 anticoagulated patients reported no bleeding complications [149]. Based on similar rates of bleeding from other studies of laparoscopic procedures reviewed by the authors, caution in chronically anticoagulated patients is warranted, particularly in those requiring bridging with low-molecular-weight heparin [148].

Laparoscopic cholecystectomy in the setting of systemic anticoagulation

- Caution in chronically anticoagulated patients is warranted even after cessation of pharmacotherapy, particularly in those bridged with low-molecular-weight heparin (Level III, Grade B)

Porcelain gallbladder

The relationship between calcification of the gallbladder wall and gallbladder cancer has been oft repeated; however, there is relatively little published data regarding the relationship between the two, with almost no published data from this decade. One of the most recent available studies, from 2000 [150], reviewed pathological findings from 25,900 cholecystectomies over 27 years; there were 150 gallbladders with cancer and 44 with calcified walls, 17 with complete intramural calcification (the classic porcelain gallbladder) and 27 with selective mucosal calcification. None of the specimens with complete intramural calcification had concomitant associated cancer, while only 2 of the 27 with selective mucosal calcification had associated cancer, correlating with a 5% incidence in calcified gallbladders (0% in true porcelain gallbladders). There is one study, from 2004, addressing calcified gallbladders in laparoscopic cholecystectomy [151] with 13 of 1,608 laparoscopic cholecystectomy specimens having calcified walls, again noting no cancer in ten gallbladders with complete intramural calcification while one of three specimens with selective mucosal calcifications had associated cancer, which suggests that patients with suspected



calcifications should be carefully studied, with open cholecystectomy recommended for those with selective mucosal calcifications.

Porcelain gallbladder

- Patients with suspected gallbladder calcifications should be carefully studied, with open cholecystectomy recommended for those with selective mucosal calcifications (Level III, Grade B)

Gallbladder polyps

Polyploid lesions of the gallbladder, which can be found in about 1-5% of adults on ultrasound in Western populations [152, 153] and 9.6% in Asian populations [154], are defined as elevations of the gallbladder mucosa. Polyploid lesions of the gallbladder can be true polyps which demonstrate neoplastic changes and may be benign, dysplastic or malignant, or can be pseudopolyps such as cholesterol polyps, inflammatory polyps, or adenomyoma, which are all benign [152, 155]. Gallbladder polyps are most frequently cholesterol polyps, which are usually small (less than 1 cm) and multiple, and tend to remain stable with regard to size and number. Patients with cholesterol polyps usually do not develop concomitant stones or symptoms [156]. A recent comparison of preoperative ultrasound findings with pathological examination of cholecystectomy specimens in Western patients suggests that size is the only reliable indicator for malignant potential, with all malignancies found in polyps greater than 6 mm [152], though non-Western populations may develop malignancies in smaller polyps [155]. There are no randomized studies to direct decisions regarding gallbladder polyps [157], and despite recent studies, the management of gallbladder polyps remains controversial. A reasonable approach would include laparoscopic cholecystectomy for larger, especially single, polyps or those with associated symptoms, with watchful waiting for small (<5 mm), asymptomatic polyps.

Gallbladder polyps

- Laparoscopic cholecystectomy should be considered for larger, especially single, polyps or

those with associated symptoms, with watchful waiting for small (<5 mm) asymptomatic polyps (Level II, Grade B)

Gallbladder cancer

The incidence of gallbladder cancer in the USA is 1.2 per 100,000; the only curative therapy is surgical resection, and except for those with early-stage disease, survival is extremely poor. Gallbladder cancer is found unexpectedly upon pathological examination in less than 1% of specimens after laparoscopic cholecystectomy [158, 159]. Laparoscopic cholecystectomy is considered curative for cancers confined to the gallbladder mucosa (T1a), while cancers which invade the muscularis (T1b) may have lymph node metastases or lymphatic invasion which prompts some authors to recommend hepatoduodenal lymph node dissection for these lesions, but an initial open versus laparoscopic approach does not influence survival [160-163]. Inadvertent opening of cancerous gallbladders during laparoscopic cholecystectomy increases the likelihood of recurrence and port-site metastases [164-166]. Cancers which are more locally advanced or those with nodal involvement should be referred to specialty centers for consideration of more extensive resection or re-resection [159].

Gallbladder cancer

- Laparoscopic cholecystectomy is considered curative for cancers confined to the gallbladder mucosa (T1a) (Level II, Grade B)
- Cancers which are more locally advanced or those with nodal involvement should be referred to specialty centers for consideration of more extensive resection or re-resection (Level II, Grade B)

POSTOPERATIVE MANAGEMENT

Length of stay

Patients undergoing uncomplicated laparoscopic cholecystectomy for symptomatic cholelithiasis may



be discharged home on the day of surgery [167]. Control of postoperative pain, nausea, and vomiting is important to successful same-day discharge [168], and admission rates despite planned same-day discharge are reported to be 1-39%; patients older than 50 years may be at increased risk for admission [168-174]. Re-admission rates range from 0 to 8%; common causes for readmission after sameday discharge include pain, intra-abdominal fluid collections, bile leaks, and bile duct stones [167, 170]. Time to discharge after surgery for patients with acute cholecystitis, bile duct stones, or in patients converted to an open procedure should be determined on an individual basis.

Length of stay

- Patients undergoing uncomplicated laparoscopic cholecystectomy for symptomatic cholelithiasis may be discharged home on the day of surgery; control of postoperative pain, nausea, and vomiting is important to successful same-day discharge (Level II, Grade B)
- Patients older than 50 years may be at increased risk for admission (Level II, Grade B)
- Time to discharge after surgery for patients with acute cholecystitis, bile duct stones, or in patients converted to an open procedure should be determined on an individual basis (Level III, Grade A)

Reduced-port and single-incision laparoscopic cholecystectomy

All parts of the *SAGES Guidelines for the Clinical Application of Laparoscopic Biliary Tract Surgery* apply to reduced-port and single-incision approaches to laparoscopic cholecystectomy. The indications, contraindications, and preoperative preparation for reduced-port and single-incision approaches are the same as those for multiport cholecystectomy. Access and equipment, are, in their essentials, the same for reduced-port and single-incision approaches and for multiport procedures. Access to the abdominal cavity in reduced-port and single-incision approaches should follow accepted standards for safe entry, including avoidance and recognition of complications.

Standard instruments may be used in single-incision or multiport procedures. With respect to specialized access devices and nonrigid instruments, there have been no trials or adequate evaluative studies yet published to offer any recommendation for these devices. Introduction of new instruments, access devices or new techniques should be done with caution and/or under study protocol, and prior to the addition of any new instrument or device, it should, to the extent possible, be proven safe, and not limit adherence to established guidelines for safe performance of laparoscopic cholecystectomy. Adequate training should be obtained on any new device or instrument prior to utilization in a patient. As with any new technique, outcomes should be continuously assessed to ensure continued patient safety as single-incision techniques are developed; to date, only studies with limited numbers of patients have been reported [175-177]. Dissection performed during single-incision procedures should follow “best practice” approaches recommended for multiport cholecystectomy including dynamic traction of the fundus of the gallbladder, dynamic lateral retraction of the gallbladder infundibulum, and identification and maintenance of the “critical view” of the cystic duct and artery to avoid inadvertent injury to the common bile duct or hepatic arteries. During initial procedures, a low threshold for using additional port sites should be maintained so as not to jeopardize a safe dissection and result.

Single-incision cholecystectomy

- The indications, contraindications, and preoperative preparation for reduced-port and single-incision approaches are the same as those for multiport cholecystectomy (Level III, Grade A)
- Access to the abdominal cavity in reduced-port and singleincision approaches should follow accepted standards for safe entry, including avoidance and recognition of complications (Level III, Grade A)
- Introduction of new instruments, access devices or new techniques should be done with cau-



tion and/or under study protocol, and prior to the addition of any new instrument or device, it should, to the extent possible, be proven safe, and not limit adherence to established guidelines for safe performance of laparoscopic cholecystectomy (Level III, Grade A)

- During initial procedures, a low threshold for using additional port sites should be maintained so as not to jeopardize a safe dissection and result (Level III, Grade A)

Disclosures Author Robert D. Fanelli is a board member of New Wave Surgical Corporation, received honoraria from Ethicon EndoSurgery for speaking/teaching, received honoraria from Boston Scientific Corporation, Inc. for speaking/teaching, and was an independent contractor for Cook Surgical, Inc. Authors Keith N. Apelgren, Keenan Berghoff, Paul Curcillo, David W. Overby, and William S. Richardson have no conflicts of interest or financial ties to disclosure.

Appendix 1: Levels of evidence

Level I	Evidence from properly conducted randomized, controlled trials
Level II	Evidence from controlled trials without randomization Or Cohort or case-control studies Or Multiple time series, dramatic uncontrolled experiments
Level III	Descriptive case series, opinions of expert panels

Appendix 2: Scale used for recommendation grading

Grade A	Based on high-level (level I or II), well-performed studies with uniform interpretation and conclusions by the expert panel
Grade B	Based on high-level, well-performed studies with varying interpretation and conclusions by the expert panel
Grade C	Based on lower-level evidence (level II or less) with inconsistent findings and/or varying interpretations or conclusions by the expert panel

Appendix 3: Literature review method, search terms, and results

Literature review method

Systematic literature searches for each topic were performed on MEDLINE during the course of the review. In general, the search strategy was limited to articles in English language, on humans, and published within the last 5 years. The abstracts were reviewed by the two committee members (D.W.O., K.N.A.). Randomized controlled trials, meta-analyses, and systematic reviews were selected for further review along with prospective and retrospective studies including studies with smaller samples, which were considered when additional evidence was lacking.

Search terms and results

A. Indications:

1. Search date: September 2009
2. Search terms: “cholecystectomy indications”
3. Limits: English language, humans, and published within the last 5 years
4. Results: 91 articles, abstracts reviewed, 6 chosen as pertinent, 1 additional earlier landmark publication included

B. Antibiotic prophylaxis:

1. Search date: July 2009
2. Search terms: “laparoscopic cholecystectomy prophylaxis antibiotics”
3. Limits: English language, humans, and published within the last 5 years
4. Results: 13 articles, abstracts reviewed, 4 chosen as pertinent

C. Abdominal access. See “Access injuries” below

D. Safe technique:

1. Search date: August 2009
2. Search terms: “laparoscopic cholecystectomy bile duct injury prevention”
3. Limits: English language, humans, and published within the last 5 years
4. Results: 33 articles, abstracts reviewed, 8 chosen as pertinent

E. Intraoperative cholangiography:

1. Search date: August 2009
2. Search terms: “intraoperative cholangiogram choledocholithiasis”
3. Limits: English language, humans, and published within the last 5 years
4. Results: 69 articles, abstracts reviewed, 12 chosen as pertinent

F. Intraoperative ultrasound:

1. Search date: August 2009
2. Search terms: “laparoscopic cholecystectomy intraoperative ultrasound”
3. Limits: English language, humans, and published within the last 5 years
4. Results: 59 articles, abstracts reviewed, 4 chosen as pertinent

G. Laparoscopic bile duct exploration, ERCP with stone extraction, and altered anatomy:

1. Search date: August 2009



2. Search terms: "laparoscopic bile duct exploration"
3. Limits: English language, humans, and published within the last 5 years
4. Results: 101 articles, abstracts reviewed, 15 chosen as pertinent

H. Laparoscopic endobiliary stent placement:

1. Search date: August 2009
2. Search terms: "laparoscopic endobiliary stent"
3. Limits: None
4. Results: 14 articles, abstracts reviewed, 4 chosen as pertinent

I. Dissection of the gallbladder from the liver bed:

1. Search date: August 2009
2. Search terms: "laparoscopic cholecystectomy dissection"
3. Limits: English language, humans, and published within the last 5 years
4. Results: 83 articles, abstracts reviewed, 5 chosen as pertinent

J. Use of drains:

1. Search date: August 2009
2. Search terms: "laparoscopic cholecystectomy drains"
3. Limits: English language, humans, and published within the last 5 years
4. Results: 9 articles, abstracts reviewed, 2 chosen as pertinent

K. Conversion to laparotomy:

1. Search date: February 2009
2. Search terms: "laparoscopic cholecystectomy conversion to laparotomy"
3. Limits: English language, humans, and published within the last 5 years
4. Results: 33 articles, abstracts reviewed, 7 chosen as pertinent

L. Access injuries:

1. Search date: August 2009
2. Search terms: "laparoscopic access complication"
3. Limits: English language, humans, and published within the last 5 years
4. Results: 90 articles, abstracts reviewed, 4 chosen as pertinent

M. Common bile duct injuries:

1. Search date: February 2009
2. Search terms: "laparoscopic cholecystectomy bile duct injury"
3. Limits: English language, humans, and published within the last 5 years
4. Additional hand searching of bibliographies
5. Results: 194 articles, abstracts reviewed, 19 chosen as pertinent

N. Biliary dyskinesia:

1. Search date: September 2009
2. Search terms: "cholecystectomy biliary dyskinesia"
3. Limits: English language, humans, and published within the last 5 years
4. Additional hand searching of bibliographies
5. Results: 40 articles, abstracts reviewed, 6 chosen as pertinent

O. Acute cholecystitis:

1. Search date: March 2009
2. Search terms: "laparoscopic cholecystectomy acute cholecystitis"

3. Limits: English language, humans, and published within the last 5 years
4. Results: 219 articles, abstracts reviewed, 38 chosen as pertinent

P. Gallstone pancreatitis:

1. Search date: April 2009
2. Search terms: "laparoscopic cholecystectomy acute pancreatitis"
3. Limits: English language, humans, and published within the last 5 years
4. Results: 77 articles, abstracts reviewed, 13 chosen as pertinent

Q. Laparoscopic cholecystectomy surgery in the setting of cirrhosis:

1. Search date: April 2009
2. Search terms: "laparoscopic cholecystectomy cirrhosis"
3. Limits: English language, humans, and published within the last 5 years
4. Additional hand searching of bibliographies
5. Results: 69 articles, abstracts reviewed, 13 chosen as pertinent

R. Laparoscopic cholecystectomy surgery in the setting of systemic anticoagulation:

1. Search date: April 2009
2. Search terms: "laparoscopic cholecystectomy anticoagulation"
3. Limits: None
4. Additional hand searching of bibliographies
5. Results: 11 articles, abstracts reviewed, 2 chosen as pertinent

S. Porcelain gallbladder:

1. Search date: April 2009
2. Search terms: "laparoscopic cholecystectomy porcelain gallbladder"
3. Limits: None
4. Additional hand searching of bibliographies
5. Results: 16 articles, abstracts reviewed, 2 chosen as pertinent

T. Gallbladder polyps:

1. Search date: April 2009
2. Search terms: "gallbladder polyps"
3. Limits: English language, humans, and published within the last 5 years
4. Results: 59 articles, abstracts reviewed, 6 chosen as pertinent

U. Gallbladder cancer:

1. Search date: June 2009
2. Search terms: "laparoscopic cholecystectomy gallbladder cancer"
3. Limits: English language, humans, and published within the last 5 years
4. Results: 108 articles, abstracts reviewed, 9 chosen as pertinent

V. Length of stay:

1. Search date: July 2009
2. Search terms: "laparoscopic cholecystectomy hospital discharge"
3. Limits: English language, humans, and published within the last 5 years
4. Results: 58 articles, abstracts reviewed, 8 chosen as pertinent

W. Single-incision cholecystectomy:

1. Search date: September 2009
2. Search terms: "single incision laparoscopic cholecystectomy"
3. Limits: English language, humans, and published within the last 5 years
4. Results: 15 articles, abstracts reviewed, 3 chosen as representative



Appendix 4

This document was prepared and revised by the SAGES Guidelines Committee:

D. Wayne Overby, MD
Keith N. Apelgren, MD
Keenan R. Beghoff, MD
Paul Curcillo, MD
Ziad Awad, MD
Ronald Clements, MD
David Edelman, MD
Erika Fellinger, MD
Stephen Haggerty, MD
Steven Heneghan, MD
Henry Lujan, MD
Sumeet Mittal, MD
Jonathan Pearl, MD
Raymond Price, MD
Patrick Reardon, MD
John Roth, MD
Alan Saber, MD

Dimitrios Stefanidis, MD
Julio Teixeria, MD
Andrew Wright, MD
Marc Zerey, MD
William Richardson, MD, Co-Chair
Robert Fanelli, MD, Chair

It was reviewed and approved by the Board of Governors of the Society of American Gastrointestinal and Endoscopic Surgeons (SAGES), January 2010.

Requests for prints should be sent to:

Society of American Gastrointestinal and Endoscopic Surgeons (SAGES)

11300 West Olympic Blvd., Suite 600

Los Angeles, CA 90064, USA Tel.: ?1 310 437 0544

Fax: ?1 310 437 0585

Email: publications@sages.org

Web: <http://www.sages.org/>

This is a revision of a SAGES publication, which was printed November 2002.

REFERENCES

1. NIH Releases (1992) Consensus statement on gallstones, bile duct stones and laparoscopic cholecystectomy. *Am Fam Physician* 46:1571–1574
2. Patel JA, Patel NA, Piper GL, Smith DE III, Malhotra G, Colella JJ (2009) Perioperative management of cholelithiasis in patients presenting for laparoscopic Roux-en-Y gastric bypass: have we reached a consensus? *Am Surg* 75:470–476 (discussion 6)
3. Gurusamy KS, Samraj K, Fusai G, Davidson BR (2008) Early versus delayed laparoscopic cholecystectomy for biliary colic. *Cochrane Database Syst Rev* CD007196
4. Halldestam I, Kullman E, Borch K (2008) Defined indications for elective cholecystectomy for gallstone disease. *Br J Surg* 95:620–626
5. Gourgiotis S, Dimopoulos N, Germanos S, Vougas V, Alfaras P, Hadjiyannakis E (2007) Laparoscopic cholecystectomy: a safe approach for management of acute cholecystitis. *JLS* 11:219–224
6. Curro G, Baccarani U, Adani G, Cucinotta E (2007) Laparoscopic cholecystectomy in patients with mild cirrhosis and symptomatic cholelithiasis. *Transplant Proc* 39:1471–1473
7. Heinrich S, Schafer M, Rousson V, Clavien PA (2006) Evidence-based treatment of acute pancreatitis: a look at established paradigms. *Ann Surg* 243:154–168
8. Choudhary A, Bechtold ML, Puli SR, Othman MO, Roy PK (2008) Role of prophylactic antibiotics in laparoscopic cholecystectomy: a meta-analysis. *J Gastrointest Surg* 12:1847–1853 (discussion 53)
9. Chang WT, Lee KT, Chuang SC et al (2006) The impact of prophylactic antibiotics on postoperative infection complication in elective laparoscopic cholecystectomy: a prospective randomized study. *Am J Surg* 191:721–725
10. Dervisoglou A, Tsioufas S, Kanellakopoulou K et al (2006) The value of chemoprophylaxis against *Enterococcus* species in elective cholecystectomy: a randomized study of cefuroxime vs ampicillin-sulbactam. *Arch Surg* 141:1162–1167
11. Steinberg JP, Braun BI, Hellinger WC et al (2009) Timing of antimicrobial prophylaxis and the risk of surgical site infections: results from the Trial to Reduce Antimicrobial Prophylaxis Errors. *Ann Surg* 250:10–16
12. Guidelines for deep venous thrombosis prophylaxis during laparoscopic surgery (2007) *Surg Endosc* 21:1007–1009.
13. Scott-Conner CEH, ed (2005) *The SAGES manual: fundamentals of laparoscopy, thoracoscopy, and GI endoscopy*, 2 edn. Birkhäuser, Springer Science & Business Media, Inc., New York
14. Ahmad G, Duffy JM, Phillips K, Watson A (2008) Laparoscopic entry techniques. *Cochrane Database Syst Rev* CD006583
15. Larobina M, Nottle P (2005) Complete evidence regarding major vascular injuries during laparoscopic access. *Surg Laparosc Endosc Percutan Tech* 15:119–123
16. Dekker SW, Hugh TB (2008) Laparoscopic bile duct injury: understanding the psychology and heuristics of the error. *ANZ J Surg* 78:1109–1114
17. Kholdebarin R, Boetto J, Harnish JL, Urbach DR (2008) Risk factors for bile duct injury during laparoscopic cholecystectomy: a case-control study. *Surg Innov* 15:114–119
18. Singh K, Ohri A (2006) Anatomic landmarks: their usefulness in safe laparoscopic cholecystectomy. *Surg Endosc* 20:1754–1758



19. Yegiyants S, Collins JC (2008) Operative strategy can reduce the incidence of major bile duct injury in laparoscopic cholecystectomy. *Am Surg* 74:985–987
20. Avgerinos C, Kelgiorgi D, Touloumis Z, Baltatzi L, Dervenis C (2009) One thousand laparoscopic cholecystectomies in a single surgical unit using the “critical view of safety” technique. *J Gastrointest Surg* 13(3):498–503
21. Debru E, Dawson A, Leibman S et al (2005) Does routine intraoperative cholangiography prevent bile duct transection? *Surg Endosc* 19:589–593
22. Waage A, Nilsson M (2006) Iatrogenic bile duct injury: a population-based study of 152 776 cholecystectomies in the Swedish Inpatient Registry. *Arch Surg* 141:1207–1213
23. Nuzzo G, Giuliante F, Giovannini I et al (2005) Bile duct injury during laparoscopic cholecystectomy: results of an Italian national survey on 56 591 cholecystectomies. *Arch Surg* 140:986–992
24. Way LW, Stewart L, Gantert W et al (2003) Causes and prevention of laparoscopic bile duct injuries: analysis of 252 cases from a human factors and cognitive psychology perspective. *Ann Surg* 237:460–469
25. Paganini AM, Guerrieri M, Sarnari J et al (2007) Thirteen years’ experience with laparoscopic transcystic common bile duct exploration for stones. Effectiveness and long-term results. *Surg Endosc* 21:34–40
26. Hamouda AH, Goh W, Mahmud S, Khan M, Nassar AH (2007) Intraoperative cholangiography facilitates simple transcystic clearance of ductal stones in units without expertise for laparoscopic bile duct surgery. *Surg Endosc* 21:955–959
27. Wenner DE, Whitwam P, Turner D, Chadha A, Degani J (2006) Laparoscopic cholecystectomy and management of biliary tract stones in a freestanding ambulatory surgery center. *JLS* 10:47–51
28. Lacitignola S, Minardi M (2008) Management of common bile duct stones: a ten-year experience at a tertiary care center. *JLS* 12:62–65
29. Bertolin-Bernades R, Sabater-Orti L, Calvete-Chornet J et al (2007) Mild acute biliary pancreatitis vs cholelithiasis: are there differences in the rate of choledocholithiasis? *J Gastrointest Surg* 11:875–879
30. Metcalfe MS, Ong T, Bruening MH, Iswariah H, WemyssHolden SA, Maddern GJ (2004) Is laparoscopic intraoperative cholangiogram a matter of routine? *Am J Surg* 187:475–481
31. Tinoco R, Tinoco A, El-Kadre L, Peres L, Sueth D (2008) Laparoscopic common bile duct exploration. *Ann Surg* 247:674–679
32. Topal B, Fieuws S, Tomczyk K et al (2009) Clinical models are inaccurate in predicting bile duct stones in situ for patients with gallbladder. *Surg Endosc* 23:38–44
33. Perry KA, Myers JA, Deziel DJ (2008) Laparoscopic ultrasound as the primary method for bile duct imaging during cholecystectomy. *Surg Endosc* 22:208–213
34. Machi J, Oishi AJ, Tajiri T, Murayama KM, Furumoto NL, Oishi RH (2007) Routine laparoscopic ultrasound can significantly reduce the need for selective intraoperative cholangiography during cholecystectomy. *Surg Endosc* 21:270–274
35. Machi J, Johnson JO, Deziel DJ et al (2009) The routine use of laparoscopic ultrasound decreases bile duct injury: a multicenter study. *Surg Endosc* 23:384–388
36. Hakamada K, Narumi S, Toyoki Y et al (2008) Intraoperative ultrasound as an educational guide for laparoscopic biliary surgery. *World J Gastroenterol* 14:2370–2376
37. Martin DJ, Vernon DR, Toouli J (2006) Surgical versus endoscopic treatment of bile duct stones. *Cochrane Database Syst Rev* (2):CD003327
38. Kharbutli B, Velanovich V (2008) Management of preoperatively suspected choledocholithiasis: a decision analysis. *J Gastrointest Surg* 12:1973–1980
39. Clayton ES, Connor S, Alexakis N, Leandros E (2006) Metaanalysis of endoscopy and surgery versus surgery alone for common bile duct stones with the gallbladder in situ. *Br J Surg* 93:1185–1191
40. Williams EJ, Green J, Beckingham I, Parks R, Martin D, Lombard M (2008) Guidelines on the management of common bile duct stones (CBDS). *Gut* 57:1004–1021
41. Bingener J, Schwesinger WH (2006) Management of common bile duct stones in a rural area of the United States: results of a survey. *Surg Endosc* 20:577–579
42. Schroepel TJ, Lambert PJ, Mathiason MA, Kothari SN (2007) An economic analysis of hospital charges for choledocholithiasis by different treatment strategies. *Am Surg* 73:472–477
43. Poulouse BK, Speroff T, Holzman MD (2007) Optimizing choledocholithiasis management: a cost-effectiveness analysis. *Arch Surg* 142:43–48 (discussion 9)
44. Poulouse BK, Arbogast PG, Holzman MD (2006) National analysis of in-hospital resource utilization in choledocholithiasis management using propensity scores. *Surg Endosc* 20:186–190
45. Topal B, Aerts R, Penninckx F (2007) Laparoscopic common bile duct stone clearance with flexible choledochoscopy. *Surg Endosc* 21:2317–2321
46. Tang CN, Tsui KK, Ha JP, Siu WT, Li MK (2006) Laparoscopic exploration of the common bile duct: 10-year experience of 174 patients from a single centre. *Hong Kong Med J* 12:191–196
47. Stromberg C, Nilsson M, Leijonmarck CE (2008) Stone clearance and risk factors for failure in laparoscopic transcystic exploration of the common bile duct. *Surg Endosc* 22:1194–1199
48. Campbell-Lloyd AJ, Martin DJ, Martin IJ (2008) Long-term outcomes after laparoscopic bile duct exploration: a 5-year follow up of 150 consecutive patients. *ANZ J Surg* 78:492–494
49. Taylor CJ, Kong J, Ghusn M, White S, Crampton N, Layani L (2007) Laparoscopic bile duct exploration: results of 160 consecutive cases with 2-year follow up. *ANZ J Surg* 77:440–445



50. Alhamdani A, Mahmud S, Jameel M, Baker A (2008) Primary closure of choledochotomy after emergency laparoscopic common bile duct exploration. *Surg Endosc* 22:2190–2195
51. Kanamaru T, Sakata K, Nakamura Y, Yamamoto M, Ueno N, Takeyama Y (2007) Laparoscopic choledochotomy in management of choledocholithiasis. *Surg Laparosc Endosc Percutan Tech* 17:262–266
52. Karaliotas C, Sgourakis G, Goumas C, Papaioannou N, Lilis C, Leandros E (2008) Laparoscopic common bile duct exploration after failed endoscopic stone extraction. *Surg Endosc* 22:1826–1831
53. Jameel M, Darmas B, Baker AL (2008) Trend towards primary closure following laparoscopic exploration of the common bile duct. *Ann R Coll Surg Engl* 90:29–35
54. O'Neill CJ, Gillies DM, Gani JS (2008) Choledocholithiasis: overdiagnosed endoscopically and undertreated laparoscopically. *ANZ J Surg* 78:487–491
55. Gersin KS, Fanelli RD (1998) Laparoscopic endobiliary stenting as an adjunct to common bile duct exploration. *Surg Endosc* 12:301–304
56. Fanelli RD, Gersin KS (2001) Laparoscopic endobiliary stenting: a simplified approach to the management of occult common bile duct stones. *J Gastrointest Surg* 5:74–80
57. Fanelli RD, Gersin KS, Mainella MT (2002) Laparoscopic endobiliary stenting significantly improves success of postoperative endoscopic retrograde cholangiopancreatography in low-volume centers. *Surg Endosc* 16:487–491
58. Costi R, Mazzeo A, Tartamella F, Manceau C, Vacher B, Valverde A (2010) Cholecystocholedocholithiasis: a case-control study comparing the short and long-term outcomes for a “laparoscopy-first” attitude with the outcome for sequential treatment (systematic endoscopic sphincterotomy followed by laparoscopic cholecystectomy). *Surg Endosc* 24(1):51–62
59. Ahmed AR, Husain S, Saad N, Patel NC, Waldman DL, O'Malley W (2007) Accessing the common bile duct after Roux-en-Y gastric bypass. *Surg Obes Relat Dis* 3:640–643
60. Neri V, Ambrosi A, Fersini A, Tartaglia N, Valentino TP (2007) Antegrade dissection in laparoscopic cholecystectomy. *JLS* 11:225–228
61. Cengiz Y, Janes A, Grehn A, Israelsson LA (2005) Randomized trial of traditional dissection with electrocautery versus ultrasonic fundus-first dissection in patients undergoing laparoscopic cholecystectomy. *Br J Surg* 92:810–813
62. Fullum TM, Kim S, Dan D, Turner PL (2005) Laparoscopic “dome-down” cholecystectomy with the LCS-5 Harmonic scalpel. *JLS* 9:51–57
63. Bessa SS, Al-Fayoumi TA, Katri KM, Awad AT (2008) Clipless laparoscopic cholecystectomy by ultrasonic dissection. *J Laparoendosc Adv Surg Tech A* 18:593–598
64. Caliskan K, Nursal TZ, Yildirim S et al (2006) Hydrodissection with adrenaline-lidocaine-saline solution in laparoscopic cholecystectomy. *Langenbecks Arch Surg* 391:359–363
65. Gurusamy KS, Samraj K, Mullerat P, Davidson BR (2007) Routine abdominal drainage for uncomplicated laparoscopic cholecystectomy. *Cochrane Database Syst Rev* (4):CD006004
66. Tzovaras G, Liakou P, Fafoulakis F, Baloyiannis I, Zacharoulis D, Hatzitheofilou C (2009) Is there a role for drain use in elective laparoscopic cholecystectomy? A controlled randomized trial. *Am J Surg* 197:759–763
67. Zhang WJ, Li JM, Wu GZ, Luo KL, Dong ZT (2008) Risk factors affecting conversion in patients undergoing laparoscopic cholecystectomy. *ANZ J Surg* 78:973–976
68. Del Rio P, Dell'Abate P, Soliani P, Sivelli R, Sianesi M (2006) Videolaparoscopic cholecystectomy for acute cholecystitis: analyzing conversion risk factors. *J Laparoendosc Adv Surg Tech A* 16:105–107
69. Kauvar DS, Brown BD, Braswell AW, Harnisch M (2005) Laparoscopic cholecystectomy in the elderly: increased operative complications and conversions to laparotomy. *J Laparoendosc Adv Surg Tech A* 15:379–382
70. Nachnani J, Supe A (2005) Pre-operative prediction of difficult laparoscopic cholecystectomy using clinical and ultrasonographic parameters. *Indian J Gastroenterol* 24:16–18
71. Simopoulos C, Botaitis S, Polychronidis A, Tripsianis G, Karayiannakis AJ (2005) Risk factors for conversion of laparoscopic cholecystectomy to open cholecystectomy. *Surg Endosc* 19:905–909
72. Visser BC, Parks RW, Garden OJ (2008) Open cholecystectomy in the laparoendoscopic era. *Am J Surg* 195:108–114
73. Al Salamah SM (2005) Outcome of laparoscopic cholecystectomy in acute cholecystitis. *J Coll Physicians Surg Pak* 15:400–403
74. Fuller J, Ashar BS, Carey-Corradro J (2005) Trocar-associated injuries and fatalities: an analysis of 1399 reports to the FDA. *J Minim Invasive Gynecol* 12:302–307
75. Shamiyeh A, Wayand W (2004) Laparoscopic cholecystectomy: early and late complications and their treatment. *Langenbecks Arch Surg* 389:164–171
76. Tantia O, Jain M, Khanna S, Sen B (2008) Iatrogenic biliary injury: 13, 305 cholecystectomies experienced by a single surgical team over more than 13 years. *Surg Endosc* 22:1077–1086
77. Karvonen J, Gullichsen R, Laine S, Salminen P, Gronroos JM (2007) Bile duct injuries during laparoscopic cholecystectomy: primary and long-term results from a single institution. *Surg Endosc* 21:1069–1073
78. Diamantis T, Tsigris C, Kiriakopoulos A et al (2005) Bile duct injuries associated with laparoscopic and open cholecystectomy: an 11-year experience in one institute. *Surg Today* 35:841–845
79. Gurusamy KS, Samraj K (2006) Early versus delayed laparoscopic cholecystectomy for acute cholecystitis. *Cochrane Database Syst Rev* (4):CD005440
80. Soderlund C, Frozanpor F, Linder S (2005) Bile duct injuries at laparoscopic cholecystectomy: a single-institution prospective study. Acute cholecystitis indicates an increased risk. *World J Surg* 29:987–993



81. Li J, Frilling A, Nadalin S, Paul A, Malago M, Broelsch CE (2008) Management of concomitant hepatic artery injury in patients with iatrogenic major bile duct injury after laparoscopic cholecystectomy. *Br J Surg* 95:460–465
82. Stewart L, Robinson TN, Lee CM, Liu K, Whang K, Way LW (2004) Right hepatic artery injury associated with laparoscopic bile duct injury: incidence, mechanism, and consequences. *J Gastrointest Surg* 8:523–530 (discussion 30–31)
83. Bektas H, Schrem H, Winny M, Klempnauer J (2007) Surgical treatment and outcome of iatrogenic bile duct lesions after cholecystectomy and the impact of different clinical classification systems. *Br J Surg* 94:1119–1127
84. Schmidt SC, Langrehr JM, Hintze RE, Neuhaus P (2005) Longterm results and risk factors influencing outcome of major bile duct injuries following cholecystectomy. *Br J Surg* 92:76–82
85. Sicklick JK, Camp MS, Lillemoe KD et al (2005) Surgical management of bile duct injuries sustained during laparoscopic cholecystectomy: perioperative results in 200 patients. *Ann Surg* 241:786–792 (discussion 93–95)
86. Nuzzo G, Giuliani F, Giovannini I et al (2008) Advantages of multidisciplinary management of bile duct injuries occurring during cholecystectomy. *Am J Surg* 195:763–769
87. Sabbaghian MS, Rich BS, Rothberger GD et al (2008) Evaluation of surgical outcomes and gallbladder characteristics in patients with biliary dyskinesia. *J Gastrointest Surg* 12:1324–1330
88. Bingener J, Richards ML, Schwesinger WH, Sirinek KR (2004) Laparoscopic cholecystectomy for biliary dyskinesia: correlation of preoperative cholecystokinin cholescintigraphy results with postoperative outcome. *Surg Endosc* 18:802–806
89. Geiger TM, Awad ZT, Burgard M et al (2008) Prognostic indicators of quality of life after cholecystectomy for biliary dyskinesia. *Am Surg* 74:400–404
90. Paaajanen H, Miilunpohja S, Joukainen S, Heikkinen J (2009) Role of quantitative cholescintigraphy for planning laparoscopic cholecystectomy in patients with gallbladder dyskinesia and chronic abdominal pain. *Surg Laparosc Endosc Percutan Tech* 19:16–19
91. Ponsky TA, DeSagun R, Brody F (2005) Surgical therapy for biliary dyskinesia: a meta-analysis and review of the literature. *J Laparoendosc Adv Surg Tech A* 15:439–442
92. Gurusamy KS, Junnarkar S, Farouk M, Davidson BR (2009) Cholecystectomy for suspected gallbladder dyskinesia. *Cochrane Database Syst Rev* (1):CD007086
93. Ainsworth AP, Adamsen S, Rosenberg J (2007) Surgery for acute cholecystitis in Denmark. *Scand J Gastroenterol* 42:648–651
94. Yamashita Y, Takada T, Hirata K (2006) A survey of the timing and approach to the surgical management of patients with acute cholecystitis in Japanese hospitals. *J Hepatobiliary Pancreat Surg* 13:409–415
95. Csiksz NG, Tseng JF, Shah SA (2008) Trends in surgical management for acute cholecystitis. *Surgery* 144:283–289
96. Campbell EJ, Montgomery DA, Mackay CJ (2008) A national survey of current surgical treatment of acute gallstone disease. *Surg Laparosc Endosc Percutan Tech* 18:242–247
97. Campbell EJ, Montgomery DA, MacKay CJ (2007) A survey of current surgical treatment of acute gallstone disease in the west of Scotland. *Scott Med J* 52:15–19
98. Siddiqui T, MacDonald A, Chong PS, Jenkins JT (2008) Early versus delayed laparoscopic cholecystectomy for acute cholecystitis: a meta-analysis of randomized clinical trials. *Am J Surg* 195:40–47
99. Yamashita Y, Takada T, Kawarada Y et al (2007) Surgical treatment of patients with acute cholecystitis: Tokyo Guidelines. *J Hepatobiliary Pancreat Surg* 14:91–97
100. Lam CM, Yuen AW, Chik B, Wai AC, Fan ST (2005) Variation in the use of laparoscopic cholecystectomy for acute cholecystitis: a population-based study. *Arch Surg* 140:1084–1088
101. Shikata S, Noguchi Y, Fukui T (2005) Early versus delayed cholecystectomy for acute cholecystitis: a meta-analysis of randomized controlled trials. *Surg Today* 35:553–560
102. Soffer D, Blackburne LH, Schulman CI et al (2007) Is there an optimal time for laparoscopic cholecystectomy in acute cholecystitis? *Surg Endosc* 21:805–809
103. Stevens KA, Chi A, Lucas LC, Porter JM, Williams MD (2006) Immediate laparoscopic cholecystectomy for acute cholecystitis: no need to wait. *Am J Surg* 192:756–761
104. Tzavaras G, Zacharoulis D, Liakou P, Theodoropoulos T, Paroutoglou G, Hatzitheofilou C (2006) Timing of laparoscopic cholecystectomy for acute cholecystitis: a prospective non randomized study. *World J Gastroenterol* 12:5528–5531
105. Chau CH, Siu WT, Tang CN et al (2006) Laparoscopic cholecystectomy for acute cholecystitis: the evolving trend in an institution. *Asian J Surg* 29:120–124
106. Wang YC, Yang HR, Chung PK, Jeng LB, Chen RJ (2006) Urgent laparoscopic cholecystectomy in the management of acute cholecystitis: timing does not influence conversion rate. *Surg Endosc* 20:806–808
107. Kolla SB, Aggarwal S, Kumar A et al (2004) Early versus delayed laparoscopic cholecystectomy for acute cholecystitis: a prospective randomized trial. *Surg Endosc* 18:1323–1327
108. Johansson M, Thune A, Nelvin L, Stiernstam M, Westman B, Lundell L (2005) Randomized clinical trial of open versus laparoscopic cholecystectomy in the treatment of acute cholecystitis. *Br J Surg* 92:44–49
109. Hadad SM, Vaidya JS, Baker L, Koh HC, Heron TP, Thompson AM (2007) Delay from symptom onset increases the conversion rate in laparoscopic cholecystectomy for acute cholecystitis. *World J Surg* 31:1298–1301 (discussion 302–303)
110. Lee AY, Carter JJ, Hochberg MS, Stone AM, Cohen SL, Pachter HL (2008) The timing of surgery for cholecystitis: a review of 202 consecutive patients at a large municipal hospital. *Am J Surg* 195:467–470



111. Casillas RA, Yegiyants S, Collins JC (2008) Early laparoscopic cholecystectomy is the preferred management of acute cholecystitis. *Arch Surg* 143:533–537
112. Daniak CN, Peretz D, Fine JM, Wang Y, Meinke AK, Hale WB (2008) Factors associated with time to laparoscopic cholecystectomy for acute cholecystitis. *World J Gastroenterol* 14:1084–1090
113. Lau H, Lo CY, Patil NG, Yuen WK (2006) Early versus delayed-interval laparoscopic cholecystectomy for acute cholecystitis: a metaanalysis. *Surg Endosc* 20:82–87
114. Asoglu O, Ozmen V, Karanlik H et al (2004) Does the complication rate increase in laparoscopic cholecystectomy for acute cholecystitis? *J Laparoendosc Adv Surg Tech A* 14:81–86
115. Macri A, Scuderi G, Saladino E et al (2006) Acute gallstone cholecystitis in the elderly: treatment with emergency ultrasonographic percutaneous cholecystostomy and interval laparoscopic cholecystectomy. *Surg Endosc* 20:88–91
116. Leveau P, Andersson E, Carlgren I, Willner J, Andersson R (2008) Percutaneous cholecystostomy: a bridge to surgery or definite management of acute cholecystitis in high-risk patients? *Scand J Gastroenterol* 43:593–596
117. Akyurek N, Salman B, Yuksel O et al (2005) Management of acute calculous cholecystitis in high-risk patients: percutaneous cholecystotomy followed by early laparoscopic cholecystectomy. *Surg Laparosc Endosc Percutan Tech* 15:315–320
118. Welschbillig-Meunier K, Pessaux P, Lebigot J et al (2005) Percutaneous cholecystostomy for high-risk patients with acute cholecystitis. *Surg Endosc* 19:1256–1259
119. Teoh WM, Cade RJ, Banting SW, Mackay S, Hassen AS (2005) Percutaneous cholecystostomy in the management of acute cholecystitis. *ANZ J Surg* 75:396–398
120. Li JC, Lee DW, Lai CW, Li AC, Chu DW, Chan AC (2004) Percutaneous cholecystostomy for the treatment of acute cholecystitis in the critically ill and elderly. *Hong Kong Med J* 10:389–393
121. Tsumura H, Ichikawa T, Hiyama E et al (2004) An evaluation of laparoscopic cholecystectomy after selective percutaneous transhepatic gallbladder drainage for acute cholecystitis. *Gastrointest Endosc* 59:839–844
122. Kirshtein B, Bayme M, Bolotin A, Mizrahi S, Lantsberg L (2008) Laparoscopic cholecystectomy for acute cholecystitis in the elderly: is it safe? *Surg Laparosc Endosc Percutan Tech* 18:334–339
123. do Amaral PC, Azaro Filho Ede M, Galvao TD et al (2006) Laparoscopic cholecystectomy for acute cholecystitis in elderly patients. *JLS* 10:479–483
124. Lowenfels AB, Maisonneuve P, Sullivan T (2009) The changing character of acute pancreatitis: epidemiology, etiology, and prognosis. *Curr Gastroenterol Rep* 11:97–103
125. Frey CF, Zhou H, Harvey DJ, White RH (2006) The incidence and case-fatality rates of acute biliary, alcoholic, and idiopathic pancreatitis in California, 1994–2001. *Pancreas* 33:336–344
126. Kimura Y, Takada T, Kawarada Y et al (2006) JPN Guidelines for the management of acute pancreatitis: treatment of gallstone-induced acute pancreatitis. *J Hepatobiliary Pancreat Surg* 13:56–60
127. Working Party of the British Society of Gastroenterology, Association of Surgeons of Great Britain and Ireland; Pancreatic Society of Great Britain and Ireland, Association of Upper GI Surgeons of Great Britain and Ireland (2005) UK guidelines for the management of acute pancreatitis. *Gut* 54(Suppl 3):iii1–9
128. Nebiker CA, Frey DM, Hamel CT, Oertli D, Kettelhack C (2009) Early versus delayed cholecystectomy in patients with biliary acute pancreatitis. *Surgery* 145:260–264
129. Sinha R (2008) Early laparoscopic cholecystectomy in acute biliary pancreatitis: the optimal choice? *HPB (Oxford)* 10:332–335
130. Chiang DT, Thompson G (2008) Management of acute gallstone pancreatitis: so the story continues. *ANZ J Surg* 78:52–54
131. Singhal T, Balakrishnan S, Grandy-Smith S, Hunt J, Asante M, El-Hasani S (2006) Gallstones: best served hot. *JLS* 10:332–335
132. Taylor E, Wong C (2004) The optimal timing of laparoscopic cholecystectomy in mild gallstone pancreatitis. *Am Surg* 70:971–975
133. Cameron DR, Goodman AJ (2004) Delayed cholecystectomy for gallstone pancreatitis: re-admissions and outcomes. *Ann R Coll Surg Engl* 86:358–362
134. Lakatos L, Mester G, Reti G, Nagy A, Lakatos PL (2004) Selection criteria for preoperative endoscopic retrograde cholangiopancreatography before laparoscopic cholecystectomy and endoscopic treatment of bile duct stones: results of a retrospective, single center study between 1996–2002. *World J Gastroenterol* 10:3495–3499
135. Jackson H, Granger S, Price R et al (2008) Diagnosis and laparoscopic treatment of surgical diseases during pregnancy: an evidence-based review. *Surg Endosc* 22:1917–1927
136. Festi D, Dormi A, Capodicasa S et al (2008) Incidence of gallstone disease in Italy: results from a multicenter, population-based Italian study (the MICOL project). *World J Gastroenterol* 14:5282–5289
137. Chen CH, Huang MH, Yang JC et al (2006) Prevalence and risk factors of gallstone disease in an adult population of Taiwan: an epidemiological survey. *J Gastroenterol Hepatol* 21:1737–1743
138. Zhang Y, Liu D, Ma Q, Dang C, Wei W, Chen W (2006) Factors influencing the prevalence of gallstones in liver cirrhosis. *J Gastroenterol Hepatol* 21:1455–1458
139. Bingener J, Cox D, Michalek J, Mejia A (2008) Can the MELD score predict perioperative morbidity for patients with liver cirrhosis undergoing laparoscopic cholecystectomy? *Am Surg* 74:156–159
140. Curro G, Iapichino G, Melita G, Lorenzini C, Cucinotta E (2005) Laparoscopic cholecystectomy in Child–Pugh class C cirrhotic patients. *JLS* 9:311–315



141. Mancero JM, D'Albuquerque LA, Gonzalez AM, Larrea FI, de Oliveira e Silva A (2008) Laparoscopic cholecystectomy in cirrhotic patients with symptomatic cholelithiasis: a case-control study. *World J Surg* 32:267–270
142. Leandros E, Albanopoulos K, Tsigris C et al (2008) Laparoscopic cholecystectomy in cirrhotic patients with symptomatic gallstone disease. *ANZ J Surg* 78:363–365
143. Schiff J, Misra M, Rendon G, Rothschild J, Schwaitzberg S (2005) Laparoscopic cholecystectomy in cirrhotic patients. *Surg Endosc* 19:1278–1281
144. Ji W, Li LT, Chen XR, Li JS (2004) Application of laparoscopic cholecystectomy in patients with cirrhotic portal hypertension. *Hepatobiliary Pancreat Dis Int* 3:270–274
145. Ji W, Li LT, Wang ZM, Quan ZF, Chen XR, Li JS (2005) A randomized controlled trial of laparoscopic versus open cholecystectomy in patients with cirrhotic portal hypertension. *World J Gastroenterol* 11:2513–2517
146. Palanivelu C, Rajan PS, Jani K et al (2006) Laparoscopic cholecystectomy in cirrhotic patients: the role of subtotal cholecystectomy and its variants. *J Am Coll Surg* 203:145–151
147. Philips JA, Lawes DA, Cook AJ et al (2008) The use of laparoscopic subtotal cholecystectomy for complicated cholelithiasis. *Surg Endosc* 22:1697–1700
148. Ercan M, Bostanci EB, Ozer I et al (2010) Postoperative hemorrhagic complications after elective laparoscopic cholecystectomy in patients receiving long-term anticoagulant therapy. *Langenbecks Arch Surg* 395(3):247–253
149. Leandros E, Gomatos IP, Mami P, Kastellanos E, Albanopoulos K, Konstadoulakis MM (2005) Elective laparoscopic cholecystectomy for symptomatic gallstone disease in patients receiving anticoagulant therapy. *J Laparoendosc Adv Surg Tech A* 15:357–360
150. Stephen AE, Berger DL (2001) Carcinoma in the porcelain gallbladder: a relationship revisited. *Surgery* 129:699–703
151. Kwon AH, Inui H, Matsui Y, Uchida Y, Hukui J, Kamiyama Y (2004) Laparoscopic cholecystectomy in patients with porcelain gallbladder based on the preoperative ultrasound findings. *Hepatogastroenterology* 51:950–953
152. Zielinski MD, Atwell TD, Davis PW, Kendrick ML, Que FG (2009) Comparison of surgically resected polypoid lesions of the gallbladder to their pre-operative ultrasound characteristics. *J Gastrointest Surg* 13:19–25
153. Kratzer W, Haenle MM, Voegtle A et al (2008) Ultrasonographically detected gallbladder polyps: a reason for concern? A seven-year follow-up study. *BMC Gastroenterol* 8:41
154. Lin WR, Lin DY, Tai DI et al (2008) Prevalence of and risk factors for gallbladder polyps detected by ultrasonography among healthy Chinese: analysis of 34 669 cases. *J Gastroenterol Hepatol* 23:965–969
155. Aldouri AQ, Malik HZ, Waytt J et al (2009) The risk of gallbladder cancer from polyps in a large multiethnic series. *Eur J Surg Oncol* 35:48–51
156. Colecchia A, Larocca A, Scaioli E et al (2009) Natural history of small gallbladder polyps is benign: evidence from a clinical and pathogenetic study. *Am J Gastroenterol* 104:624–629
157. Gurusamy KS, Abu-Amara M, Farouk M, Davidson BR (2009) Cholecystectomy for gallbladder polyp. *Cochrane Database Syst Rev* (1):CD007052
158. Duffy A, Capanu M, Abou-Alfa GK et al (2008) Gallbladder cancer (GBC): 10-year experience at Memorial Sloan-Kettering Cancer Centre (MSKCC). *J Surg Oncol* 98:485–489
159. Shih SP, Schulick RD, Cameron JL et al (2007) Gallbladder cancer: the role of laparoscopy and radical resection. *Ann Surg* 245:893–901
160. You DD, Lee HG, Paik KY, Heo JS, Choi SH, Choi DW (2008) What is an adequate extent of resection for T1 gallbladder cancers? *Ann Surg* 247:835–838
161. Kang CM, Lee WJ, Choi GH et al (2007) Does “clinical” R0 have validity in the choice of simple cholecystectomy for gallbladder carcinoma? *J Gastrointest Surg* 11:1309–1316
162. Kang CM, Choi GH, Park SH et al (2007) Laparoscopic cholecystectomy only could be an appropriate treatment for selected clinical R0 gallbladder carcinoma. *Surg Endosc* 21:1582–1587
163. Cucinotta E, Lorenzini C, Melita G, Iapichino G, Curro G (2005) Incidental gall bladder carcinoma: does the surgical approach influence the outcome? *ANZ J Surg* 75:795–798
164. Kwon AH, Imamura A, Kitade H, Kamiyama Y (2008) Unsuspected gallbladder cancer diagnosed during or after laparoscopic cholecystectomy. *J Surg Oncol* 97:241–245
165. Chan KM, Yeh TS, Jan YY, Chen MF (2006) Laparoscopic cholecystectomy for early gallbladder carcinoma: long-term outcome in comparison with conventional open cholecystectomy. *Surg Endosc* 20:1867–1871
166. Yamamoto H, Hayakawa N, Kitagawa Y et al (2005) Unsuspected gallbladder carcinoma after laparoscopic cholecystectomy. *J Hepatobiliary Pancreat Surg* 12:391–398
167. Tenconi SM, Boni L, Colombo EM, Dionigi G, Rovera F, Cassinotti E (2008) Laparoscopic cholecystectomy as day-surgery procedure: current indications and patients' selection. *Int J Surg* 6(Suppl 1):S86–S88
168. Psaila J, Agrawal S, Fountain U et al (2008) Day-surgery laparoscopic cholecystectomy: factors influencing same-day discharge. *World J Surg* 32:76–81
169. Chauhan A, Mehrotra M, Bhatia PK, Baj B, Gupta AK (2006) Day care laparoscopic cholecystectomy: a feasibility study in a public health service hospital in a developing country. *World J Surg* 30:1690–1695 (discussion 6–7)
170. Sherigar JM, Irwin GW, Rathore MA, Khan A, Pillow K, Brown MG (2006) Ambulatory laparoscopic cholecystectomy outcomes. *JSL* 10:473–478



171. Kavanagh T, Hu P, Minogue S (2008) Daycase laparoscopic cholecystectomy: a prospective study of post-discharge pain, analgesic and antiemetic requirements. *Ir J Med Sci* 177:111–115
172. Kasem A, Paix A, Grandy-Smith S, El-Hasani S (2006) Is laparoscopic cholecystectomy safe and acceptable as a day case procedure? *J Laparoendosc Adv Surg Tech A* 16:365–368
173. Chok KS, Yuen WK, Lau H, Lee F, Fan ST (2004) Outpatient laparoscopic cholecystectomy in Hong Kong Chinese—an outcome analysis. *Asian J Surg* 27:313–316
174. Bueno Lledo J, Planells Roig M, Arnau Bertomeu C et al (2006) Outpatient laparoscopic cholecystectomy: a new gold standard for cholecystectomy. *Rev Esp Enferm Dig* 98:14–24
175. Hodgett SE, Hernandez JM, Morton CA, Ross SB, Albrink M, Rosemurgy AS (2009) Laparoendoscopic single site (LESS) cholecystectomy. *J Gastrointest Surg* 13:188–192
176. Chow A, Purkayastha S, Aziz O, Paraskeva P (2010) Singleincision laparoscopic surgery for cholecystectomy: an evolving technique. *Surg Endosc* 24(3):709–714
177. Ersin S, Firat O, Sozbilen M (2010) Single-incision laparoscopic cholecystectomy: is it more than a challenge? *Surg Endosc* 24(1):68–71



4º CURSO TEÓRICO-PRÁTICO DE CIRURGIA HEPATO-BILIO-PANCREÁTICA

Thomas E. Starzl Symposium on HPB Surgery

Organização e Segurança em Cirurgia HBP
Porto, 2, 3 e 4 Fevereiro 2012



ORGANIZAÇÃO



UNIDADE
HEPATO-BILIO-PANCREÁTICA E ESPLÉNICA
Serviço de Cirurgia Geral do CHSJ

SECRETARIADO



curso.cirurgiaHBP@eruditus.pt
www.eruditus.pt
tel | +351 214 430 901
mob | +351 917 289 264
mob | +351 918 027 877

PROGRAMA PRELIMINAR

2 Fevereiro 2012

CURSO PRÉ-CONGRESSO

Cultura de Segurança para Equipas Cirúrgicas

3 e 4 Fevereiro 2012

TEMAS

Organização e acreditação
Como atingir zero mortalidade
Cirurgia em maus fígados
Registos
Transplante
Tumores das vias biliares

4 Fevereiro 2012

CURSO PRÁTICO

Centro de Cirurgia Experimental

CONVIDADOS ESTRANGEIROS

Ben-Hur Ferraz Neto
Darius Mirza
Evaristo Varo
Jose Luis Garcia Sabrido
Masatoshi Makuuchi
Paul McMaster

INSCRIÇÃO

4º Curso Teórico-Prático Hepato-Bilio-Pancreático
200€ médicos | 150€ enfermeiros

Curso Segurança (inscritos no 4º Curso HBP) | 50€
**Número limitado a 50 participantes*

Curso Prático (inscritos no 4º Curso HBP) | 100€
**Número limitado 20 participantes*

