



# Salicylic acid signal transduction: the initiation of biosynthesis, perception and transcriptional reprogramming

Carolyn Seyfferth and Kenichi Tsuda\*

Department of Plant Microbe Interactions, Max Planck Institute for Plant Breeding Research, Cologne, Germany

## Edited by:

Hua Lu, University of Maryland at Baltimore County, USA

## Reviewed by:

Murray Grant, University of Exeter, UK

Zhixiang Chen, Purdue University, USA

## \*Correspondence:

Kenichi Tsuda, Department of Plant Microbe Interactions, Max Planck Institute for Plant Breeding Research, Carl-von-Linné-Weg 10, D-50829 Cologne, Germany  
e-mail: tsuda@mpipz.mpg.de

The phytohormone salicylic acid (SA) is a small phenolic compound that regulates diverse physiological processes, in particular plant resistance against pathogens. Understanding SA-mediated signaling has been a major focus of plant research. Pathogen-induced SA is mainly synthesized via the isochorismate pathway in chloroplasts, with ICS1 (ISOCHORISMATE SYNTHASE 1) being a critical enzyme. Calcium signaling regulates activities of a subset of transcription factors thereby activating nuclear *ICS1* expression. The produced SA triggers extensive transcriptional reprogramming in which NPR1 (NON-EXPRESSOR OF PATHOGENESIS-RELATED GENES 1) functions as the central coactivator of TGA transcription factors. Recently, two alternative but not exclusive models for SA perception mechanisms were proposed. The first model is that NPR1 homologs, NPR3 and NPR4, perceive SA thereby regulating NPR1 protein accumulation. The second model describes that NPR1 itself perceives SA, triggering an NPR1 conformational change thereby activating SA-mediated transcription. Besides the direct SA binding, NPR1 is also regulated by SA-mediated redox changes and phosphorylation. Emerging evidence show that pathogen virulence effectors target SA signaling, further strengthening the importance of SA-mediated immunity.

**Keywords:** calcium, ICS1, NPR1, plant immunity, salicylic acid, SA perception, transcriptional reprogramming

## INTRODUCTION

The phytohormone salicylic acid (SA) is a small phenolic compound that functions as an important signaling molecule during plant immunity (Vlot et al., 2009; Robert-Seilaniantz et al., 2011; Pieterse et al., 2012). Since constitutive SA accumulation is often associated with stunted plant growth, resulting in reduction of plant fitness (Ishihara et al., 2008; Pajerowska-Mukhtar et al., 2012; Chandran et al., 2014), SA biosynthesis and SA-mediated signaling are tightly controlled.

The plant immune system comprises multiple layers, such as pattern-triggered immunity (PTI) and effector-triggered immunity (ETI; Jones and Dangl, 2006; Tsuda and Katagiri, 2010). PTI is triggered by recognition of common microbial components (MAMPs, microbe-associated molecular patterns), such as bacterial flagellin or the fungal cell wall component chitin (Boller and Felix, 2009; Macho and Zipfel, 2014). MAMP recognition stimulates generation of reactive oxygen species, intracellular calcium influx, transient activation of mitogen-activated protein kinases (MAPKs), and the production of SA (Tsuda et al., 2008a,b; Tsuda and Katagiri, 2010). Virulent pathogens, for example, the bacterial pathogen *Pseudomonas syringae* pv. *tomato* DC3000 (*Pto* DC3000), however, can suppress PTI in *Arabidopsis* and tomato by effectors, injected via bacterial secretion systems into the plant cell (Lohou et al., 2013; Xin and He, 2013). Recent studies identified various effectors that interfere with SA signaling (Uppalapati

et al., 2007; Djamei et al., 2011; Caillaud et al., 2013; Jiang et al., 2013; Rabe et al., 2013; Gimenez-Ibanez et al., 2014; Liu et al., 2014), highlighting the importance of SA signaling for plant immunity. To regain resistance, plants have acquired intracellular receptors [resistance (R) proteins], which induce the second layer of defense after effector recognition, termed ETI (Eitas and Dangl, 2010; Bonardi and Dangl, 2012; Jacob et al., 2013). Activation of ETI also induces SA accumulation and MAPK activation, which are also important for resistance against pathogens during ETI (Tsuda et al., 2013). Additionally, SA has vital roles in establishing systemic acquired resistance (SAR), a form of long-term and broad-spectrum resistance throughout the entire plant after local pathogen infection (Wang et al., 2006; Fu and Dong, 2013).

In this review, we summarize SA signal transduction from regulation of biosynthesis, perception, to transcriptional reprogramming during plant immunity. We also discuss compensation mechanisms that would provide robust immunity once SA signaling is compromised, for example, by pathogen effector attack. SA signaling pathway is highly interconnected with other phytohormone signaling such as mediated by jasmonates (JA), ethylene, and abscisic acid (Robert-Seilaniantz et al., 2011; Pieterse et al., 2012; Derksen et al., 2013). For example, JA and ethylene signaling negatively regulate SA biosynthesis at the transcriptional level (Chen et al., 2009; Zheng et al., 2012). However, discussions on these are beyond the scope of this review.

## THE BIOSYNTHESIS OF SA IN PLANTS

### BIOSYNTHETIC PATHWAYS

Two major SA biosynthetic pathways in plants were identified: the isochorismate (IC) and the phenylalanine ammonia-lyase (PAL) pathways. Both pathways commonly utilize chorismate, the end product of the shikimate pathway, to produce SA (Dempsey et al., 2011). IC synthase (ICS) and PAL are critical enzymes for these pathways, respectively. Homologs of *ICS* and *PAL* genes are present throughout the plant kingdom, including *Arabidopsis*, tobacco, tomato, populus, sunflower, and pepper (Wildermuth et al., 2001; Cochrane et al., 2004; Uppalapati et al., 2007; Catinot et al., 2008; Yuan et al., 2009; Sadeghi et al., 2013; Kim and Hwang, 2014), suggesting the importance of these SA biosynthesis pathways to survive during the course of evolution. In *Arabidopsis*, mutations in *ICS1* lead to an almost complete loss of pathogen-induced SA accumulation (Wildermuth et al., 2001). However, *Arabidopsis* quadruple *PAL* mutants, in which *PAL* activity is reduced to 10%, also show lower SA accumulation (50%) compared to the wild type upon pathogen infection (Huang et al., 2010). Thus, while contribution of the *PAL* pathway is evident, the *IC* pathway is the major route for SA biosynthesis during plant immunity.

In chloroplasts, *ICS* catalyzes the conversion of chorismate into *IC* (Wildermuth et al., 2001; Strawn et al., 2007; Garcion et al., 2008), which is further converted to SA (Dempsey et al., 2011). In some bacteria, conversion of *IC* to SA is catalyzed by *IC* pyruvate lyases (*IPLs*; Dempsey et al., 2011). However, plant genomes encode no homologous genes to bacterial *IPLs*. Expression of bacterial enzymes catalyzing this conversion together with *ICS* in chloroplasts leads to constitutive accumulation of SA (Verberne et al., 2000; Mauch et al., 2001). Thus, it is conceivable that plants have yet-determined gene(s) whose product(s) possess *IPL* activity in chloroplasts. However, metabolic enzymes such as the acyl acid amido synthetase *GH3.12* [also known as *PBS3/WIN3/GDG1* (*AVRPPHB* SUSCEPTIBLE 3/*HOPW1-INTERACTING 3/GH3-LIKE* DEFENSE GENE 1); Nobuta et al., 2007; Zhang et al., 2007; Okrent et al., 2009; Westfall et al., 2010, 2012] and the acyltransferase *EPS1* (*ENHANCED PSEUDOMONAS* SUSCEPTIBILITY 1; Zheng et al., 2009) are involved in SA accumulation, perhaps by providing SA precursors or regulatory molecules for SA biosynthesis. Thus, SA biosynthesis may be more complex in plants compared to bacteria. SA export from chloroplasts is mediated by the *MATE*-transporter *EDS5* (*ENHANCED DISEASE SUSCEPTIBILITY 5*; Serrano et al., 2013). This export seems important for SA accumulation and distribution in the cell since SA accumulation is compromised in *eds5* mutants (Nawrath et al., 2002; Ishihara et al., 2008).


### REGULATION OF SA BIOSYNTHESIS

Salicylic acid biosynthesis is tightly regulated since constitutive SA accumulation has negative impacts on plant fitness (Ishihara et al., 2008; Pajerowska-Mukhtar et al., 2012; Chandran et al., 2014). Accumulating evidence show that transcriptional control of *ICS1* by calcium signaling is key for the initiation of SA biosynthesis (Figure 1). The concentration of calcium ions ( $\text{Ca}^{2+}$ ) in the cytosol transiently increases upon immune receptor activation through  $\text{Ca}^{2+}$  channels. Elevation of intracellular  $\text{Ca}^{2+}$ , called

$\text{Ca}^{2+}$  signature, is decoded by  $\text{Ca}^{2+}$  sensor proteins, such as calmodulin (*CaM*) and  $\text{Ca}^{2+}$ -dependent protein kinases (*CDPKs*; Dodd et al., 2010; Boudsocq and Sheen, 2013; Poovaiah et al., 2013; Schulz et al., 2013). Binding of *CaM* regulates target protein activities thereby relaying  $\text{Ca}^{2+}$  signatures to downstream responses. During *Arabidopsis* immunity, the *CaM*-binding transcription factor *CBP60g* (*CALMODULIN BINDING PROTEIN 60g*) and its homolog *SARD1* (*SYSTEMIC ACQUIRED RESISTANCE DEFICIENT 1*) control *ICS1* transcription (Wang et al., 2009, 2011; Zhang et al., 2010; Wan et al., 2012). *CaM*-binding is required for *CBP60g* function, whereas *SARD1* does not appear to be a *CaM*-binding protein (Wang et al., 2009). Despite this difference, *CBP60g* and *SARD1* are partially redundant for *ICS1* expression and SA accumulation during immunity. However, dual regulation of *ICS1* transcription by *CBP60g* and *SARD1* seems important for temporal dynamics of SA biosynthesis: *CBP60g* mainly contributes to SA biosynthesis at early stages after *P. syringae* infection while *SARD1* does at late stages (Wang et al., 2011). Another close homolog of *CBP60g*, *CBP60a*, negatively regulates *ICS1* expression upon *CaM*-binding (Truman et al., 2013). Conceivably, upon pathogen attack, *CBP60g* and *SARD1* bind to the *ICS1* promoter and activate its expression, at least partly by removing the negative regulator *CBP60a* from the *ICS1* promoter.

Unlike *CaM*, *CDPKs* have both intrinsic  $\text{Ca}^{2+}$  sensing and responding sites thereby allowing individual *CDPK* proteins to relay  $\text{Ca}^{2+}$  signatures to downstream components via phosphorylation events. Recently, the *CDPKs*, *CPK4*, 5, 6, and 11, were shown to re-localize to the nucleus, and to interact with and phosphorylate the *WRKY* transcription factors, *WRKY8*, 28, and 48, during ETI mediated by the plasma membrane-associated immune receptors *RPS2* (*RESISTANCE TO P.SYRINGAE 2*) or *RPM1* (*RESISTANCE TO P.SYRINGAE PV MACULICULA 1*; Gao et al., 2013). Mutants in *WRKY8* or *WRKY48* are compromised in pathogen-induced *ICS1* expression. Furthermore, *WRKY28* directly interacts with the *ICS1* promoter (van Verk et al., 2011), which might be regulated through phosphorylation by *CPK4*, 5, 6, or 11. Collectively, these results suggest that during ETI, these *CDPKs* relay  $\text{Ca}^{2+}$  signatures to activate *ICS1* transcription via *WRKY* transcription factors.

Besides *ICS1* regulation, calcium signaling also affects the maintenance of SA accumulation through transcriptional regulation of *EDS1* (*ENHANCED DISEASE SUSCEPTIBILITY 1*; Du et al., 2009), encoding a central regulator of the positive feedback loop of SA accumulation (Feys et al., 2001). A *CaM*-binding transcription factor, *CAMTA3/SR1* (*CALMODULIN BINDING TRANSCRIPTION ACTIVATOR 3/SIGNAL-RESPONSIVE GENE 1*), binds to the *EDS1* promoter to repress its transcription, and mutants of *CAMTA3/SR1* show elevated SA levels and enhanced immunity against *P. syringae* and the fungal pathogen *Botrytis cinerea*. Combinatorial mutant analysis indicates that *CAMTA3/SR1* and its homologs *CAMTA1/2* also suppress expression of *CBP60g*, *SARD1*, and *ICS1* (Kim et al., 2013). Thus, the three *CAMTA* homologs coordinately suppress SA accumulation, but it remains unknown if the *CAMTA* transcription factors directly target the promoters of *CBP60g*, *SARD1*, and *ICS1*. It was recently shown



**FIGURE 1 | Regulation of SA accumulation by calcium signaling.** MAMP or effector recognition increases intracellular Ca<sup>2+</sup> concentrations thereby regulating calcium sensor proteins, such as CaM and CDPKs. The CaM-binding transcription factors CBP60g and CBP60a are positive and negative regulators of *ICS1* transcription, respectively. A homolog of CBP60a/g, SARD1, is not a CaM-binding protein but functions redundantly with CBP60g for *ICS1* transcription. WRKY28, whose DNA-binding activity is

regulated by the CDPKs CPK5 and CPK11, also contributes to *ICS1* expression. *ICS1* mediates SA production in chloroplasts, by conversion of chorismate into the SA-precursor isochorismate. SA may be transported through the MATE-transporter EDS5 into the cytosol. The EDS1/PAD4 complex contributes to the positive feedback loop of SA accumulation. Repression of *EDS1* transcription by the Ca<sup>2+</sup>/CaM-binding transcription factor CAMTA3 represents a fine-tuning mechanism for SA accumulation.

that a CAMTA3/SR1-interacting protein links CAMTA3/SR1 to ubiquitin-mediated protein degradation thereby enhancing *EDS1* expression and immunity against *P. syringae* (Zhang et al., 2014).

In summary, these results clearly indicate the importance of Ca<sup>2+</sup> signaling in regulation of SA accumulation during immunity through transcriptional regulation of genes involved in SA biosynthesis and maintenance. However, how plants spatiotemporally

coordinate positive and negative regulators of SA biosynthesis and accumulation remains to be investigated.

### SA PERCEPTION


Identification of SA receptor(s) has been one of the major research interests for the last two decades. Considering its diverse functions in environmental stress response, plants may have multiple SA receptors. Indeed, biochemical approaches identified a num-



ber of SA-interacting proteins, and activities of these proteins were affected by SA-binding (Chen and Klessig, 1991; Chen et al., 1993; Durner and Klessig, 1995; Du and Klessig, 1997; Slaymaker et al., 2002; Kumar and Klessig, 2003; Forouhar et al., 2005; Park et al., 2009; Tripathi et al., 2010; Tian et al., 2012; Moreau et al., 2013). However, these SA-binding proteins do not fully explain SA response including SA-mediated transcriptional reprogramming. Recently, the three NPR (NON-EXPRESSOR of PATHOGENESIS-RELATED GENES) family members, NPR1, NPR3, and NPR4, were identified as *bona fide* SA receptors in *Arabidopsis* (Fu et al., 2012; Wu et al., 2012). In this section, we discuss how these NPR proteins function as SA receptors.

NPR1 is a master regulator of SA-mediated transcriptional reprogramming and immunity, functioning as a transcriptional coactivator (Pajerowska-Mukhtar et al., 2013). NPR1 comprises a BTB/POZ (broad-complex, tramtrack, and bric-à-brac/poxvirus and zinc-finger) domain, an ankyrin repeat domain, and a nuclear localization sequence. Mutations in *NPR1* lead to an almost

complete loss of SA-mediated transcriptional reprogramming and great susceptibility to (hemi)-biotrophic pathogens (Shah et al., 1997; Volko et al., 1998; Dong, 2004). Therefore, it was not surprising but sensational that Wu et al. (2012) found NPR1 to be a *bona fide* SA receptor (**Figure 2A**). Using an equilibrium method, they showed that *Arabidopsis* NPR1 directly binds SA ( $K_d = 140$  nM), but not inactive structural analogs, through Cys<sup>521/529</sup> via the transition metal copper. Consistently, Cys<sup>521/529</sup> were previously identified as key amino acid residues for *Arabidopsis* NPR1 function (Rochon et al., 2006). Biochemical approaches indicate that SA-binding triggers a conformational change in NPR1. Further protein deletion analyses suggest that the C-terminal transactivation domain of NPR1 is intramolecularly inhibited by the N-terminal BTB/POZ domain and that SA-binding releases the transactivation domain from BTB/POZ suppression. Thus, the study established a model with NPR1 as an SA receptor that also functions as a master signal transducer of SA signaling. However, Cys<sup>521/529</sup> are not conserved among



plant species, raising an issue of the evolutionary significance of the SA perception mechanism via NPR1. In addition, another study showed that NPR1 does not bind SA in a conventional non-equilibrium  $^3\text{H}$ -SA binding assay (Yan and Dong, 2014). Instead, Fu et al. (2012) identified two homologs of NPR1, NPR3 and NPR4, as SA receptors (**Figure 2B**; Fu et al., 2012). NPR1 is subject to degradation via the 26S proteasome pathway in the absence of SA (Spoel et al., 2009). Once SA increases upon pathogen infection, NPR1 is stabilized. However, full induction of SA-responsive genes also requires NPR1 turnover. Thus, regulation of NPR1 protein level is critical for SA response. Fu et al. (2012) found that NPR3 and NPR4 interact with NPR1 and are required for NPR1 degradation (Fu et al., 2012). NPR4 has a high SA affinity ( $K_d = 46$  nM) whereas NPR3 shows a low affinity ( $K_d = 981$  nM), suggesting differential regulations of NPR1 by NPR3 and NPR4. Interestingly, SA disrupts NPR1–NPR4 interaction, but facilitates NPR1–NPR3 interaction. These observations support a model in which NPR3 and NPR4 create an NPR1 protein concentration gradient in order to regulate NPR1–mediated transcription: in the absence of SA, NPR4-mediated NPR1 degradation prevents NPR1 accumulation whereas high SA levels also prevent NPR1 accumulation due to NPR3. Thus, NPR1-mediated signaling is active only at intermediate SA levels. This model is consistent with the observation that NPR1 protein highly accumulates at sites surrounding the infection site in a leaf. These regions are supposed to contain intermediate SA levels, while the infection site may have too high SA levels. Although this model is attractive, further validation is required.


Collectively, two alternative but not exclusive SA perception mechanisms in plant cells were identified, but further research is still required to address fundamental questions. For example, the subcellular location(s) of SA perception have not been addressed yet. The nuclear NPR1 pool is necessary for SA-mediated transcription (Mou et al., 2003). Consistently, NPR3 and NPR4 are nuclear proteins, and therefore SA is likely perceived by them in the nucleus to regulate nuclear NPR1 amount. On the other hand, the cytosolic NPR1 pool may regulate cross-talk between SA- and JA-mediated transcriptional reprogramming (Spoel et al., 2003), suggesting that SA is also perceived in the cytosol. Since SA perception by nuclear NPR3 and NPR4 does not explain this observation, cytosolic NPR1 activity may be regulated by the direct SA binding.

### SA-MEDIATED TRANSCRIPTIONAL REPROGRAMMING

NPR1 controls expression of more than 95% of the responsive genes to the SA-analog benzothiadiazole (BTH; Wang et al., 2006). Functional regulation of NPR1 is not only mediated by the direct SA binding, but also by SA-triggered redox changes (Mou et al., 2003). In the absence of SA, NPR1 is present as an oligomer formed through intermolecular disulfide bonds. SA triggers changes in the cellular redox potential, thereby reducing cysteine residues in NPR1 through the thioredoxins TRXh3 and TRXh5, resulting in monomerization of NPR1 (Tada et al., 2008). Mutations in the cysteine residues (Cys<sup>82</sup> or Cys<sup>216</sup>) lead to constitutive monomerization and nuclear accumulation of NPR1, resulting in activation of *PR1* expression (Mou et al., 2003). Nuclear accumulation of NPR1 triggered by SA can be explained

by stabilization of nuclear NPR1 or translocation of the NPR1 monomer from the cytosol to the nucleus. Thus, SA-triggered NPR1 monomerization and nuclear accumulation are important steps for NPR1-mediated transcription. However, forced nuclear localization of NPR1 is not sufficient for transcriptional reprogramming, as the presence of SA is additionally required for full *PR1* induction (Kinkema et al., 2000; Spoel et al., 2003). This can be explained by the observation that SA-binding triggers the NPR1 conformational change thereby allowing NPR1 to regulate gene expression (Wu et al., 2012). Additional regulation of NPR1 involves phosphorylation (Spoel et al., 2009). SA triggers phosphorylation of NPR1 at the N-terminus (Ser<sup>11/15</sup>) in the nucleus via yet-determined kinase(s). NPR1 phosphorylation contributes to its recruitment to a ubiquitin ligase, resulting in proteasome-mediated NPR1 degradation. This degradation is required for the proper transcriptional control by NPR1, perhaps by allowing fresh NPR1 to reinitiate the next cycle of transcription.

NPR1 regulates transcription of SA-responsive genes through interactions with specific transcription factors (**Figure 3**). Identified major transcription factors belong to a subclass of the basic leucine zipper transcription factor family, TGA (Gatz, 2013). The *Arabidopsis* genome encodes 10 TGA transcription factors, which are structurally divided into five subgroups and all bind the consensus DNA sequence TGACG. Yeast-two-hybrid analyses with NPR1 and TGA transcription factors show interaction specificity for clade II TGAs (TGA2/TGA5/TGA6) and TGA3 (clade III; Zhou et al., 2000; Hepworth et al., 2005). Genetic analysis reveal that TGA2, TGA5, and TGA6 repress *PR1* transcription in the absence of SA, but on the other hand are required for



**FIGURE 3 | SA-mediated transcriptional regulation of *PR1* through NPR1.** In the absence of SA, repression of *PR1* expression can be achieved by repressor complexes (blue), such as the TGA2–NPR1–NIMIN1 complex through the co-repressor TOPLESS or the CBNAC–SNI1 complex. SA triggers a conformational change of NPR1 and dissociation of NIMIN1, resulting in forming activator complexes (pink) including TGA transcription factors and SSN2. The DNA repair proteins BRCA2 and RAD51 are also involved in SA-mediated transcription.

*PR1* induction in the presence of SA (Zhang et al., 2003). In the absence of SA, TGA2 binds to the *PR1* promoter thereby repressing its transcription (Rochon et al., 2006; Boyle et al., 2009). An NPR1-interacting protein, NIMIN1 (NPR1/NIM1-INTERACTING PROTEIN 1), can form a ternary complex with TGA2 through NPR1 at least in yeast (Weigel et al., 2005). Transcriptional repression by TGA2 may be achieved through NIMIN1 interacting with a transcriptional co-repressor, TOPLESS (Braun et al., 2011). Conceivably, SA allows NPR1 to form a different complex with TGA2 and other TGA factors, such as TGA3 thereby activating *PR1* transcription (Johnson et al., 2003). The NIMIN1–NPR1–TGA2 complex is dissociated in the presence of SA in yeast (Hermann et al., 2013). Thus, NIMIN1 dissociation from the NPR1–TGA transcriptional complex by SA may contribute to activation of the NPR1–TGA transcriptional complex. This transcriptional activation may be relayed through specific mediator subunits, such as the Mediator subunit MED15, since *med15* mutants are insensitive to SA (Canet et al., 2012).

A suppressor screen of *npr1* identified SNI1 (SUPPRESSOR OF *NPR1* INDUCIBLE 1) as another repressor of SA-responsive genes (e.g., *PR1*) in unstressed conditions (Li et al., 1999; Mosher et al., 2006). SNI1-mediated transcriptional repression may be achieved through the CaM-binding NAC (NAM, ATAF1,2, CUC2) transcription repressor CBNAC, since SNI1 directly interacts with CBNAC and enhances CBNAC-binding activity to the *PR1* promoter (Kim et al., 2012). Upon SA treatment, SNI1 is dissociated from the *PR1* promoter and replaced by the DNA repair protein SSN2 (Song et al., 2011). Although SSN2 contains a DNA-binding domain, its binding to the *PR1* promoter requires NPR1 and the transcription factor TGA7. These results suggest that SA triggers NPR1 activation through nuclear accumulation and conformational change, resulting in the formation of a TGA7–NPR1–SSN2 complex that activates *PR1* transcription. Additional DNA repair proteins, such as BRCA2A (BREAST CANCER 2A) and RAD51D, are also functionally associated with SA-mediated transcription (Durrant et al., 2007; Wang et al., 2010; Song et al., 2011). Interestingly, SA and *Pseudomonas* infection cause DNA damage, such as DNA double strand breaks, suggesting that DNA damage response is an intrinsic component of SA-mediated transcription during plant immunity (Yan et al., 2013; Song and Bent, 2014).

Besides functional regulation of transcription factors by NPR1 through complex formation, NPR1 also controls expression of transcription factors, such as WRKY transcription factors, which are required for SA-mediated transcriptional reprogramming (Wang et al., 2006; Pajerowska-Mukhtar et al., 2012). The *Arabidopsis* genome encodes 74 WRKY factors which bind the specific DNA sequence (C/TTGACT/C), termed the W-box (Rushton et al., 2010). WRKY factors form a complex interconnected regulatory network, containing recurring regulatory patterns, such as both positive and negative feedback and feedforward loops. This WRKY network ensures rapid and efficient signal amplification and allows tight control to limit the plant immune response. Furthermore, the presence of multiple W-boxes in the *NPR1* promoter suggests regulation of *NPR1* expression by WRKY factors, which is indeed supported by NPR1 promoter analysis (Yu et al., 2001). Thus, WRKY transcriptional regulatory networks

downstream of NPR1 amplify and fine-tune SA-mediated transcriptional reprogramming.

## COMPENSATION OF SA SIGNALING

The importance of SA signaling during immunity is reflected by the fact that pathogen effectors target it for virulence, either by preventing SA accumulation (Djamei et al., 2011; Rabe et al., 2013; Liu et al., 2014) or by dampen SA signaling and transcriptional regulation, using the antagonistic interaction between SA and JA signaling (Uppalapati et al., 2007; Caillaud et al., 2013; Jiang et al., 2013; Gimenez-Ibanez et al., 2014). It is reasonable to assume that plants have evolved compensatory mechanism(s) to circumvent weakened SA signaling upon effector attack, thereby ensuring robust immune response (Tsuda and Katagiri, 2010). For example, although it is believed that SA and JA signaling antagonize each other, a recent study suggests the compensation of SA accumulation by JA (Kim et al., 2014). The MAMP *flg22* induces SA accumulation in an *ICS1*-dependent manner (Tsuda et al., 2008b). Additionally, a component of the SA amplification loop, PAD4, is required for full induction of SA (Zhou et al., 1998; Tsuda et al., 2008b). In agreement with the antagonistic relationship between JA and SA, single mutation in the JA biosynthesis gene *DDE2* leads to higher SA accumulation upon *flg22* treatment. However, combined mutation in *DDE2* and *PAD4* diminishes SA accumulation comparable to that in *sid2*, suggesting that although JA suppresses SA accumulation through PAD4, it also supports SA accumulation once *PAD4* is compromised. Thus, JA signaling represents a compensation mechanism for SA accumulation during PTI.

In addition to JA, MAPK signaling also compensates SA signaling to secure transcriptional regulation of SA-responsive genes in ETI (Tsuda et al., 2013). Activation of the *Arabidopsis* MAPKs MPK3 and MPK6 is transient during PTI, but sustained during RPS2- and RPM1-mediated ETI (Tsuda et al., 2013), or upon *B. cinerea* infection (Han et al., 2010). While transient activation of MPK3 and MPK6 is not sufficient to overcome SA-dependency of a subset of SA-responsive genes such as *PR1*, prolonged activation of MPK3 and MPK6 facilitates their transcriptional regulation independent of SA. Furthermore, this compensation mechanism does not require NPR1 since *NPR1* mutation does not affect *PR1* induction mediated by prolonged MAPK activation. It can be assumed that prolonged MAPK activation bypasses the requirement of NPR1 to regulate transcription factor(s) involved in SA response. Although transcription factors shared by SA and the MAPK cascade are not known, large-scale protein target identifications of MPK3 and MPK6 would help to identify candidates (Popescu et al., 2009; Hoehenwarter et al., 2013). Among them, TGA transcription factors are reasonable candidates (Wang and Fobert, 2013). However, how this quantitative MAPK activation leads to qualitatively different transcriptional outputs still remains to be determined. One possible answer lays in temporal regulation of transcription factor(s). Hereby, the MAPKs first activate expression of transcription factor(s), and later on phosphorylate the accumulated transcription factor(s), representing a feedforward loop for activation of the transcription factor(s). In this case, only prolonged MAPK activation ensures activation of the transcription factor(s). Indeed, the MAPKs regulate expression of



a diverse transcription factor set (Mao et al., 2011; Li et al., 2012; Meng et al., 2013; Tsuda et al., 2013; Frei dit Frey et al., 2014), but whether the MAPKs also phosphorylate them is a future issue.

## CONCLUSIONS AND PERSPECTIVES

Over the past decade a number of researches have shed light into our understanding of SA-mediated signaling, through the discoveries of calcium signaling as the major switch for SA biosynthesis, NPR family members as SA receptors, and the mechanism for NPR1-mediated transcriptional reprogramming. However, many questions are still unanswered, starting with identification of plant *IPL* gene(s) to further validate the IC pathway as the major route for SA biosynthesis in plants. The controversy for SA perception should also be solved in the future. In addition, information for temporal and spatial dynamics of SA biosynthesis and SA-mediated transcriptional reprogramming is missing. For this, systems approaches using time-series genomics data sets and tissue-specific analysis will help our conception (Mine et al., 2014). Most studies are based on experiments using the model plant *Arabidopsis*. Analysis of different plant species is necessary to understand evolutionary conservation and diversification of SA signal transduction. Finally, identification of the molecular components in MAPK-mediated SA/NPR1-independent gene regulation of SA-responsive genes in ETI will shed light on the molecular mechanism of SA compensation.

## ACKNOWLEDGMENTS

We thank Akira Mine, Shajahan Anver, and Sajjad Khani for critical reading of the manuscript. This work was supported by the Max Planck Society (Carolin Seyfferth, Kenichi Tsuda) and a grant from the German Research Council SFB670 (Kenichi Tsuda).

## REFERENCES

- Boller, T., and Felix, G. (2009). A renaissance of elicitors: perception of microbe-associated molecular patterns and danger signals by pattern-recognition receptors. *Annu. Rev. Plant Biol.* 60, 379–406. doi: 10.1146/annurev.arplant.57.032905.105346
- Bonardi, V., and Dangl, J. L. (2012). How complex are intracellular immune receptor signaling complexes? *Front. Plant Sci.* 3:237. doi: 10.3389/fpls.2012.00237
- Boudsocq, M., and Sheen, J. (2013). CDPKs in immune and stress signaling. *Trends Plant Sci.* 18, 30–40. doi: 10.1016/j.tplants.2012.08.008
- Boyle, P., Le Su, E., Rochon, A., Shearer, H. L., Murmu, J., Chu, J. Y., et al. (2009). The BTB/POZ domain of the *Arabidopsis* disease resistance protein NPR1 interacts with the repression domain of TGA2 to negate its function. *Plant Cell* 21, 3700–3713. doi: 10.1105/tpc.109.069971
- Braun, P., Carvunis, A. R., Charlotteaux, B., Dreze, M., Ecker, J. R., Hill, D. E., et al. (2011). Evidence for network evolution in an *Arabidopsis* interactome map. *Science* 333, 601–607. doi: 10.1126/science.1203877
- Caillaud, M. C., Asai, S., Rallapalli, G., Piquerez, S., Fabro, G., and Jones, J. D. (2013). A downy mildew effector attenuates salicylic acid-triggered immunity in *Arabidopsis* by interacting with the host mediator complex. *PLoS Biol.* 11:e1001732. doi: 10.1371/journal.pbio.1001732
- Canet, J. V., Dobon, A., and Tornero, P. (2012). Non-recognition-of-BTH4, an *Arabidopsis* mediator subunit homolog, is necessary for development and response to salicylic acid. *Plant Cell* 24, 4220–4235. doi: 10.1105/tpc.112.103028
- Catinot, J., Buchala, A., Abou-Mansour, E., and Metraux, J. P. (2008). Salicylic acid production in response to biotic and abiotic stress depends on isochlorismate in *Nicotiana benthamiana*. *FEBS Lett.* 582, 473–478. doi: 10.1016/j.febslet.2007.12.039
- Chandran, D., Rickert, J., Huang, Y., Steinwand, M. A., Marr, S. K., and Wildermuth, M. C. (2014). Atypical E2F transcriptional repressor DEL1 acts at the intersection of plant growth and immunity by controlling the hormone salicylic acid. *Cell Host Microbe* 15, 506–513. doi: 10.1016/j.chom.2014.03.007
- Chen, H., Xue, L., Chintamanani, S., Germain, H., Lin, H., Cui, H., et al. (2009). ETHYLENE INSENSITIVE3 and ETHYLENE INSENSITIVE3-LIKE1 repress SALICYLIC ACID INDUCTION DEFICIENT2 expression to negatively regulate plant innate immunity in *Arabidopsis*. *Plant Cell* 21, 2527–2540. doi: 10.1105/tpc.108.065193
- Chen, Z., and Klessig, D. F. (1991). Identification of a soluble salicylic acid-binding protein that may function in signal transduction in the plant disease-resistance response. *Proc. Natl. Acad. Sci. U.S.A.* 88, 8179–8183. doi: 10.1073/pnas.88.18.8179
- Chen, Z., Silva, H., and Klessig, D. F. (1993). Active oxygen species in the induction of plant systemic acquired resistance by salicylic acid. *Science* 262, 1883–1886. doi: 10.1126/science.8266079
- Cochrane, F. C., Davin, L. B., and Lewis, N. G. (2004). The *Arabidopsis* phenylalanine ammonia lyase gene family: kinetic characterization of the four PAL isoforms. *Phytochemistry* 65, 1557–1564. doi: 10.1016/j.phytochem.2004.05.006
- Dempsey, D. A., Vlot, A. C., Wildermuth, M. C., and Klessig, D. F. (2011). Salicylic Acid biosynthesis and metabolism. *Arabidopsis Book* 9, e0156. doi: 10.1199/tab.0156
- Derksen, H., Rampitsch, C., and Daayf, F. (2013). Signaling cross-talk in plant disease resistance. *Plant Sci.* 207, 79–87. doi: 10.1016/j.plantsci.2013.03.004
- Djamei, A., Schipper, K., Rabe, F., Ghosh, A., Vincon, V., Kahnt, J., et al. (2011). Metabolic priming by a secreted fungal effector. *Nature* 478, 395–398. doi: 10.1038/nature10454
- Dodd, A. N., Kudla, J., and Sanders, D. (2010). The language of calcium signaling. *Annu. Rev. Plant Biol.* 61, 593–620. doi: 10.1146/annurev-arplant-070109-104628
- Dong, X. (2004). NPR1, all things considered. *Curr. Opin. Plant Biol.* 7, 547–552. doi: 10.1016/j.pbi.2004.07.005
- Du, H., and Klessig, D. F. (1997). Identification of a soluble, high-affinity salicylic acid-binding protein in tobacco. *Plant Physiol.* 113, 1319–1327.
- Du, L., Ali, G. S., Simons, K. A., Hou, J., Yang, T., Reddy, A. S., et al. (2009). Ca<sup>2+</sup>/calmodulin regulates salicylic-acid-mediated plant immunity. *Nature* 457, 1154–1158. doi: 10.1038/nature07612
- Durner, J., and Klessig, D. F. (1995). Inhibition of ascorbate peroxidase by salicylic acid and 2,6-dichloroisonicotinic acid, two inducers of plant defense responses. *Proc. Natl. Acad. Sci. U.S.A.* 92, 11312–11316. doi: 10.1073/pnas.92.24.11312
- Durrant, W. E., Wang, S., and Dong, X. (2007). *Arabidopsis* SN11 and RAD51D regulate both gene transcription and DNA recombination during the defense response. *Proc. Natl. Acad. Sci. U.S.A.* 104, 4223–4227. doi: 10.1073/pnas.0609357104
- Eitas, T. K., and Dangl, J. L. (2010). NB-LRR proteins: pairs, pieces, perception, partners, and pathways. *Curr. Opin. Plant Biol.* 13, 472–477. doi: 10.1016/j.pbi.2010.04.007
- Feys, B. J., Moisan, L. J., Newman, M. A., and Parker, J. E. (2001). Direct interaction between the *Arabidopsis* disease resistance signaling proteins, EDS1 and PAD4. *EMBO J.* 20, 5400–5411. doi: 10.1093/emboj/20.19.5400
- Forouhar, F., Yang, Y., Kumar, D., Chen, Y., Fridman, E., Park, S. W., et al. (2005). Structural and biochemical studies identify tobacco SAMP2 as a methyl salicylate esterase and implicate it in plant innate immunity. *Proc. Natl. Acad. Sci. U.S.A.* 102, 1773–1778. doi: 10.1073/pnas.0409227102
- Frei dit Frey, N., Garcia, A. V., Bigeard, J., Zaag, R., Bueso, E., Garmier, M., et al. (2014). Functional analysis of *Arabidopsis* immune-related MAPKs uncovers a role for MPK3 as negative regulator of inducible defences. *Genome Biol.* 15, R87. doi: 10.1186/gb-2014-15-6-r87
- Fu, Z. Q., and Dong, X. (2013). Systemic acquired resistance: turning local infection into global defense. *Annu. Rev. Plant Biol.* 64, 839–863. doi: 10.1146/annurev-arplant-042811-105606
- Fu, Z. Q., Yan, S., Saleh, A., Wang, W., Ruble, J., Oka, N., et al. (2012). NPR3 and NPR4 are receptors for the immune signal salicylic acid in plants. *Nature* 486, 228–232. doi: 10.1038/nature11162
- Gao, X., Chen, X., Lin, W., Chen, S., Lu, D., Niu, Y., et al. (2013). Bifurcation of *Arabidopsis* NLR immune signaling via Ca<sup>2+</sup>-dependent protein kinases. *PLoS Pathog.* 9:e1003127. doi: 10.1371/journal.ppat.1003127
- Garcion, C., Lohmann, A., Lamodièrre, E., Catinot, J., Buchala, A., Doermann, P., et al. (2008). Characterization and biological function of the ISOCHORISMATE SYNTHASE2 gene of *Arabidopsis*. *Plant Physiol.* 147, 1279–1287. doi: 10.1104/pp.108.119420

- Gatz, C. (2013). From pioneers to team players: TGA transcription factors provide a molecular link between different stress pathways. *Mol. Plant Microbe Interact.* 26, 151–159. doi: 10.1094/MPMI-04-12-0078-IA
- Gimenez-Ibanez, S., Boter, M., Fernandez-Barbero, G., Chini, A., Rathjen, J. P., and Solano, R. (2014). The bacterial effector HopX1 targets JAZ transcriptional repressors to activate jasmonate signaling and promote infection in *Arabidopsis*. *PLoS Biol.* 12:e1001792. doi: 10.1371/journal.pbio.1001792
- Han, L., Li, G. J., Yang, K. Y., Mao, G., Wang, R., Liu, Y., et al. (2010). Mitogen-activated protein kinase 3 and 6 regulate *Botrytis cinerea*-induced ethylene production in *Arabidopsis*. *Plant J.* 64, 114–127. doi: 10.1111/j.1365-3113.2010.04318.x
- Hepworth, S. R., Zhang, Y., Mckim, S., Li, X., and Haughn, G. W. (2005). BLADE-ON-PETIOLE-dependent signaling controls leaf and floral patterning in *Arabidopsis*. *Plant Cell* 17, 1434–1448. doi: 10.1105/tpc.104.030536
- Hermann, M., Maier, F., Masroor, A., Hirth, S., Pfitzner, A. J., and Pfitzner, U. M. (2013). The *Arabidopsis* NIMIN proteins affect NPR1 differentially. *Front. Plant Sci.* 4:88. doi: 10.3389/fpls.2013.00088
- Hoehenwarter, W., Thomas, M., Nukarinen, E., Egelhofer, V., Rohrig, H., Weckwerth, W., et al. (2013). Identification of novel in vivo MAP kinase substrates in *Arabidopsis thaliana* through use of tandem metal oxide affinity chromatography. *Mol. Cell. Proteomics* 12, 369–380. doi: 10.1074/mcp.M112.020560
- Huang, J., Gu, M., Lai, Z., Fan, B., Shi, K., Zhou, Y. H., et al. (2010). Functional analysis of the *Arabidopsis* PAL gene family in plant growth, development, and response to environmental stress. *Plant Physiol.* 153, 1526–1538. doi: 10.1104/pp.110.157370
- Ishihara, T., Sekine, K. T., Hase, S., Kanayama, Y., Seo, S., Ohashi, Y., et al. (2008). Overexpression of the *Arabidopsis thaliana* EDS5 gene enhances resistance to viruses. *Plant Biol.* 10, 451–461. doi: 10.1111/j.1438-8677.2008.00050.x
- Jacob, F., Vernaldi, S., and Maekawa, T. (2013). Evolution and conservation of plant NLR functions. *Front. Immunol.* 4:297. doi: 10.3389/fimmu.2013.00297
- Jiang, S., Yao, J., Ma, K. W., Zhou, H., Song, J., He, S. Y., et al. (2013). Bacterial effector activates jasmonate signaling by directly targeting JAZ transcriptional repressors. *PLoS Pathog.* 9:e1003715. doi: 10.1371/journal.ppat.1003715
- Johnson, C., Boden, E., and Arias, J. (2003). Salicylic acid and NPR1 induce the recruitment of trans-activating TGA factors to a defense gene promoter in *Arabidopsis*. *Plant Cell* 15, 1846–1858. doi: 10.1105/tpc.012211
- Jones, J. D., and Dangl, J. L. (2006). The plant immune system. *Nature* 444, 323–329. doi: 10.1038/nature05286
- Kim, D. S., and Hwang, B. K. (2014). An important role of the pepper phenylalanine ammonia-lyase gene (PAL1) in salicylic acid-dependent signalling of the defence response to microbial pathogens. *J. Exp. Bot.* 65, 2295–2306. doi: 10.1093/jxb/eru109
- Kim, H. S., Park, H. C., Kim, K. E., Jung, M. S., Han, H. J., Kim, S. H., et al. (2012). A NAC transcription factor and SN1 cooperatively suppress basal pathogen resistance in *Arabidopsis thaliana*. *Nucleic Acids Res.* 40, 9182–9192. doi: 10.1093/nar/gks683
- Kim, Y., Park, S., Gilmour, S. J., and Thomashow, M. F. (2013). Roles of CAMTA transcription factors and salicylic acid in configuring the low-temperature transcriptome and freezing tolerance of *Arabidopsis*. *Plant J.* 75, 364–376. doi: 10.1111/tpj.12205
- Kim, Y., Tsuda, K., Igarashi, D., Hillmer, R. A., Sakakibara, H., Myers, C. L., et al. (2014). Mechanisms underlying robustness and tunability in a plant immune signaling network. *Cell Host Microbe* 15, 84–94. doi: 10.1016/j.chom.2013.12.002
- Kinkema, M., Fan, W., and Dong, X. (2000). Nuclear localization of NPR1 is required for activation of PR gene expression. *Plant Cell* 12, 2339–2350. doi: 10.1105/tpc.12.12.2339
- Kumar, D., and Klessig, D. F. (2003). High-affinity salicylic acid-binding protein 2 is required for plant innate immunity and has salicylic acid-stimulated lipase activity. *Proc. Natl. Acad. Sci. U.S.A.* 100, 16101–16106. doi: 10.1073/pnas.0307162100
- Li, G., Meng, X., Wang, R., Mao, G., Han, L., Liu, Y., et al. (2012). Dual-level regulation of ACC synthase activity by MPK3/MPK6 cascade and its downstream WRKY transcription factor during ethylene induction in *Arabidopsis*. *PLoS Genet.* 8:e1002767. doi: 10.1371/journal.pgen.1002767
- Li, X., Zhang, Y., Clarke, J. D., Li, Y., and Dong, X. (1999). Identification and cloning of a negative regulator of systemic acquired resistance, SN11, through a screen for suppressors of npr1-1. *Cell* 98, 329–339. doi: 10.1016/S0092-8674(00)81962-5
- Liu, T., Song, T., Zhang, X., Yuan, H., Su, L., Li, W., et al. (2014). Unconventionally secreted effectors of two filamentous pathogens target plant salicylate biosynthesis. *Nat. Commun.* 5, 4686. doi: 10.1038/ncomms5686
- Lohou, D., Lonjon, F., Genin, S., and Vaillau, F. (2013). Type III chaperones & Co in bacterial plant pathogens: a set of specialized bodyguards mediating effector delivery. *Front. Plant Sci.* 4:435. doi: 10.3389/fpls.2013.00435
- Macho, A. P., and Zipfel, C. (2014). Plant PRRs and the activation of innate immune signaling. *Mol. Cell* 54, 263–272. doi: 10.1016/j.molcel.2014.03.028
- Mao, G., Meng, X., Liu, Y., Zheng, Z., Chen, Z., and Zhang, S. (2011). Phosphorylation of a WRKY transcription factor by two pathogen-responsive MAPKs drives phytoalexin biosynthesis in *Arabidopsis*. *Plant Cell* 23, 1639–1653. doi: 10.1105/tpc.111.084996
- Mauch, F., Mauch-Mani, B., Gaille, C., Kull, B., Haas, D., and Reimann, C. (2001). Manipulation of salicylate content in *Arabidopsis thaliana* by the expression of an engineered bacterial salicylate synthase. *Plant J.* 25, 67–77. doi: 10.1046/j.1365-3113x.2001.00940.x
- Meng, X., Xu, J., He, Y., Yang, K. Y., Mordorski, B., Liu, Y., et al. (2013). Phosphorylation of an ERF transcription factor by *Arabidopsis* MPK3/MPK6 regulates plant defense gene induction and fungal resistance. *Plant Cell* 25, 1126–1142. doi: 10.1105/tpc.112.109074
- Mine, A., Sato, M., and Tsuda, K. (2014). Toward a systems understanding of plant-microbe interactions. *Front. Plant Sci.* 5:423. doi: 10.3389/fpls.2014.00423
- Moreau, M., Westlake, T., Zampogna, G., Popescu, G., Tian, M., Noutsos, C., et al. (2013). The *Arabidopsis* oligopeptidases TOP1 and TOP2 are salicylic acid targets that modulate SA-mediated signaling and the immune response. *Plant J.* 76, 603–614. doi: 10.1111/tpj.12320
- Mosher, R. A., Durrant, W. E., Wang, D., Song, J., and Dong, X. (2006). A comprehensive structure-function analysis of *Arabidopsis* SN11 defines essential regions and transcriptional repressor activity. *Plant Cell* 18, 1750–1765. doi: 10.1105/tpc.105.039677
- Mou, Z., Fan, W., and Dong, X. (2003). Inducers of plant systemic acquired resistance regulate NPR1 function through redox changes. *Cell* 113, 935–944. doi: 10.1016/S0092-8674(03)00429-X
- Nawrath, C., Heck, S., Parinithawong, N., and Mettraux, J. P. (2002). EDS5, an essential component of salicylic acid-dependent signaling for disease resistance in *Arabidopsis*, is a member of the MATE transporter family. *Plant Cell* 14, 275–286.
- Nobuta, K., Okrent, R. A., Stoutemyer, M., Rodibaugh, N., Kempema, L., Wildermuth, M. C., et al. (2007). The GH3 acyl adenylase family member PBS3 regulates salicylic acid-dependent defense responses in *Arabidopsis*. *Plant Physiol.* 144, 1144–1156. doi: 10.1104/pp.107.097691
- Okrent, R. A., Brooks, M. D., and Wildermuth, M. C. (2009). *Arabidopsis* GH3.12 (PBS3) conjugates amino acids to 4-substituted benzoates and is inhibited by salicylate. *J. Biol. Chem.* 284, 9742–9754. doi: 10.1074/jbc.M806662200
- Pajerowska-Mukhtar, K. M., Emerine, D. K., and Mukhtar, M. S. (2013). Tell me more: roles of NPRs in plant immunity. *Trends Plant Sci.* 18, 402–411. doi: 10.1016/j.tplants.2013.04.004
- Pajerowska-Mukhtar, K. M., Wang, W., Tada, Y., Oka, N., Tucker, C. L., Fonseca, J. P., et al. (2012). The HSF-like transcription factor TBF1 is a major molecular switch for plant growth-to-defense transition. *Curr. Biol.* 22, 103–112. doi: 10.1016/j.cub.2011.12.015
- Park, S. W., Liu, P. P., Forouhar, F., Vlot, A. C., Tong, L., Tietjen, K., et al. (2009). Use of a synthetic salicylic acid analog to investigate the roles of methyl salicylate and its esterases in plant disease resistance. *J. Biol. Chem.* 284, 7307–7317. doi: 10.1074/jbc.M807968200
- Pieterse, C. M., Van Der Does, D., Zamioudis, C., Leon-Reyes, A., and Van Wees, S. C. (2012). Hormonal modulation of plant immunity. *Annu. Rev. Cell Dev. Biol.* 28, 489–521. doi: 10.1146/annurev-cellbio-092910-154055
- Poovaiah, B. W., Du, L., Wang, H., and Yang, T. (2013). Recent advances in calcium/calmodulin-mediated signaling with an emphasis on plant-microbe interactions. *Plant Physiol.* 163, 531–542. doi: 10.1104/pp.113.220780
- Popescu, S. C., Popescu, G. V., Bachan, S., Zhang, Z., Gerstein, M., Snyder, M., et al. (2009). MAPK target networks in *Arabidopsis thaliana* revealed using functional protein microarrays. *Genes Dev.* 23, 80–92. doi: 10.1101/gad.1740009
- Rabe, F., Ajami-Rashidi, Z., Doehlemann, G., Kahmann, R., and Djamei, A. (2013). Degradation of the plant defence hormone salicylic acid by the biotrophic



- fungus *Ustilago maydis*. *Mol. Microbiol.* 89, 179–188. doi: 10.1111/mmi.12269
- Robert-Seilaniantz, A., Grant, M., and Jones, J. D. (2011). Hormone crosstalk in plant disease and defense: more than just jasmonate-salicylate antagonism. *Annu. Rev. Phytopathol.* 49, 317–343. doi: 10.1146/annurev-phyto-073009-114447
- Rochon, A., Boyle, P., Wignes, T., Fobert, P. R., and Despres, C. (2006). The coactivator function of *Arabidopsis* NPR1 requires the core of its BTB/POZ domain and the oxidation of C-terminal cysteines. *Plant Cell* 18, 3670–3685. doi: 10.1105/tpc.106.046953
- Rushton, P. J., Somssich, I. E., Ringler, P., and Shen, Q. J. (2010). WRKY transcription factors. *Trends Plant Sci.* 15, 247–258. doi: 10.1016/j.tplants.2010.02.006
- Sadeghi, M., Dehghan, S., Fischer, R., Wenzel, U., Vilcinskas, A., Kavousi, H. R., et al. (2013). Isolation and characterization of isochorismate synthase and cinnamate 4-hydroxylase during salinity stress, wounding, and salicylic acid treatment in *Carthamus tinctorius*. *Plant Signal. Behav.* 8:e27335. doi: 10.4161/psb.27335
- Schulz, P., Herde, M., and Romeis, T. (2013). Calcium-dependent protein kinases: hubs in plant stress signaling and development. *Plant Physiol.* 163, 523–530. doi: 10.1104/pp.113.222539
- Serrano, M., Wang, B., Aryal, B., Garcion, C., Abou-Mansour, E., Heck, S., et al. (2013). Export of salicylic acid from the chloroplast requires the multidrug and toxin extrusion-like transporter EDS5. *Plant Physiol.* 162, 1815–1821. doi: 10.1104/pp.113.218156
- Shah, J., Tsui, F., and Klessig, D. F. (1997). Characterization of a salicylic acid-insensitive mutant (sail) of *Arabidopsis thaliana*, identified in a selective screen utilizing the SA-inducible expression of the *tms2* gene. *Mol. Plant Microbe Interact.* 10, 69–78. doi: 10.1094/MPMI.1997.10.1.69
- Slaymaker, D. H., Navarre, D. A., Clark, D., del Pozo, O., Martin, G. B., and Klessig, D. F. (2002). The tobacco salicylic acid-binding protein 3 (SABP3) is the chloroplast carbonic anhydrase, which exhibits antioxidant activity and plays a role in the hypersensitive defense response. *Proc. Natl. Acad. Sci. U.S.A.* 99, 11640–11645. doi: 10.1073/pnas.182427699
- Song, J., and Bent, A. F. (2014). Microbial pathogens trigger host DNA double-strand breaks whose abundance is reduced by plant defense responses. *PLoS Pathog.* 10:e1004030. doi: 10.1371/journal.ppat.1004030
- Song, J., Durrant, W. E., Wang, S., Yan, S., Tan, E. H., and Dong, X. (2011). DNA repair proteins are directly involved in regulation of gene expression during plant immune response. *Cell Host Microbe* 9, 115–124. doi: 10.1016/j.chom.2011.01.011
- Spoel, S. H., Koornneef, A., Claessens, S. M., Korzelius, J. P., Van Pelt, J. A., Mueller, M. J., et al. (2003). NPR1 modulates cross-talk between salicylate- and jasmonate-dependent defense pathways through a novel function in the cytosol. *Plant Cell* 15, 760–770. doi: 10.1105/tpc.009159
- Spoel, S. H., Mou, Z., Tada, Y., Spivey, N. W., Genschik, P., and Dong, X. (2009). Proteasome-mediated turnover of the transcription coactivator NPR1 plays dual roles in regulating plant immunity. *Cell* 137, 860–872. doi: 10.1016/j.cell.2009.03.038
- Strawn, M. A., Marr, S. K., Inoue, K., Inada, N., Zubieta, C., and Wildermuth, M. C. (2007). *Arabidopsis* isochorismate synthase functional in pathogen-induced salicylate biosynthesis exhibits properties consistent with a role in diverse stress responses. *J. Biol. Chem.* 282, 5919–5933. doi: 10.1074/jbc.M605193200
- Tada, Y., Spoel, S. H., Pajerowska-Mukhtar, K., Mou, Z., Song, J., Wang, C., et al. (2008). Plant immunity requires conformational changes [corrected] of NPR1 via S-nitrosylation and thioredoxins. *Science* 321, 952–956. doi: 10.1126/science.1156970
- Tian, M., Von Dahl, C. C., Liu, P. P., Friso, G., Van Wijk, K. J., and Klessig, D. F. (2012). The combined use of photoaffinity labeling and surface plasmon resonance-based technology identifies multiple salicylic acid-binding proteins. *Plant J.* 72, 1027–1038. doi: 10.1111/tpj.12016
- Tripathi, D., Jiang, Y. L., and Kumar, D. (2010). SABP2, a methyl salicylate esterase is required for the systemic acquired resistance induced by acibenzolar-S-methyl in plants. *FEBS Lett.* 584, 3458–3463. doi: 10.1016/j.febslet.2010.06.046
- Truman, W., Sreekanta, S., Lu, Y., Bethke, G., Tsuda, K., Katagiri, F., et al. (2013). The CALMODULIN-BINDING PROTEIN60 family includes both negative and positive regulators of plant immunity. *Plant Physiol.* 163, 1741–1751. doi: 10.1104/pp.113.227108
- Tsuda, K., Glazebrook, J., and Katagiri, F. (2008a). The interplay between MAMP and SA signaling. *Plant Signal. Behav.* 3, 359–361. doi: 10.4161/psb.3.6.5702
- Tsuda, K., Sato, M., Glazebrook, J., Cohen, J. D., and Katagiri, F. (2008b). Interplay between MAMP-triggered and SA-mediated defense responses. *Plant J.* 53, 763–775. doi: 10.1111/j.1365-313X.2007.03369.x
- Tsuda, K., and Katagiri, F. (2010). Comparing signaling mechanisms engaged in pattern-triggered and effector-triggered immunity. *Curr. Opin. Plant Biol.* 13, 459–465. doi: 10.1016/j.pbi.2010.04.006
- Tsuda, K., Mine, A., Bethke, G., Igarashi, D., Botanga, C. J., Tsuda, Y., et al. (2013). Dual regulation of gene expression mediated by extended MAPK activation and salicylic acid contributes to robust innate immunity in *Arabidopsis thaliana*. *PLoS Genet.* 9:e1004015. doi: 10.1371/journal.pgen.1004015
- Uppalapati, S. R., Ishiga, Y., Wangdi, T., Kunkel, B. N., Anand, A., Mysore, K. S., et al. (2007). The phytotoxin coronatine contributes to pathogen fitness and is required for suppression of salicylic acid accumulation in tomato inoculated with *Pseudomonas syringae* pv. *tomato* DC3000. *Mol. Plant Microbe Interact.* 20, 955–965. doi: 10.1094/MPMI-20-8-0955
- van Verk, M. C., Bol, J. F., and Linthorst, H. J. (2011). WRKY transcription factors involved in activation of SA biosynthesis genes. *BMC Plant Biol.* 11:89. doi: 10.1186/1471-2229-11-89
- Verberne, M. C., Verpoorte, R., Bol, J. F., Mercado-Blanco, J., and Linthorst, H. J. (2000). Overproduction of salicylic acid in plants by bacterial transgenes enhances pathogen resistance. *Nat. Biotechnol.* 18, 779–783. doi: 10.1038/77347
- Vlot, A. C., Dempsey, D. A., and Klessig, D. F. (2009). Salicylic Acid, a multifaceted hormone to combat disease. *Annu. Rev. Phytopathol.* 47, 177–206. doi: 10.1146/annurev.phyto.050908.135202
- Volko, S. M., Boller, T., and Ausubel, F. M. (1998). Isolation of new *Arabidopsis* mutants with enhanced disease susceptibility to *Pseudomonas syringae* by direct screening. *Genetics* 149, 537–548.
- Wan, D., Li, R., Zou, B., Zhang, X., Cong, J., Wang, R., et al. (2012). Calmodulin-binding protein CBP60g is a positive regulator of both disease resistance and drought tolerance in *Arabidopsis*. *Plant Cell Rep.* 31, 1269–1281. doi: 10.1007/s00299-012-1247-7
- Wang, D., Amornsiripanitch, N., and Dong, X. (2006). A genomic approach to identify regulatory nodes in the transcriptional network of systemic acquired resistance in plants. *PLoS Pathog.* 2:e123. doi: 10.1371/journal.ppat.0020123
- Wang, L., and Fobert, P. R. (2013). *Arabidopsis* clade I TGA factors regulate apoplastic defences against the bacterial pathogen *Pseudomonas syringae* through endoplasmic reticulum-based processes. *PLoS ONE* 8:e77378. doi: 10.1371/journal.pone.0077378
- Wang, L., Tsuda, K., Sato, M., Cohen, J. D., Katagiri, F., and Glazebrook, J. (2009). *Arabidopsis* CaM binding protein CBP60g contributes to MAMP-induced SA accumulation and is involved in disease resistance against *Pseudomonas syringae*. *PLoS Pathog.* 5:e1000301. doi: 10.1371/journal.ppat.1000301
- Wang, L., Tsuda, K., Truman, W., Sato, M., Nguyen Le, V., Katagiri, F., et al. (2011). CBP60g and SARD1 play partially redundant critical roles in salicylic acid signaling. *Plant J.* 67, 1029–1041. doi: 10.1111/j.1365-313X.2011.04655.x
- Wang, S., Durrant, W. E., Song, J., Spivey, N. W., and Dong, X. (2010). *Arabidopsis* BRCA2 and RAD51 proteins are specifically involved in defense gene transcription during plant immune responses. *Proc. Natl. Acad. Sci. U.S.A.* 107, 22716–22721. doi: 10.1073/pnas.1005978107
- Weigel, R. R., Pfitzner, U. M., and Gatz, C. (2005). Interaction of NIMIN1 with NPR1 modulates PR gene expression in *Arabidopsis*. *Plant Cell* 17, 1279–1291. doi: 10.1105/tpc.104.027441
- Westfall, C. S., Herrmann, J., Chen, Q., Wang, S., and Jez, J. M. (2010). Modulating plant hormones by enzyme action: the GH3 family of acyl acid amido synthetases. *Plant Signal. Behav.* 5, 1607–1612. doi: 10.4161/psb.5.12.13941
- Westfall, C. S., Zubieta, C., Herrmann, J., Kapp, U., Nanao, M. H., and Jez, J. M. (2012). Structural basis for prereceptor modulation of plant hormones by GH3 proteins. *Science* 336, 1708–1711. doi: 10.1126/science.1221863
- Wildermuth, M. C., Dewdney, J., Wu, G., and Ausubel, F. M. (2001). Isochorismate synthase is required to synthesize salicylic acid for plant defence. *Nature* 414, 562–565. doi: 10.1038/35107108
- Wu, Y., Zhang, D., Chu, J. Y., Boyle, P., Wang, Y., Brindle, I. D., et al. (2012). The *Arabidopsis* NPR1 protein is a receptor for the plant defense hormone salicylic acid. *Cell Rep.* 1, 639–647. doi: 10.1016/j.celrep.2012.05.008
- Xin, X. F., and He, S. Y. (2013). *Pseudomonas syringae* pv. *tomato* DC3000: a model pathogen for probing disease susceptibility and hormone signaling

- in plants. *Annu. Rev. Phytopathol.* 51, 473–498. doi: 10.1146/annurev-phyto-082712-102321
- Yan, S., and Dong, X. (2014). Perception of the plant immune signal salicylic acid. *Curr. Opin. Plant Biol.* 20C, 64–68. doi: 10.1016/j.pbi.2014.04.006
- Yan, S., Wang, W., Marques, J., Mohan, R., Saleh, A., Durrant, W. E., et al. (2013). Salicylic acid activates DNA damage responses to potentiate plant immunity. *Mol. Cell* 52, 602–610. doi: 10.1016/j.molcel.2013.09.019
- Yu, D., Chen, C., and Chen, Z. (2001). Evidence for an important role of WRKY DNA binding proteins in the regulation of NPR1 gene expression. *Plant Cell* 13, 1527–1540. doi: 10.1105/tpc.13.7.1527
- Yuan, Y., Chung, J. D., Fu, X., Johnson, V. E., Ranjan, P., Booth, S. L., et al. (2009). Alternative splicing and gene duplication differentially shaped the regulation of isochorismate synthase in *Populus* and *Arabidopsis*. *Proc. Natl. Acad. Sci. U.S.A.* 106, 22020–22025. doi: 10.1073/pnas.0906869106
- Zhang, L., Du, L., Shen, C., Yang, Y., and Poovaiah, B. W. (2014). Regulation of plant immunity through ubiquitin-mediated modulation of  $Ca^{2+}$ -calmodulin-AtSR1/CAMTA3 signaling. *Plant J.* 78, 269–281. doi: 10.1111/tj.12473
- Zhang, Y., Tessaro, M. J., Lassner, M., and Li, X. (2003). Knockout analysis of *Arabidopsis* transcription factors TGA2, TGA5, and TGA6 reveals their redundant and essential roles in systemic acquired resistance. *Plant Cell* 15, 2647–2653. doi: 10.1105/tpc.014894
- Zhang, Y., Xu, S., Ding, P., Wang, D., Cheng, Y. T., He, J., et al. (2010). Control of salicylic acid synthesis and systemic acquired resistance by two members of a plant-specific family of transcription factors. *Proc. Natl. Acad. Sci. U.S.A.* 107, 18220–18225. doi: 10.1073/pnas.1005225107
- Zhang, Z., Li, Q., Li, Z., Staswick, P. E., Wang, M., Zhu, Y., et al. (2007). Dual regulation role of GH3.5 in salicylic acid and auxin signaling during *Arabidopsis*-*Pseudomonas syringae* interaction. *Plant Physiol.* 145, 450–464. doi: 10.1104/pp.107.106021
- Zheng, X. Y., Spivey, N. W., Zeng, W., Liu, P. P., Fu, Z. Q., Klessig, D. F., et al. (2012). Coronatine promotes *Pseudomonas syringae* virulence in plants by activating a signaling cascade that inhibits salicylic acid accumulation. *Cell Host Microbe* 11, 587–596. doi: 10.1016/j.chom.2012.04.014
- Zheng, Z., Qualley, A., Fan, B., Dudareva, N., and Chen, Z. (2009). An important role of a BAHD acyl transferase-like protein in plant innate immunity. *Plant J.* 57, 1040–1053. doi: 10.1111/j.1365-313X.2008.03747.x
- Zhou, J. M., Trifa, Y., Silva, H., Pontier, D., Lam, E., Shah, J., et al. (2000). NPR1 differentially interacts with members of the TGA/OBF family of transcription factors that bind an element of the PR-1 gene required for induction by salicylic acid. *Mol. Plant Microbe Interact.* 13, 191–202. doi: 10.1094/MPMI.2000.13.2.191
- Zhou, N., Tootle, T. L., Tsui, F., Klessig, D. F., and Glazebrook, J. (1998). PAD4 functions upstream from salicylic acid to control defense responses in *Arabidopsis*. *Plant Cell* 10, 1021–1030. doi: 10.1105/tpc.10.6.1021

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 15 October 2014; accepted: 24 November 2014; published online: 09 December 2014.

Citation: Seyfferth C and Tsuda K (2014) Salicylic acid signal transduction: the initiation of biosynthesis, perception and transcriptional reprogramming. *Front. Plant Sci.* 5:697. doi: 10.3389/fpls.2014.00697

This article was submitted to *Plant-Microbe Interaction*, a section of the journal *Frontiers in Plant Science*.

Copyright © 2014 Seyfferth and Tsuda. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.