

Sarcopenia and Sarcopenic Obesity in Patients Undergoing Orthopedic Surgery

Hyung-Min Ji, MD, Jun Han, MD, Dong San Jin, MD*, Hyunseok Suh, MD,
Yoon-Sok Chung, MD[†], Ye-Yeon Won, MD

Department of Orthopaedics, Ajou University School of Medicine, Suwon, Korea,

**Joint Centre, Beijing Mary Orthopedic Hospital, Beijing, China,*

[†]Department of Endocrinology and Metabolism, Ajou University School of Medicine, Suwon, Korea

Background: The purpose of this retrospective study was to determine the prevalence of sarcopenia and sarcopenic obesity among patients who underwent orthopedic surgery (OS).

Methods: A total of 222 patients were reviewed immediately after or prior to OS. In the control group, 364 patients from outpatient departments (OPDs) who did not have any OS were enrolled. Whole-body dual-energy X-ray absorptiometry was used to analyze body composition. Skeletal muscle mass was adjusted for height squared, total body weight, and height and fat mass (residuals). Obesity was defined as body mass index (BMI) > 25.0 kg/m².

Results: The prevalence of sarcopenia in the OS group was 25.7%, 44.1%, and 26.6%, respectively, according to the 3 different criteria. The prevalence was significantly lower in the OPD group (6.0%, 33.1%, and 14.8%, respectively). The highest rates of sarcopenia with height-adjusted definition were seen in patients with a femoral neck fracture. In the multivariate analysis, factors associated with sarcopenia were male gender, older age, and lower BMI (odds ratio [OR]: 28.38, 1.03, and 1.83, respectively) when muscle mass was adjusted for height, whereas male gender, older age, and higher BMI were associated with sarcopenia (OR: 1.04, 2.57, and 1.83, respectively) when adjusted for weight. When residuals were used as a cutoff, decreased BMI and total hip bone mineral density (0.1 g/cm²) were independent risk factors associated with sarcopenia (OR: 1.09 and 1.05). The prevalence of sarcopenic obesity ranged from 1.8% to 21.2%.

Conclusions: Our study demonstrated a high prevalence of sarcopenia among OS patients.

Keywords: Sarcopenia, Prevalence, Retrospective studies, Body composition, Orthopedics

Sarcopenia is a syndrome characterized by progressive and generalized loss of skeletal muscle mass and strength due to senescence, which may lead to physical disability, poor quality of life, and even increased mortality.¹⁾ The unprecedentedly rapid aging process in Asian countries has drawn much attention to the geriatric condition.¹⁾

Recently, a number of researches have been pub-

lished on the prevalence and related risk factors of sarcopenia in various cohorts in Asia.²⁻⁹⁾ To the best of our knowledge, however, the prevalence of sarcopenia among adults undergoing orthopedic surgery (OS) has never been reported. In the field of OS, bone mineral density (BMD) and bone quality have been the primary focus; however, recent studies have revealed that both bone and muscle are endocrine organs^{10,11)} that are closely associated in both formation and function.¹²⁾ Moreover, sarcopenia and osteoporosis constitute a hazardous duo that causes frailty in the elderly.¹³⁾ A recent study has demonstrated the synergistic effects of medications for osteoporosis and mechanical stimuli generated by the muscle, which resulted in the enhanced cortical thickness.¹⁴⁾ Such evidence might

Received September 30, 2015; Accepted February 19, 2016

Correspondence to: Ye-Yeon Won, MD

Department of Orthopaedics, Ajou University School of Medicine, 164
World cup-ro, Yeongtong-gu, Suwon 16499, Korea

Tel: +82-31-219-5220, Fax: +82-31-219-4537

E-mail: thrtkr@ajou.ac.kr

Copyright © 2016 by The Korean Orthopaedic Association

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Clinics in Orthopedic Surgery • pISSN 2005-291X eISSN 2005-4408

reveal how sarcopenia adversely affects musculoskeletal health and function in the intraoperative and postoperative settings, especially among the elderly who undergo a number of OS; thus, data on the prevalence of sarcopenia in this population may be invaluable. Moreover, the exact definition of sarcopenia in the orthopaedic field is still debatable, although the European Working Group on Sarcopenia in Older People devised a working clinical definition and consensus-based diagnostic criteria for sarcopenia in 2010.¹⁾ According to these criteria, one must have low muscle mass in order to be diagnosed as sarcopenic. However, a variety of methods are currently used to define low muscle mass¹⁵⁻¹⁷⁾ for orthopedic patients in the absence of any consensus on the ideal criterion of low muscle mass. Recently, the combined impact of muscle loss and fat gain on the functional disability in older persons has been reported,¹⁸⁾ and the prevalence of a condition known as sarcopenic obesity has become of great interest.

The purpose of this study was to identify the prevalence of sarcopenia and sarcopenic obesity among adult patients who had common OS of specific types, namely intra- and extracapsular proximal femoral fracture surgery and total hip/knee replacement (THR/TKR) and to compare the prevalence of sarcopenia in them with that in nonorthopedic patients at our and other outpatient departments (OPDs).

METHODS

Subjects

We retrospectively reviewed electronic medical records of 222 patients who underwent OS at Ajou University Hospital between February 2011 and April 2013 and those of 364 patients who did not have any orthopedic proce-

dures in various OPDs including Orthopedics, Internal Medicine, and Family Medicine. The study was conducted after obtaining approval from Ajou University Hospital Institutional Review Board (IRB No. AJIRB-MED-MDB-14-172) and informed consent was waived by the IRB. Ajou University Hospital is located in an urban area with a population of more than 1 million and has served as the sole tertiary hospital for over 20 years. Initially, 401 OPD patients were considered for inclusion; however, 9.2% (37 patients) of them were excluded due to surgery on the lower extremity. All OPD patients underwent dual-energy X-ray absorptiometry (DXA) for routine BMD assessment, and most of them were prescribed medications for osteoporosis. All of the OS patients underwent whole-body DXA (Lunar iDXA; GE Healthcare, Waukesha, WI, USA) 2 weeks prior to or after surgery. Patients who underwent elective procedures such as THR and TKR underwent preoperative whole-body DXA routinely whereas patients who suffered from fragility hip fractures received DXA postoperatively when they were ambulatory and free of severe pain. Patients who did not receive whole-body DXA were excluded from the study; 23% (66 patients) of all OS patients (288 patients) who did not complete whole-body DXA during the study period were excluded.

Patient Demographics

Ultimately, 586 patients (117 males and 469 females) were included in the study (Table 1). There were significantly more women in the OPD group. The average age of the patients was 66.1 ± 13.0 years. Patients undergoing OS were significantly older by 6.4 years than the OPD patients and taller, whereas body mass index (BMI) was slightly higher in the OPD group. The total hip BMD was higher in the OPD group than in the OS group. Male patients in

Table 1. Patient Demographics of OS Patients and OPD Patients

Characteristic	TKR (n = 79)	THR (n = 38)	ITF surgery (n = 53)	FNF surgery (n = 52)	Overall OS patients (n = 222)	OPD patients (n = 364)	p-value*
Age (yr)	70.3 ± 6.2	60.9 ± 13.3	77.5 ± 8.9	69.4 ± 10.7	70.2 ± 10.8	63.6 ± 13.6	<0.001
Female (%)	91.1	69.1	62.3	69.2	68.0	86.8	<0.001
Height (cm)	153.7 ± 6.9	163.2 ± 10.6	157.0 ± 9.2	159.3 ± 7.2	157.4 ± 8.9	154.8 ± 8.7	0.001
Weight (kg)	63.2 ± 9.9	62.5 ± 11.2	56.1 ± 10.1	56.0 ± 9.0	59.7 ± 10.5	60.0 ± 13.7	0.766
Body mass index (kg/m ²)	26.8 ± 3.1	23.4 ± 3.4	22.6 ± 3.8	21.9 ± 3.8	24.1 ± 4.0	24.9 ± 4.2	0.015
Total hip BMD (g/cm ²)	0.815 ± 0.107	0.887 ± 0.153	0.662 ± 0.145	0.723 ± 0.136	0.771 ± 0.153	0.826 ± 0.159	<0.001

Values are presented as mean ± standard deviation.

OS: orthopedic surgery, OPD: outpatient department, TKR: total knee replacement, THR: total hip replacement, ITF: intertrochanteric fracture, FNF: femoral neck fracture, BMD: bone mineral density.

*Unpaired Student *t*-test and chi-square test.

the OS group were older (67.8 ± 12.8 year vs. 53.9 ± 18.1 year, $p < 0.001$), shorter (166.3 ± 7.4 cm vs. 169.4 ± 8.2 cm, $p = 0.032$), and lighter (62.3 ± 11.2 kg vs. 78.5 ± 20.8 kg, $p < 0.001$) than those in the OPD group. The average BMI of the OS group was lower than that of the OPD group (22.1 ± 3.4 kg/m² vs. 27.2 ± 5.6 kg/m², $p < 0.001$). The total hip BMD of the OS group was significantly lower than that of the OPD group (0.845 ± 0.144 g/cm² vs. 0.979 ± 0.177 g/cm², $p = 0.011$). Women were older in the OS group than in the OPD group (71.3 ± 9.5 year vs. 65.1 ± 12.2 year, $p < 0.001$) and their total hip BMD was lower in the OS group than in the OPD group (0.737 ± 0.146 g/cm² vs. 0.803 ± 0.142 g/cm², $p < 0.001$), while there was no significant difference in weight and BMI between the groups (58.6 ± 10.0 kg vs. 57.2 ± 9.6 kg, $p = 0.185$; 24.9 ± 4.0 kg/m² vs. 24.6 ± 3.8 kg/m², $p = 0.365$).

Body Composition and BMD

Body composition and BMD were measured by whole-body DXA. The accuracy error for BMD measurements was determined according to the standardized protocols from the International Society for Clinical Densitometry.¹⁹ All BMD measurements were performed by an experienced technician at the lumbar spine and the proximal femur. The total hip BMD was used for final analysis. Appendicular skeletal mass (ASM) was defined as the sum of muscle mass in the arms and legs, assuming that all nonfat and nonbone tissue is skeletal muscle. The total fat mass was recorded in grams and as a percentage of the total body weight.

Definitions of Sarcopenia and Sarcopenic Obesity

Three different measures of ASM were applied for defining sarcopenia: ASM relative to height squared, ASM relative to body weight, and ASM relative to height and total fat mass (residuals). Class I sarcopenia was defined as when the height- or weight-adjusted ASM was between 1 and 2 standard deviations (SDs) below the gender-specific mean for young adults. Class II sarcopenia was defined as when the ASM was below 2 SD.¹⁶ Height-adjusted ASM was calculated using the equation (ASM/height², kg/m²) previously proposed by Baumgartner et al.¹⁵ The cut-off value for class II sarcopenia among Koreans was defined as 6.58 kg/m² for men and 4.59 kg/m² for women.²⁰ Body weight-adjusted ASM was expressed as a percentage of lean body mass relative to body weight, skeletal muscle mass index (ASM/weight \times 100 [%]),¹⁶ and the cut-off value of Koreans was defined as 29.1% for men and 23.0% for women.²⁰ The ASM relative to height and total fat mass was determined after adjusting for height and fat mass using a linear

regression model.¹⁷ The residuals of the regression model were analysed, and the 20th percentile of the distribution of residuals of all individuals was used as the cut-off point, which was -0.98 for men and -0.91 for women. Obesity among Koreans was defined as having a BMI of > 25.0 kg/m².

Background Data

The potential demographic risk factors for sarcopenia were assessed: OS, age, gender, BMI, and total hip BMD. All subjects were Korean. The patients were categorized into 3 groups according to age (8–50 years, 51–70 years, and > 70 years) to account for the different levels of physical activity and their potential influence on sarcopenia.

Statistical Analysis

We calculated the distribution of relative ASM adjusted by height, weight, and residuals (mean, SD), the percentages of patients with class I and II sarcopenia adjusted for height and weight, and also the ratio of patients with sarcopenia when the residual method was used. We used unpaired Student *t*-test to compare characteristics between the OS and OPD patients. Univariate logistic regression models were used to identify the independent relative risk for class II sarcopenia. Stepwise multivariate logistic regression analysis was performed to evaluate the independent effects of age, gender, OS, BMI, and the total femoral BMD, where *p*-values were all less than 0.15. In the multivariate analysis, a forward method was used and age was considered a continuous variable. The level of significance was set at $p \leq 0.05$. The unadjusted odds ratio (OR), 95% confidence interval (CI), and *p*-value estimated for each risk factor were also determined. The chi-square test and Fisher exact test were used to compare ratios when indicated. All analyses were performed using two-tailed tests on SPSS ver. 16.0 (SPSS Inc., Chicago, IL, USA).

RESULTS

Prevalence of Sarcopenia According to Different Criteria

When using the height-adjusted ASM, the average relative lean mass in OS patients was 6.33 ± 1.06 kg/m² for males and 5.85 ± 0.92 kg/m² for females, whereas the value in OPD patients was 7.89 ± 1.47 kg/m² for males and 5.88 ± 0.88 kg/m² for females. The prevalence of sarcopenia in the OS group was significantly higher irrespective of the criteria (Table 2). The subjects were divided into 2 groups based on gender and then subdivided into 3 different age groups to elucidate the effects of age and gender on the prevalence of class II sarcopenia (Table 2). The residual method was the only measure in which the prevalence was

Table 2. Prevalence of Sarcopenia among OS Patients and OPD Patients Based on the ASM/Height, ASM/Weight, and Residual Methods

Variable	ASM/height ²			ASM/weight			Residual		
	OS	OPD	p-value	OS	OPD	p-value	OS	OPD	p-value
Whole							26.6	14.8	<0.001* [†]
Normal	51.8	70.1		14.0	21.8		-	-	-
Type I	22.5	23.9		41.9	45.1		-	-	-
Type II	25.7	6.0	<0.001* [†]	44.1	33.1	0.011*	-	-	-
Male									
18–50 yr (n = 23)	25.0	10.5	0.453*	50.0	47.4	1.000*	25.0	16.7	1.000*
51–70 yr (n = 47)	56.7 [†]	0.0 [†]	<0.001* [†]	53.3	52.9	1.000*	23.3 [†]	0.0 [†]	0.039* [†]
>70 yr (n = 47)	71.4	41.7	0.064	57.1	41.7	0.354	30.3	8.3	0.240*
Total (n = 117)	62.3 [†]	14.6 [†]	<0.001 [†]	55.1	47.9	0.459	26.9 [†]	8.5 [†]	0.016 [†]
Female									
18–50 yr (n = 28)	40.0 [†]	0.0 [†]	0.026* [†]	60.0	17.4	0.082*	40.0	4.3	0.073*
51–70 yr (n = 233)	5.4	1.7	0.152*	44.6 [†]	24.9 [†]	0.007 [†]	21.4	13.6	0.202
>70 yr (n = 208)	9.8	10.3	0.894	34.8	43.1	0.223	27.2	21.6	0.346

Values are presented as percentage.

OS: orthopedic surgery, OPD: outpatient department, ASM: appendicular skeletal mass.

*Fisher exact test, otherwise chi-square test. [†]Statistical significance.

higher in both male and female patients.

Comparison of Prevalence of Sarcopenia According to Different Types of Surgeries

The highest rate of class II sarcopenia with height-adjusted definition was seen in patients who were diagnosed with femoral neck fracture (FNF) (Fig. 1). When adjusted for weight, the rate was highest in patients who received THR (52.6%). When using the residual method, the prevalence of sarcopenia was highest in the FNF patients (32.7%).

Prevalence of Obesity and Sarcopenic Obesity According to Different Criteria

There was no significant difference between the height-adjusted ASM and weight-adjusted ASM with regard to the prevalence of obesity and sarcopenic obesity. However, there were significantly more patients with sarcopenic obesity when the residual method was applied ($p = 0.049$) (Table 3).

Risk Factors According to Different Criteria for Sarcopenia

When the ASM was adjusted for height, males and elderly patients were more likely to be diagnosed as sarcopenic in

the multivariate analysis (Table 4). When the ASM was adjusted for weight, older age, male sex, and lower BMI were independent risk factors. When the diagnosis of sarcopenia was determined by the residual method, lower BMI and BMD were independent risk factors.

DISCUSSION

Patients with sarcopenia are at a greater risk for falls²¹⁾ and incident disability.²²⁾ Therefore, there are significant implications of sarcopenia in patients undergoing OS, especially those performed on the lower extremities among the elderly. Our study demonstrated that there were significantly more patients with sarcopenia in the OS group than in the OPD group. The prevalence ranged from 25.7% to 44.1% in the OS group according to different criteria whereas this was between 6.0% and 33.1% in the OPD group. In all age- and gender-matched subgroups (except for women > 70 years of age), the prevalence was also higher in the OS patients based on all 3 criteria even though no statistical significance was found in all subgroups. To the best of our knowledge, these results are unprecedented; hence, we were unable to find any comparative studies. This discrepancy between the OS group and the OPD groups was veri-




Fig. 1. Bar chart depicting the prevalence of type I and II sarcopenia among the different types of surgeries when adjusted for height (A), weight (B), and residuals (C). ASM: appendicular skeletal mass, TKR: total knee replacement, THR: total hip replacement.

Table 3. Prevalence of Obesity and Sarcopenic Obesity Based on the ASM/Height, ASM/Weight, and Residual Methods

Variable	Obesity	ASM/height ²		ASM/weight		Residual	
		NSO	SO	NSO	SO	NSO	SO
Outpatient department (n = 364)	41.8	41.5	0.3	22.6	19.2	38.2	3.6
Total knee replacement (n = 79)	74.7	73.4	1.3	39.3	35.4	65.8	8.9
Total hip replacement (n = 38)	28.9	28.9	0.0	10.5	18.4	21.0	7.9
Intertrochanteric fracture surgery (n = 53)	28.3	24.5	3.8	17.0	11.3	22.6	5.7
Femoral neck fracture surgery (n = 52)	21.2	19.3	1.9	9.7	11.5	15.4	5.8
Overall orthopedic surgery patients (n = 222)	43.2	41.4	1.8	22.0	21.2	36.0	7.2

Values are presented as percentage.
 ASM: appendicular skeletal mass, NSO: nonsarcopenic obesity, SO: sarcopenic obesity.

Table 4. Univariate and Multivariate Analysis of Risk Factors for Sarcopenia by Different Criteria

Variable	Univariate analysis			Multivariate analysis		
	OR*	95% CI [†]	p-value	OR*	95% CI [†]	p-value
ASM/height ²						
Age [‡]	2.79	1.65–4.71	< 0.001	1.03	1.00–1.07	0.015
Man	11.32	6.70–19.14	< 0.001	28.38	10.69–75.32	< 0.001
OS [§]	5.38	3.17–9.09	< 0.001	-	-	-
BMI	1.71	1.52–1.91	< 0.001	1.83	1.55–2.18	< 0.001
BMD [¶]	1.38	1.17–1.63	< 0.001	-	-	-
ASM/weight						
Age [‡]	1.43	1.01–2.03	0.046	2.57	1.58–4.18	< 0.001
Man	2.14	1.42–3.23	< 0.001	1.04	1.02–1.05	< 0.001
OS [§]	1.59	1.13–2.24	0.008	-	-	-
BMI	1.13	1.08–1.18	< 0.001	1.83	1.55–2.18	< 0.001
BMD [¶]	1.46	0.84–1.73	0.648	-	-	-
Residual						
Age [‡]	1.73	1.12–2.67	0.013	-	-	-
OS [§]	1.98	1.31–3.01	0.001	-	-	-
BMI	1.15	1.17–1.63	< 0.001	1.09	1.02–1.79	0.007
BMD [¶]	1.44	1.24–1.67	< 0.001	1.05	1.11–1.54	0.001

OR: odds ratio, CI: confidence interval, ASM: appendicular skeletal mass, OS: orthopedic surgery, BMI: body mass index, BMD: bone mineral density.

*OR estimated from univariate and stepwise multivariate logistic regression analyses. [†]CI of the estimated OR. [‡]Whenever age increased by 1 year. [§]OS patients compared with outpatient department patients. ^{||}Whenever BMI decreased by 1.0 kg/m². [¶]Whenever BMD decreased by 0.1 g/cm².

fied by our results of multivariate regression analysis. In the univariate logistic regression analysis, OS patients were 5.38, 1.59, and 1.98 times more likely to have sarcopenia when ASM was adjusted for height, weight, and residuals, respectively. In the multivariate analysis, however, OS was not an independent risk factor after adjusting for variables, whereas age, gender, and BMI were all independent risk factors when the ASM was adjusted for height and weight with BMD being the only independent factor when the residual method was used. These differences in patient demographics between the 2 groups resulted in significantly different values of prevalence. The demographics of the OS patients from this study were quite similar to those in recently reported studies on TKR,²³⁾ THR,²⁴⁾ and hip fracture surgery,²⁵⁾ which were conducted on Korean patients, and so were the demographics of the OPD patients in our study when compared with the data from a nationwide survey on nutritional status.²⁰⁾

However, the prevalence of sarcopenia among elderly patients from a national survey in our country was 12.4% in men and 0.1% in women based on the height-adjusted criteria and 9.7% in men and 11.8% in women based on the weight-adjusted criteria.²⁰⁾ These values are clearly lower than ours: 41.7% of men and 10.3% of women in the OPD group were sarcopenic based on the height-adjusted criteria, whereas 41.7% of males and 43.1% of females were sarcopenic based on the weight-adjusted criteria. These discrepancies might be attributable to the fact that all of the OPD patients underwent DXA for various underlying conditions that could affect the bone quantity and quality leading to osteopenia and osteoporosis, such as rheumatologic or metabolic disorder, although we did not investigate this further.

There were clear differences in the prevalence of sarcopenia among different types of OS when applying different criteria. The highest prevalence of class II sarco-

penia based on the height-adjusted definition was seen in patients with FNF and intertrochanteric fracture (ITF). The prevalence was higher in the FNF group than the ITF group. Although the male-to-female ratio and average age were higher in the ITF group, BMI was lower by 0.7 kg/m² in the FNF group, which resulted in the higher prevalence. The prevalence was the lowest among patients who had TKR, which was even lower than that of the OPD patients. The female predominance in TKR patients might explain such a low sarcopenia prevalence. In contrast, when adjusted for weight, the prevalence of class II sarcopenia was highest in patients who had THR (52.6%), which was slightly higher than patients who had FNF surgery (48.1%) or TKR (44.3%). The OR of gender and BMI in case of weight-adjusted sarcopenia (1.04 and 1.83, respectively) using the multivariate regression models were relatively smaller compared to those using the height-adjusted definition (28.38 and 1.83, respectively). Therefore, it seems logical to assume that there would be another factor for the prediction of sarcopenia when adjusting for weight. Another nationwide study performed in Korea showed that men and women in the lowest skeletal muscle mass index quintile were more likely to have metabolic syndrome by 4.62 times and 3.55 times, respectively, than patients in the highest quintile.²⁶⁾ Thus, sarcopenia based on the weight-adjusted definition may be associated with metabolic syndrome. Sarcopenia based on the residual method (ASM adjusted for height and total fat mass) was reported to be closely related to lower extremity physical function.¹⁷⁾ Recent studies on the prevalence of sarcopenia demonstrated that the residual method could be more useful in identifying overweight senior population with sarcopenia than the height-adjusted method.^{27,28)} Loss of BMD involving the whole hip, which is one of the well-known risk factors for sarcopenia, was an independent risk factor for sarcopenia only when applying the residual method in the multiple regression analysis and a significant difference in prevalence between the OS and OPD patients in both men and women was also observed only when the residual method was applied. Our findings and abovementioned studies suggest that the residual method should be included when clinicians attempt to evaluate muscle volume before and after OS.

This study has its limitations. First, only elderly Ko-

rean patients were included in the study; thus, such a high prevalence of sarcopenia as in our OS patients might not be found in other ethnic groups. However, the risk factors for sarcopenia and the associated OR reported in this study were comparable to those in studies involving different ethnicities.^{17,27,28)} Therefore, our findings might be cautiously considered clinically relevant for clinicians outside Korea. Second, the body composition data from whole-body DXA bears fundamental shortcomings; sarcopenia cannot be diagnosed based on the relative muscle volume alone, as muscle strength and functional capacity also play a role.¹⁾ Besides, recent studies have demonstrated that fat degeneration of the thigh muscles is far more important for predicting incident disability and future hip fractures than muscle mass alone.^{22,29)} Our study did not address serum markers such as tumor necrosis factor (TNF- α), interleukin (IL-6), IL-1, albumin, or 25-hydroxy vitamin D levels³⁰⁾ because the primary focus of the study was to evaluate the muscle volume and to highlight and raise awareness among clinicians of such high prevalence of sarcopenia. We hope that our study would serve as a foundation or reference for future studies. Third, the timing of DXA scans was not standardized among OS patients: DXA scans were performed prior to surgery in patients with conditions other than hip fracture, whereas patients with hip fracture underwent DXA scans when they were able to ambulate with little pain after surgery. All of these limitations necessitate further prospective studies among OS patients.

In conclusion, we demonstrated that the prevalence of sarcopenia among Korean adults receiving OS on a lower extremity is relatively high. Male gender, older age, and BMI were universally significant risk factors while total hip BMD mattered only for sarcopenia based on the definition of ASM adjusted by height and fat mass. These risk factors should be taken into consideration pre- and postoperatively in light of the clinical significance of sarcopenia for musculoskeletal health among the elderly.

CONFLICT OF INTEREST

No potential conflict of interest relevant to this article was reported.

REFERENCES

1. Cruz-Jentoft AJ, Baeyens JP, Bauer JM, et al. Sarcopenia: European consensus on definition and diagnosis: report of the European Working Group on Sarcopenia in Older People. *Age Ageing*. 2010;39(4):412-23.

2. Hida T, Ishiguro N, Shimokata H, et al. High prevalence of sarcopenia and reduced leg muscle mass in Japanese patients immediately after a hip fracture. *Geriatr Gerontol Int.* 2013;13(2):413-20.
3. Kaido T, Ogawa K, Fujimoto Y, et al. Impact of sarcopenia on survival in patients undergoing living donor liver transplantation. *Am J Transplant.* 2013;13(6):1549-56.
4. Kim JK, Choi SR, Choi MJ, et al. Prevalence of and factors associated with sarcopenia in elderly patients with end-stage renal disease. *Clin Nutr.* 2014;33(1):64-8.
5. Kim SH, Kim TH, Hwang HJ. The relationship of physical activity (PA) and walking with sarcopenia in Korean males aged 60 years and older using the Fourth Korean National Health and Nutrition Examination Survey (KNHANES IV-2, 3), 2008-2009. *Arch Gerontol Geriatr.* 2013;56(3):472-7.
6. Lee WJ, Liu LK, Peng LN, Lin MH, Chen LK; ILAS Research Group. Comparisons of sarcopenia defined by IWGS and EWGSOP criteria among older people: results from the I-Lan longitudinal aging study. *J Am Med Dir Assoc.* 2013;14(7):528.e1-7.
7. Lin CC, Lin WY, Meng NH, et al. Sarcopenia prevalence and associated factors in an elderly Taiwanese metropolitan population. *J Am Geriatr Soc.* 2013;61(3):459-62.
8. Liu LK, Lee WJ, Liu CL, et al. Age-related skeletal muscle mass loss and physical performance in Taiwan: implications to diagnostic strategy of sarcopenia in Asia. *Geriatr Gerontol Int.* 2013;13(4):964-71.
9. Miyakoshi N, Hongo M, Mizutani Y, Shimada Y. Prevalence of sarcopenia in Japanese women with osteopenia and osteoporosis. *J Bone Miner Metab.* 2013;31(5):556-61.
10. Fukumoto S, Martin TJ. Bone as an endocrine organ. *Trends Endocrinol Metab.* 2009;20(5):230-6.
11. Pratesi A, Tarantini F, Di Bari M. Skeletal muscle: an endocrine organ. *Clin Cases Miner Bone Metab.* 2013;10(1):11-4.
12. DiGirolamo DJ, Kiel DP, Esser KA. Bone and skeletal muscle: neighbors with close ties. *J Bone Miner Res.* 2013;28(7):1509-18.
13. Park JH, Park KH, Cho S, et al. Concomitant increase in muscle strength and bone mineral density with decreasing IL-6 levels after combination therapy with alendronate and calcitriol in postmenopausal women. *Menopause.* 2013;20(7):747-53.
14. Poole KE, Treece GM, Ridgway GR, Mayhew PM, Borggrefe J, Gee AH. Targeted regeneration of bone in the osteoporotic human femur. *PLoS One.* 2011;6(1):e16190.
15. Baumgartner RN, Koehler KM, Gallagher D, et al. Epidemiology of sarcopenia among the elderly in New Mexico. *Am J Epidemiol.* 1998;147(8):755-63.
16. Janssen I, Heymsfield SB, Ross R. Low relative skeletal muscle mass (sarcopenia) in older persons is associated with functional impairment and physical disability. *J Am Geriatr Soc.* 2002;50(5):889-96.
17. Newman AB, Kupelian V, Visser M, et al. Sarcopenia: alternative definitions and associations with lower extremity function. *J Am Geriatr Soc.* 2003;51(11):1602-9.
18. Rolland Y, Lauwers-Cances V, Cristini C, et al. Difficulties with physical function associated with obesity, sarcopenia, and sarcopenic-obesity in community-dwelling elderly women: the EPIDOS (EPIDemiologie de l'OSteoporose) Study. *Am J Clin Nutr.* 2009;89(6):1895-900.
19. Shepherd JA, Lu Y, Wilson K, et al. Cross-calibration and minimum precision standards for dual-energy X-ray absorptiometry: the 2005 ISCD Official Positions. *J Clin Densitom.* 2006;9(1):31-6.
20. Kim YS, Lee Y, Chung YS, et al. Prevalence of sarcopenia and sarcopenic obesity in the Korean population based on the Fourth Korean National Health and Nutritional Examination Surveys. *J Gerontol A Biol Sci Med Sci.* 2012;67(10):1107-13.
21. Moreland JD, Richardson JA, Goldsmith CH, Clase CM. Muscle weakness and falls in older adults: a systematic review and meta-analysis. *J Am Geriatr Soc.* 2004;52(7):1121-9.
22. Visser M, Goodpaster BH, Kritchevsky SB, et al. Muscle mass, muscle strength, and muscle fat infiltration as predictors of incident mobility limitations in well-functioning older persons. *J Gerontol A Biol Sci Med Sci.* 2005;60(3):324-33.
23. Chang MJ, Kang YG, Chang CB, Seong SC, Kim TK. The patterns of limb length, height, weight and body mass index changes after total knee arthroplasty. *J Arthroplasty.* 2013;28(10):1856-61.
24. Kang BJ, Ha YC, Hwang SC, Lee YK, Koo KH. Midterm results of large diameter Biolox forte ceramic head on delta ceramic liner articulation in total hip arthroplasty. *J Arthroplasty.* 2014;29(12):2412-4.
25. Lee YK, Ha YC, Yoon BH, Koo KH. Incidence of second hip fracture and compliant use of bisphosphonate. *Osteoporos Int.* 2013;24(7):2099-104.
26. Kim TN, Yang SJ, Yoo HJ, et al. Prevalence of sarcopenia and sarcopenic obesity in Korean adults: the Korean sarcopenic obesity study. *Int J Obes (Lond).* 2009;33(8):885-92.
27. Figueiredo CP, Domiciano DS, Lopes JB, et al. Prevalence of sarcopenia and associated risk factors by two diagnostic cri-

- teria in community-dwelling older men: the Sao Paulo Ageing & Health Study (SPAH). *Osteoporos Int.* 2014;25(2):589-96.
28. Estrada M, Kleppinger A, Judge JO, Walsh SJ, Kuchel GA. Functional impact of relative versus absolute sarcopenia in healthy older women. *J Am Geriatr Soc.* 2007;55(11):1712-9.
29. Lang T, Cauley JA, Tylavsky F, et al. Computed tomographic measurements of thigh muscle cross-sectional area and attenuation coefficient predict hip fracture: the health, aging, and body composition study. *J Bone Miner Res.* 2010;25(3):513-9.
30. Pahor M, Manini T, Cesari M. Sarcopenia: clinical evaluation, biological markers and other evaluation tools. *J Nutr Health Aging.* 2009;13(8):724-8.