

Schwarz–Christoffel Mapping

Tobin A. Driscoll
University of Delaware

Lloyd N. Trefethen
Oxford University



PUBLISHED BY THE PRESS SYNDICATE OF THE UNIVERSITY OF CAMBRIDGE
The Pitt Building, Trumpington Street, Cambridge, United Kingdom

CAMBRIDGE UNIVERSITY PRESS
The Edinburgh Building, Cambridge CB2 2RU, UK
40 West 20th Street, New York, NY 10011-4211, USA
477 Williamstown Road, Port Melbourne, VIC 3207, Australia
Ruiz de Alarcón 13, 28014 Madrid, Spain
Dock House, The Waterfront, Cape Town 8001, South Africa

<http://www.cambridge.org>

© Cambridge University Press 2002

This book is in copyright. Subject to statutory exception
and to the provisions of relevant collective licensing agreements,
no reproduction of any part may take place without
the written permission of Cambridge University Press.

First published 2002

Printed in the United Kingdom at the University Press, Cambridge

Typeface Times Roman 10/13 pt. *System* L^AT_EX 2_ε [TB]

A catalog record for this book is available from the British Library.

Library of Congress Cataloging in Publication Data

Driscoll, Tobin A. (Tobin Allen), 1969–

Schwarz–Christoffel mapping / Tobin A. Driscoll, Lloyd N. Trefethen.

p. cm.—(Cambridge monographs on applied and computational mathematics ; v. 8)

ISBN 0-521-80726-3

1. Conformal mapping. I. Trefethen, Lloyd N. (Lloyd Nicholas) II. Title.
III. Cambridge monographs on applied and computational mathematics ; 8.

QA360 .D75 2002
516.3'6—dc21 2001043099

ISBN 0 521 80726 3 hardback

Contents

Figures	<i>page</i> xi
Preface	xv
1 Introduction	1
1.1 The Schwarz–Christoffel idea	1
1.2 History	4
2 Essentials of Schwarz–Christoffel mapping	9
2.1 Polygons	9
2.2 The Schwarz–Christoffel formula	10
2.3 Polygons with one or two vertices	12
2.4 Triangles	16
2.5 Rectangles and elliptic functions	18
2.6 Crowding	20
3 Numerical methods	23
3.1 Side-length parameter problem	23
3.2 Quadrature	27
3.3 Inverting the map	29
3.4 Cross-ratio parameter problem	30
3.5 Mapping using cross-ratios	36
3.6 Software	39
4 Variations	41
4.1 Mapping from the disk	42
4.2 Mapping from a strip	44
4.3 Mapping from a rectangle	47
4.4 Exterior maps	51

4.5	Periodic regions and fractals	55
4.6	Reflections and other transformations	57
4.7	Riemann surfaces	58
4.8	Gearlike regions	60
4.9	Doubly connected regions	64
4.10	Circular-arc polygons	70
4.11	Curved boundaries	73
5	Applications	75
5.1	Why use Schwarz–Christoffel maps?	75
5.2	Piecewise-constant boundary conditions	77
5.3	Alternating Dirichlet and Neumann conditions	83
5.4	Oblique derivative boundary conditions	87
5.5	Generalized parameter problems	99
5.6	Free-streamline flows	101
5.7	Mesh generation	105
5.8	Polynomial approximation and matrix iterations	108
5.9	Symmetric multiply connected domains	111
	Appendix: Using the SC Toolbox	115
	Bibliography	121
	Index	131

Figures

1.1	Notational conventions	<i>page</i> 2
1.2	Action of a term in the SC product	3
1.3	The effect of prevertices on side lengths	4
1.4	Schwarz's plot of the conformal map of a square	6
2.1	Interior angles for a vertex at infinity	10
2.2	Map from the disk to a half-plane	13
2.3	Map from the half-plane to a strip	13
2.4	Map from a disk to a strip	14
2.5	A different map of the half-plane to a strip	14
2.6	Map from a doubly slit plane using reflection	15
2.7	Other maps obtained by reflection	15
2.8	Maps from the half-plane to wedges	16
2.9	Maps to triangles with two infinite vertices	16
2.10	Maps to triangles with one infinite vertex	17
2.11	Map from the half-plane to a rectangle	18
2.12	Map of a generalized quadrilateral	19
2.13	Illustration of crowding	20
3.1	Location of the last vertex by intersection	24
3.2	Maps to regions with 100 vertices	26
3.3	Endless descent in the side-length parameter problem	27
3.4	Embeddings with different crowding localities	31
3.5	Triangulation and quadrilaterals in the CRDT scheme	32
3.6	Summary of the CRDT idea	34
3.7	Preprocessing step for CRDT	35

3.8	Map to a “maze” using CRDT	37
3.9	Map to the maze from a multiply elongated region	38
4.1	Examples of disk maps	43
4.2	More examples of disk maps	44
4.3	Elementary factor in the strip map	45
4.4	Divergence adjustment of the strip map	46
4.5	Examples of strip maps	47
4.6	More examples of strip maps	48
4.7	Comparing the rectangle, half-plane, and strip	49
4.8	Examples of rectangle maps	50
4.9	More examples of rectangle maps	51
4.10	Examples of exterior maps	53
4.11	More examples of exterior maps	54
4.12	Map to a periodic channel	55
4.13	Map to a self-similar spiral	56
4.14	Map to a symmetric doubly connected region by reflection	57
4.15	Map to a symmetrically periodic channel by reflection	58
4.16	Maps to certain regions with circular arcs	59
4.17	Map to a three-sheeted Riemann surface	61
4.18	Maps to gearlike regions	62
4.19	Logarithms of the gearlike regions	63
4.20	Notation for the doubly connected map	64
4.21	First steps toward the doubly connected map	65
4.22	Convergence of a doubly connected SC factor	67
4.23	Examples of doubly connected SC maps	70
4.24	Maps to circular-arc polygons	73
5.1	Solving Laplace’s equation with two boundary values	77
5.2	Components of a piecewise-constant harmonic function	79
5.3	Examples of four-boundary-value harmonic functions	81
5.4	More piecewise-constant Dirichlet examples	82
5.5	Solutions to the D/N/D/N Laplace problem	84
5.6	Solutions of three-value alternating D/N problems	84
5.7	Riemann surface in the four-value alternating D/N problem	87
5.8	Examples of alternating D/N solutions	88
5.9	More examples of alternating D/N solutions	89
5.10	Oblique derivative problem on an L-shaped region	90
5.11	Solution of the L-shaped oblique derivative problem	91
5.12	More solutions to the L-shaped oblique problem	92

5.13	Examples from a space of solutions to an oblique derivative problem	95
5.14	More examples of solutions to oblique derivative problems	96
5.15	Illustration of the Hall effect	97
5.16	Illustration of reflected Brownian motion	98
5.17	Resistor trimming problem	100
5.18	Multiple resistance measurements for locating cracks	100
5.19	Wake flow past a polygonal obstacle	101
5.20	Potential and computational planes for wake flow	102
5.21	Elementary factor in the free-streamline map	103
5.22	Examples of wake flows	106
5.23	Notation for approximation discussion	109
5.24	Level curves of Faber polynomials	110
5.25	Green's function for a symmetric multiply connected region	113
5.26	More examples of Green's functions	114

1.1 The Schwarz–Christoffel idea

The idea behind the Schwarz–Christoffel (SC) transformation and its variations is that a conformal transformation f may have a derivative that can be expressed as

$$f' = \prod f_k \tag{1.1}$$

for certain canonical functions f_k . A surprising variety of conformal maps can be fitted into this basic framework. In fact, virtually all conformal transformations whose analytic forms are known are Schwarz–Christoffel maps, albeit sometimes disguised by an additional change of variables.

Geometrically speaking, the significance of (1.1) is that

$$\arg f' = \sum \arg f_k.$$

In the classical transformation, each $\arg f_k$ is designed to be a step function, so the resulting $\arg f'$ is piecewise constant with specific jumps (i.e., f maps the real axis onto a polygon). To be specific, let P be the region in the complex plane \mathbf{C} bounded by a polygon Γ with vertices w_1, \dots, w_n , given in counter-clockwise order, and interior angles $\alpha_1\pi, \dots, \alpha_n\pi$. For now, we assume that P is bounded and without cusps or slits, so that $\alpha_k \in (0, 2)$ for each k . Let f be a conformal map of the upper half-plane H^+ onto P , and let $z_k = f^{-1}(w_k)$ be the k th **prevertex**.¹ We shall assume $z_n = \infty$ without loss of generality, for if infinity is not already a prevertex, we can simply introduce its image (which lies

¹ The Carathéodory–Osgood theorem [Hen74] guarantees a continuous extension of f to the boundary. Hence the prevertices are well defined.




Figure 1.1. Notational conventions for the Schwarz–Christoffel transformation. In this case, z_1 and z_2 are mathematically distinct but graphically difficult to distinguish. As with all figures in this book, everything shown is not just schematic but also quantitatively correct.

on Γ) as a new vertex with interior angle π . The other prevertices z_1, \dots, z_{n-1} are real. Figure 1.1 illustrates these definitions.

As with all conformal maps, the main effort is in getting the boundary right. By the Schwarz reflection principle, which was invented for this purpose, f can be analytically continued across the segment (z_k, z_{k+1}) . In particular, f' exists on this segment, and we see that $\arg f'$ must be constant there. Furthermore, $\arg f'$ must undergo a specific jump at $z = z_k$, namely

$$\left[\arg f'(z) \right]_{z_k^-}^{z_k^+} = (1 - \alpha_k)\pi = \beta_k\pi. \quad (1.2)$$

The angle $\beta_k\pi$ is the **turning angle** at vertex k . We now identify a function f_k that is analytic in H^+ , satisfies (1.2), and otherwise has $\arg f_k$ constant on \mathbf{R} :

$$f_k = (z - z_k)^{-\beta_k}. \quad (1.3)$$

Any branch consistent with H^+ will work; to be definite, we pick the branch with $f_k(z) > 0$ if $z > z_k$ on \mathbf{R} . The action of f_k on the real line is sketched in Figure 1.2.

The preceding argument suggests the form

$$f'(z) = C \prod_{k=1}^{n-1} f_k(z)$$

for some constant C . We will prove the following fundamental theorem of Schwarz–Christoffel mapping in section 2.2.




Figure 1.2. Action of a term (1.3) in the SC product. In either case, the argument of the image jumps by $\beta_k \pi$ at z_k .

Theorem 1.1. *Let P be the interior of a polygon Γ having vertices w_1, \dots, w_n and interior angles $\alpha_1 \pi, \dots, \alpha_n \pi$ in counterclockwise order. Let f be any conformal map from the upper half-plane H^+ to P with $f(\infty) = w_n$. Then*

$$f(z) = A + C \int^z \prod_{k=1}^{n-1} (\zeta - z_k)^{\alpha_k - 1} d\zeta \quad (1.4)$$

for some complex constants A and C , where $w_k = f(z_k)$ for $k = 1, \dots, n - 1$.

The lower integration limit is left unspecified, as it affects only the value of A .

The formula also applies to polygons that have slits ($\alpha = 2$) or vertices at infinity ($-2 \leq \alpha \leq 0$). Indeed, arbitrary real exponents can meaningfully appear in (1.4), although the resulting region may overlap itself and not be bounded by a polygon in the usual sense of the term; see section 4.7.

Formula (1.4) can be adapted to maps from different regions (such as the unit disk), to exterior maps, to maps with branch points, to doubly connected regions, to regions bounded by circular arcs, and even to piecewise analytic boundaries. These and other variations are the subject of Chapter 4.

But there is a major difficulty we have not yet mentioned: without knowledge of the prevertices z_k , we cannot use (1.4) to compute values of the map. In view of how we arrived at (1.4), the image $f(\mathbf{R} \cup \{\infty\})$ of the extended real line will necessarily be *some* polygon whose interior angles match those of P , no matter what real values of z_k are used; that much is forced by the parameters α_k . (Here we are broadening the usual idea of “polygon” to allow




Figure 1.3. The effect of prevertices on side lengths. The region on the left is the “target,” whereas the region on the right illustrates the type of distortion that may occur if the prevertices are chosen incorrectly.

self-intersections.) The prevertices, however, influence the side lengths of $f(\mathbf{R} \cup \{\infty\})$, as illustrated in Figure 1.3. Determining their correct values for a given polygon is the Schwarz–Christoffel **parameter problem**, and its solution is the first step in using the SC formula.² In sections 2.3–2.5 we will consider some of the classical cases for which the parameter problem can be solved explicitly.

In the majority of practical problems, there is no analytic solution for the prevertices, which depend nonlinearly on the side lengths of Γ . Numerical computation is also usually needed to evaluate the integral in (1.4) and to invert the map. Thus, much of the potential of SC mapping went unrealized until computers became readily available in the last quarter of the twentieth century. Numerical issues are discussed in Chapter 3.

1.2 History

The roots of conformal mapping lie early in the nineteenth century. Gauss considered such problems in the 1820s. The Riemann mapping theorem was first stated in Riemann’s celebrated doctoral dissertation of 1851: any simply connected region in the complex plane can be conformally mapped onto any other, provided that neither is the entire plane.³ The Schwarz–Christoffel formula was discovered soon afterwards, independently by Christoffel in 1867 and Schwarz in 1869.

² Sometimes the constants A and C are included as unknowns in the parameter problem. However, they can be found easily once the prevertices are known, for they just describe a scaling, rotation, and translation of the image.

³ Riemann’s proof, based on the Dirichlet principle, was later pointed out by Weierstrass to be incomplete. Rigorous proofs did not appear until the work of Koebe, Osgood, Carathéodory, and Hilbert early in the twentieth century.

Elwin Bruno Christoffel (1829–1900) was born in the German town of Montjoie (now Monschau) and was studying mathematics in Berlin under Dirichlet and others when Riemann’s dissertation appeared.⁴ Christoffel completed his doctoral degree in 1856 and in 1862 succeeded Dedekind as a professor of mathematics at the Swiss Federal Institute of Technology in Zurich. It was in Zurich that he published the first paper on the Schwarz–Christoffel formula, with the Italian title, “Sul problema delle temperature stazionarie e la rappresentazione di una data superficie” [Chr67]. Christoffel’s motivation was the problem of heat conduction, which he approached by means of the Green’s function. This paper presented the discovery that, in the case of a polygonal domain, the Green’s function could be obtained via a conformal map from the half-plane, as in (1.4). In subsequent papers he extended these ideas to exteriors of polygons and to curved boundaries [Chr70a, Chr70b, Chr71].

Hermann Amandus Schwarz (1843–1921) grew up nearly a generation after Christoffel but also very much under the influence of Riemann. In the late 1860s he was living in Halle, where his discovery of the Schwarz–Christoffel formula apparently came independently of Christoffel’s. His three papers on the subject [Sch69a, Sch69b, Sch90] cover much of the same territory as Christoffel’s, including the generalizations to curved boundaries (section 4.11) and to circular polygons (section 4.10), but the emphasis is quite different—more numerical and more concerned with particular cases such as triangles in [Sch69b] and quadrilaterals in [Sch69a].⁵ Schwarz even published the world’s first plot of a Schwarz–Christoffel map, reproduced in Figure 1.4. Schwarz’s papers included his famous reflection principle: if an analytic function f , extended continuously to a straight or circular boundary arc, maps the boundary arc to another straight or circular arc, then f can be analytically continued across the arc by reflection.

In 1869 Christoffel moved briefly to the Gewerbeakademie in Berlin, and Schwarz succeeded him in Zurich. By this time the two were well aware of each other’s work; the phrase *Schwarz–Christoffel transformation* is now nearly universal (although the order of the names is reversed in some of the literature of the former Soviet Union).

In the 130 years since its discovery, the Schwarz–Christoffel formula has had an extensive impact in theoretical complex analysis, especially as a constructive

⁴ For extensive biographical information on Christoffel, the reader is referred to the sesquicentennial volume [BF81], particularly Pfluger’s paper therein on Christoffel’s work on the SC formula.

⁵ Schwarz also credits Weierstrass for proving the existence of a solution for the unknown parameters (which Schwarz proved for $n = 4$) in the general case.




Figure 1.4. Schwarz's 1869 plot of the conformal map of a square onto a disk, reproduced from [Sch69b].

tool for proving the Riemann mapping theorem and related results. Its practical implementation—the main subject of this book—lagged far behind. Schwarz himself was the first to point out the importance of the parameter problem (discussed in the preceding section). This problem limited practical use to simple special cases, until the invention of computers.

Algorithmic discussions of the computation of Schwarz–Christoffel maps to prescribed polygons appear in several books, including those of Kantorovich and Krylov [KK64] and Gaier [Gai64]. Algorithms and in some cases computer programs have also appeared in numerous technical articles over the years, but in most of the earlier cases the authors were unaware of each other's work, and the quality of the result was wanting. Crucial issues that were often neglected included efficient evaluation of the SC integral and the need to impose necessary ordering conditions on the prevertices while solving the parameter problem. The most generally applicable computer programs for the classical problem are those of Trefethen [Tre80] (SCPACK) and Driscoll [Dri96] (SC Toolbox). The former was developed around 1980, and the latter began development in 1993. Both have been widely disseminated in the public domain.

Here is a list, more or less chronological, of contributors to constructive SC mapping of whom we are aware.

Gauss (1820s): Idea of conformal mapping

Riemann (1851): Riemann mapping theorem

Christoffel [Chr67, Chr70a, Chr70b, Chr71]: Discovery of SC formula and variants

Schwarz [Sch69a, Sch69b, Sch90]: Discovery of SC formula and variants

Kantorovich & Krylov [KK64] (first published 1936)

- Polozkii** (1955)
- Filchakov** ([Fil61], 1968, 1969, 1975)
- Binns** [Bin61, Bin62, Bin64]
- Pisacane & Malvern** [PM63]
- Savenkov** (1963, 1964)
- Gaier** [Gai64]: Book on numerical conformal mapping
- Haeusler** (1966)
- Lawrenson & Gupta** [LG68]: Adaptive quadrature, equations solver for parameters
- Beigel** (1969)
- Hoffman** (1971, 1974)
- Gaier** [Gai72]: “Crowding” phenomenon
- Howe** [How73]
- Vecheslavov, Tolstobrova & Kokoulin** [VT73, VK74]: Doubly connected regions
- Foster & Anderson** [FA74, And75]
- Cherednichenko & Zhelankina** [CZ75]
- Squire** [Squ75]
- Meyer** [Mey79]: Comparison of algorithms
- Nicolaide** [Nic77]
- Prochazka** [HP78, Pro83]: FORTRAN package KABBAV
- Davis et al.** [Dav79, ADHE82, SD85]: Curved boundaries
- Hopkins & Roberts** [HR79]: Solution by Kufarev’s method
- Reppe** [Rep79]: First fully robust algorithm
- Binns, Rees & Kahan** [BRK79]
- Volkov** [Vol79, Vol87, Vol88]
- Trefethen** [Tre80, Tre84, Tre89, Tre93]: Robust algorithm, SCPACK, generalized parameter problems
- Brown** [Bro81]
- Tozoni** [Toz83]
- Hoekstra** (1983, [Hoe86]): Curved boundaries, doubly connected regions
- Sridhar & Davis** [SD85]: Strip maps
- Floryan & Zemach** [Flo85, Flo86, FZ87]: Channel flows, periodic regions
- Bjørstad & Grosse** [BG87]: Software for circular-arc polygons
- Dias, Elcrat & Trefethen** [ET86, DET87, DE92]: Free-streamline flows
- Däppen** [Döp87, Döp88]: Doubly connected regions
- Costamagna** ([Cos87, Cos01]): Applications in electricity and magnetism
- Howell & Trefethen** [How90, HT90, How93, How94]: Integration methods, elongated regions, circular-arc polygons
- Pearce** [Pea91]: Gearlike domains

Chaudhry [Cha92, CS92]: Piecewise smooth boundaries

Gutlyanskii & Zaidan [GZ94]: Kufarev's method

Driscoll [Dri96]: SC Toolbox for MATLAB

Hu [Hu95, Hu98]: Doubly connected regions (FORTRAN package DSCPACK)

Driscoll & Vavasis [DV98]: CRDT algorithm based on cross-ratios

Jamili (1999): Doubly connected regions

For more background information on conformal mapping in general and Schwarz–Christoffel mapping in particular, see [AF97, BF81, Hen74, Neh52, SL91, TD98, vS59, Wal64].