

## Score to identify the severity of adult patients with influenza A (H1N1) 2009 virus infection at hospital admission

A. Capelastegui · J. M. Quintana · A. Bilbao · P. P. España ·  
O. Garin · J. Alonso · J. Astray · R. Cantón · J. Castilla · A. Castro ·  
M. Delgado-Rodríguez · P. Godoy · F. González-Candelas ·  
V. Martín · J. M. Mayoral · T. Pumarola · S. Tamames · N. Soldevila ·  
M. Baricot · A. Domínguez · The CIBERESP Cases and Controls in  
Pandemic Influenza Working Group, Spain

Received: 27 January 2012 / Accepted: 21 March 2012 / Published online: 24 April 2012  
© Springer-Verlag 2012

**Abstract** The objective of this paper was to develop a prognostic index for severe complications among hospitalized patients with influenza A (H1N1) 2009 virus infection. We conducted a prospective observational cohort study of 618 inpatients with 2009 H1N1 virus infection admitted to 36 Spanish hospitals between July 2009 and February 2010. Risk factors evaluated included host-related factors and clinical data at admission. We developed a composite index of severe in-hospital complications (SIHC), which included: mortality, mechanical ventilation, septic shock, acute respiratory distress syndrome, and requirement for resuscitation

maneuvers. Six factors were independently associated with SIHC: age >45 years, male sex, number of comorbidities, pneumonia, dyspnea, and confusion. From the  $\beta$  parameter obtained in the multivariate model, a weight was assigned to each factor to compute the individual influenza risk score. The score shows an area under the receiver operating characteristic (ROC) curve of 0.77. The SIHC rate was 1.9 % in the low-risk group, 10.3 % in the intermediate-risk group, and 29.6 % in the high-risk group. The odds ratio for complications was 21.8 for the high-risk group compared with the low-risk group. This easy-to-score influenza A

A. Capelastegui · P. P. España  
Servicio de Respiratorio, Hospital Galdakao,  
Galdakao, Bizkaia, Spain

J. M. Quintana  
Unidad de Investigación, Hospital Galdakao,  
Galdakao, Bizkaia, Spain

A. Bilbao  
Unidad de Investigación, Hospital Basurto,  
Bilbao, Bizkaia, Spain

O. Garin · J. Alonso  
Institut Municipal de Investigació Mèdica,  
Barcelona, Spain

J. Astray  
Área de Epidemiología, Comunidad de Madrid,  
Madrid, Spain

R. Cantón  
Hospital Universitario Ramón y Cajal,  
Madrid, Spain

J. Castilla  
Instituto de Salud Pública de Navarra,  
Pamplona, Spain

A. Castro  
CIBER Enfermedades Respiratorias,  
Madrid, Spain

M. Delgado-Rodríguez  
Universidad de Jaen,  
Jaen, Spain

P. Godoy  
Departament de Salut, Generalitat of Catalonia,  
Barcelona, Spain

F. González-Candelas  
Centro Superior de Investigación en Salud Pública,  
Universitat de València,  
Valencia, Spain

V. Martín  
Instituto de Biomedicina, Universidad de León,  
Leon, Spain

J. M. Mayoral  
Servicio de Vigilancia de Andalucía,  
Sevilla, Spain

(H1N1) 2009 virus infection risk index accurately stratifies patients hospitalized for H1N1 virus infection into low-, intermediate-, and high-risk groups for SIHC.

### Abbreviations

ARDS	Acute respiratory distress syndrome
AUC	Area under the receiver operating characteristic curve
BMI	Body mass index
CI	Confidence intervals
ED	Emergency department
ICU	Intensive care unit
OR	Odds ratio
Ref.	Reference group
RT-PCR	Real-time polymerase chain reaction
SD	Standard deviation
SIHC	Severe in-hospital complications

### Introduction

The influenza A (H1N1) virus pandemic in 2009 shocked the health systems of many countries and raised great social alarm. Much of this alarm was due to the information emitted on the epidemiological and clinical characteristics of the infection: a high capacity for contagion and rapid spread [1, 2]. Unlike seasonal influenza, the infection affected mostly young

T. Pumarola  
Red Española de Investigación en Patología Infecciosa,  
Madrid, Spain

S. Tamames  
Dirección General de Salud Pública e Investigación,  
Desarrollo e Innovación, Junta de Castilla y León,  
Valladolid, Spain

J. M. Quintana · A. Bilbao · O. Garin · J. Alonso · J. Astray ·  
R. Cantón · J. Castilla · M. Delgado-Rodríguez · P. Godoy ·  
F. González-Candelas · V. Martín · N. Soldevila · M. Baricot ·  
A. Domínguez  
CIBER Epidemiología y Salud Pública,  
Barcelona, Spain

A. Domínguez  
Universitat de Barcelona,  
Barcelona, Spain

J. M. Quintana  
Universidad del País Vasco/EHU,  
Leioa, Spain

A. Capelastegui (✉)  
Service of Pneumology, Hospital de Galdakao-Usansolo,  
48960 Galdakao, Bizkaia, Spain  
e-mail: ALBERTO.CAPELASTEGUISAIZ@OSAKIDETZA.NET

people and fewer people aged >65 years [3]. Initially, it was regarded as a potentially severe disease, with a high mortality rate, above all in young people without comorbidities, who required intensive care [4–6]. Obesity and pregnancy were identified as risk factors [4, 7–9].

In this context, extraordinary measures were taken with respect to the care of hospitalized patients. For example, the isolation of patients together with thorough cleaning and protection of health care personnel, both steps aimed to limit contagion. Molecular microbiological determinations were used to provide an early diagnosis of the virus. The specific treatment of the virus was standardized for all infected patients admitted. Extraordinary resources were provided in order to increase intensive care unit (ICU) staff. However, we lacked specific tools for the early identification of adult patients hospitalized with a bad prognosis.

Although a large amount of information about the epidemiology and clinical management of influenza A (H1N1) 2009 virus infection has been obtained in a remarkably short period, a major gap exists in understanding disease severity and identifying at-risk populations. Most studies have focused on epidemiological aspects of the general population and on patients admitted to ICUs. Selecting the most seriously ill patients exclusively according to ICU admittance involves significant bias, due to the variability in ICU selection criteria [10, 11], especially if the initial alarm raised by the pandemic and the emphasis on critical care is taken into account. In addition, the information available on risk factors groups children and adults together. It is reasonable to suppose that serious risk factors could differ between children and adults and, in addition, the hospital care of children and adults is structured differently. No in-depth studies have been carried out to identify risk factors of severe evolution for adult patients hospitalized for influenza A (H1N1) 2009 virus infection in order to manage these patients.

The aim of this study was to identify risk factors present at admission in adult patients hospitalized due to influenza A (H1N1) 2009 virus infection during the period 2009–2010 that were independently associated with worse outcomes and, thereby, develop a prognostic influenza severity index.

### Patients and methods

#### Setting and study design

A multicenter matched case–control study was carried out in 36 hospitals and primary care centers from seven Spanish regions (Andalusia, Catalonia, Castile and Leon, Madrid, Navarre, the Basque Country, and Valencia). Cases and controls were recruited between July 2009 and February 2010. A case was defined as a patient admitted to hospital

for >24 h with influenza A (H1N1) virus infection laboratory-confirmed by real-time polymerase chain reaction (RT-PCR). Twenty-five patients aged  $\geq 18$  years were recruited from each of the 36 study hospitals, and were chosen by the systematic sampling of all patients admitted with laboratory-confirmed influenza A (H1N1) 2009 virus infection [12]. For the purpose of this article, only the prospective cohort of patients admitted to any of the participant hospitals were included (the cases) and the controls are not included. Therefore, for this study, based in our selection criteria, we included only adult cases ( $\geq 18$  years) who were hospitalized selected from the 36 hospitals studied. We excluded patients who had nosocomial infection, defined as pandemic virus infection in a patient that appears  $\geq 48$  h after admission for another cause. All information collected was treated as confidential, in strict observance of legislation on observational studies. The study was approved by the Ethics Committees of the hospitals involved. Written informed consent was obtained from all patients included in the study.

#### Data collection

A structured questionnaire was administered to patients by specifically trained personnel. We collected information on sociodemographic characteristics, pre-existing medical conditions, vaccinations, toxic habits, previous medications, exposure to social environments which could contribute to contagion, and the adoption of measures to prevent influenza. Variables on pre-existing medical conditions and vaccination were completed and verified by review of the medical records.

#### Possible predictive variables considered

The following demographic variables and pre-existing medical conditions were considered for this study: age, sex, previous hospital admission, history of pneumonia in the previous two years, obesity [body mass index (BMI) $\geq 30$ ], morbid obesity (BMI $\geq 40$ ), pregnancy in women aged 15–49 years, smoking, alcoholism, comorbidities (chronic obstructive pulmonary disease, asthma, other chronic respiratory diseases, cardiovascular disease, renal failure, diabetes, liver disease, HIV infection, disabling neurological disease, rheumatologic diseases, cancer, immunodeficiency, asplenia), vaccination, previous treatment, and clinical data at admission. For each vaccine, a case was considered to be vaccinated if they had received the vaccine at least 15 days before the onset of symptoms.

#### Outcomes

Severe in-hospital complications (SIHC) were the primary outcome of interest. This was a composite variable including: hospital mortality or requirement for mechanical

ventilation or the presence of septic shock or acute respiratory distress syndrome (ARDS) or a requirement for resuscitation maneuvers during hospitalization. Shock was defined as systolic blood pressure below 90 mmHg without anti-hypertensive drugs and the need for vasopressive agents. Mechanical ventilation was included when it was needed for at least 24 h and always after influenza A diagnosis in the Emergency Department (ED). We did not include patients who received non-invasive positive pressure ventilation.

#### Statistical analysis

Univariate and multivariate logistic regression models were then constructed to identify the statistical significance of each risk factor. The dependent variable was SIHC, and the independent variables were factors with a significance of  $p < 0.15$  in the univariate analysis. The odds ratio (OR) and 95 % confidence intervals (95 % CI) were calculated. The possible interaction between variables was also examined. The predictive accuracy of the model was determined by calculating the area under the receiver operating characteristic curve (AUC) for discrimination [13] and by comparing predicted and observed SIHC using the Hosmer–Lemeshow test for calibration [14]. Multilevel analysis with generalized estimated equations was carried out to determine whether the statistical significance of each predictive variable remained after adjusting for the Spanish regions.

To develop the influenza risk score, we first assigned a weight to each risk factor in relation to each  $\beta$  parameter based on the multivariate logistic regression model. Then, we added the weights of each of the risk factors presented by a patient. The predictive accuracy of the influenza risk score was determined by means of the AUC [13] and its calibration was tested by the Hosmer–Lemeshow test [14]. In addition, we attempted to validate the risk score by  $K$ -fold cross-validation [15, 16], which uses part of the available data to fit the model, and a different part to test it. That is, the model is validated in a random subsample which was not involved in the development of the model. This process is repeated sequentially for all partitions of the original sample. Thus, we split the data into  $K=10$  roughly equal-sized parts, we fitted the model with  $K-1$  parts of the data, and validated it by predicting the remaining  $k$ th part of the data. This procedure was repeated for each  $K$ th part, until the ten groups were all used in the validation, meaning that all cases were used once in the validation of the risk score [15].

Once the influenza risk score was developed, we created three categories (low, intermediate, and high risk) in relation to the predicted SIHC. The performance of the index score categories was studied using a logistic regression model and the AUC, and the Cochran–Armitage test for trend was used

**Table 1** Characteristics and outcomes of patients hospitalized with influenza A (H1N1) virus infection ( $N=618$ )

	<i>n</i> (%)
<b>Characteristics</b>	
Age, years, mean (SD)	48.60 (15.7)
Age, groups, years	
≤45 years	275 (44.5)
46–65 years	242 (39.2)
>65 years	101 (16.3)
Female	320 (51.8)
Hospitalized during the last year	133 (22.1)
Previous hospitalization by influenza A (H1N1) virus infection	3 (0.5)
Pneumonia in the last two years	93 (16.3)
Obesity	
No	414 (78)
$30 \leq \text{BMI} \leq 40$	96 (18.1)
$\text{BMI} \geq 40$	21 (4)
Pregnant women	48 (7.8)
Toxic habit	
Smoking	
No	308 (50.4)
Yes	176 (28.8)
Ex-smoker	127 (20.8)
Alcoholism	44 (7.1)
Drugs	13 (2.1)
<b>Comorbidities</b>	
Chronic obstructive pulmonary disease	72 (11.7)
Asthma	103 (16.8)
Others chronic pulmonary disease	74 (13.2)
Chronic respiratory insufficiency	54 (8.8)
Cardiovascular disease	89 (14.5)
Renal failure	40 (6.5)
Diabetes	89 (14.5)
Liver disease	33 (5.4)
AIDS	19 (3.1)
AIDS/symptomatic infection by HIV	21 (3.4)
Disabling neurological disease	20 (3.3)
Cognitive deterioration	8 (1.3)
Neuromuscular disease	8 (2.1)
Convulsive event	12 (3.1)
Rheumatologic disease	21 (3.5)
Neoplasia	62 (10.1)
Immunodeficiency	10 (1.6)
Asplenia	4 (0.7)
<b>No. of comorbidities</b>	
0	240 (38.8)
1	172 (27.8)
2	115 (18.6)
>2	91 (14.7)

**Table 1** (continued)

	<i>n</i> (%)
<b>Vaccination</b>	
Pandemic vaccine	10 (1.8)
Seasonal influenza vaccine in last year	174 (28.8)
23-valent pneumococcal vaccine in last 5 years	35 (6.3)
7-valent conjugated pneumococcal vaccine in last 5 years	10 (1.8)
<b>Previous treatment</b>	
Previous antibiotics	229 (37.1)
Length of antibiotic treatment, days, mean (SD)	1.14 (4.9)
Systemic corticosteroids in last 90 days	94 (18)
Inhaled corticosteroids	134 (21.8)
<b>Clinical status at admission</b>	
Multilobar and/or bilateral involvement	35 (5.7)
Pneumonia	174 (28.2)
Confusion	37 (6.2)
Fever	565 (92.8)
Dyspnea	404 (67.9)
<b>Outcomes</b>	
Hospital mortality	5 (1.3)
Shock	22 (5.4)
Mechanical ventilation	33 (5.3)
Acute respiratory distress syndrome	19 (4.9)
Resuscitation maneuvers	5 (1.3)

SD, standard derivation

Data are given as frequency (percentage) unless otherwise stated. Percentages exclude patients with missing data

to study the trend of the proportion of SIHC according to risk categories.

All effects were considered to be significant at  $p < 0.05$ , unless otherwise stated. All statistical analyses were performed using SAS for Windows statistical software, version 9.1 (SAS Institute, Inc., Carey, NC) and R<sup>®</sup> version 2.13.0 software.

## Results

A total of 618 patients hospitalized with influenza A (H1N1) 2009 virus infection were included. Patient characteristics and outcomes are shown in Table 1. The mean age was 48.6 years [standard deviation (SD), 15.7], and 44.5 % were aged <46 years. Almost 40 % had no comorbidity, while 14.7 % had  $\geq 3$  comorbidities. Of the 320 women included, 48 (15 %) were pregnant. A total of 117 patients (22 %) were obese ( $\text{BMI} \geq 30$ ), of which 21 (17.9 %) were morbidly obese ( $\text{BMI} \geq 40$ ). The SIHC rate was 9.9 % (61/618).

In the univariate analyses, several host-related factors, such as age, gender, smoking, comorbidities, and clinical

data at admission, were significantly associated with the likelihood of SIHC (Table 2). Obesity was not associated with SIHC (OR, 1.39; 95 % CI, 0.72–2.66;  $p=0.3275$ ), and no pregnant woman developed SIHC.

In the multivariate analysis, six factors were independently associated with SIHC: age, sex, number of comorbidities, and the presence of pneumonia, confusion, and dyspnea at admission (Table 3). The logistic model showed good discrimination, with an AUC value of 0.77. The model was also well calibrated, with a Hosmer–Lemeshow  $p$ -value of 0.9618. The statistical significance of each predictive variable remained after adjustment for Spanish regions.

Based on the multivariate logistic model, a weight was assigned to each risk factor in relation to each  $\beta$  parameter

(Table 3). By adding up the weights assigned to each predictive variable, an individual influenza risk score was given to each patient, ranging from 0 to 8, with a higher score corresponding to a higher likelihood of SIHC. The risk score was significantly associated with the likelihood of developing SIHC (OR, 2.11; 95 % CI, 1.68–2.64;  $p<0.0001$ ), and was well calibrated (Hosmer–Lemeshow  $o$ -value, 0.9603). The influenza risk score showed good discrimination (AUC, 0.76; 95 % CI, 0.71–0.82), in addition to the good results showed by the  $K$ -fold cross-validation, which had an AUC (95 % CI) of 0.74 (0.68–0.80) (Fig. 1).

Three risk categories were assigned using the influenza risk score (Table 4): low risk (0–2 points); intermediate risk (3–4 points); high risk (>4 points). The percentage of SIHC

**Table 2** Risk factors significantly associated with severe in-hospital complication in the univariate analyses ( $N=618$ )

Characteristics	Severe in-hospital complication, $n$ (%)	Odds ratio (95 % CI)	$p$ -value
Age, in groups			
≤45 years	15 (5.5)	Ref.	
46–65 years	32 (13.2)	2.6 (1.4–5)	0.003
>65 years	14 (13.9)	2.8 (1.3–6)	0.009
Sex			
Female	21 (6.6)	Ref.	
Male	40 (13.4)	2.2 (1.3–3.8)	0.005
Smoking			
No	20 (6.5)	Ref.	
Yes	22 (12.5)	2.1 (1.1–3.9)	0.03
Ex-smoker	17 (13.4)	2.2 (1.1–4.4)	0.02
No. of comorbidities			
0	19 (7.9)	Ref.	
1	15 (8.7)	1.1 (0.6–2.3)	0.77
2	9 (7.8)	1 (0.4–2.3)	0.98
>2	18 (19.8)	2.9 (1.4–5.8)	0.003
Clinical data at admission			
Multilobar and/or bilateral involvement			
No	54 (9.3)	Ref.	
Yes	7 (20)	2.5 (1–5.9)	0.04
Pneumonia			
No	37 (8.3)	Ref.	
Yes	24 (13.8)	1.8 (1–3)	0.04
Confusion			
No	49 (8.8)	Ref.	
Yes	10 (27)	3.9 (1.8–8.5)	0.0007
Fever			
No	9 (20.5)	Ref.	
Yes	49 (8.7)	0.4 (0.2–0.8)	0.01
Dyspnea			
No	6 (3.1)	Ref.	
Yes	53 (13.1)	4.7 (2–11)	0.0005

CI, confidence interval; Ref., reference group

Percentages exclude patients with missing data. Only factors with a significance of  $p<0.15$  in the univariate analysis are presented



**Table 3** Risk factors significantly associated with severe in-hospital complication in the multivariate analyses ( $N=618$ )

Risk factors	$\beta$ parameter	Odds ratio (95 % CI)	$p$ -value	Weight
Intercept	-4.91			
Age (years)				
>45 vs. $\leq 45$	0.76	2.1 (1.1–4.2)	0.03	1
Sex				
Male vs. female	0.88	2.4 (1.3–4.4)	0.004	1
No. of comorbidities				
$\geq 3$ vs. $< 3$	0.80	2.2 (1.2–4.4)	0.02	1
Pneumonia	0.65	1.9 (1.1–3.5)	0.03	1
Confusion	1.35	3.9 (1.6–9.1)	0.002	2
Dyspnea	1.50	4.5 (1.95–11)	0.0009	2
AUC		0.77		
Hosmer–Lemeshow $p$ -value <sup>a</sup>		0.9618		

CI, confidence interval;  $\beta$  parameter=estimated  $\beta$  coefficient; AUC, area under the receiver operating characteristic curve  
All risk factors were examined jointly  
<sup>a</sup>A significant value for the Hosmer–Lemeshow statistic indicates a significant deviation between predicted and observed outcomes


ranged from 1.9 % (95 % CI, 0.1–3.7) in patients classified as low risk to 29.6 % (95 % CI, 19.7–39.5) in patients classified as high risk (trend test,  $p<0.001$ ). The OR for the high-risk group was 21.8 compared with the low-risk group. The risk categories showed good discrimination, with an AUC value of 0.74.

## Discussion

This study was able to derive and validate a 2009 H1N1 virus infection influenza risk score with acceptable validity, discriminative ability, and generalizability, using data from a large cohort of inpatients from 36 geographically distinct Spanish hospitals. The risk score has various strengths. We developed a clinical prediction tool with six variables using information on predictive factors, including host-related factors and clinical conditions. The tool can be readily assessed and computed by physicians using information readily available in the ED.

Our results suggest that males aged  $>45$  years and those patients with more than two underlying chronic conditions have an increased risk. However, the greatest risk factors are associated with clinical factors at hospital admission, when a patient with dyspnea and confusion, and, for example, pneumonia, would have a probability of almost 30 % of developing SIHC. The main value of this predictive score is its ability to identify patients who need additional monitoring and more aggressive treatment after the first ED evaluation, either in the ICU, intermediate care units, or specialized regular wards, depending on the severity. The ED is the natural setting for the use of this severity score, but it could also be useful in outpatient services, as an adjunct to clinical judgment. It is easy and quick to apply. Patients with a score of  $>2$  points should be taken to hospital and patients with a score of  $>4$  points should be hospitalized on an emergency basis. However, in patients with a score of  $<3$ , the chances of severe complications would be slim and the patient could be managed at home.

**Fig. 1** Receiver operating characteristic curve of predicting severe in-hospital complication according to the individual influenza risk score (**a**) and according to the 10-fold cross-validation model (**b**). **a** Area under the curve (AUC) [95 % confidence interval (CI)], 0.76 (0.71–0.82); **b** AUC (95 % CI), 0.74 (0.68–0.80)



**Table 4** Validation of the influenza risk score: severe in-hospital complication by index score categories

Risk group (points)	No. with SIHC/ no. at risk	Percentage (95 % CI) <sup>a</sup>	Odds ratio (95 % CI)
Low risk (0–2)	4/211	1.9 (0.1–3.7)	Ref.
Intermediate risk (3–4)	30/291	10.3 (6.8–13.8)	6 (2.1–17.1)
High risk (>4)	24/81	29.6 (19.7–39.5)	21.8 (7.3–65.3)
AUC			0.74

AUC, area under the receiver operating characteristic curve; CI, confidence interval; Ref., reference group

The low-risk group was considered as the reference group

<sup>a</sup> $p < 0.001$  for the Cochran–Armitage test for trend

The strengths of the study include the relatively large cohort of patients recruited from different settings during the influenza A (H1N1) 2009 pandemic and the amount of clinical information collected. The main limitation of this study comes from its original design as a case–control study, where only 25 patients were recruited from each participating center and followed until discharge. Likewise, some laboratory parameters were not collected. Although we performed cross-validation, external validation of our score is still required.

Many of the predictors incorporated in our severity score have been established as risk factors for influenza A (H1N1)-associated complications in earlier studies. Comorbidities have been associated with an increased risk of complications both in seasonal influenza [17, 18] and in the influenza A (H1N1) 2009 pandemic [3, 7, 19–21]. We found that the risk of SIHC only increases in patients with >2 comorbidities. In contrast to seasonal influenza, where the greatest risk occurs in patients aged  $\geq 70$  years [18, 22], in both this study and other reports [19], severe outcomes occurred in patients aged >45 years.

We found that men had a higher risk of SIHC than women: similar results were reported by a Chinese study [7]. Another previous study showed that male sex was an independent risk factor for prolonged RT-PCR positivity in cases infected by influenza A (H1N1) virus [23]. Men have been found to be at a higher risk than women for death due to pneumonia [24, 25] and for sepsis [26]. Patterns of inflammation, coagulation, and fibrinolysis biomarkers in men may explain the reduced survival [27].

A preliminary report of 32 patients with influenza A (H1N1) 2009 virus infection hospitalized in a Spanish ICU showed that pneumonia was associated with a relatively high case–fatality rate [28]. The rate of patients hospitalized in wards and the ICU with pneumonia in the 2009 pandemic was higher [3, 4, 8, 19, 29]. We sought to confirm that pneumonia was an independent risk factor for SIHC.

Severe obesity has been identified as a risk factor for influenza A (H1N1) 2009 virus infection [4, 6–8]. However, in accordance with other studies [3, 30], we found that obesity was not associated with a higher risk of SIHC. In fact, obesity has not been identified as a risk factor for seasonal influenza complications [31, 32]. However, the prevalence of obesity in our series was similar to that of the general Spanish population [33], while a previous Spanish study found a high prevalence of obesity in patients hospitalized for 2009 H1N1 virus infection [34]. Obese patients may have been under-represented in our sample because the recruiting process of hospitalized cases collected 25 cases in each center during the pandemic period, but without trying to be representative of all cases seen in each hospital. Further investigation is needed in order to clarify the association between obesity and severe influenza.

In previous influenza epidemics and pandemics, pregnancy has been associated with an increased risk of severe disease [35, 36]. Likewise, recent reports suggest that there is an increased hospitalization rate and severity of illness in pregnant women infected by influenza A (H1N1) 2009 virus [8, 9, 19], while a Chinese study found that pregnancy was an independent risk factor associated with severe illness [7]. However, in our series, none of the pregnant women hospitalized for 2009 H1N1 virus infection developed SIHC, similar to the results of a previous Spanish study [34] and a Canadian study [20] that did not identify pregnancy as a risk factor for ICU admission or death. Preventive measures carried out in Spain, together with a fast diagnosis, early evaluation, and early antiviral treatment, may explain the relatively low rate of severity in pregnant women infected with the influenza A (H1N1) 2009 virus.

We did not use ICU admission as a criterion for determining SIHC because the decision to admit a patient to the ICU depends on individual clinical judgment and local hospital practices, differences that could account for much of the variability in ICU admission [10, 11]. On the other hand, the risk of death from influenza A infection is not the same as the need for inpatient care. We considered in-hospital death, mechanical ventilation, septic shock, ARDS, and resuscitation maneuvers as endpoints, given their more objective nature as variables [10]. This is a major strength of our study.

## Conclusions

Although the 2009 pandemic is over, gaining deeper knowledge of influenza A (H1N1) 2009 remains essential, as it will plan for the next, unavoidable pandemic and because, as in the 2010–2011 seasonal influenza in Spain [37], the influenza A (H1N1) virus might be one factor responsible for future seasonal influenza epidemics.

We identified risk factors for severe in-hospital complications (SIHC) in patients hospitalized due to influenza A (H1N1) virus infection and developed a clinical severity score that is very easy and simple to apply. The use of this score at diagnosis, or at diagnostic presumption, even with ambulatory patients, could assist decisions on care management. Early identification of the sickest patients could allow earlier interventions and, thus, potentially improve outcomes.

**The other members of the CIBERESP Cases and Controls in Pandemic Influenza Working Group are:** *Andalucía:* Ernestina Azor (Médico Centinela), Miguel Angel Bueno (Complejo Hospitalario de Jaén), Manuel Carnero (Hospital Virgen de la Victoria), Jerónimo Carrillo (Médico Centinela), Fernández-Crehuet Joaquín (Hospital Virgen de la Victoria), Víctor Fuentes (Hospital Costa del Sol), Virtudes Gallardo (Servicio de Epidemiología), M<sup>a</sup> Luisa Gómez (Complejo Hospitalario de Jaén), Reyes López (Hospital Infanta Elena de Huelva), José Ramón Maldonado (Hospital de Torrecárdenas), Marcial Mariscal (Complejo Hospitalario de Jaén), Belén Martínez (Complejo Hospitalario de Jaén), Jose M<sup>a</sup> Mayoral (Servicio de Epidemiología), Áurea Morillo (Hospital Virgen del Rocío), Rosa Moyano (Médico Centinela), José M<sup>a</sup> Navarro (Laboratorio de Referencia de Gripe), Juan Antonio Navarro (Médico Centinela), Salvador Oña (Hospital Carlos Haya), Esteban Pérez (Servicio de Epidemiología), M<sup>a</sup> José Pérez (Hospital Virgen de Valme), Mercedes Pérez (Laboratorio de Referencia de Gripe), Juan Pedro Quesada (Complejo Hospitalario de Jaén), Jorge del Diego Salas (Hospital Virgen de la Victoria), María Sillero (Complejo Hospitalario de Jaén), M<sup>a</sup> Carmen Ubago (Hospital Virgen de las Nieves), Manuel Vázquez (Médico Centinela), Francisco Zafra (Médico Centinela), Manuel Zarzuela (Hospital Puerta del Mar). *Comunidad Valenciana:* José Blanquer (Hospital Clínico de Valencia), María Morales (Hospital Doctor Peset), Red Centinela Sanitaria de la Comunidad Valenciana. *Castilla y León:* Demetrio Carriedo (Complejo Asistencial de León), José Javier Castrodeza (Dirección General de Salud Pública, Desarrollo e Innovación), Florentino Díez (Complejo Asistencial de León), Isabel Fernández (Complejo Asistencial de León), Silvia Fernandez (Complejo Asistencial de León), Ignacio Ledesma (Complejo Asistencial de León), Raul Ortiz de Lejarazu (Centro Nacional de Gripe de Valladolid), Juan Ortiz de Saracho (Hospital de El Bierzo), Alberto Pérez (Dirección General de Salud Pública, Desarrollo e Innovación), Ana Pueyo (Complejo Asistencial de Burgos), Pedro Redondo (Servicio Territorial de Sanidad y Bienestar Social de León), Juan Ortiz (Hospital del Bierzo), José Luis Viejo (Complejo Asistencial de Burgos). *Cataluña:* Alvar Agustí (Hospital Clínic de Barcelona), Ferrán Barbé (Hospital Arnau de Vilanova), Lluís Blanch (Hospital de Sabadell), Xavier Bonfill (Hospital de Sant Pau), Eva Borràs (Dirección General de Salud Pública), Carlos Bravo (Hospital Vall d'Hebrón), Francesc Calafell (Universitat Pompeu Fabra), Joan Cayla (Agencia de Salud Publica de Barcelona), Ignacio Garcia (Hospital Germans Trias i Pujol), Juan Jose Garcia (Hospital Sant Joan de Deu), Joaquim Gea (Hospital del Mar), Ned Hayes (Hospital Clínic, CRESIB), Juan Pablo Horcajada (Hospital del Mar), Joaquín López-Contreras (Hospital de Sant Pau), Ana Martínez (Dirección General de Salud Pública), Fernando Moraga (Hospital Vall d'Hebrón), Gemma Navarro (Hospital de Sabadell), Virginia Pomar (Hospital de Sant Pau), María Teresa Puig (Hospital de Sant Pau), Tomas Pumarola (Laboratorio de Referencia de Gripe), Antoni Rosell (Hospital de Bellvitge), Juan Ruiz (Hospital Germans Trias i Pujol), Marc Saez (Universidad de Girona), Núria Torner (Dirección General de Salud Pública), Antoni Torres (Hospital Clínic de Barcelona), Cecilia Tortajada (Agencia de Salud Publica de Barcelona), Antoni Trilla (Hospital Clínic de Barcelona), Ana Vilella (Hospital Clínic de

Barcelona). *Madrid:* Carlos Álvarez (Hospital 12 de Octubre), Fernando Baquero (Hospital Universitario Ramón y Cajal), Rafael Cantón (Hospital Universitario Ramón y Cajal), Esther Córdoba (Area de Epidemiología de la Comunidad de Madrid), María Enríquez (Hospital 12 de Octubre), Juan Carlos Galán (Hospital Universitario Ramón y Cajal), Juan García (Area de Epidemiología de la Comunidad de Madrid), Ricard Génova (Area de Epidemiología de la Comunidad de Madrid), Elisa Gil (Area de Epidemiología de la Comunidad de Madrid), Susana Jiménez (Area de Epidemiología de la Comunidad de Madrid), Josefina López (Area de Epidemiología de la Comunidad de Madrid), Fernando Martín (Area de Epidemiología de la Comunidad de Madrid), María Luisa Martínez (Area de Epidemiología de la Comunidad de Madrid), José Ramón Paño (Hospital Universitario La Paz), Francisco Pozo (Hospital 12 de octubre), Ana Robustillo (Hospital Universitario Ramón y Cajal), Elena Rodríguez (Area de Epidemiología de la Comunidad de Madrid), María Romero (Hospital Universitario La Paz), Silvia Sánchez (Area de Epidemiología de la Comunidad de Madrid), M<sup>a</sup> Angeles Valdeón (Hospital Universitario Ramón y Cajal), Cenegundis Valdés (Area de Epidemiología de la Comunidad de Madrid). *Navarra:* Patricia Fanlo (Hospital Virgen del Camino), Francisco Gil (Hospital Virgen del Camino), Antonia Martínez (Instituto de Salud Pública), Leyre Martínez (Instituto de Salud Pública), María Ruiz (Hospital Virgen del Camino). *Pais Vasco:* Urko Aguirre (Hospital Galdakao), José María Antóñana (Hospital de Cruces), Javier Arístegui (Hospital Basurto), Itziar Astigarraga (Hospital de Cruces), Amaia Bilbao (Fundación Vasca de Innovación e Investigación Sanitarias), Alberto Caspelastegui (Hospital Galdakao), Gustavo Cilla (Hospital Donostia), Antonio Escobar (Hospital Basurto), Pedro Pablo España (Hospital Galdakao), Felipe Esteban (Hospital Txagorritxu), Carmen Garaizar (Fundación Vasca de Innovación e Investigación Sanitarias), Susana García (Hospital Galdakao), Javier Korta (Hospital Donostia), José Luis Lobo (Hospital Txagorritxu), Emilio Pérez Trallero (Hospital Donostia), Jose Ignacio Pijoan (Hospital de Cruces), José María Quintana (Hospital Galdakao), Mikel Santiago (Hospital de Cruces), Cristina Sarasqueta (Hospital Donostia), Eva Tato (Hospital Txagorritxu).

**Sources of support** This study was supported by the Ministerio de Ciencia e Innovación Instituto de Salud Carlos III, Programa de Investigación Sobre Gripe A/H1N1 (grant GR09/0030), Agency for the Management of Grants for University Research (AGAUR grant number 2009/SGR 42), Department of Health of the Basque Country, Osakidetza, and the thematic networks—CIBERESP and CIBERES—of the Instituto de Salud Carlos III.

**Conflicts of interest** All authors declare that there are no conflicts of interest for any of the authors.

All authors had full access to the data.

## References

1. Miller MA, Viboud C, Balinska M et al (2009) The signature features of influenza pandemics—implications for policy. *N Engl J Med* 360:2595–2598
2. Writing Committee of the WHO Consultation on Clinical Aspects of Pandemic (H1N1) 2009 Influenza, Bautista E, Chotpitayasunondh T et al (2010). Clinical aspects of pandemic 2009 influenza A (H1N1) virus infection. *N Engl J Med* 362:1708–1719
3. Jain S, Kamimoto L, Bramley AM et al (2009) Hospitalized patients with 2009 H1N1 influenza in the United States, April–June 2009. *N Engl J Med* 361:1935–1944



4. Kumar A, Zarychanski R, Pinto R et al (2009) Critically ill patients with 2009 influenza A(H1N1) infection in Canada. *JAMA* 302:1872–1879
5. Perez-Padilla R, de la Rosa-Zamboni D, Ponce de Leon S et al (2009) Pneumonia and respiratory failure from swine-origin influenza A (H1N1) in Mexico. *N Engl J Med* 361:680–689
6. Domínguez-Cherit G, Lapinsky SE, Macías AE et al (2009) Critically ill patients with 2009 influenza A(H1N1) in Mexico. *JAMA* 302:1880–1887
7. Yu H, Feng Z, Uyeki TM et al (2011) Risk factors for severe illness with 2009 pandemic influenza A (H1N1) virus infection in China. *Clin Infect Dis* 52:457–465
8. ANZIC Influenza Investigators, Webb SA, Pettilä V et al (2009) Critical care services and 2009 H1N1 influenza in Australia and New Zealand. *N Engl J Med* 361:1925–1934
9. Louie JK, Acosta M, Jamieson DJ et al; California Pandemic (H1N1) Working Group (2010) Severe 2009 H1N1 influenza in pregnant and postpartum women in California. *N Engl J Med* 362:27–35
10. Angus DC, Marrie TJ, Obrosky DS et al (2002) Severe community-acquired pneumonia: use of intensive care services and evaluation of American and British Thoracic Society Diagnostic criteria. *Am J Respir Crit Care Med* 166:717–723
11. McQuillan P, Pilkington S, Allan A et al (1998) Confidential inquiry into quality of care before admission to intensive care. *BMJ* 316:1853–1858
12. Domínguez A, Alonso J, Astray J et al (2011) Risk factors of influenza (H1N1) 2009 hospitalization and effectiveness of pharmaceutical and nonpharmaceutical interventions in its prevention: a case-control study. *Rev Esp Salud Publica* 85:3–15
13. Hanley JA, McNeil BJ (1982) The meaning and use of the area under a receiver operating characteristic (ROC) curve. *Radiology* 143:29–36
14. Hosmer DW, Lemeshow S (1989) *Applied logistic regression*. Wiley Interscience, New York
15. Hastie T, Tibshirani R, Friedman J (2009) *The elements of statistical learning: data mining, inference, and prediction*. Springer, New York
16. Sxteyerberg EW (2009) *Clinical prediction models*. Springer, New York
17. Thompson WW, Shay DK, Weintraub E et al (2004) Influenza-associated hospitalizations in the United States. *JAMA* 292:1333–1340
18. Thompson WW, Shay DK, Weintraub E et al (2003) Mortality associated with influenza and respiratory syncytial virus in the United States. *JAMA* 289:179–186
19. Louie JK, Acosta M, Winter K et al (2009) Factors associated with death or hospitalization due to pandemic 2009 influenza A(H1N1) infection in California. *JAMA* 302:1896–1902
20. Campbell A, Rodin R, Kropp R et al (2010) Risk of severe outcomes among patients admitted to hospital with pandemic (H1N1) influenza. *CMAJ* 182:349–55
21. Donaldson LJ, Rutter PD, Ellis BM et al (2009) Mortality from pandemic A/H1N1 2009 influenza in England: public health surveillance study. *BMJ* 339:b5213
22. Schanzer DL, Tam TW, Langley JM et al (2007) Influenza-attributable deaths, Canada 1990–1999. *Epidemiol Infect* 135:1109–1116
23. Cao B, Li XW, Mao Y et al (2009) Clinical features of the initial cases of 2009 pandemic influenza A (H1N1) virus infection in China. *N Engl J Med* 361:2507–2517
24. Kaplan V, Angus DC, Griffin MF et al (2002) Hospitalized community-acquired pneumonia in the elderly: age- and sex-related patterns of care and outcome in the United States. *Am J Respir Crit Care Med* 165:766–772
25. Fine MJ, Auble TE, Yealy DM et al (1997) A prediction rule to identify low-risk patients with community-acquired pneumonia. *N Engl J Med* 336:243–250
26. Martin GS, Mannino DM, Eaton S et al (2003) The epidemiology of sepsis in the United States from 1979 through 2000. *N Engl J Med* 348:1546–1554
27. Reade MC, Yende S, D’Angelo G et al (2009) Differences in immune response may explain lower survival among older men with pneumonia. *Crit Care Med* 37:1655–1662
28. Rello J, Rodríguez A, Ibañez P et al (2009) Intensive care adult patients with severe respiratory failure caused by Influenza A (H1N1)v in Spain. *Crit Care* 13:R148
29. Vasoo S, Singh K, Trenholme GM (2010) Predicting need for hospitalization of patients with pandemic (H1N1) 2009, Chicago, Illinois, USA. *Emerg Infect Dis* 16:1594–1597
30. Díaz E, Rodríguez A, Martín-Loeches I et al (2011) Impact of obesity in patients infected with 2009 influenza A(H1N1). *Chest* 139:382–386
31. Fiore AE, Shay DK, Broder K et al; Centers for Disease Control and Prevention (2009) Prevention and control of seasonal influenza with vaccines: recommendations of the Advisory Committee on Immunization Practices (ACIP), 2009. *MMWR Recomm Rep* 58(RR-8):1–52
32. Ho YC, Wang JL, Wang JT et al (2009) Prognostic factors for fatal adult influenza pneumonia. *J Infect* 58:439–445
33. Rodríguez-Martín A, Novalbos Ruiz JP, Martínez Nieto JM et al (2009) Life-style factors associated with overweight and obesity among Spanish adults. *Nutr Hosp* 24:144–151
34. Santa-Olalla Peralta P, Cortes-García M, Vicente-Herrero M et al (2010) Risk factors for disease severity among hospitalised patients with 2009 pandemic influenza A (H1N1) in Spain, April–December 2009. *Euro Surveill* 15(38). pii: 19667
35. Harris JW (1919) Influenza occurring in pregnant women. *JAMA* 72:978–980
36. Rasmussen SA, Jamieson DJ, Bresee JS (2008) Pandemic influenza and pregnant women. *Emerg Infect Dis* 14:95–100
37. Instituto de Salud Carlos III. Weekly report of the Spanish Influenza Surveillance System. Week 06/2011. In Spanish. Available online at: <http://vgripe.isciii.es/gripe/documentos/20102011/boletines/grn062011.pdf>