



Research Journal of **Microbiology**

ISSN 1816-4935



Academic
Journals Inc.

www.academicjournals.com

Screening and Isolation of Novel Glutaminase Free L-asparaginase from Fungal Endophytes

¹A. Nagarajan, ²N. Thirunavukkarasu, ³T.S. Suryanarayanan and ¹Sathyanarayana N. Gummadi

¹Department of Biotechnology, Laboratory of Applied and Industrial Microbiology, Bhupat and Jyoti Mehta School of Biosciences, Indian Institute of Technology Madras, Chennai, 600036, India

²Department of Botany, Ramakrishna Mission Vivekananda College, Chennai, 600004, India

³Vivekanandha Institute of Tropical Mycology, Ramakrishna Mission Vidyapith, Chennai, 600004, India

Corresponding Authors: S.N. Gummadi, Department of Biotechnology, Indian Institute of Technology Madras, Chennai, 600036, India Tel: 914422574114 Fax: 914422574102 and T.S. Suryanarayanan, Vivekanandha Institute of Tropical Mycology, Ramakrishna Mission Vidyapith, Chennai, 600004, India

ABSTRACT

L-asparaginase (E.C.3.5.1.1) has been commonly used for the treatment of acute lymphoblastic leukemia in adults and children. It is also used in food industry to reduce acrylamide formation during the preparation of fried food items containing starch at high temperatures. Several microorganisms from the diverse group of bacteria and yeast were reported to be used for L-asparaginase production however, many of the strains also coproduce L-glutaminase which is highly undesirable as it results in cellular stress and neurotoxicity. Thus identification of new sources for the production of glutaminase free L-asparaginase needs to be explored. In this study, we screened endophytic fungi isolated from trees of moist deciduous and semi evergreen forests of the Western Ghats and plants growing in Rono Hills, Arunachal Pradesh, India for the production of glutaminase free L-asparaginase. Using a simple agar plate assay, we found that 33 strains were positive for the L-asparaginase activity among which 19 strains showed glutaminase free L-asparaginase activity. Our results show that: *Alternaria* sp. endophytic in the leaf of *Withania somnifera* and growing in the moist deciduous forest of the Western Ghats showed maximum enzyme activity. Optimization of process parameters reveal that maximum L-asparaginase production was observed at 96 h of fermentation and high concentration of glucose in the medium as the carbon source inhibited enzyme production in *Alternaria* sp. This is the first report on production of glutaminase free L-asparaginase by fungal endophyte *Alternaria* sp.

Key words: L-asparaginase, endophytes, production

INTRODUCTION

L-asparaginase (E.C.3.5.1.1) is an enzyme which is present in diverse types of microorganisms, plants and animals (Mohan Kumar *et al.*, 2013). L-asparaginase has been commercially used to prevent acrylamide formation in fried food products (Pedreschi *et al.*, 2008; Mohan Kumar *et al.*, 2013). L-asparaginase from bacterial sources has been reported to be used especially as a therapeutic agent in the treatment of acute lymphoblastic leukemia in children (Muller and Boos, 1998; Pieters *et al.*, 2011). This enzyme depletes malignant cells by preventing the formation of essential growth factors for tumor development. L-asparaginase production by various

microorganisms including, *Erwinia carotovora* (Maladkar *et al.*, 1993), *Escherichia coli* (Cedar and Schwartz, 1968; Wei and Liu, 1998), *Aspergillus* sp. (Sarquis *et al.*, 2004) and also from marine derived fungal endophytes such as *Fusarium* sp., *Phomopsis* sp., *Trichoderma* sp. and *Sargasam wightii* (Thirunavukkarasu *et al.*, 2011) has been studied extensively. A recent report revealed that clinical *E. coli* L-asparaginase is an important drug in treatment of patients with Acute Lymphoblastic Leukemia (ALL) (Rytting, 2012). Though it is a strong candidate for treatment of ALL, several secondary complications hinder the use of this enzyme in tumor treatment including side effects such as breathing problems, neural disorders, pancreatitis and also affects reproductive fertility (Duval *et al.*, 2002). *Pyrococcus furiosus* and its mutants MTCC 5580-5582 produces L-asparaginase which was found to be highly stable (Kundu *et al.*, 2013). *Aspergillus tamari* and *Aspergillus terreus* have been used for production of L-asparaginase but the yields were very low due to the presence of glutaminase and urease (Sarquis *et al.*, 2004). *Aspergillus niger* and *Aspergillus oryzae* have been preferred because of their high yields (Laan *et al.*, 2008; Eisele *et al.*, 2011). L-asparaginase from *Bacillus licheniformis* (RAM 8) with low glutaminase levels was optimized for the removal of acrylamide that is formed in baked and fried food (Mahajan *et al.*, 2012). Although the production of L-asparaginase has been reported from various organisms, the major constraint in commercialization is L-glutaminase coproduction with L-asparaginase which leads to toxicity. Very few reports are available on glutaminase free L-asparaginase produced by microorganisms. An intercellular glutaminase free L-asparaginase from few microorganisms such as *Pectobacterium carotovorum* MTCC 1428 (Kumar *et al.*, 2011), *Vibrio succinogenes* (Distasio *et al.*, 1982), *Pseudomonas stutzeri* (Manna *et al.*, 1995), *Pyrococcus furiosus* and its mutants MTCC 5580-5582 (Kundu *et al.*, 2013) has been reported. Endophytic fungi reside inside living tissues of all groups of plants without causing any disease in them. These fungi are known to produce novel bioactive compounds (Gunatilaka, 2006; Weber, 2009; Suryanarayanan *et al.*, 2009; Debbab *et al.*, 2011). These fungi have also been studied for their use in biological control of plant diseases (Vega, 2008; Rocha *et al.*, 2011). Our recent studies and those by others indicate that endophytic fungi are also a source of novel industrial enzymes (Nagaraju *et al.*, 2009; Rajulu *et al.*, 2011; Suryanarayanan *et al.*, 2012; Robl *et al.*, 2013). In an earlier study, it was reported that many of the endophytic fungi associated with marine algae produce L-asparaginase (Thirunavukkarasu *et al.*, 2011). In this study, we screened endophytic fungi isolated from trees of moist deciduous and semi evergreen forests of the Western Ghats and plants growing in Rono Hills, Arunachal Pradesh, India for the production of glutaminase free L-asparaginase.

MATERIALS AND METHODS

L-asparagine and L-glutamine was procured from Sigma-Aldrich. All chemicals used in the production medium and protein analysis were of analytical grade and of the highest purity from SRL, Hi-Media and Merck, India.

Isolation of endophytes: From each plant species, mature healthy leaves or outer bark were collected, washed thoroughly in running tap water and cut into segments (0.5 cm²) and surface sterilized by serial washing in 70% ethanol and sodium hypochlorite (Suryanarayanan *et al.*, 1998). They were then plated on antibiotic-amended Potato Dextrose Agar (PDA) medium in petri dishes, sealed using Parafilm™ and incubated in a light chamber with 12 h light: 12 h dark cycle for 28 days at 26±1°C (Bills and Polishook, 1992). The tissue segments were observed periodically

and the endophytic fungi growing out of them were cultured in PDA slants and were identified using standard taxonomic keys (Ellis, 1971, 1976; Subramanian, 1971; Barnett and Hunter, 1972; Von Arx, 1974; Sutton, 1980; Onions *et al.*, 1981; Ellis and Ellis, 1987).

Plate assay for L-asparaginase: A modified Czapek Dox (CD) medium (glucose 2 g L⁻¹, L-asparagine 10 g L⁻¹ or L-glutamine 10 g L⁻¹, KH₂PO₄ 1.52 g L⁻¹, KCl 0.52 g L⁻¹, MgSO₄.7H₂O 0.52 g L⁻¹, FeSO₄.7H₂O 0.01 g L⁻¹ and agar 20 g L⁻¹) was used for plate assay (Thirunavukkarasu *et al.*, 2011). A 2.5% stock solution of phenol red was prepared in ethanol (pH 6.2) and 3 mL of this was added to 1000 mL of Czapek Dox medium. A mycelial plug (5 mm diameter) cut from the growing margin of the colony of an endophyte or phelophyte was placed in a petri dish containing 20 mL of this medium. After 72 h of incubation at 26±1°C, the appearance of a pink zone around the fungal colony in an otherwise yellow medium indicated L-asparaginase activity (Gulati *et al.*, 1997). In order to study the effect of glucose on growth and enzyme production, fungal endophytes were inoculated on modified CD media with and without 2% (w/v) glucose. Fungal growth and the enzyme zone were measured as described earlier.

Shake flask experiments for enzyme production: Fungal endophyte strains producing glutaminase free L-asparaginase were maintained and sub-cultured every 7 days in PDA medium and incubated at room temperature. The production of L-asparaginase was optimized in modified CD medium (Kumar *et al.*, 2010). Modified CD medium contained glucose 2.0 g L⁻¹, L-asparagine 10.0 g L⁻¹ or L-glutamine 10.0 g L⁻¹, KH₂PO₄ 1.52 g L⁻¹, KCl 0.52 g L⁻¹, MgSO₄.7H₂O 0.52 g L⁻¹, (NH₄)₂SO₄ 5.0 g L⁻¹, FeSO₄.7H₂O 0.01 g L⁻¹ and the initial pH was set to 6.2 (Thirunavukkarasu *et al.*, 2011; Kumar *et al.*, 2011). Seven days old fungal endophyte mycelium plug was cut (~5 mm diameter) using cork borer and inoculated in 50 mL modified CD medium in a 250 mL Erlenmeyer flask. The culture flask was incubated at 30°C and 180 rpm in an orbital shaker for 120 h. Samples were withdrawn at regular intervals of 24 h and used for the measurement of L-asparaginase production. In order to study the effect of carbon and nitrogen source, varying concentrations of glucose (1.0, 2.0 and 5.0 g L⁻¹) and L-asparagine (5.0, 10.0 and 20.0 g L⁻¹) was incorporated in CD medium and growth was carried out for five days. The samples were collected for every 24 h and assayed as mentioned above.

Enzyme assays: The culture filtrates were centrifuged at 10,000×g for 10 min at 4°C and their protein contents were determined by Lowry method (Lowry *et al.*, 1951). The enzyme assay reaction mixture contained 0.5 mL of 0.1 M potassium phosphate buffer pH 8.0, 0.1 mL of 40 mM L-asparagine and 100 µg of extracellular protein (culture filtrate of each endophyte) and final volume was made upto 2.0 mL with distilled water. Reaction mixture was incubated at 37°C in stirred water bath for 30 min and enzymatic reaction was stopped by adding 0.5 mL of 1.5 M Trichloro Acetic Acid (TCA). The reaction mixture was centrifuged at 10,000×g for 5 min at 4°C to remove precipitates. The ammonia released was determined using colorimetric method by adding 200 µL Nessler's reagent into a tube containing 100 µL supernatant and 1700 µL distilled water. This mixture was vortexed and incubated at room temperature for 20 min and absorbance was measured at 436 nm (Kumar *et al.*, 2010). Ammonium sulphate was used to plot the standard curve to determine ammonia released in the reaction mixture (Mahajan *et al.*, 2012). One unit (IU) of L-asparaginase activity is determined as the amount of enzyme needed to liberate 1 µmol of ammonia per min at 37°C when L-asparagine is used as substrate (Imada *et al.*, 1973; Kumar *et al.*, 2010). Glutaminase activity was measured by direct nesslerization as described by

Mashburn and Wriston (1964), using L-glutamine as the substrate. One Unit (IU) of L-glutaminase activity is calculated as the amount of enzyme needed to liberate 1 μ mol of ammonia in presence of L-glutamine per min at 37°C. Specific activity of protein is defined as enzyme activity per milligram of protein used in the assay.

Statistical analysis: All the experiments are performed at least four times and one way analysis of variance (ANOVA) was performed in Minitab 16. A $p < 0.05$ was considered significant.

RESULTS AND DISCUSSION

Screening of fungal endophytes for glutaminase free L-asparaginase: Thirty three different fungal endophyte strains were collected from plant hosts in Western Ghats and Rono Hills in India. The samples were critically scrutinized for the production of L-asparaginase by plate assay method. The plate assay method involves the use of Czapeck Dox medium with L-asparagine as a nitrogen source and phenol red as a pH indicator to monitor the changes in pH. If the organism produces L-asparaginase, it could grow on a normal plate containing L-asparagine and degrades it into aspartic acid and ammonia. Release of ammonia in the medium changes the pH towards alkaline which is indicated by a visible change in medium colour to pink. This pink zone was initially used to screen the fungal endophytes (Gulati *et al.*, 1997; Thirunavukkarasu *et al.*, 2011). For example *C. lunata* showed pink zone only when grown in plate containing L-asparagine and not in plate containing L-glutamine (Fig. 1a), suggesting that the strain produces glutaminase free L-asparaginase. While another strain, *L. theobromae* showed pink zone when grown both on L-asparagine as well as in L-glutamine plate implying that this strain produces both the enzymes (Fig. 1b).

In order to screen the endophytes that produces glutaminase free L-asparaginase, all 33 fungal endophytes were allowed to grow on the modified CD medium containing L-glutamine as a nitrogen source and the colour change was monitored. Endophytes that produced pink colour zone on L-glutamine medium containing plates (Fig. 1b) were eliminated. Of the fungi studied, only a species of *Periconia* and *S. intermedia* lacked asparaginase activity (Table 1). Twelve fungi, viz., *Acremonium* sp., *Alternaria* sp. 2, *Aspergillus* sp., *Botrytis* sp., *C. cladosporioides*, *Corynespora* sp., *Cylindrocladium* sp., *Fusarium* sp., *L. theobromae*, *Pestalotiopsis* sp. 1 and 2 and *Sordaria* sp., were positive for both the enzymes (Table 1). Some strains producing only glutaminase free L-asparaginase (Fig. 1a) were selected and plate assay was performed. Nineteen fungal endophytes were shown to be producing glutaminase free L-asparaginase which were further examined for the highest activity of L-asparaginase production by measuring the colony diameter (mm) and diameter of the pink colour zone (mm) which indicates the enzyme production quantitatively (Fig. 2).

Effect of glucose on colony growth and enzyme activity by plate assay: Previous reports showed that concomitant addition of L-asparagine and glucose inhibits growth and activity (Geckil *et al.*, 2006). Few reports showed that presence of glucose in media marginally favours activity and is a good choice compared to other substrates because of its low cost, making it an economically viable process (Kumar *et al.*, 2010). The existing reports are contradictory regarding the effect of glucose and nitrogen. Hence, to check the effect of glucose on growth and




Fig. 1(a-c): Plate assay for screening fungal endophytes producing L-asparaginase, (a) *Curvularia lunata* grown on plates containing L-asparagine and L-glutamine showing pink zone only when L-asparagine was used as the substrate, (b) *Lasiodiplodia theobromae* showing pink zone on both plates containing L-asparagine and L-glutamine as substrate and (c) Glutaminase free L-asparaginase showing pink colour zones around the fungal endophyte colony

L-asparaginase production, plate assay was performed in the presence and absence of glucose. Among the 19 fungal endophytes screened and isolated that exhibited glutaminase free L-asparaginase activity, 10 strains which showed relatively higher enzyme activity zone were taken and grown in the presence and absence of media containing 2 g L^{-1} glucose. Results showed that the presence of glucose inhibited colony growth and enzyme activity in most of the endophytes (Fig. 3a and b). *Curvularia lunata*, *Phomopsis* sp. and *Dreschlera* sp. showed an increase in both colony growth and enzyme activity in the presence of glucose. Presence of glucose for *Corynespora* sp. did not have much effect on the colony growth but the enzyme activity zone was drastically reduced (Fig. 3b) (Narayana *et al.*, 2008). As a widely used carbon source, effect of glucose on L- asparaginase activity has been well studied over the years. Glucose is reported

Table 1: Isolated novel fungal endophytes screened for glutaminase free L-asparaginase production (plate assay)

Endophytes	Host	Organ	Location	L-asparaginase	L-glutaminase
<i>Acremonium</i> sp.	<i>Lagerstroemia microcarpa</i>	Leaf	Moist deciduous	+	+
<i>Alternaria alternata</i>	<i>Macaranga peltata</i>	Leaf	Evergreen	+	-
<i>Alternaria brassicicola</i>	<i>Litsea floribunda</i>	Leaf	Evergreen	+	-
<i>Alternaria</i> sp.	<i>Terminalia paniculata</i>	Leaf	Moist deciduous	+	-
<i>Alternaria</i> sp.	<i>Syzygium densiflorum</i>	Leaf	Evergreen	+	+
<i>Alternaria</i> sp. 1	<i>Withania somnifera</i>	Leaf	Moist deciduous	+	-
<i>Alternaria</i> sp. 2	<i>Withania somnifera</i>	Leaf	Moist deciduous	+	-
<i>Arthriniump.</i>	<i>Elaeocarpus serratus</i>	Bark	Evergreen	+	-
<i>Aspergillus</i> sp.	<i>Cassia fistula</i>	Leaf	Moist deciduous	+	+
<i>Botrytis</i> sp.	<i>Daphniphyllum neilgherrense</i>	Bark	Evergreen	+	+
<i>Capnodium</i> sp.	<i>Ligustrum roxburghii</i>	Leaf	Evergreen	+	-
<i>Chaetomium</i>	<i>Phoebe lanceolata</i>	Bark	Evergreen	+	-
<i>Cladosporium cladospriosides</i>	<i>Diospyros montana</i>	Leaf	Moist deciduous	+	+
<i>Corynespora</i> sp.	<i>Withania somnifera</i>	Leaf	Moist deciduous	+	-
<i>Corynespora</i> sp.	<i>Eunonymus undulates</i>	Leaf	Evergreen	+	-
<i>Corynespora</i> sp. 2	<i>Ligustrum roxburghii</i>	Leaf	Evergreen	+	+
<i>Curvularia lunata</i>	<i>Litsea floribunda</i>	Leaf	Evergreen	+	-
<i>Cylindrocladium</i>	<i>Cycas</i> sp.	Leaf	Ronohills	+	+
<i>Dreschlera</i> sp.	<i>Syzygium densiflorum</i>	Leaf	Evergreen	+	-
<i>Fusarium</i> sp.	<i>Calicarba arborea</i>	Leaf	Ronohills	+	+
<i>Humicola</i> sp.	<i>Eurya nitida</i>	Leaf	Evergreen	+	-
<i>Lasiodiplodia theobromae</i>	<i>Litsea floribunda</i>	Leaf	Evergreen	+	+
<i>Lasiodiplodia theobromae</i>	<i>Cordia oblique</i>	Leaf	Moist deciduous	+	-
<i>Periconia</i> sp.	<i>Elaeocarpus serratus</i>	Bark	Evergreen	-	-
<i>Pestalotiopsis</i> sp.	<i>Terminalia bellerica</i>	Leaf	Moist deciduous	+	+
<i>Pestalotiopsis</i> sp.	<i>Litsea cubeba</i>	Leaf	Ronohills	+	+
<i>Phomopsis</i> sp.	<i>Diospyros montana</i>	Leaf	Moist deciduous	+	-
<i>Phomopsis</i> sp.	<i>Careya arborea</i>	Leaf	Moist deciduous	+	-
<i>Phyllosticta</i> sp.	<i>Eurya nitida</i>	Leaf	Evergreen	+	-
<i>Pithomyces</i> sp.	<i>Eurya nitida</i>	Bark	Evergreen	+	-
<i>Sordaria</i> sp.	<i>Cordia wallichii</i>	Bark	Moist deciduous	+	+
<i>Sporormiella intermedia</i>	<i>Butea monosperma</i>	Leaf	Moist deciduous	-	-
<i>Torulomyces</i> sp.	<i>Turpinia nepalensis</i>	Bark	Evergreen	+	-

to induce the production of L-asparaginase in several microbes including certain strains of *Serratia marcescens* (Sukumaran *et al.*, 1979) and *Bacillus circulans* (Hymavathi *et al.*, 2010). Glucose acts as a repressor and inhibits L-asparaginase production in a few organisms such as *Streptomyces albidoflavus* (Narayana *et al.*, 2008) and *Serratia marcescens* (Heinemann and Howard, 1969). It has been reported that when 0.4% of glucose was used, L-asparaginase production was not inhibited but by further increase in the glucose concentration upto 1%, pH of the medium changed from alkaline (pH 7.5-8.0) to acidic (pH 5.2-6.9) and L-asparaginase production was inhibited (Heinemann and Howard, 1969).

L-asparaginase production by shake flask: L-asparaginase producing microorganisms either produce this enzyme constitutively or upon induction. The physico-chemical conditions for L-asparaginase production differ among various microorganisms (Savitri and Azmi, 2003). In this study, endophytes produced L-asparaginase when grown only in the presence of L-asparagine.

L-asparaginase produced was determined by colorimetric estimation of released ammonia at 436 nm. Ten fungal endophytes, which exhibited highest activity in plate assay were taken for production studies in shake flask. Among the 10 fungal endophytes, WS/*Alternaria* sp. 2 exhibited the highest activity of 1.17 ± 0.04 U mg^{-1} at 96 h, followed by TPM/*Alternaria* sp. 0.5 ± 0.06 U mg^{-1}




Fig. 2: Colony growth and enzyme activity for fungal endophytes. Plate assay method was used to calculate the colony growth and enzyme activity by measuring the colony zone diameter (mm) and enzyme activity zone diameter (mm) in the plates. Experiment was repeated thrice and the error bars represent standard error of mean




Fig. 3(a-b): Continue




Fig. 3(a-b): Effect of glucose on (a) Colony growth and (b) Enzyme activity by plate assay in L-asparaginase producing fungal endophytes. There is a significant decline in L-asparaginase activity of DIOM/*Phomopsis* sp., WS/*Alternaria* sp., 2 and EU/*Corynespora* sp. in the presence of glucose ($p < 0.05$, by ANOVA). In WS/*Alternaria* sp. 1, *Alternaria alternata* and WS/*Alternaria brassicicola*, the mean colony growth decreases significantly ($p < 0.05$, by ANOVA) in the presence of glucose. Experiment was repeated thrice and the error bars represent standard error of mean

(Fig. 4). Though other class of microorganisms has been reported to produce L-asparaginase activity with higher and comparable values, this is the first report to show glutaminase free L-asparaginase activity by fungal endophytes.

Effect of carbon source on L-asparaginase production: Glucose is commonly used as the primary carbon source for most of the microorganisms producing primary and secondary metabolites. Since we have already confirmed that glucose play a key role in enzyme activity by plate assay method, it is essential to optimize the glucose concentration for maximizing the enzyme production in shake flasks. WS/*Alternaria* sp. 2, was grown in CD media containing 1, 2 and 5 g L⁻¹ of glucose, respectively and samples were collected for every 24 till 120 h (Fig. 5a). Our results showed that culture grown in 1 g L⁻¹ of glucose exhibited a maximum activity of 2.1 U mg⁻¹ at 96h, whereas culture grown in 2 and 5 g L⁻¹ for 96 h showed 0.34 and 0.31 U mg⁻¹, respectively (Fig. 5b). Thus it is confirmed that 1 g L⁻¹ was optimal for L-asparaginase production in WS/*Alternaria* sp. 2. The enzyme production beyond 96 h decreases drastically which could be due to several reasons. It was observed that the production of the L-asparaginase was significantly reduced in the presence of glucose at higher concentrations, where it act as repressor for L-asparaginase in *Enterobacter aerogenes* (Mukherjee *et al.*, 2000) and similar trend was




Fig. 4: Specific activity of glutaminase free L-asparaginase in shake flask experiments. *WS/Alternaria* sp. 2 shows maximum specific activity. Experiment was repeated thrice and the error bars represent standard error of mean




Fig. 5(a-b): Effect of glucose on specific activity of glutaminase free L-asparaginase produced in shake flask method, (a) *WS/Alternaria* sp. was inoculated in the modified CD medium containing various concentration of glucose and (b) Specific activity at varying glucose concentrations




Fig. 6: Effect of L-asparagine on specific activity of glutaminase free L- asparaginase in shake flask at varying concentrations of L-asparagine 5, 10 and 20 g L⁻¹. Experiment was repeated thrice and the error bars represent standard error of mean

observed in *Fusarium* sp. (Thirunavukkarasu *et al.*, 2011), *Serrtia marcescens* (Heinemann *et al.*, 1970) and *Erwinia aroideae* (Liu and Zajic, 1972).

Effect of nitrogen concentration on L-asparaginase production: To study the effect of nitrogen source concentration on L-asparaginase production by *WS/Alternaria* sp. 2, different concentrations of L-asparagine (5, 10 and 20 g L⁻¹) was supplemented in the media. Culture grown in 10 g L⁻¹ of L-asparagine showed maximum activity of 1.8 U mg⁻¹ at 120 h, followed by 1.3 and 0.9 U mg⁻¹ for cultures grown on 5 and 20 g L⁻¹, respectively (Fig. 6). The time dependent increase in the enzyme activity shows that L-asparaginase expression increases in the presence of L-asparagine, suggesting that it is nitrogen regulated and inducible as observed in other microbes (Sarquis *et al.*, 2004). Effect of varying concentration (0.5-2.0%) of L-asparagine as a sole nitrogen source on L-asparaginase production was studied in *Streptomyces* ABR2 and the optimum concentration was determined to be 1.0% (Mostafa and Salama, 1979; Sudhir *et al.*, 2012). Another report showed that 1% (w/v) of L-asparagine in the media exhibited maximum production of L-asparaginase by *Aspergillus terreus* MTCC 1782. The activity of L-asparaginase decreases with increase in concentration of L-asparagine above 1% which might be due to substrate inhibition (Shaffer *et al.*, 1988; Baskar and Renganathan, 2011).

CONCLUSION

In this study, several endophytes have been identified that are capable of producing glutaminase free L-asparaginase. Among all the screened endophytes, *WS/Alternaria* sp. 2, produced maximum glutaminase free L-asparaginase. The optimal concentrations of glucose and L-asparagine were found to be 10 and 1 g L⁻¹, respectively. Under these conditions, the strain produced a maximum specific activity of 1.65 U mg⁻¹. This is the first report on the production of glutaminase free L-asparaginase by endophytic fungi *Alternaria* sp. Although, preliminary, the

study gains importance as it identifies a novel eukaryotic (and hence possibly more human-compatible than a bacterial source of the enzyme) source of such a desirable enzyme for therapeutic uses. Further media components screening and optimization in shake flasks and bioreactors are needed to enhance the productivity.

REFERENCES

- Barnett, H.L. and B.B. Hunter, 1972. Illustrated Genera of Imperfect Fungi. 3rd Edn., Burgess Publishing Co., Minneapolis, MN., USA., Pages: 241.
- Baskar, G. and S. Renganathan, 2011. Optimization of media components and operating conditions for exogenous production of fungal L-asparaginase. *Chiang Mai J. Sci.*, 38: 270-279.
- Bills, G.F. and J.D. Polishook, 1992. Recovery of endophytic fungi from *Chamaecyparis hyoides*. *Sydowia*, 44: 1-12.
- Cedar, H. and J.H. Schwartz, 1968. Production of L-asparaginase II by *Escherichia coli*. *J. Bacteriol.*, 96: 2043-2048.
- Debbab, A., A.H. Aly and P. Proksch, 2011. Bioactive secondary metabolites from endophytes and associated marine derived fungi. *Fungal Diversity*, 49: 1-12.
- Distasio, J.A., A.M. Salazar, M. Nadji and D.L. Durden, 1982. Glutaminase-free asparaginase from *Vibrio succinogenes*: An antilymphoma enzyme lacking hepatotoxicity. *Int. J. Cancer*, 30: 343-347.
- Duval, M., S. Suciú, A. Ferster, X. Riolland and B. Nelken *et al.*, 2002. Comparison of *Escherichia coli*-asparaginase with *Erwinia*-asparaginase in the treatment of childhood lymphoid malignancies: Results of a randomized European organisation for research and treatment of cancer-children's leukemia group phase 3 trial. *Blood*, 99: 2734-2739.
- Eisele, N., D. Linke, K. Bitzer, S. Na'amnieh, M. Nimtz and R.G. Berger, 2011. The first characterized asparaginase from a basidiomycete, *Flammulina velutipes*. *Bioresour. Technol.*, 102: 3316-3321.
- Ellis, M.B., 1971. Dematiaceous Hyphomycetes. Commonwealth Mycological Institute, Kew, Surrey, England.
- Ellis, M.B., 1976. More Dematiaceous Hyphomycetes. 1st Edn., Commonwealth Mycological Inst., Kew, Surrey, UK., Pages: 507.
- Ellis, M.B. and J.P. Ellis, 1987. Microfungi on Land Plants: An Identification Handbook. Croom Helm, London.
- Geckil, H., S. Gencer, B. Ates, U. Ozer, M. Uckun and I. Yilmaz, 2006. Effect of *Vitreoscilla* hemoglobin on production of a chemotherapeutic enzyme, L-asparaginase, by *Pseudomonas aeruginosa*. *Biotechnol. J.*, 1: 203-208.
- Gulati, R., R.K. Saxena and R. Gupta, 1997. A rapid plate assay for screening L-asparaginase producing micro-organisms. *Lett. Applied Microbiol.*, 24: 23-26.
- Gunatilaka, A.A.L., 2006. Natural products from plant-associated microorganisms: Distribution, structural diversity, bioactivity and implication of their occurrence. *J. Nat. Prod.*, 69: 509-526.
- Heinemann, B. and A.J. Howard, 1969. Production of tumor-inhibitory L-asparaginase by submerged growth of *Serratia marcescens*. *Applied Microbiol.*, 18: 550-554.
- Heinemann, B., A.J. Howard and H.J. Palocz, 1970. Influence of dissolved oxygen levels on production of L-asparaginase and prodigiosin by *Serratia marcescens*. *Applied Environ. Microbiol.*, 19: 800-804.

- Hymavathi, M., T. Sathish, P. Brahmaiah and R.S. Prakasham, 2010. Impact of carbon and nitrogen sources on L-asparaginase production by isolated *Bacillus circulans* (MTCC 8574): Application of saturated Plackett-Burman design. *Chem. Biochem. Eng. Q.*, 24: 473-480.
- Imada, A., S. Igarasi, K. Nakahama and M. Isono, 1973. Asparaginase and glutaminase activities of micro-organisms. *J. Gen. Microbiol.*, 76: 85-99.
- Kumar, S., V.V. Dasu and K. Pakshirajan, 2010. Localization and production of novel L-asparaginase from *Pectobacterium carotovorum* MTCC 1428. *Process Biochem.*, 45: 223-229.
- Kumar, S., V.V. Dasu and K. Pakshirajan, 2011. Purification and characterization of glutaminase-free L-asparaginase from *Pectobacterium carotovorum* MTCC 1428. *Bioresour. Technol.*, 102: 2077-2082.
- Kundu, B., S. Bansal and P. Mishra, 2013. Mutants of L-asparaginase. US Patent No. 20130330316 A1. <http://www.google.com/patents/US20130330316>.
- Laan, V.D.J.M., M.C. Stor, D.I. Lange and L. Mohrmann, 2008. *Asparaginase* enzyme variants and uses thereof. US Patent No. WO2008128974. <http://www.google.com/patents/WO2008128974A1?cl=en>
- Liu, F.S. and J.E. Zajic, 1972. L-Asparaginase synthesis by *Erwinia aroideae*. *Applied Environ. Microbiol.*, 23: 667-668.
- Lowry, O.H., N.J. Rosebrough, A.L. Farr and R.J. Randall, 1951. Protein measurement with the folin phenol reagent. *J. Biol. Chem.*, 193: 265-275.
- Mahajan, R.V., S. Saran, K Kameswaran, V. Kumar and R.K. Saxena, 2012. Efficient production of L-asparaginase from *Bacillus licheniformis* with low-glutaminase activity: Optimization, scale up and acrylamide degradation studies. *Bioresour. Technol.*, 125: 11-16.
- Maladkar, N.K., V.K. Singh and S.R. Naik, 1993. Fermentative production and isolation of L-asparaginase from *Erwinia carotovora*, EC-113. *Hindustan Antibiot. Bull.*, 35: 77-86.
- Manna, S., A. Sinha, R. Sadhukhan and S.L. Chakrabaty, 1995. Purification, characterization and antitumor activity of Lasparaginase isolated from *Pseudomonas stutzeri* MB-405. *Curr. Microbiol.*, 30: 291-298.
- Mashburn, L.T. and J.C. Wriston Jr., 1964. Tumor inhibitory effect of L-asparaginase from *Escherichia coli*. *Arch. Biochem. Biophys.*, 105: 450-453.
- Mohan Kumar, N.S., R. Ramasamy and H.K. Manonmani, 2013. Production and optimization of L-asparaginase from *Cladosporium* sp. using agricultural residues in solid state fermentation. *Ind. Crops Prod.*, 43: 150-158.
- Mostafa, S.A. and M.S. Salama, 1979. L-Asparaginase-producing *Streptomyces* from the soil of Kuwait. *Zentralblatt Bakteriologie Parasitenkunde Infektionskrankheiten Hygiene*, 134: 325-334.
- Mukherjee, J., S. Majumdar and T. Scheper, 2000. Studies on nutritional and oxygen requirements for production of l-asparaginase by *Enterobacter aerogenes*. *Applied Microb. Biotechnol.*, 53: 180-184.
- Muller, H.J. and J. Boos, 1998. Use of L-asparaginase in childhood ALL. *Crit. Rev. Oncol. Hematol.*, 282: 97-113.
- Nagaraju, D., M.B. GovindaRajulu, N.E. El Gueddari, T.S. Suryanarayanan and B.M. Moerschbacher, 2009. Identification and characterization of chitinolytic enzymes from endophytic fungi. *Sugars in Norwich-Royal Society*, London.

- Narayana, K.J.P., K.G. Kumar and M. Vijayalakshmi, 2008. L-asparaginase production by *Streptomyces albidoflavus*. Indian J. Microbiol., 48: 331-336.
- Onions, A.H.A., D. Allsopp and H.O.W. Eggins, 1981. Smith's Introduction to Industrial Mycology. 7th Edn., Edward Arnold (Publishers) Ltd., London, ISBN: 0 7131 2811 9, pp: 398.
- Pedreschi, F., K. Kaack and K. Granby, 2008. The effect of asparaginase on acrylamide formation in French fries. Food Chem., 109: 386-392.
- Pieters, R., S.P. Hunger, J. Boos, C. Rizzari and L. Silverman *et al.*, 2011. L-asparaginase treatment in acute lymphoblastic leukemia: A focus on *Erwinia* asparaginase. Cancer, 117: 238-249.
- Rajulu, M.B.G., N. Thirunavukkarasu, T.S. Suryanarayanan, J.P. Ravishankar, N.E. El Gueddari and B.M. Moerschbacher, 2011. Chitinolytic enzymes from endophytic fungi. Fungal Diversity, 47: 43-53.
- Robl, D., P. Da Delabona, C.M. Mergel, J.D. Rojas and P. dos Costa *et al.*, 2013. The capability of endophytic fungi for production of hemicellulases and related enzymes. BMC Biotechnol., Vol. 13. 10.1186/1472-6750-13-94
- Rocha, A.C., D. Garcia, A.P. Uetanabaro, R.T. Carneiro, I.S. Araujo, C.R. Mattos and A. Goes-Neto, 2011. Foliar endophytic fungi from *Hevea brasiliensis* and their antagonism on *Microcyclus ulei*. Fungal Diversity, 47: 75-84.
- Rytting, M.E., 2012. Role of L-asparaginase in acute lymphoblastic leukemia: Focus on adult patients. Blood Lymphatic Cancer: Targets Therapy, 2: 117-124.
- Sarquis, M.I.M., E.M.M. Oliveira, A.S. Santos and G.L. Da Costa, 2004. Production of L-asparaginase by filamentous fungi. Mem. Inst. Oswaldo Cruz., 99: 489-492.
- Savitri, N.A. and W. Azmi, 2003. Microbial L-asparaginase: A potent antitumour enzyme. Indian J. Biotechnol., 2: 184-194.
- Shaffer, P.M., H.N. Arst, L. Estberg, L. Fernando, T. Ly and M. Sitter, 1988. An asparaginase of *Aspergillus nidulans* is subject to oxygen repression in addition to nitrogen metabolite repression. Mol. Gen. Genet. MGG, 212: 337-341.
- Subramanian, C.V., 1971. Hyphomycetes: An Account of Indian Species, Except Cercosporae. Indian Council of Agricultural Research, New Delhi, India, Pages: 930.
- Sudhir, A.P., B.R. Dave, K.A. Trivedi and R.B. Subramanian, 2012. Production and amplification of an L-asparaginase gene from actinomycete isolate *Streptomyces* ABR2. Ann. Microbiol., 62: 1609-1614.
- Sukumaran, C.P., D.V. Singh and P.R. Mahadevan, 1979. Synthesis of L-asparaginase by *Serratia marcescens* (Nima). J. Biosci., 1: 263-269.
- Suryanarayanan, T.S., V. Kumaresan and J.A. Johnson, 1998. Foliar fungal endophytes from two species of the mangrove *Rhizophora*. Can. J. Microbiol., 44: 1003-1006.
- Suryanarayanan, T.S., N. Thirunavukkarasu, M.B. Govindarajulu, F. Sasse, R. Jansen and T.S. Murali, 2009. Fungal endophytes and bioprospecting. Fungal Biol. Rev., 23: 9-19.
- Suryanarayanan, T.S., N. Thirunavukkarasu, M.B. Govindarajulu and V. Gopalan, 2012. Fungal endophytes: An untapped source of biocatalysts. Fungal Diversity, 54: 19-30.
- Sutton, B.C., 1980. The Coelomycetes: Fungi Imperfecti with Pycnidia Acervuli and Stromata. Commonwealth Mycological Institute, Kew, Surrey, England, ISBN: 9780851984469, Pages: 696.

- Thirunavukkarasu, N., T.S. Suryanarayanan, T.S. Murali, J.P. Ravishankar and S.N. Gummadi, 2011. L-asparaginase from marine derived fungal endophytes of seaweeds. *Mycosphere*, 2: 147-155.
- Vega, F.E., 2008. Insect pathology and fungal endophytes. *J. Invertebrate Pathol.*, 98: 277-279.
- Von Arx, J.A., 1974. The Genera of Fungi Sporulation in Pure Culture. 2nd Edn., J. Cramer in der A.-R.-Gantner-Verlag-Kommanditgesellschaft, Vaduz, Liechtenstein, ISBN-13: 9783768206938, Pages: 315.
- Weber, D., 2009. Endophytic Fungi, Occurrence and Metabolites. In: *The Mycota XV Physiology and Genetics*, Anke, T. and D. Weber (Eds.). Springer, New York, pp: 153-195.
- Wei, D.Z. and H. Liu, 1998. Promotion of L-asparaginase production by using n-dodecane. *Biotechnol. Techniques*, 12: 129-131.