

This article was downloaded by: [CRA]

On: 21 March 2013, At: 23:42

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Archives Of Phytopathology And Plant Protection

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gapp20>

Screening of plant-derived antifungal substances useful for the control of seedborne pathogens

Catello Pane ^a, Domenica VILLECCO ^a, Graziana ROSCIGNO ^b, Enrica De Falco ^b & Massimo ZACCARDELLI ^a

^a Consiglio per la Ricerca e la Sperimentazione in Agricoltura - Centro di Ricerca per l'Orticoltura, Pontecagnano, Italy

^b Dipartimento di Farmacia, Università degli Studi di Salerno, Fisciano, Italy

Version of record first published: 20 Mar 2013.

To cite this article: Catello Pane , Domenica VILLECCO , Graziana ROSCIGNO , Enrica De Falco & Massimo ZACCARDELLI (2013): Screening of plant-derived antifungal substances useful for the control of seedborne pathogens, Archives Of Phytopathology And Plant Protection, DOI:10.1080/03235408.2013.771458

To link to this article: <http://dx.doi.org/10.1080/03235408.2013.771458>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Screening of plant-derived antifungal substances useful for the control of seedborne pathogens

Catello Pane^{a*}, Domenica Vilecco^a, Graziana Roscigno^b, Enrica De Falco^b and Massimo Zaccardelli^a

^aConsiglio per la Ricerca e la Sperimentazione in Agricoltura – Centro di Ricerca per l'Orticoltura, Pontecagnano, Italy; ^bDipartimento di Farmacia, Università degli Studi di Salerno, Fisciano, Italy

(Received 25 January 2013; final version received 28 January 2013)

The effect of plant essential oils, *Solanum chilense* and *Aster sedifolius* raw saponins, plant-derived colourants and *Brassica carinata* glucosinolate-containing tissues, for the control of seedborne fungi *Alternaria dauci*, *Alternaria radicina*, *Colletotrichum lindemuthianum* and *Ascochyta rabiei*, was evaluated. *In vitro* assays indicated that oregano essential oil, bark chestnut-derived colourant and *B. carinata* meals, showed a large spectrum of activity against all pathogens. *S. chilense* raw saponins were effective against *A. dauci* and *A. radicina*, while those from *A. sedifolius* also inhibited significantly *C. lindemuthianum*. Screening showed the antifungal potential of natural compounds.

Keywords: dyeing plants; essential oils; glucosinolates; saponins; smart agriculture

Introduction

Seedborne pathogens cause diseases where the seeds are the main source of inoculum (Nome et al. 2002). The control of pathogens associated with the seeds is very problematic, because the infections departing from coat and embryo are difficult to control using only agronomic practices. Therefore, prevention of Seedborne diseases is based on the use of certified plant material free from pathogens and/or using resistant cultivars. However, because plant resistance against these pathogens is very erratic, the control of these microbes was currently entrusted to chemical fungicides. Nevertheless, the preference of the consumers for safe and organic productions has stimulated the research of sustainable alternative pesticides (Saba and Messina 2003). For this reason, plant-derived compounds with antifungal activity, such as essential oils (Isman 2000), saponins (Barile et al. 2006) and glucosinolates (Brown and Morra 1997) are promising tools to totally replace or integrate chemicals for disease management. Recently, plants used for dye extraction have been shown to possess antimicrobial activity own to the presence of a large amount of tannins, anthraquinones or naphthoquinones (Singh et al. 2005; Chengaiah et al. 2010). Seed application of plant-derived molecules may be considerate an integrated pest management strategy able to improve pathogen control and to reduce unsustainable use of traditional fungicides (Ntalli and Menkissoglu-Spiroudi 2011).

*Corresponding author. Email: catello.pane@entecra.it

This research was focused on environmentally compatible control of four Seedborne pathogens: *Alternaria dauci* (J.G. Kühn) J.W. Groves & Skolko, *Alternaria radicina* Meier, Drechsler & E.D. Eddy, *Colletotrichum lindemuthianum* (Sacc. & Magnus) Lams.-Scrib. and *Ascochyta rabiei* (Pass.) Labr. *A. dauci* and *A. radicina* are related pathogens causing severe blights on leaves, petioles and stems of carrot (*Daucus carota* Blotter). *A. radicina* can cause a further distinctive black rot symptom, consisting in damping-off and rotting of roots, crowns and seedlings of carrots that become unmarketable. *C. lindemuthianum* and *A. rabiei* are the causal agents of anthracnose of common bean (*Phaseolus vulgaris* L.) and leaf blight of chickpea (*Cicer arietinum* L.), respectively. The research of alternative to chemicals for the control of these diseases already proposed the use of seed treatment with physical methods, resistance inducers and antagonistic microorganisms (Tinivella et al. 2009; Koch et al. 2010). Aim of this study was to show a further screening of other phytochemicals, such as essential oils, natural dyeing, raw saponins and *Brassica carinata* L. meals containing glucosinolates for their *in vitro* antifungal activity against the described Seedborne pathogens.

Results and discussion

This study provided new data about *in vitro* inhibition activity of some plant-derived substances against Seedborne pathogens *A. radicina*, *A. dauci*, *A. rabiei* and *C. lindemuthianum*. *In vitro* assays indicate that plant essential oils showed variable effects on mycelial growth of the tested fungi. Oregano oil showed the largest spectrum of activity, because inhibited all tested fungi. On the contrary, rosemary oil does not affect any pathogen used in this work. *A. radicina* was more sensible to essential oils than *A. dauci*. A number of essential oils, including hyssop, oregano, basil, caraway, sage, marjoram and balm, were able to suppress black rot agent, whereas *A. dauci* was inhibited only by oregano and thyme oil (Table 1). About *C. lindemuthianum* and *A. rabiei*, results indicate that hyssop and oregano-derived essential oils inhibited both pathogens. Moreover, oils distilled from basil, caraway, sage and thyme blocked mycelial growth of *C. lindemuthianum*, while essential oils from lemon balm and verbena blocked *A. rabiei* growth. Essential oils generally were reported as very efficacious in order to reduce *in vitro* development of several fungi belonging to genus *Alternaria* (Tian et al. 2011), *Colletotrichum* (Chang et al. 2008; Arrebola et al. 2010; Maqbool et al. 2011) and *Ascochyta* (Zerroug et al. 2011). In agreement with our results, oregano essential oil revealed, by *in vitro* assays, a broad-spectrum antifungal activity vs. a plethora of fungi (Portillo-Ruiz et al. 2005).

Among over 10 dyeing plants extracts assayed in this study, *Castanea sativa* L. bark extract partially inhibited *in vitro* growth of all the four pathogens tested in this study, probably due to the high content of tannins. *Rubia tinctorum* L. inhibited completely the radial growth of *A. dauci* and *A. rabiei* (Table 1) and these results are in accordance with the literature data inherent the antimicrobial activity of root extract of this plant (Manojlovic et al. 2005; Kalyoncu et al. 2006). A similar case was showed by the extracts from *in vitro* grown shoots of *Quillaja saponaria* that proved effectiveness against *Botrytis cinerea* (Ribera et al. 2008). The absence of inhibition of the other tested extracts can be attributed to the low dose adopted as reported from Prusty et al. (2010). So the natural dyes require more in-depth investigation into the effect of dose and dye structure on antimicrobial property.

Raw saponin-containing tissues showed encouraging specie-specific antifungal properties. In plate assays, raw saponins contained in tissue-meals of *Solanum chilense*

Table 1. Effect of different plant-derived products (essential oils and dyeing extracts) on mycelial growth of *A. dauci*, *A. radicina*, *A. rabiei* and *C. lindemuthianum*. Signs – and + were used to indicate inhibition rank.

Plant-derived products	Concentration used	Mycelial growth inhibition ^a			
		<i>A. dauci</i>	<i>A. radicina</i>	<i>A. rabiei</i>	<i>C. lindemuthianum</i>
Basil oil	1 vol.%	–	+++	–	+++
Caraway oil	1 vol.%	–	+++	–	+++
Hyssop oil	1 vol.%	–	+++	+++	+++
Majoram oil	1 vol.%	–	+++	–	–
Melissa oil	1 vol.%	–	+++	+++	–
Oregano oil	1 vol.%	+++	+++	+++	+++
Rosmary oil	1 vol.%	–	–	–	–
Sage oil	1 vol.%	–	+++	–	+++
Thyme oil	1 vol.%	+++	–	–	+++
Verbena oil	1 vol.%	–	–	+++	–
Bark chestnut dye	1 vol.%	+	+	+	+
Clover dye	1 vol.%	–	–	–	–
Chrysanthemum dye	1 vol.%	–	–	–	–
Heather dye	1 vol.%	–	–	–	–
Wild fennel dye	1 vol.%	–	–	–	–
Inula dye	1 vol.%	–	–	–	–
Pomegranate dye	1 vol.%	–	–	–	–
Mint dye	1 vol.%	–	–	–	–
Madder dye	1 vol.%	–	–	–	–
Elder dye	1 vol.%	–	–	–	–

Note: ^aInhibition rank: 100% (+++), >50% (++) , <50% (+) significant growth reduction compared with control. Not inhibited (–).

reduced significantly radial growth rate of *A. dauci* and *A. radicina*, while those derived from *Aster sedifolius* inhibited significantly *A. radicina*, *C. lindemuthianum* and *A. rabiei*. Saponins are a class of natural compounds isolated from plants that can exhibit antimicrobial properties independently from their purity degree, as in the case of saponin-rich fraction of *Camellia oleifera* cake (Hu et al. 2012). In our work, finally, *B. carinata* glucosinolate-derived isothiocyanates were also significantly effective in *in vitro* control of all the four assayed pathogens (Figure 1) confirming the high antimicrobial activity showed by these molecules (Tierens et al. 2001).

Findings of the present work suggest that plant-derived products with antifungal properties can have a large potential for practical application in disease management, since they are composed of natural molecules compatible with environmental sustainable cropping systems. Results of the present screening gave encouraging indications regarding the potential utilisation of the assayed natural products in the protection against pathogens transmissible through the seeds. Here, we observed that the broad-spectrum efficacy of *B. carinata* seed meals, as-well-as that of some essential oils is in accordance with a number of literature data (Sridhar et al. 2003; Galletti et al. 2008; Lalitha et al. 2011). Similarly, raw saponins exhibited a great potential to implement the control of Seedborne diseases (Stuardo and San Martín 2008). All these substances with antifungal properties can be suitable biofungicides for direct seed applications, by coating or tanning, to improve plant protection against pathogens (Wulff et al. 2011). As matter of the fact, essential oils are reported to be promising for the protection of the seeds against




Figure 1. Effect of *S. chilense* (SC), *A. sedifolius* (AS) and *B. carinata* (BC) tissue meals on radial growth rate of *A. dauci*, *A. radicina*, *A. rabiei* and *C. lindemuthianum*, compared to the control (CTRL). Different letters indicate differences statistically significant. Data averages (± 1 SE) of three replicates.

infestation by specialised fungi, because they have penetration action (Lalitha et al. 2011). It is the case, for example, of the various oil seed treatments that have successfully controlled the cumin rot root disease caused by *Fusarium* spp. (Hashem et al. 2010). Among other non-chemical seed treatments, thyme oil has been proposed to reduce Seedborne diseases caused by *C. lindemuthianum* on bean and *Ascochyta* spp. on pea (Tinivella et al. 2009), *A. dauci* and *A. radicina* on carrot (Koch et al. 2010), *Phoma valerianellae* on lamb's lettuce (Schmitt et al. 2009) and *Alternaria brassicola* on cabbage (Amein et al. 2011). Likewise, seed treatments with *Brassica* spp. meals have also reported to control *Rhizoctonia solani* in cabbage (Chung et al. 2002). Previously, *B. carinata* seed meals have been proposed as geodisinfectant in a greenhouse lettuce system (Pane et al. 2012). While this product containing glucosinolates was also proposed, together some essential oils as foliar spray to improve processing tomato plant conditions in open field (Pane et al. 2013). The current research showed oregano essential oil, *B. carinata* seed meal and bark chestnut dye the most promising non-chemical fungicides that can be proposed against the examined Seedborne pathogens. However, as a perspective of this research, the practical applications of these natural products in disease control will have to be tested to evaluate the actual potential of the treatments.

Experimental

Fungal pathogens

A. radicina (strain ER1462), *A. dauci* (strain ER1530), *A. rabiei* (strain 1405AR) and *C. lindemuthianum* (strain ISCI9) were originally isolated from their specific hosts. The fungal isolates were stored at 4 °C in tubes containing potato–dextrose–agar (PDA, Difco).

In vitro plate assays

To evaluate the ability of plant-derived substances to inhibit growth of fungal plant pathogens *A. radicina*, *A. dauci*, *A. rabiei* and *C. lindemuthianum*, *in vitro* plate tests were performed. 10 essential oils of aromatic plants and 10 water extracts of dyeing plants were separately tested by submerging a plug (5 mm in diameter) removed from the edge of the growing mycelia in water suspensions. Essential oil emulsions were prepared at 1% concentration (vol./vol.). While to produce the dyeing plant extract solution 5 g of dried material in 800 ml of water for 60 min at 70 °C at neutral pH, and concentrated by rotavapor, until 1:4 in volume. After overnight incubation at 25 °C, the plug was transferred in the centre of a PDA Petri plates (diameter 90 mm). Assay with glucosinolates was carried out including, in inverted hermetically closed PDA plates and under the plug, a capsule containing 100 mg of *B. carinata* seed meals wetted with 500 µl of sterile distilled water. Raw saponins, instead, were tested amending PDA with *A. sedifolius* or *S. chilense* tissue meals at a rate of 0.5 g ml⁻¹. In all cases, control plates were prepared without the addition of the tested substances. For each treatment, plates were inoculated in triplicate and incubated at 25 °C. The diameter of the mycelia was measured daily until fungi reached the edge of control plates; data were expressed as growth rate (mm d⁻¹). Instead, in the case of essential oils and dyeing plant extracts, only *in vitro* inhibition levels were indicated as 100% (+++), >50% (++) , <50% (+) significant growth reduction compared to control, and not inhibited (-).

Statistical analysis

Data from the plate inhibition assays of the fungi with raw saponins and glucosinolates were processed with the analysis of variance (ANOVA). When the ANOVA was significant ($P \leq 0.05$), means were separated with Duncan's test.

Acknowledgements

This research was financed with funding programme 'Pro. Vi. Se. Bio.', research title Agro. Se. Bio., and 'Extravalore' provided by the Italian Ministry of Agricultural Policy. We are grateful to Dr. Maria Cammareri, (University of Naples) for providing plant tissues containing raw saponins.

References

- Amein T, Wright S, Wikström M, Koch E, Schmitt A, Stephan D, Jahn M, Tinivella F, Gullino ML, Forsberg G, Werner S, van der Wolf J, Groot S, 2011. Evaluation of non-chemical seed treatment methods for control of *Alternaria brassicicola* on cabbage seeds. J Plant Dis Plant Protec. 118:214–221.
- Arrebola E, Sivakumar D, Bacigalupo R, Korsten L. 2010. Combined application of antagonist *Bacillus amyloliquefaciens* and essential oils for the control of peach postharvest diseases. Crop Prot. 29:369–377.
- Barile E, Bonanomi G, Antignani V, Zolfaghari B, Ebrahim Sajjadi S, Scala F, Lanzotti V. 2006. Saponins from *Allium minutiflorum* with antifungal activity. Phytochem. 68:596–603.
- Brown PD, Morra MJ. 1997. Control of soil-borne plant pests using glucosinolate-containing plants. Adv Agron. 61:167–231.
- Chang HT, Cheng YH, Wu CL, Chang ST, Chang TT, Su YC. 2008. Antifungal activity of essential oil and its constituents from *Calocedrus macrolepis* var. *formosana* Florin leaf against plant pathogenic fungi. Biores Technol. 99:6266–6270.
- Chengaiyah B, Rao KM, Kumar KM, Alagusundaram M, Chetty CM. 2010. Medicinal importance of natural dyes – A review. Int J PharmTech Res. 2:144–154.

- Chung WC, Huang JW, Huang HC, Jen JF. 2002. Effect of ground *Brassica* seed meal on control of *Rhizoctonia* damping-off of cabbage. *Can J Plant Pathol.* 24:211–218.
- Galletti S, Sala E, Leoni O, Burzi PL, Cerato C. 2008. *Trichoderma* spp. tolerance to *Brassica carinata* seed meal for a combined use in biofumigation. *Biol Cont.* 45:319–327.
- Hashem M, Moharam AM, Zaied AA, Saleh FEM. 2010. Efficacy of essential oils in the control of cumin root rot disease caused by *Fusarium* spp. *Crop Prot.* 29:1111–1117.
- Hu JL, Nie SP, Huang DF, Li C, Xie MY, Wan Y. 2012. Antimicrobial activity of saponin-rich fraction from *Camellia oleifera* cake and its effect on cell viability of mouse macrophage RAW 264.7. *J Sci Food Agric.* 92:2443–2449.
- Isman MB. 2000. Plant essential oils for pest and disease management. *Crop Prot.* 19:603–608.
- Kalyoncu F, Cetin B, Saglam H. 2006. Antimicrobial activity of common madder (*Rubia tinctorum* L.). *Phytother Res.* 20:490–492.
- Koch E, Schemm A, Stephan D, Kromphardt C, Jahn M, Krauthausen HJ, Forsberg G, Werner S, Amein T, Wright SAI, et al. 2010. Evaluation of non-chemical seed treatment methods for the control of *Alternaria dauci* and *A. radicina* on carrots seeds. *Eur J Plant Pathol.* 127:99–112.
- Lalitha V, Kiran B, Raveesha KA. 2011. Antifungal and antibacterial potentiality of six essential oils extracted from plant source. *Int J Eng Sci Technol.* 3:3029–3037.
- Manojlovic NT, Solujic S, Sukdolak S, Milosev M. 2005. Antifungal activity of *Rubia tinctorum*, *Rhamnus frangula* and *Coloplaca cerina*. *Fitoterapia.* 76:244–246.
- Maqbool M, Ali A, Alderson PG, Tengku Muda Mohamed M, Siddiqui Y, Zahid N. 2011. Post-harvest application of gum arabic and essential oils for controlling anthracnose and quality of banana and papaya during cold storage. *Postharvest Biol Tec.* 62:71–76.
- Nome SF, Barreto D, Docampo DM. 2002. Seedborne pathogens. In: McDonald M, Contreras S, editors. *Proceedings International Seed Seminar: Trade, Production and Technology; 2002 Oct 15th and 16th. Santiago (Chile): Pontificia Universidad Católica de Chile, Facultad de Agronomía e Ingeniería Forestal, Departamento de Ciencias Vegetales.* p. 114–126.
- Ntalli NG, Menkissoglu-Spiroudi U. 2011. Pesticides of botanical origin: a promising tool in plant protection. In: Stoytcheva M, editor. *Pesticides – formulations, effects, fate.* Rijeca: InTech; p. 3–24.
- Pane C, Vilecco D, Pentangelo A, Lahoz E, Zaccardelli M. 2012. Integration of soil solarization with *Brassica carinata* seed meals amendment in a green house lettuce production system. *Acta Agric Scand Sect B Soil Plant Sci.* 62:291–299.
- Pane C, Rongai D, Zaccardelli M. Forthcoming 2013. Foliar spray application of glucosinolates and essential oils on processing tomato in open field production system. *Agric Sci.*
- Portillo-Ruiz MC, Viramontes-Ramos S, Muñoz-Castellanos LN, Gastélum-Franco MG, Nevárez-Moorillón GV. 2005. Antifungal activity of Mexican oregano (*Lippia berlandieri* Shauer). *J Food Prot.* 68:2713–2717.
- Prusty AK, Das T, Nayak A, Das NB. 2010. Colourimetric analysis and antimicrobial study of natural dyes and dyed silk. *J Cleaner Prod.* 18:1750–1756.
- Ribera A, Cotoras M, Zúñiga GE. 2008. Effect of extracts from *in vitro*-grown shoots of *Quillaja saponaria* Mol. on *Botrytis cinerea* Pers. *World J Microbiol Biotechnol.* 24:1803–1811.
- Saba A, Messina F. 2003. Attitudes towards organic foods and risk/benefit perception associated with pesticides. *Food Qual Pref.* 14:637–645.
- Schmitt A, Koch E, Stephan D, Kromphardt C, Jahn M, Krauthausen HJ, Forsberg G, Werner S, Amein T, Wright SAI. 2009. Evaluation of non-chemical seed treatment methods for the control of *Phoma valerianellae* on lamb's lettuce seeds. *J Plant Dis Prot.* 116:200–207.
- Singh R, Jain A, Panwar S, Gupta D, Khare SK. 2005. Antimicrobial activity of some natural dyes. *Dyes Pigm.* 66:99–102.
- Sridhar SR, Rajagopal RV, Rajavel R, Masilamani S, Narasimhan S. 2003. Antifungal activity of some essential oils. *J Agric Food Chem.* 51:7596–7599.
- Stuardo M, San Martín R. 2008. Antifungal properties of quinoa (*Chenopodium quinoa* qeite: Willd.) alkali treated saponins against *Botrytis cinerea*. *Ind Crop Prod.* 27:296–302.
- Tian J, Ban X, Zeng H, He J, Huang B, Wang Y. 2011. Chemical composition and antifungal activity of essential oil from *Cicuta virosa* L. var. *latisecta* Celak. *Int J Food Microbiol.* 145:464–470.
- Tierens KFMJ, Thomma BPHJ, Brouwer M, Schmidt J, Kistner K, Porzel A, Mauch-Mani B, Cammue BPA, Broekaert WF. 2001. Study of the role of antimicrobial glucosinolate-derived isothiocyanates in resistance of *Arabidopsis* to microbial pathogens. *Plant Physiol.* 125:1688–1699.

- Timivella F, Hirata LM, Celan MA, Wright SAI, Amein T, Schmitt A, Koch E, van der Wolf JM, Groot SPC, Stephan D, et al. 2009. Control of seed-borne pathogens on legumes by microbial and other alternative seed treatments. *Eur J Plant Pathol.* 123:139–151.
- Wulff EG, Zida E, Torp J, Lund OS. 2011. *Yucca schidigera* extract: a potential biofungicide against seedborne pathogens of sorghum. *Plant. Pathol.* 61:331–338.
- Zerroug MM, Laouer H, Nicklin J. 2011. The effect of essential oil of *Saccocalyx satureioides* Coss. et Dur. on the growth of and the production of solanapyrone a by *Ascochyta rabiei* (Pass.) Labr. *Adv Environ Biol.* 5:501–506.