



UNIVERSITÀ  
DEGLI STUDI  
DI PADOVA



# Search for massive long-lived particles decaying semileptonically at $\sqrt{s} = 13 \text{ TeV}$

**Davide Zuliani\***

University and INFN of Padova

On behalf of the LHCb Collaboration

\* for questions/comments: [davide.zuliani@cern.ch](mailto:davide.zuliani@cern.ch)

# Outline



- LHCb experiment
- Search for massive long-lived particles decaying semileptonically at  $\sqrt{s} = 13$  TeV
- Conclusions

# LHCb experiment

## A General Purpose Forward Detector

JINST 3 (2008) S08005  
 Int. J. Mod. Phys. A 30, 1530022 (2015)  
 CERN-LPCC-2018-04

- LHCb, originally designed for  $b$ - and  $c$ -hadron physics, is now considered a **general purpose forward detector**
- Unique phase space region ( $2 < \eta < 5$ ) **complementary to General Purpose Detectors (ATLAS & CMS)**





- **Excellent track momentum resolution:** 0.4% at 5 GeV to 1.0% at 200 GeV
- Impact Parameter resolution  $\sigma_{IP} = 20 \mu\text{m}$  for high- $p_T$  tracks, lifetime resolution of  $\sigma_\tau = 0.2 \text{ ps}$
- Muon ID efficiency: 97% with 1-3%  $\mu \rightarrow \pi$  misidentification
- Electron ID efficiency: 90% with 5%  $h \rightarrow e$  misidentification

# LHCb experiment

## VELO material map

JINST 13 (2018) 06, P06008

- Material map of VERtEx LOcator (VELO) is fundamental for LLP searches:
  - Displacement up to 20 cm
  - Thin VELO envelope (RF foil) - background dominated by
    - heavy flavour decays at  $< 5$  mm
    - material interactions at  $> 5$  mm
- Precise material VELO map by secondary interactions of hadrons produced in beam-gas collisions



- So far only performed analyses on Run 1 and Run 2 data with LLPs decaying within the VELO
- Searches could be extended to LLPs decaying downstream of the VELO (displacement up to 200 cm)
  - much worse momentum resolution

# LLPs searches at LHCb so far...

- Displaced leptons:
  - Dark photon
  - Low-mass di-muon resonances
  - Majorana neutrino
  - LLPs decaying to  $e^\pm \mu^\mp \nu$
  - Light boson from  $b \rightarrow s$  decays
- Displaced jets:
  - HNL in  $W^\pm \rightarrow \mu^\pm \mu^\pm$  jet
  - LLP  $\rightarrow$  jet jet
  - LLP  $\rightarrow \mu +$  jets**

[PRL 124\(2020\) 041801](#)

[LHCb-PAPER-2020-013](#)

[PRL 112 \(2014\) 131082](#)


[EPJC 81 \(2021\) 261](#)

[PRD 95 \(2017\) 071101](#)

[EPJC 81 \(2021\) 248](#)

[EPJC 77 \(2017\) 812](#)

[LHCb-PAPER-2021-028](#)



# Search for massive long-lived particles decaying semileptonically at $\sqrt{s} = 13$ TeV


## Theory context

- Supersymmetry (SUSY) is one of the most popular extension of Standard Model (SM)
  - Solves the hierarchy problem
  - Unifies gauge couplings at Planck scale
  - Dark Matter candidates
- Subset of models for Minimal Supersymmetric Standard Model (MSSM) addresses long-lived particles
  - Main signature: measurable flight distance and displaced vertices
- If considering R-parity violation (RPV) processes a MSSM long-lived particle can decay into SM particles
- In this analysis the minimal Super GRAvity (mSUGRA) theoretical model has been considered, with RPV
  - A “neutralino”  $\tilde{\chi}_1^0$  can decay into a muon and two quarks:  $\tilde{\chi}_1^0 \rightarrow \mu^+ q_i q_j (\mu^- \bar{q}_i \bar{q}_j)$

# Search for massive long-lived particles decaying semileptonically at $\sqrt{s} = 13$ TeV

## LLPs production mode

- In this analysis two productions mode have been considered:



- The Higgs-like decay analysis covers  $h^0$  masses from 30 to 200 GeV/c<sup>2</sup>
- LLPs masses are in the range [10,  $\sim m(h^0)/2$ ] GeV/c<sup>2</sup> and lifetimes in the range [5,200] ps
- The direct production mode address LLPs masses in the range [10, 90] GeV/c<sup>2</sup> and lifetimes in the range [5,200] ps
- Lifetime range well above b-hadron lifetime and vertices still within LHCb VELO
- Mass range to avoid SM b-quark states and to consider LHCb forward acceptance
- Relevant backgrounds:  $b\bar{b}$  and  $c\bar{c}$  direct production and  $Z$ ,  $W$ , Higgs and top decays

# Search for massive long-lived particles decaying semileptonically at $\sqrt{s} = 13$ TeV

## Signal Selection











- The LLP signature is a displaced vertex (charged particles tracks) + isolated muon with high  $p_T$
- Muons are expected to be more isolated than muons in hadron decays (high LLP masses)
- PV selection:
  - Small radial distance from beam axis,  $R_{xy} < 0.3$  mm
- Muon selection:
  - Online trigger selects muon with  $p_T > 10$  GeV/c
  - Offline selection: impact parameter  $IP^\mu > 0.25$  mm and  $p_T > 12$  GeV/c
  - Isolation: sum of energy tracks around muon (with muon) in a cone of radius  $R_{\eta\phi} = 0.3$
- LLP selection:
  - Once PVs are found, geometric veto on displaced vertices (remember VELO material map)
  - A LLP candidate is formed by 3 or more tracks + muon, and  $m_{inv}(\text{LLP}) > 4.5$  GeV/c<sup>2</sup>



# Search for massive long-lived particles decaying semileptonically at $\sqrt{s} = 13$ TeV

## Signal Selection

- Distribution for simulated  $b\bar{b}$  background samples and  $h^0 \rightarrow \tilde{\chi}_1^0 \tilde{\chi}_1^0$  with  $m(h^0) = 125$  GeV/ $c^2$  and  $m(\tilde{\chi}_1^0) = 40$  GeV/ $c^2$
- Shapes are consistent with  $b\bar{b}$  composition of background
- Background estimation used as cross-checks (see later)



# Search for massive long-lived particles decaying semileptonically at $\sqrt{s} = 13$ TeV


## MVA analysis

- A multivariate analysis (MVA) based on Boosted Decision Trees is used to further purify the data sample
- Ten MVA input variables are selected to perform signal-background separation
  - $p_T$  and IP of the muon
  - Ratio of energies released in ECAL and HCAL normalized to muon energy
  - $p_T$  and  $\eta$  of the LLP candidate
  - Number of tracks forming the LLP candidate
  - Vertex distance  $R_{xy}$
  - Uncertainties of the vertex:  $\sigma_R, \sigma_z$
- Muon-isolation variable and reconstructed LLP mass are used to get the LLPs yield
- Signal MVA training samples are obtained from simulation
- Background training sample is obtained from data
  - No bias on MVA performance

# Search for massive long-lived particles decaying semileptonically at $\sqrt{s} = 13$ TeV

## Signal Extraction

- After MVA application no background events survive
- Background is obtained with a data-driven method
- Muon isolation is used to find a signal and a background region:
  - Signal region = muon-isolation variable  $< 1.2$
  - Background region = muon-isolation variable  $> 1.2$
- 80% of signal events are included in the signal region
- Fit to reconstructed LLP mass from background region
  - Constraints on fit on signal region



# Search for massive long-lived particles decaying semileptonically at $\sqrt{s} = 13$ TeV

## Efficiencies and systematics

- Detector efficiency is estimated from simulations
- Several factor to be taken into account:
  - Efficiency increases with LLP mass (more particles are produced)
  - Loss of particles outside the detector when LLPs come from heavier states
  - Lower boost of heavier LLPs results in shorter flight distance (cut on radial distance  $R_{xy}$ )
  - With increasing LLPs lifetime  $\rightarrow$  material region VETO
  - Loss in efficiency due to MVA selection
- Systematic uncertainties take into account several aspects
  - Muon reconstruction
  - Vertex reconstruction
  - MVA discrimination
  - ...

Direct production mode


$m_{\tilde{\chi}_1^0}$	$\tau_{\tilde{\chi}_1^0}$	$\epsilon_{presel}$	$\epsilon$	$N_b$		$N_s$		$\chi^2/ndf$
10	10	0.61	0.13	2767.9	$\pm 88.2$	-141.8	$\pm 69.7$	1.69
20	10	0.66	0.23	43.9	$\pm 40.1$	-4.2	$\pm 5.0$	0.67
30	10	2.29	0.47	15.7	$\pm 5.8$	3.3	$\pm 5.2$	0.90
40	10	2.49	0.52	1.1	$\pm 1.4$	5.9	$\pm 2.8$	0.96
60	10	3.81	1.97	45.1	$\pm 5.6$	-8.0	$\pm 4.3$	0.80
90	10	2.52	1.68	30.8	$\pm 2.2$	-9.8	$\pm 5.0$	1.04
30	5	1.44	0.21	11.0	$\pm 2.5$	-1.0	$\pm 2.7$	0.67
30	20	2.64	0.66	13.8	$\pm 4.4$	3.2	$\pm 4.2$	0.65
30	30	2.52	0.74	5.6	$\pm 2.2$	2.4	$\pm 2.1$	0.41
30	50	2.25	0.81	16.5	$\pm 16.1$	-1.8	$\pm 3.2$	0.69
30	100	1.68	0.61	9.9	$\pm 7.4$	-1.7	$\pm 3.1$	1.10
30	200	1.06	0.29	38.0	$\pm 6.3$	0.0	$\pm 2.3$	0.79

Source	Contribution [%]
Integrated luminosity	2.0
Parton luminosity gluons fusion (quarks)	6.0 (3.0)
Simulation statistics	2.0-4.0
Muon reconstruction	2.0-3.7
$p_T^\mu$	1.0
IP $^\mu$	1.0
Vertex reconstruction	2.0
Beam line uncertainty ( $R_{xy}$ )	0.9
Muon isolation	1.7
MVA	1.7-16
Mass calibration	1.4
Total	7.3-18.9

# Search for massive long-lived particles decaying semileptonically at $\sqrt{s} = 13$ TeV

95% CL upper limits


- No excess is found
- 95% CL upper limits are computed on  $\sigma(\text{LLPs}) \times \mathcal{B}(\text{LLPs} \rightarrow q\bar{q}\mu)$  for both production modes
- Statistical and systematic uncertainties are included as nuisance parameters



# Search for massive long-lived particles decaying semileptonically at $\sqrt{s} = 13$ TeV

## 95% CL upper limits


- No excess is found
- 95% CL upper limits are computed on  $\sigma(\text{LLPs}) \times \mathcal{B}(\text{LLPs} \rightarrow q\bar{q}\mu)$  for both production modes
- Statistical and systematic uncertainties are included as nuisance parameters



# Search for massive long-lived particles decaying semileptonically at $\sqrt{s} = 13$ TeV

95% CL upper limits

- No excess is found
- 95% CL upper limits are computed on  $\sigma(\text{LLPs}) \times \mathcal{B}(\text{LLPs} \rightarrow q\bar{q}\mu)$  for both production modes
- Statistical and systematic uncertainties are included as nuisance parameters



# Conclusions

## Wrap up

- LHCb is by all means a **general purpose forward detector**
- LHCb is a unique place to study LLPs:
  - Detection of low-mass particles and soft signatures
  - Studies on b- and c-decays
  - Phase space region complementary to ATLAS and CMS
- Here we studied massive LLPs decaying semileptonically into a muon and two quarks
  - Two production modes (direct production and Higgs like boson decay)
  - Different mass and lifetime ranges
  - Upper limits on  $\sigma(\text{LLPs}) \times \mathcal{B}(\text{LLPs} \rightarrow q\bar{q}\mu)$  with sensitivity of the order  $O(1 \text{ pb})$
- If you're interested in LLPs searches at LHCb go check [yesterday's presentation](#)



**Thank you for  
your attention!**