

 Open access • Journal Article • DOI:10.1103/PHYSREVLTT.102.231801

Search for resonant diphoton production with the D0 detector — [Source link](#)

[V. M. Abazov](#), [Brad Abbott](#), [M. Abolins](#), [B. S. Acharya](#) ...+510 more authors

Institutions: [Joint Institute for Nuclear Research](#), [University of Oklahoma](#), [Michigan State University](#), [Tata Institute of Fundamental Research](#) ...+73 more institutions

Published on: 12 Jun 2009 - [Physical Review Letters](#) (American Physical Society)

Topics: [Higgs boson](#), [Standard Model](#), [Tevatron](#), [Supersymmetry](#) and [Particle decay](#)

Related papers:

- [PYTHIA 6.4 Physics and Manual](#)
- [The upgraded DØ detector](#)
- [Search for a Higgs boson in diphoton final states with the D0 detector in 9.6 fb\(-1\) of p\(p\)over-bar collisions at root s=1.96 TeV](#)
- [Search for a Fermiophobic and Standard Model Higgs Boson in Diphoton Final States](#)
- [Measurement of direct photon pair production cross sections in p over\(p,γ\) collisions at sqrt\(s\) = 1.96 TeV](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/search-for-resonant-diphoton-production-with-the-d0-detector-543r24ptvi>

Search for Resonant Diphoton Production with the D0 Detector

V. M. Abazov,³⁶ B. Abbott,⁷⁵ M. Abolins,⁶⁵ B. S. Acharya,²⁹ M. Adams,⁵¹ T. Adams,⁴⁹ E. Aguilo,⁶ M. Ahsan,⁵⁹ G. D. Alexeev,³⁶ G. Alkhazov,⁴⁰ A. Alton,^{64,*} G. Alverson,⁶³ G. A. Alves,² M. Anastasoia,³⁵ L. S. Ancu,³⁵ T. Andeen,⁵³ B. Andrieu,¹⁷ M. S. Anzelc,⁵³ M. Aoki,⁵⁰ Y. Arnoud,¹⁴ M. Arov,⁶⁰ M. Arthaud,¹⁸ A. Askew,^{49,†} B. Åsman,⁴¹ A. C. S. Assis Jesus,³ O. Atramentov,⁴⁹ C. Avila,⁸ J. BackusMayes,⁸² F. Badaud,¹³ L. Bagby,⁵⁰ B. Baldin,⁵⁰ D. V. Bandurin,⁵⁹ P. Banerjee,²⁹ S. Banerjee,²⁹ E. Barberis,⁶³ A.-F. Barfuss,¹⁵ P. Bargassa,⁸⁰ P. Baringer,⁵⁸ J. Barreto,² J. F. Bartlett,⁵⁰ U. Bassler,¹⁸ D. Bauer,⁴³ S. Beale,⁶ A. Bean,⁵⁸ M. Begalli,³ M. Biegel,⁷³ C. Belanger-Champagne,⁴¹ L. Bellantoni,⁵⁰ A. Bellavance,⁵⁰ J. A. Benitez,⁶⁵ S. B. Beri,²⁷ G. Bernardi,¹⁷ R. Bernhard,²³ I. Bertram,⁴² M. Besançon,¹⁸ R. Beuselinck,⁴³ V. A. Bezzubov,³⁹ P. C. Bhat,⁵⁰ V. Bhatnagar,²⁷ G. Blazey,⁵² F. Blekman,⁴³ S. Blessing,⁴⁹ K. Bloom,⁶⁷ A. Boehnlein,⁵⁰ D. Boline,⁶² T. A. Bolton,⁵⁹ E. E. Boos,³⁸ G. Borissoy,⁴² T. Bose,⁷⁷ A. Brandt,⁷⁸ R. Brock,⁶⁵ G. Brooijmans,⁷⁰ A. Bross,⁵⁰ D. Brown,¹⁹ X. B. Bu,⁷ N. J. Buchanan,⁴⁹ D. Buchholz,⁵³ M. Buehler,⁸¹ V. Buescher,²² V. Bunichev,³⁸ S. Burdin,^{42,‡} T. H. Burnett,⁸² C. P. Buszello,⁴³ P. Calfayan,²⁵ B. Calpas,¹⁵ S. Calvet,¹⁶ J. Cammin,⁷¹ M. A. Carrasco-Lizarraga,³³ E. Carrera,⁴⁹ W. Carvalho,³ B. C. K. Casey,⁵⁰ H. Castilla-Valdez,³³ S. Chakrabarti,⁷² D. Chakraborty,⁵² K. M. Chan,⁵⁵ A. Chandra,⁴⁸ E. Cheu,⁴⁵ D. K. Cho,⁶² S. Choi,³² B. Choudhary,²⁸ L. Christofek,⁷⁷ T. Christoudias,⁴³ S. Cihangir,⁵⁰ D. Claes,⁶⁷ J. Clutter,⁵⁸ M. Cooke,⁵⁰ W. E. Cooper,⁵⁰ M. Corcoran,⁸⁰ F. Couderc,¹⁸ M.-C. Cousinou,¹⁵ S. Crépe-Renaudin,¹⁴ V. Cuplov,⁵⁹ D. Cutts,⁷⁷ M. Ćwiok,³⁰ H. da Motta,² A. Das,⁴⁵ G. Davies,⁴³ K. De,⁷⁸ S. J. de Jong,³⁵ E. De La Cruz-Burelo,³³ C. De Oliveira Martins,³ K. DeVaughan,⁶⁷ F. Déliot,¹⁸ M. Demarteau,⁵⁰ R. Demina,⁷¹ D. Denisov,⁵⁰ S. P. Denisov,³⁹ S. Desai,⁵⁰ H. T. Diehl,⁵⁰ M. Diesburg,⁵⁰ A. Dominguez,⁶⁷ T. Dorland,⁸² A. Dubey,²⁸ L. V. Dudko,³⁸ L. Duflot,¹⁶ S. R. Dugad,²⁹ D. Duggan,⁴⁹ A. Duperrin,¹⁵ S. Dutt,²⁷ J. Dyer,⁶⁵ A. Dyshkant,⁵² M. Eads,⁶⁷ D. Edmunds,⁶⁵ J. Ellison,⁴⁸ V. D. Elvira,⁵⁰ Y. Enari,⁷⁷ S. Eno,⁶¹ P. Ermolov,^{38,§§} M. Escalier,¹⁵ H. Evans,⁵⁴ A. Evdokimov,⁷³ V. N. Evdokimov,³⁹ A. V. Ferapontov,⁵⁹ T. Ferbel,^{61,71} F. Fiedler,²⁴ F. Filthaut,³⁵ W. Fisher,⁵⁰ H. E. Fisk,⁵⁰ M. Fortner,⁵² H. Fox,⁴² S. Fu,⁵⁰ S. Fuess,⁵⁰ T. Gadfort,⁷⁰ C. F. Galea,³⁵ C. Garcia,⁷¹ A. Garcia-Bellido,⁷¹ V. Gavrilov,³⁷ P. Gay,¹³ W. Geist,¹⁹ W. Geng,^{15,65} C. E. Gerber,⁵¹ Y. Gershtein,^{49,†} D. Gillberg,⁶ G. Ginter,⁷¹ B. Gómez,⁸ A. Goussiou,⁸² P. D. Grannis,⁷² H. Greenlee,⁵⁰ Z. D. Greenwood,⁶⁰ E. M. Gregores,⁴ G. Grenier,²⁰ Ph. Gris,¹³ J.-F. Grivaz,¹⁶ A. Grohsjean,²⁵ S. Grünendahl,⁵⁰ M. W. Grünewald,³⁰ F. Guo,⁷² J. Guo,⁷² G. Gutierrez,⁵⁰ P. Gutierrez,⁷⁵ A. Haas,⁷⁰ N. J. Hadley,⁶¹ P. Haefner,²⁵ S. Hagopian,⁴⁹ J. Haley,⁶⁸ I. Hall,⁶⁵ R. E. Hall,⁴⁷ L. Han,⁷ K. Harder,⁴⁴ A. Harel,⁷¹ J. M. Hauptman,⁵⁷ J. Hays,⁴³ T. Hebbeker,²¹ D. Hedin,⁵² J. G. Hegeman,³⁴ A. P. Heinson,⁴⁸ U. Heintz,⁶² C. Hensel,^{22,§} K. Herner,⁷² G. Hesketh,⁶³ M. D. Hildreth,⁵⁵ R. Hirosky,⁸¹ T. Hoang,⁴⁹ J. D. Hobbs,⁷² B. Hoeneisen,¹² M. Hohlfield,²² S. Hossain,⁷⁵ P. Houben,³⁴ Y. Hu,⁷² Z. Hubacek,¹⁰ N. Huske,¹⁷ V. Hynek,⁹ I. Iashvili,⁶⁹ R. Illingworth,⁵⁰ A. S. Ito,⁵⁰ S. Jabeen,⁶² M. Jaffré,¹⁶ S. Jain,⁷⁵ K. Jakobs,²³ C. Jarvis,⁶¹ R. Jesik,⁴³ K. Johns,⁴⁵ C. Johnson,⁷⁰ M. Johnson,⁵⁰ D. Johnston,⁶⁷ A. Jonckheere,⁵⁰ P. Jonsson,⁴³ A. Juste,⁵⁰ E. Kajfasz,¹⁵ D. Karmanov,³⁸ P. A. Kasper,⁵⁰ I. Katsanos,⁷⁰ V. Kaushik,⁷⁸ R. Kehoe,⁷⁹ S. Kermiche,¹⁵ N. Khalatyan,⁵⁰ A. Khanov,⁷⁶ A. Kharchilava,⁶⁹ Y. N. Kharzheev,³⁶ D. Khatidze,⁷⁰ T. J. Kim,³¹ M. H. Kirby,⁵³ M. Kirsch,²¹ B. Klima,⁵⁰ J. M. Kohli,²⁷ J.-P. Konrath,²³ A. V. Kozelov,³⁹ J. Kraus,⁶⁵ T. Kuhl,²⁴ A. Kumar,⁶⁹ A. Kupco,¹¹ T. Kurča,²⁰ V. A. Kuzmin,³⁸ J. Kvita,⁹ F. Lacroix,¹³ D. Lam,⁵⁵ S. Lammers,⁷⁰ G. Landsberg,⁷⁷ P. Lebrun,²⁰ W. M. Lee,⁵⁰ A. Leflat,³⁸ J. Lellouch,¹⁷ J. Li,^{78,§§} L. Li,⁴⁸ Q. Z. Li,⁵⁰ S. M. Lietti,⁵ J. K. Lim,³¹ J. G. R. Lima,⁵² D. Lincoln,⁵⁰ J. Linnemann,⁶⁵ V. V. Lipaev,³⁹ R. Lipton,⁵⁰ Y. Liu,⁷ Z. Liu,⁶ A. Lobodenko,⁴⁰ M. Lokajicek,¹¹ P. Love,⁴² H. J. Lubatti,⁸² R. Luna-Garcia,^{33,||} A. L. Lyon,⁵⁰ A. K. A. Maciel,² D. Mackin,⁸⁰ R. J. Madaras,⁴⁶ P. Mättig,²⁶ A. Magerkurth,⁶⁴ P. K. Mal,⁸² H. B. Malbouisson,³ S. Malik,⁶⁷ V. L. Malyshev,³⁶ Y. Maravin,⁵⁹ B. Martin,¹⁴ R. McCarthy,⁷² M. M. Meijer,³⁵ A. Melnitchouk,⁶⁶ L. Mendoza,⁸ P. G. Mercadante,⁵ M. Merkin,³⁸ K. W. Merritt,⁵⁰ A. Meyer,²¹ J. Meyer,^{22,§} J. Mitrevski,⁷⁰ R. K. Mommsen,⁴⁴ N. K. Mondal,²⁹ R. W. Moore,⁶ T. Moulík,⁵⁸ G. S. Muanza,¹⁵ M. Mulhearn,⁷⁰ O. Mundal,²² L. Mundim,³ E. Nagy,¹⁵ M. Naimuddin,⁵⁰ M. Narain,⁷⁷ H. A. Neal,⁶⁴ J. P. Negret,⁸ P. Neustroev,⁴⁰ H. Nilsen,²³ H. Nogima,³ S. F. Novaes,⁵ T. Nunnemann,²⁵ D. C. O'Neil,⁶ G. Obrant,⁴⁰ C. Ochando,¹⁶ D. Onoprienko,⁵⁹ N. Oshima,⁵⁰ N. Osman,⁴³ J. Osta,⁵⁵ R. Otec,¹⁰ G. J. Otero y Garzón,¹ M. Owen,⁴⁴ M. Padilla,⁴⁸ P. Padley,⁸⁰ M. Pangilinan,⁷⁷ N. Parashar,⁵⁶ S.-J. Park,^{22,§} S. K. Park,³¹ J. Parsons,⁷⁰ R. Partridge,⁷⁷ N. Parua,⁵⁴ A. Patwa,⁷³ G. Pawloski,⁸⁰ B. Penning,²³ M. Perfilov,³⁸ K. Peters,⁴⁴ Y. Peters,²⁶ P. Pétroff,¹⁶ M. Petteni,⁴³ R. Piegaia,¹ J. Piper,⁶⁵ M.-A. Pleier,²² P. L. M. Podesta-Lerma,^{33,||} V. M. Podstavkov,⁵⁰ Y. Pogorelov,⁵⁵ M.-E. Pol,² P. Polozov,³⁷ B. G. Pope,⁶⁵ A. V. Popov,³⁹ C. Potter,⁶ W. L. Prado da Silva,³ H. B. Prosper,⁴⁹ S. Protopopescu,⁷³ J. Qian,⁶⁴ A. Quadt,^{22,§} B. Quinn,⁶⁶ A. Rakitine,⁴² M. S. Rangel,² K. Ranjan,²⁸ P. N. Ratoff,⁴² P. Renkel,⁷⁹ P. Rich,⁴⁴ M. Rijssenbeek,⁷² I. Ripp-Baudot,¹⁹ F. Rizatdinova,⁷⁶ S. Robinson,⁴³ R. F. Rodrigues,³ M. Rominsky,⁷⁵ C. Royon,¹⁸

P. Rubinov,⁵⁰ R. Ruchti,⁵⁵ G. Safronov,³⁷ G. Sajot,¹⁴ A. Sánchez-Hernández,³³ M. P. Sanders,¹⁷ B. Sanghi,⁵⁰ G. Savage,⁵⁰ L. Sawyer,⁶⁰ T. Scanlon,⁴³ D. Schaile,²⁵ R. D. Schamberger,⁷² Y. Scheglov,⁴⁰ H. Schellman,⁵³ T. Schliephake,²⁶ S. Schlobohm,⁸² C. Schwanenberger,⁴⁴ R. Schwienhorst,⁶⁵ J. Sekaric,⁴⁹ H. Severini,⁷⁵ E. Shabalina,⁵¹ M. Shamim,⁵⁹ V. Shary,¹⁸ A. A. Shchukin,³⁹ R. K. Shivpuri,²⁸ V. Siccaldi,¹⁹ V. Simak,¹⁰ V. Sirotenko,⁵⁰ P. Skubic,⁷⁵ P. Slattery,⁷¹ D. Smirnov,⁵⁵ G. R. Snow,⁶⁷ J. Snow,⁷⁴ S. Snyder,⁷³ S. Söldner-Rembold,⁴⁴ L. Sonnenschein,¹⁷ A. Sopczak,⁴² M. Sosebee,⁷⁸ K. Soustruznik,⁹ B. Spurlock,⁷⁸ J. Stark,¹⁴ V. Stolin,³⁷ D. A. Stoyanova,³⁹ J. Strandberg,⁶⁴ S. Strandberg,⁴¹ M. A. Strang,⁶⁹ E. Strauss,⁷² M. Strauss,⁷⁵ R. Ströhmer,²⁵ D. Strom,⁵³ L. Stutte,⁵⁰ S. Sumowidagdo,⁴⁹ P. Svoisky,³⁵ A. Sznajder,³ A. Tanasijczuk,¹ W. Taylor,⁶ B. Tiller,²⁵ F. Tissandier,¹³ M. Titov,¹⁸ V. V. Tokmenin,³⁶ I. Torchiani,²³ D. Tsybychev,⁷² B. Tuchming,¹⁸ C. Tully,⁶⁸ P. M. Tuts,⁷⁰ R. Unalan,⁶⁵ L. Uvarov,⁴⁰ S. Uvarov,⁴⁰ S. Uzunyan,⁵² B. Vachon,⁶ P. J. van den Berg,³⁴ R. Van Kooten,⁵⁴ W. M. van Leeuwen,³⁴ N. Varelas,⁵¹ E. W. Varnes,⁴⁵ I. A. Vasilyev,³⁹ P. Verdier,²⁰ L. S. Vertogradov,³⁶ M. Verzocchi,⁵⁰ D. Vilanova,¹⁸ F. Villeneuve-Seguié,⁴³ P. Vint,⁴³ P. Vokac,¹⁰ M. Voutilainen,^{67,**} R. Wagner,⁶⁸ H. D. Wahl,⁴⁹ M. H. L. S. Wang,⁵⁰ J. Warchol,⁵⁵ G. Watts,⁸² M. Wayne,⁵⁵ G. Weber,²⁴ M. Weber,^{50,††} L. Welty-Rieger,⁵⁴ A. Wenger,^{23,‡‡} N. Wermes,²² M. Wetstein,⁶¹ A. White,⁷⁸ D. Wicke,²⁶ M. R. J. Williams,⁴² G. W. Wilson,⁵⁸ S. J. Wimpenny,⁴⁸ M. Wobisch,⁶⁰ D. R. Wood,⁶³ T. R. Wyatt,⁴⁴ Y. Xie,⁷⁷ C. Xu,⁶⁴ S. Yacoub,⁵³ R. Yamada,⁵⁰ W.-C. Yang,⁴⁴ T. Yasuda,⁵⁰ Y. A. Yatsunenko,³⁶ Z. Ye,⁵⁰ H. Yin,⁷ K. Yip,⁷³ H. D. Yoo,⁷⁷ S. W. Youn,⁵³ J. Yu,⁷⁸ C. Zeitnitz,²⁶ S. Zelitch,⁸¹ T. Zhao,⁸² B. Zhou,⁶⁴ J. Zhu,⁷² M. Zielinski,⁷¹ D. Zieminska,⁵⁴ L. Zivkovic,⁷⁰ V. Zutshi,⁵² and E. G. Zverev³⁸

(D0 Collaboration)

¹Universidad de Buenos Aires, Buenos Aires, Argentina²LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil³Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil⁴Universidade Federal do ABC, Santo André, Brazil⁵Instituto de Física Teórica, Universidade Estadual Paulista, São Paulo, Brazil⁶University of Alberta, Edmonton, Alberta, Canada,

Simon Fraser University, Burnaby, British Columbia, Canada,

York University, Toronto, Ontario, Canada,

and McGill University, Montreal, Quebec, Canada

⁷University of Science and Technology of China, Hefei, People's Republic of China⁸Universidad de los Andes, Bogotá, Colombia⁹Center for Particle Physics, Charles University, Prague, Czech Republic¹⁰Czech Technical University, Prague, Czech Republic¹¹Center for Particle Physics, Institute of Physics, Academy of Sciences of the Czech Republic, Prague, Czech Republic¹²Universidad San Francisco de Quito, Quito, Ecuador¹³LPC, Université Blaise Pascal, CNRS/IN2P3, Clermont, France¹⁴LPSC, Université Joseph Fourier Grenoble 1, CNRS/IN2P3, Institut National Polytechnique de Grenoble, Grenoble, France¹⁵CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France¹⁶LAL, Université Paris-Sud, IN2P3/CNRS, Orsay, France¹⁷LPNHE, IN2P3/CNRS, Universités Paris VI and VII, Paris, France¹⁸CEA, Irfu, SPP, Saclay, France¹⁹IPHC, Université Louis Pasteur, CNRS/IN2P3, Strasbourg, France²⁰IPNL, Université Lyon 1, CNRS/IN2P3, Villeurbanne, France and Université de Lyon, Lyon, France²¹III. Physikalisches Institut A, RWTH Aachen University, Aachen, Germany²²Physikalisches Institut, Universität Bonn, Bonn, Germany²³Physikalisches Institut, Universität Freiburg, Freiburg, Germany²⁴Institut für Physik, Universität Mainz, Mainz, Germany²⁵Ludwig-Maximilians-Universität München, München, Germany²⁶Fachbereich Physik, University of Wuppertal, Wuppertal, Germany²⁷Panjab University, Chandigarh, India²⁸Delhi University, Delhi, India²⁹Tata Institute of Fundamental Research, Mumbai, India³⁰University College Dublin, Dublin, Ireland³¹Korea Detector Laboratory, Korea University, Seoul, Korea³²SungKyunKwan University, Suwon, Korea³³CINVESTAV, Mexico City, Mexico³⁴FOM-Institute NIKHEF and University of Amsterdam/NIKHEF, Amsterdam, The Netherlands

- ³⁵Radboud University Nijmegen/NIKHEF, Nijmegen, The Netherlands
³⁶Joint Institute for Nuclear Research, Dubna, Russia
³⁷Institute for Theoretical and Experimental Physics, Moscow, Russia
³⁸Moscow State University, Moscow, Russia
³⁹Institute for High Energy Physics, Protvino, Russia
⁴⁰Petersburg Nuclear Physics Institute, St. Petersburg, Russia
⁴¹Lund University, Lund, Sweden,
Royal Institute of Technology and Stockholm University, Stockholm, Sweden,
and Uppsala University, Uppsala, Sweden
⁴²Lancaster University, Lancaster, United Kingdom
⁴³Imperial College, London, United Kingdom
⁴⁴University of Manchester, Manchester, United Kingdom
⁴⁵University of Arizona, Tucson, Arizona 85721, USA
⁴⁶Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA
⁴⁷California State University, Fresno, California 93740, USA
⁴⁸University of California, Riverside, California 92521, USA
⁴⁹Florida State University, Tallahassee, Florida 32306, USA
⁵⁰Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA
⁵¹University of Illinois at Chicago, Chicago, Illinois 60607, USA
⁵²Northern Illinois University, DeKalb, Illinois 60115, USA
⁵³Northwestern University, Evanston, Illinois 60208, USA
⁵⁴Indiana University, Bloomington, Indiana 47405, USA
⁵⁵University of Notre Dame, Notre Dame, Indiana 46556, USA
⁵⁶Purdue University Calumet, Hammond, Indiana 46323, USA
⁵⁷Iowa State University, Ames, Iowa 50011, USA
⁵⁸University of Kansas, Lawrence, Kansas 66045, USA
⁵⁹Kansas State University, Manhattan, Kansas 66506, USA
⁶⁰Louisiana Tech University, Ruston, Louisiana 71272, USA
⁶¹University of Maryland, College Park, Maryland 20742, USA
⁶²Boston University, Boston, Massachusetts 02215, USA
⁶³Northeastern University, Boston, Massachusetts 02115, USA
⁶⁴University of Michigan, Ann Arbor, Michigan 48109, USA
⁶⁵Michigan State University, East Lansing, Michigan 48824, USA
⁶⁶University of Mississippi, University, Mississippi 38677, USA
⁶⁷University of Nebraska, Lincoln, Nebraska 68588, USA
⁶⁸Princeton University, Princeton, New Jersey 08544, USA
⁶⁹State University of New York, Buffalo, New York 14260, USA
⁷⁰Columbia University, New York, New York 10027, USA
⁷¹University of Rochester, Rochester, New York 14627, USA
⁷²State University of New York, Stony Brook, New York 11794, USA
⁷³Brookhaven National Laboratory, Upton, New York 11973, USA
⁷⁴Langston University, Langston, Oklahoma 73050, USA
⁷⁵University of Oklahoma, Norman, Oklahoma 73019, USA
⁷⁶Oklahoma State University, Stillwater, Oklahoma 74078, USA
⁷⁷Brown University, Providence, Rhode Island 02912, USA
⁷⁸University of Texas, Arlington, Texas 76019, USA
⁷⁹Southern Methodist University, Dallas, Texas 75275, USA
⁸⁰Rice University, Houston, Texas 77005, USA
⁸¹University of Virginia, Charlottesville, Virginia 22901, USA
⁸²University of Washington, Seattle, Washington 98195, USA
(Received 14 January 2009; published 12 June 2009)

We present a search for a narrow resonance in the inclusive diphoton final state using $\sim 2.7 \text{ fb}^{-1}$ of data collected with the D0 detector at the Fermilab Tevatron $p\bar{p}$ Collider. We observe good agreement between the data and the background prediction, and set the first 95% C.L. upper limits on the production cross section times the branching ratio for decay into a pair of photons for resonance masses between 100 and 150 GeV. This search is also interpreted in the context of several models of electroweak symmetry breaking with a Higgs boson decaying into two photons.

At a hadron collider, diphoton ($\gamma\gamma$) production allows detailed studies of the standard model (SM) [1], as well as searches for new phenomena, such as new heavy resonances [2], extra spatial dimensions, or cascade decays of heavy new particles [3]. Within the SM, continuum $\gamma\gamma + X$ production is characterized by a steeply falling $\gamma\gamma$ mass ($M_{\gamma\gamma}$) spectrum, on top of which a heavy resonance decaying into $\gamma\gamma$ can potentially be observed. In particular, this is considered one of the most promising discovery channels for a light SM Higgs boson at the LHC [4], despite the small branching ratio of $B(H \rightarrow \gamma\gamma) \sim 0.2\%$ for $110 < M_H < 140$ GeV [5,6]. At the Tevatron, the dominant SM Higgs boson production mechanism is gluon fusion (GF) ($gg \rightarrow H$), followed by associated production with a W or Z boson ($q\bar{q}' \rightarrow VH$, $V = W, Z$), and vector boson fusion (VBF) ($VV \rightarrow H$) [7–9]. While the SM Higgs production rate at the Tevatron is not sufficient to observe it in the $\gamma\gamma$ mode, the Hgg and $H\gamma\gamma$ couplings, being loop-mediated, are particularly sensitive to new physics effects. Furthermore, in some models beyond the SM [2], $B(H \rightarrow \gamma\gamma)$ can be enhanced significantly relative to the SM prediction.

In this Letter, we present a search for a narrow resonance in the $M_{\gamma\gamma}$ spectrum using a data sample collected by the D0 detector [10] at the Fermilab Tevatron Collider. The selection of an inclusive $\gamma\gamma + X$ sample and the use of the $M_{\gamma\gamma}$ spectrum make the results of this search quasi-model independent. We use the SM Higgs boson (H_{SM}) with $H \rightarrow \gamma\gamma$ as a reference model, resulting in the first such search at the Tevatron, and a forerunner to similar planned searches at the LHC. Additionally, we consider other models of electroweak symmetry breaking (EWSB) with enhanced $B(H \rightarrow \gamma\gamma)$ [2], as a consequence of suppressed Higgs couplings to either (i) all fermions (known as bosonic or fermiophobic Higgs boson, H_f), (ii) only down-type fermions (H_u , as it gives mass to up-type fermions), or (iii) only top and bottom quarks (known as electroweak Higgs boson, H_{EW}). All models considered have SM-like production cross sections, with the exception of the H_f and H_{EW} models, where GF is absent or has a negligibly small cross section.

The subdetectors most relevant to this analysis are the central tracking system, composed of a silicon microstrip tracker (SMT) and a central fiber tracker (CFT) embedded in a 2 T solenoidal magnetic field, the central preshower (CPS), and the liquid-argon and uranium sampling calorimeter. The CPS is located immediately before the inner layer of the calorimeter and is formed of one radiation length of absorber followed by several layers of scintillating strips. The calorimeter consists of a central section with coverage of $|\eta| < 1.1$ [11], and two end calorimeters covering up to $|\eta| \simeq 4.2$. The electromagnetic (EM) section of the calorimeter is segmented into four longitudinal layers (EM*i*, $i = 1, 4$) with transverse segmentation of $\Delta\eta \times \Delta\phi = 0.1 \times 0.1$ [11], except in EM3, where it is $0.05 \times$

0.05 . The calorimeter is well suited for a precise measurement of electron and photon energies, providing a resolution of $\sim 3.6\%$ at energies of ~ 50 GeV. The data used in this analysis were collected using triggers requiring at least two clusters of energy in the EM calorimeter and correspond to an integrated luminosity of $2.7 \pm 0.2 \text{ fb}^{-1}$ [12].

Events are selected by requiring at least two photon candidates with transverse momentum $p_T > 25$ GeV and $|\eta| < 1.1$, for which the trigger requirements are fully efficient. The photon p_T is computed with respect to the reconstructed event primary vertex (PV) with the highest number of associated tracks, which is required to be within 60 cm of the geometrical center of the detector along the beam axis. The PV reconstruction efficiency in $\gamma\gamma + X$ events is $\sim 98\%$, with $\sim 95\%$ probability to match the true vertex. Photons are selected from EM clusters reconstructed within a cone with radius $\mathcal{R} = \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2} = 0.2$ by requiring (i) $\geq 97\%$ of the cluster energy is deposited in the EM calorimeter, (ii) the calorimeter isolation variable $I = [E_{\text{tot}}(0.4) - E_{\text{EM}}(0.2)]/E_{\text{EM}}(0.2) < 0.1$, where $E_{\text{tot}}(\mathcal{R})$ ($E_{\text{EM}}(\mathcal{R})$) is the total (EM) energy in a cone of radius \mathcal{R} , (iii) the energy-weighted shower width in the $r - \phi$ plane in EM3 is $< \sqrt{14}$ cm, and (iv) the scalar sum of the p_T of all tracks ($p_{\text{Trk}}^{\text{sum}}$) originating from the primary vertex in an annulus of $0.05 < \mathcal{R} < 0.4$ around the cluster is < 2 GeV. To suppress electrons misidentified as photons, the EM clusters are required to not be spatially matched to tracker activity, either a reconstructed track, or a density of hits in the SMT and CFT consistent with that of an electron [13]. To suppress jets misidentified as photons, a neural network (NN) is trained using a set of variables sensitive to differences between photons and jets in the tracker activity and in the energy deposits in the calorimeter and CPS: $p_{\text{Trk}}^{\text{sum}}$, the numbers of cells above a threshold in EM1 within $\mathcal{R} < 0.2$ and $0.2 < \mathcal{R} < 0.4$ of the EM cluster, the number of CPS clusters within $\mathcal{R} < 0.1$ of the EM cluster, and the squared-energy-weighted width of the energy deposit in the CPS. The NN is trained using $\gamma\gamma$ and dijet Monte Carlo (MC) samples and its performance is verified using a data sample of $Z \rightarrow \ell^+ \ell^- \gamma$ ($\ell = e, \mu$) events. Figure 1(a) compares the NN output (O_{NN}) spectrum for photons and jets. Photon candidates are required to have $O_{\text{NN}} > 0.1$, which is $\sim 98\%$ efficient for real photons and rejects $\sim 50\%$ of misidentified jets. Finally, $M_{\gamma\gamma}$, computed from the two highest p_T photons, is required to be > 60 GeV. In total, 5608 events are selected in data.

All MC samples used in this analysis are generated using PYTHIA [14] with CTEQ6L [15] parton distribution functions (PDFs), and processed through a GEANT-based [16] simulation of the D0 detector and the same reconstruction software as the data. Signal samples are generated separately for GF, VH, and VBF production and normalized using the theoretical cross sections [7–9] and branching ratio predictions from HDECAY [5].




FIG. 1 (color online). (a) Normalized O_{NN} spectrum for photons and jets. (b) $M_{\gamma\gamma}$ spectrum in data (points) compared to the total background parametrization (solid line), including the DDP contribution derived via a sideband fit, and the total background prediction (dashed line) inside the search region for $M_H = 130$ GeV. The inset figure compares the data to the total background prediction inside the search region including its 1 standard deviation (s.d.) uncertainty band, as well as the expected H_{SM} signal scaled by a factor of 50. (c) Observed and expected 95% C.L. upper limits on $\sigma \times B$ as a function of M_H . Also shown are the predictions for $\sigma \times B$ in the different EWSB scenarios discussed in the text.

This analysis is affected by instrumental backgrounds such as $\gamma + \text{jet}$, dijet, and $Z/\gamma^* \rightarrow e^+e^-$ (ZDY) production, with jets or electrons misidentified as photons, as well as an irreducible background from direct diphoton production (DDP). All backgrounds, except for ZDY, are estimated directly from data.

The ZDY background is estimated using the MC simulation, normalized to the next-to-next-to-leading-order cross section [17]. The selection efficiencies determined by the MC simulation are corrected to the corresponding values measured in the data. On average each electron has a 2% probability to satisfy the photon selection criteria, mainly due to the inefficiency of the track-match veto requirements. The total contribution from ZDY is estimated to be 88 ± 10 events.

Backgrounds due to $\gamma + \text{jet}$ and dijet events are directly estimated from data by using a 4×4 matrix background estimation method [18]. After final event selection, a tightened O_{NN} requirement ($O_{NN} > 0.75$) is used to classify the events into four categories depending on whether the two highest- p_T photons, only the leading photon, only the trailing photon, or neither of the two photons satisfy this requirement. The corresponding

numbers of events, after subtraction of the estimated ZDY contributions, are denoted as N_{pp} , N_{pf} , N_{fp} , and N_{ff} . The different relative efficiency of the $O_{NN} > 0.75$ requirement between real photons and jets allows the estimation of the sample composition by solving a linear system of equations: $(N_{pp}, N_{pf}, N_{fp}, N_{ff})^T = \mathcal{E}(N_{\gamma\gamma}, N_{\gamma j}, N_{j\gamma}, N_{jj})^T$, where $N_{\gamma\gamma}$ (N_{jj}) is the number of $\gamma\gamma$ (dijet) events and $N_{\gamma j}$ ($N_{j\gamma}$) is the number of $\gamma + \text{jet}$ events with the leading (trailing) cluster as the photon. The 4×4 matrix \mathcal{E} contains the efficiency terms (parametrized as a function of $|\eta|$), estimated in photon and jet MC samples and validated in data. The estimated sample composition is $N_{\gamma\gamma} = 3155 \pm 125(\text{stat})$, $N_{\gamma j + j\gamma} = 1680 \pm 149(\text{stat})$, and $N_{jj} = 685 \pm 93(\text{stat})$. The shape of the $M_{\gamma\gamma}$ spectrum for the sum of the $\gamma + \text{jet}$ and dijet backgrounds is obtained from an independent control data sample by requiring $O_{NN} < 0.1$ for one of the photon candidates, and is parametrized with an exponential function. The resulting shape is found to be in excellent agreement with that derived by directly applying the 4×4 matrix method bin by bin in the final selected sample, but has smaller statistical fluctuations, especially in the high $M_{\gamma\gamma}$ region.

TABLE I. Numbers of selected events in data, expected backgrounds, expected H_{SM} signal, and signal acceptance (for each production mechanism: GF, VH, VBF), in the search region for different M_H values. The expected signal includes contributions from GF, VH, and VBF processes, the latter two representing $\sim 21\%$ – 24% of the total signal.

M_H (GeV)	100	110	120	130	140	150
$Z/\gamma^* \rightarrow e^+e^-$	55 ± 7	17 ± 3	6 ± 2	5 ± 1	4 ± 1	3 ± 1
$\gamma\gamma$	742 ± 62	481 ± 42	324 ± 34	236 ± 30	161 ± 28	124 ± 22
$\gamma j + jj$	540 ± 66	319 ± 39	204 ± 25	133 ± 16	89 ± 11	61 ± 8
Total background	1337 ± 29	817 ± 26	534 ± 19	374 ± 12	254 ± 7	188 ± 5
Data	1385	827	544	357	270	202
H_{SM} signal	1.62 ± 0.11	1.61 ± 0.11	1.51 ± 0.10	1.26 ± 0.08	0.90 ± 0.06	0.54 ± 0.04
Acceptance (%)	19.9, 18.8, 20.3	20.4, 19.9, 21.6	21.0, 20.6, 22.3	21.5, 21.2, 22.9	21.8, 22.0, 23.5	22.1, 22.2, 24.1

After subtraction of the ZDY, γ + jet, and dijet background contributions, the $M_{\gamma\gamma}$ spectrum is examined for the presence of a narrow resonance. For each assumed M_H value (between 100 and 150 GeV, in steps of 5 GeV), the search region is defined to be $(M_H - 15 \text{ GeV}, M_H + 15 \text{ GeV})$, where 15 GeV corresponds to about five times the expected $M_{\gamma\gamma}$ resolution. The DDP background is estimated by performing a sideband fit to the $M_{\gamma\gamma}$ spectrum in the 70 to 200 GeV range (this excludes the search region) using an exponential function [see Fig. 1(b)]. Such a parametrization has been validated using a next-to-leading-order calculation for this process [19].

Systematic uncertainties affecting the normalization and shape of the $M_{\gamma\gamma}$ spectrum are estimated for both signal and backgrounds. Uncertainties affecting the ZDY background normalization include integrated luminosity (6.1%), electron misidentification rate (14.3%), and ZDY cross section (3.9%). Such uncertainties are propagated, via the 4×4 matrix method, to the estimated normalization of the γ + jet and dijet background contributions, affected in addition by the uncertainty on the $O_{\text{NN}} > 0.75$ selection efficiency for photons (2%) and jets (10%). The uncertainty in the shape of the γ + jet and dijet $M_{\gamma\gamma}$ spectrum is given by the statistics of the control data sample used to parametrize it. The above uncertainties, as well as the statistical uncertainties of the sideband fitting method, result in systematic uncertainties in the normalization and shape of the DDP background contribution. Uncertainties affecting the signal normalization include integrated luminosity (6.1%), acceptance due to the photon identification efficiency (6.8%), and PDFs (1.7%–2.2%) [15]. Finally, the location of the peak in the $M_{\gamma\gamma}$ spectrum for signal is affected by the uncertainty in the relative data-to-MC photon energy scale (0.6%).

Table I shows the number of events in the data, expected background, and expected H_{SM} signal in six different search regions. The inset in Fig. 1(b) illustrates the $M_{\gamma\gamma}$ spectrum in the search region for $M_H = 130 \text{ GeV}$, found to be in good agreement with the background prediction. The $M_{\gamma\gamma}$ spectrum in the search region is used to derive upper limits on the production cross section times branching ratio for $H \rightarrow \gamma\gamma$ ($\sigma \times B$) as a function of M_H . The SM prediction for the ratio of the production cross sections for the three signal production mechanisms is assumed. Limits are calculated at the 95% C.L. using the modified frequentist approach with a Poisson log-likelihood ratio test statistic [20,21]. The impact of systematic uncertainties is incorporated via convolution of the Poisson probability distributions for signal and background with Gaussian distributions corresponding to the different sources of systematic uncertainty. The correlations in systematic uncertainties are maintained between signal and backgrounds.

The resulting limits on $\sigma \times B$ are given in Table II, and displayed in Fig. 1(c). Although the SM Higgs boson is

TABLE II. Observed and expected 95% C.L. upper limits on $\sigma \times B$ (in fb) for different M_H values. The expected limit is defined as the median of the distribution of limits in background-only pseudoexperiments.

M_H (GeV)	100	105	110	115	120	125	130	135	140	145	150
Exp. $\sigma \times B$	88	78	71	63	58	53	49	45	41	39	36
Obs. $\sigma \times B$	78	110	112	76	50	46	42	53	56	66	52

used as a reference model, the fact that the signal acceptance is found to be almost independent of the production mechanism (see Table I) makes the estimated limits applicable to other models of new physics with a narrow resonance decaying into $\gamma\gamma$. In the context of models of EWSB with enhanced $B(H \rightarrow \gamma\gamma)$, the current search excludes a H_f boson with $M_H < 101 \text{ GeV}$ at 95% C.L., improving (slightly) upon previous results at the Tevatron [22]. While none of the other EWSB scenarios explored can currently be excluded, the expected sensitivity is within less than a factor of 4 of the prediction for the H_u model for $M_H < 110 \text{ GeV}$, and only a factor ~ 20 above the SM prediction for $115 \leq M_H \leq 130 \text{ GeV}$. As a result, this search contributes to the overall sensitivity of the SM Higgs boson search at the Tevatron from the combination of multiple channels [23]. Assuming the same integrated luminosity in all channels and a single Tevatron experiment, this analysis is expected to improve the combined upper limit on the SM Higgs production cross section by $\sim 5\%$ for $115 \leq M_H \leq 130 \text{ GeV}$. Finally, this search is used to derive 95% C.L. upper limits on $B(H \rightarrow \gamma\gamma)$ between 14.1% and 33.9% for M_H in the range 100–150 GeV, in the case of models where the Higgs boson does not couple to the top quark. Conversely, for models where the GF production mode is available, this inclusive search allows improvement of the upper limits on $B(H \rightarrow \gamma\gamma)$ to 3.4%–7.2% in the same mass range. These represent the most stringent limits on $B(H \rightarrow \gamma\gamma)$ for M_H in the range 100–150 GeV to date, significantly improving upon previous results from LEP and the Tevatron [22].

In summary, we have performed an inclusive search for a narrow resonance with mass between 100 and 150 GeV decaying into $\gamma\gamma$ at the Tevatron. This channel is used to increase the overall sensitivity of the SM Higgs boson search program at the Tevatron [23] and allows the probe of new physics models predicting an enhanced rate for $H \rightarrow \gamma\gamma$.

We thank the staffs at Fermilab and collaborating institutions, and acknowledge support from the DOE and NSF (U.S.); CEA and CNRS/IN2P3 (France); FASI, Rosatom and RFBR (Russia); CNPq, FAPERJ, FAPESP and FUNDUNESP (Brazil); DAE and DST (India); Colciencias (Colombia); CONACyT (Mexico); KRF and KOSEF (Korea); CONICET and UBACyT (Argentina); FOM (The Netherlands); STFC (United Kingdom); MSMT and GACR (Czech Republic); CRC Program,

CFI, NSERC and WestGrid Project (Canada); BMBF and DFG (Germany); SFI (Ireland); The Swedish Research Council (Sweden); CAS and CNSF (China); and the Alexander von Humboldt Foundation (Germany).

*Visitor from Augustana College, Sioux Falls, SD, USA.

[†]Visitor from Rutgers University, Piscataway, NJ, USA.

[‡]Visitor from The University of Liverpool, Liverpool, UK.

[§]Visitor from II. Physikalisches Institut, Georg-August-University, Göttingen, Germany.

^{||}Visitor from Centro de Investigacion en Computacion-IPN, Mexico City, Mexico.

[¶]Visitor from ECFM, Universidad Autonoma de Sinaloa, Culiacán, Mexico.

**Visitor from Helsinki Institute of Physics, Helsinki, Finland.

^{††}Visitor from Universität Bern, Bern, Switzerland.

^{‡‡}Visitor from Universität Zürich, Zürich, Switzerland.

^{§§}Deceased.

- [1] C. Balazs, E. L. Berger, P. Nadolsky, and C.-P. Yuan, *Phys. Rev. D* **76**, 013009 (2007), and references therein.
- [2] S. Mrenna and J. Wells, *Phys. Rev. D* **63**, 015006 (2000), and references therein.
- [3] See, e.g., G. F. Giudice and R. Rattazzi, *Phys. Rep.* **322**, 419 (1999); M. C. Kumar, P. Mathews, V. Ravindran, and A. Tripathi, *Phys. Lett. B* **672**, 45 (2009), and references therein.
- [4] G. Aad *et al.* (ATLAS Collaboration), arXiv:0901.0512; G. L. Bayatian *et al.* (CMS Collaboration), *J. Phys. G* **34**, 095 (2007).
- [5] A. Djouadi, J. Kalinowski, and M. Spira, *Comput. Phys. Commun.* **108**, 56 (1998).
- [6] Throughout this Letter we adopt units where $c = 1$.
- [7] S. Catani *et al.*, *J. High Energy Phys.* **07** (2003) 028.
- [8] U. Aglietti *et al.*, arXiv:hep-ph/0610033.
- [9] K. A. Assamagan *et al.*, arXiv:hep-ph/0406152.
- [10] V. M. Abazov *et al.* (D0 Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **565**, 463 (2006).
- [11] Pseudorapidity is defined as $\eta = -\ln[\tan(\theta/2)]$, where θ is the polar angle relative to the proton beam direction. ϕ is defined as the azimuthal angle in the plane transverse to the proton beam direction.
- [12] T. Andeen *et al.*, Report No. FERMILAB-TM-2365, 2007.
- [13] V. M. Abazov *et al.* (D0 Collaboration), *Phys. Lett. B* **659**, 856 (2008).
- [14] T. Sjöstrand *et al.*, *Comput. Phys. Commun.* **135**, 238 (2001); we use PYTHIA version v6.323.
- [15] J. Pumplin *et al.*, *J. High Energy Phys.* **07** (2002) 012; D. Stump *et al.*, *J. High Energy Phys.* **10** (2003) 046.
- [16] R. Brun and F. Carminati, CERN Program Library Long Writeup Report No. W5013, 1993; we use GEANT version v3.21.
- [17] R. Hamberg, W. L. van Neerven, and T. Matsuura, *Nucl. Phys.* **B359**, 343 (1991); **644**, 403 (2002).
- [18] D. Acosta *et al.* (CDF Collaboration), *Phys. Rev. Lett.* **95**, 022003 (2005).
- [19] T. Binoth *et al.*, *Eur. Phys. J. C* **16**, 311 (2000).
- [20] T. Junk, *Nucl. Instrum. Methods Phys. Res., Sect. A* **434**, 435 (1999); A. Read, CERN Report No. 2000-005, 2000.
- [21] W. Fisher, Report No. FERMILAB-TM-2386-E, 2006.
- [22] V. M. Abazov *et al.* (D0 Collaboration), *Phys. Rev. Lett.* **101**, 051801 (2008), and references therein.
- [23] TEVNPH Working Group, for the CDF and D0 Collaborations, arXiv:0903.4001.