



PERGAMON

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

SCIENCE @ DIRECT®

Deep-Sea Research I 50 (2003) 457–494

DEEP-SEA RESEARCH  
PART I

[www.elsevier.com/locate/dsr](http://www.elsevier.com/locate/dsr)

# Seasonal and interannual variability of benthic foraminiferal faunas at 550 m depth in the Bay of Biscay

C. Fontanier<sup>a,\*</sup>, F.J. Jorissen<sup>b,c</sup>, G. Chaillou<sup>a</sup>, C. David<sup>a</sup>, P. Anschutz<sup>a</sup>, V. Lafon<sup>a</sup>

<sup>a</sup>Department of Geology and Oceanography, Bordeaux University, CNRS UMR 58 05 EPOC, Avenue des Facultés, 33405 Talence Cedex, France

<sup>b</sup>Department for the Study of Recent and Fossil Bio-Indicators, Angers University, UPRES EA 2644, 2 Boulevard Lavoisier, 49045 Angers Cedex, France

<sup>c</sup>Laboratory for the Study of Marine Bio-indicators (LEBIM), 85350 Ile d'Yeu, France

Received 25 February 2002; received in revised form 27 November 2002; accepted 12 December 2002

## Abstract

Live benthic foraminiferal faunas were sampled 10 times between October 1997 and April 2000 at a 550 m depth open-slope station in the Bay of Biscay. Duplicate cores for 5 samplings allow distinguishing between spatial and temporal variability of the foraminiferal faunas. Although spatial patchiness of the foraminiferal faunas is substantial, especially in the 63–150 µm fraction, the temporal variability appears to be larger. The foraminiferal patterns are compared with surface water primary production as assessed by the study of available SeaWiFS satellite images. In the study area, the primary production regime is marked by a pulselike and prolonged spring bloom and possibly a short fall bloom. Such periods of elevated chlorophyll-*a* concentration are followed, after a delay of about 4–6 weeks, by a strong frequency increase of the most opportunistic taxa of benthic foraminifera. Surprisingly, no change of bottom and interstitial water oxygenation and of redox conditions within the sediment is recorded. The small taxa *Epistominella exigua*, *Reophax guttiferus*, *Bolivina spathulata*, *Cassidulina carinata* and *Nuttallides pusillus* appear to respond first to a labile organic matter input, by a reproductive event marked by a strong patchy spatial distribution hypothetically resulting of the spatial heterogeneity of organic matter deposits. *Uvigerina peregrina* and *Uvigerina mediterranea*, the most opportunistic larger taxa, strongly dominate the > 150 µm fraction during eutrophic periods (spring and fall blooms). Intermediate and deep infaunal taxa seem to depend less on fresh organic matter input, even if a small frequency increases are recorded in the > 150 µm fraction during the most productive periods; *Globobulimina affinis* and *Melonis barleeanus* show reproductive events in rather shallow sediment layers in the more oligotrophic periods of the year. A conceptual model explains the increasing delay in the response to important phytoplankton bloom periods for the successive benthic ecosystem compartments.

© 2003 Published by Elsevier Science Ltd.

**Keywords:** Benthic foraminifera; Microhabitat; Seasonality; Patchiness; Opportunism; Organic flux

## 1. Introduction

Mid-latitude primary production regimes are marked by a seasonal and interannual alternation

\*Corresponding author.

E-mail address: [c.fontanier@epoc.u-bordeaux.fr](mailto:c.fontanier@epoc.u-bordeaux.fr)  
(C. Fontanier).

of algal blooms and low productivity periods (Pfannkuche and Thiel, 1987; Parsons and Lalli, 1988; Berger and Wefer, 1990). This intermittence and seasonality of the surface water primary production is responsible for important fluctuations of the exported organic matter flux (Billet et al., 1983; Pfannkuche and Thiel, 1987; Berger and Wefer, 1990; Lohrenz et al., 1992). Since the biomass of deep-sea foraminifera is closely related to the flux of organic carbon to the seafloor (Altenbach, 1985, 1988; Altenbach and Sarnthein, 1989; Herguera and Berger, 1991; Jorissen et al., 1998; Loubere and Fariduddin, 1999; De Rijk et al., 2000; Fontanier et al., 2002; Morigi et al., 2001), the interannual and seasonal primary production oscillations could thus result in important short-term variability of the standing stocks and composition of the foraminiferal faunas. Several observations of highly opportunistic behaviour of foraminiferal taxa illustrate the capability of this group of organisms to rapidly adapt to changing trophic conditions (Gooday, 1988, 1993; Gooday and Lambshead, 1989; Barmawidjaja et al., 1992; Silva et al., 1996; Ohga and Kitazato, 1997; Jannink et al., 1998; Kitazato et al., 2000). Even in some abyssal environments (e.g. Porcupine Seabight) intermittence of food input appears to be a determinant factor, causing instantaneous and patchy bursts of benthic foraminiferal faunas that colonise freshly deposited phytodetritus (Gooday, 1988, 1993; Lambshead and Gooday, 1990; Turley et al., 1993). A similar reproductive response to artificial food input has recently been shown in laboratory experiments (Heinz et al. 2001, 2002).

Although the impact of phytodetritus deposits on foraminiferal faunas has been clearly shown many important questions about deep-sea foraminiferal ecology remain partially unanswered:

- (1) What are the delays between primary production in the euphotic zone the resulting exported organic matter input at the sediment–water interface and the response of the foraminiferal faunas in terms of density composition and microhabitat?
- (2) What are the requirements of the various foraminiferal species in terms of quality and quantity of exported organic compounds?
- (3) How important is foraminiferal patchiness in the upper sediment layers; is this patchiness related to the micro-distribution of organic matter?
- (4) Is the vertical distribution of foraminiferal species in the sediment modified by organic matter input?
- (5) Is the foraminiferal response to food input restricted to the sediment surface, or is there a faunal response at depth in the sediment as well?

Whereas most opportunistic taxa rely on fresh phytodetritus, other, less critical taxa may feed on more altered organic substances, such as zooplankton faecal pellets or organic matter in a more or less advanced state of decay. It has been speculated that some taxa may also feed either on bacteria involved in the aerobic or anaerobic degradation of more refractory organic compounds, or on their breakdown products (Lee, 1979; Thomsen and Altenbach, 1993; Kitazato, 1994; Jorissen et al., 1995).

A previous foraminiferal study along a five station bathymetric transect in the Bay of Biscay substantiated the determining role of exported organic matter flux on foraminiferal density, composition, and microhabitat (Fontanier et al., 2002). The foraminiferal density reflects the exported organic matter flux, and shows a clear decrease from 150 to 2000 m depth. Foraminiferal microhabitats, which appear to be linked to principal redox fronts, deepen from the outer continental shelf to the lower slope stations. In the deepest, most oligotrophic stations, infaunal taxa tend to disappear. Finally, Fontanier et al. (2002) propose a species specific adaptation of the TROX microhabitat model of Jorissen et al. (1995). In the present paper, we study the temporal variability of live benthic foraminiferal faunas (63–150 µm, >150 µm) at a single upper bathyal station (550 m depth) from an open-slope environment in the Bay of Biscay (Fig. 1), which is supposed to be under the influence of a typical temperate, mid latitude seasonal primary production regime. For this purpose, 13 sampling cruises took place from October 1997 to April 2000. On-line data archives of the Joint Research Centre (SeaWIFS data) allow estimates of surface water chlorophyll-*a* for




Fig. 1. Study area, bathymetry and geographical position of station B.

the duration of the study period. For all studied cores, geochemical analyses have been performed in order to determine redox conditions in the upper sediment layer. The main objectives of this study are to determine whether the foraminiferal faunas respond to the seasonal and interannual fluctuations of the exported organic matter flux. More precisely, we will concentrate on the seasonal changes of density, composition and vertical distribution of foraminiferal faunas, and their possible relationship with the qualitative and quantitative seasonal variations of the exported organic matter flux.

## 2. Study area

### 2.1. Hydrographical setting

The Bay of Biscay is a semi-enclosed basin where water mass circulation is disconnected from

the general North Atlantic Drift. The current velocity of the various water masses is generally weak (less than  $10 \text{ cm s}^{-1}$ , Tréguer et al., 1979), and the waters which enter from the north, along the Irish shelf-break, leave the area near Cape Finisterre only after 2 years. The surface waters patterns are strongly dependent on the seasonal variations of the thermocline and mixed layer (Tréguer et al., 1979), and the surface currents (velocity and directions) are widely influenced by local wind regimes (Boucher, 1985). Hydrological patterns of water masses in the southeast part of the Bay of Biscay (our study area) are presented in a previous paper (Fontanier et al., 2002).

The continental slope bordering the French shelf deepens gradually, and is interrupted by two large canyons (Cap-Ferret Canyon, Capbreton Canyon). Vertical fluxes represent the main sedimentary component in open-slope environments, whereas lateral advection may dominate

sedimentary processes in the canyons (Heussner et al., 1999). The present paper concentrates on a 550 m deep open-slope station, where the impact of laterally advected particles is supposed to be minimal, and where the linkage between surface water primary production, the vertical particle flux, and benthic life should be rather straightforward. Our station B is positioned within the Northern Atlantic Central Waters (NACW), which has a salinity of 35.60 and a temperature of about 11°C (Ogawa and Tauzin, 1973). The sediment consists predominantly of fine-grained silty mud.

## 2.2. Primary production patterns in the north-eastern Atlantic

In temperate latitudes, the alternation between mixing and stratification of the water column, and the daily input of sunlight appear to be the main factors controlling the seasonal dynamics of phytoplankton (e.g. Harris, 1986; Berger and Wefer, 1990). The moderately strong seasonality found in the Bay of Biscay is characterised by an annual sinusoidal primary production curve, dominated by opportunistic species, with a succession of phytoplankton to zooplankton. The consequences of this temporal variability are seasonal occurrences of zooplankton grazers and predators, a fluctuating exploitation of organic debris, and, possibly, seasonal changes in the growth rate of benthic organisms (Bruland et al., 1989; Jumars et al., 1989).

The Bay of Biscay is characterised by phytoplankton blooms occurring in spring, summer and autumn (Tréguer et al., 1979; Froidefond et al., 1996; Laborde et al., 1999). Spring blooms are considered as the highest chlorophyll-*a* concentration events throughout the year. The phytoplankton assemblages are mainly composed of diatoms and coccolithophorids (Tréguer et al., 1979; Fernandez et al., 1995). Wroblewski (1989) suggests a general duration of about 2 weeks for the spring bloom. Fernandez (1990), who studies the Central Cantabrian Sea (southern part of the Bay of Biscay), observes high chlorophyll-*a* and primary production rates in March, predominantly due to microflagellates. In a more recent

study Laborde et al. (1999) describe a high productivity period, which lasts for about 2 months. In view of the earlier literature, it seems obvious that this relatively long period is the result of a close succession of several individual bloom events, involving different phytoplankton groups.

Minor upwelling events occur during early summer, in relation to NNW coast-parallel winds and to riverine discharge (Froidefond et al., 1996). Large coccolithophorids blooms are the result; they have been observed on the shelf-break of the Bay of Biscay by satellite imagery as well as by in situ measurements (Holligan et al., 1983; Fernandez et al., 1993; Beaufort and Heussner, 1999). Because this system has probably a very strong interannual variability, it is difficult to determine an average primary production value. Nevertheless, total primary production estimates based on satellite imagery (Antoine et al., 1996) suggest a slight increase in early summer.

In autumn, cooling of the surface waters is generally effective without causing the total disappearance of water column stratification and of the nutricline. Nevertheless, Tréguer et al. (1979) have recorded a chlorophyll-*a* increase in October, which they describe as an autumn bloom. In this period, primary production is rather patchy on a basin scale, and strongly influenced by the hydrological structures. Lohrenz et al. (1992) speculate that primary production increases are generally caused by wind-induced mixing, resulting in nutrient injection from subsurface waters into the nutrient-depleted surface waters. Recently, Sellmer et al. (1998) showed an autumn bloom at the BIOTRANS site (47°N/20°W), occurring around the nutricline, at about 30–50 m depth, mainly composed of autotrophic dinoflagellates, with smaller amount of diatoms.

Very few quantitative data are available on primary production in general, and more specifically, on the individual bloom events occurring in the study area. Tréguer et al. (1979) estimate a primary production between 0.4 and 1.9 g C/m<sup>2</sup>/day for the spring bloom of 1973. Primary production measurements during autumn 1972 indicate a bloom with values of 0.3–0.4 g C/m<sup>2</sup>/day (Le Corre and Tréguer, 1976). This range of values agrees with recent data obtained in the Cap-Ferret

region during five ECOFER cruises: 0.7–1.2  $\text{gC/m}^2/\text{day}$  in spring (May 1990 and 1991) and 0.3  $\text{gC/m}^2/\text{day}$  in autumn (October 1990; Laborde et al., 1999). Total annual primary production has been estimated between 145 and 170  $\text{gC/m}^2/\text{yr}$  (Laborde et al., 1999).

### 3. Material and methods

In order to complete the rather scarce information about seasonal and interannual changes of phytoplankton biomass in the surface waters from the Bay of Biscay, we used on-line data archives of the Joint Research Center (European Commission) to estimate chlorophyll-*a* concentrations (SeaWiFS data) in the study area for the duration of our sampling period (October 1997–April 2000; Fig. 2). Unfortunately, weather conditions strongly affect the availability of images. Especially during autumn and winter, satellite monitoring is only rarely possible, because of the dense cloud cover.

In order to obtain a good overview of the temporal variability of the benthic foraminiferal assemblages, 13 sampling cruises were organised between October 1997 and April 2000. For meteorological reasons, sampling was impossible

during 3 cruises, and consequently, station B ( $43^\circ 49' 98\text{N}$ ,  $2^\circ 23' 04\text{W}$ , water depth 550 m, Fig. 1) was sampled 10 times (Table 1). Cores were sampled with a Barnett multi-tube corer (Barnett et al., 1984), which allows sampling of first decimeters of the sediment, the overlying bottom waters, and an undisturbed sediment–water interface. Free waters were collected immediately after core recovery for dissolved  $\text{O}_2$  measurements by the Winkler titration method (Strickland and Parsons, 1972). Profiles of pore water  $\text{O}_2$  were obtained on board with a cathode-type mini-electrode (Revsbech and Jørgensen, 1986; Helder and Bakker, 1985; Revsbech, 1983). The temperature was maintained stable with an insulating device. This operation was completed in duplicate within 30 min after core recovery. Subsequently, the core used for  $\text{O}_2$  profiling was sliced into thin horizontal sections (every 0.5 cm for the top 2 cm, 1 or 2 cm below) within 1 h and 30 min. For every level a sub-sample was centrifuged under  $\text{N}_2$  at 5000 rpm for 20 min in order to collect pore waters. Two aliquots of water were filtered ( $0.2\ \mu\text{m}$ ) and frozen at  $-25^\circ\text{C}$  for nutrient analyses. Interstitial water compounds were analysed by techniques adapted for small volumes of samples (Anschutz et al., 1999; Hyacinthe et al., 2001). Nitrate and nitrite were measured by flow




Fig. 2. Chlorophyll-*a* concentrations in the surface waters in our study area between October 1997 and April 2000 (estimates based on SEAWIFS images). Vertical dot–dash lines indicate when the sampling cruises took place.

Table 1

Sampling dates, bottom water oxygen concentration, depth in the sediment of the zero oxygen level and semi-quantitative analysis performed on the sedimentary residual parts of the first quarter of sediment for 15 cores at station B for the 10 sampling cruises; Phytoplankton and zooplankton components were observed and described according to the following classes: ++ abundant, + common, – rare, — absent

| Cores station B            | Date       | O <sub>2</sub> concentration (μmol/l) | Oxygen penetration depth (mm) | Whitish filamentous and/or amorphous phytodetrital aggregates | Radiolarians |
|----------------------------|------------|---------------------------------------|-------------------------------|---|--------------|
| OB1B/OB1B <sup>bis</sup>   | 26/10/1997 | 217                                   | 17                            | + +/–   | –/–          |
| OB2B/OB2B <sup>bis</sup>   | 30/01/1998 | 216                                   | 24                            | +   | —            |
| OB3B                       | 07/06/1998 | 212                                   | 19                            | –/+   | –/–          |
| OB4B                       | 23/07/1998 | 208                                   | 18                            | +   | —            |
| OB5B                       | 17/10/1998 | 205                                   | 21                            | –   | —            |
| OB6B                       | 08/12/1998 | 212                                   | 20                            | –   | —            |
| OB7B                       | 23/01/1999 | 220                                   | 26                            | +   | —            |
| OB8B/OB8B <sup>bis</sup>   | 19/04/1999 | 207                                   | 20                            | –/–   | –/+          |
| OB9B/OB9B <sup>bis</sup>   | 24/06/1999 | 215                                   | 21                            | + /+ +  | + /+         |
| OB10B/OB10B <sup>bis</sup> | 25/04/2000 | 221                                   | 23                            | + + /+ +  | –/–          |

injection analysis (FIA) according to Anderson (1979).

For faunal analysis, one entire 72 cm<sup>2</sup> core was sliced horizontally for each station, usually every 0.25 cm for the first cm of sediment, every half cm between 1 and 4 cm depth, and every cm between 4 and 10 cm. For evaluating the amplitude of spatial heterogeneity, all available duplicate samples were analysed as well. This concerns samples OB1B<sup>bis</sup> (the top 2 cm of a second core, partially sampled in 0.5 cm intervals), OB2B<sup>bis</sup> (top 5 cm sampled in 2.5 cm intervals), OB8B<sup>bis</sup> (sampled down to 5 cm according to the standard sampling protocol), OB9B<sup>bis</sup> and OB10B<sup>bis</sup> (both completely sampled). Except for OB10 (two consecutive multi-corer deployments), each duplicates pair was recovered from the same multi-corer deployment. Since in all cores more than 95% of the fauna is found in the top 5 cm, OB2B<sup>bis</sup>, OB8B<sup>bis</sup> and OB9B<sup>bis</sup> can be considered as fully reliable duplicates. Census data of core OB1B (> 150 μm fraction) have already been presented in Fontanier et al. (2002). Nevertheless because of some taxonomic improvements, we present slightly modified census data here (Appendix B).

We follow the same sediments storage and preparation as presented in Fontanier et al. (2002). Foraminifera belonging to the > 150 μm

fraction were studied along the 10 cm long cores. Because of its extremely time-consuming character, we limited our study of the 63–150 μm fraction to the first half cm of the sediment. In cases where the sediment surface is oblique, the volume of the top half cm layer may be slightly variable, and therefore, absolute density values for the superficial samples are less reliable. During picking, semi-quantitative analyses were performed on the residual parts of both fractions to evaluate some remarkable sedimentary components (phytodetritus and zooplankton compounds; Table 1).

The Rose Bengal staining technique (Walton, 1952; Bernhard, 1988) is routinely used to recognise foraminifera that were alive at the time of sampling. A problem of this method is that Rose Bengal may also stain the dead foraminiferal protoplasm, which may be partially preserved below the zero oxygen level within the sediment for a considerable period of time after the death of the organism (Bernhard, 1988; Corliss and Emerson, 1990). As a consequence, we use the same strict staining criteria as those detailed in Fontanier et al. (2002). Non-transparent agglutinated and miliolid taxa were broken in order to inspect the test interior. Fragments of the very fragile arborescent agglutinating foraminiferal fragments (such as *Hyperammina* spp., and *Glomospira* spp.)

were not included in the quantitative analyses, because the orange-reddish colour of their test makes it particularly difficult to appreciate whether the organism was alive or dead at the time of sampling.

Our taxonomical framework is given in Appendix A. All foraminiferal census data are listed in Appendices B and C. The total density of the live foraminiferal fauna total density has been determined by summing up the number of individuals for all levels between 0 and 10 cm depth for the >150 µm fraction, but only for the 0–0.5 cm interval for the 63–150 µm fraction. The total density per core is expressed as number of individuals found at and below a 72 cm<sup>2</sup> sediment surface. In all graphs depicting the vertical distribution of the foraminifera in the >150 µm fraction (Figs. 5a–m), the faunal densities have been standardised for a 50 cm<sup>3</sup> sediment volume.

The average living depth (ALD<sub>x</sub>, Jorissen et al., 1995) seems the best way to describe the overall vertical distribution of the total foraminifera fauna or of individual taxa, and to get a general idea about the microhabitat patterns. After a first classification with four main microhabitats proposed by Corliss and Chen (1988), it has been argued that only species living on elevated substrates can be considered as “epifaunal” (Buzas et al., 1993). Therefore, in the soft bottom communities described in this study, we recognise only three different microhabitat categories: shallow infaunal, intermediate infaunal and deep infaunal taxa. The ALD<sub>x</sub> is calculated with the following formula:

$$\text{ALD}_x = \sum_{i=0,x} (n_i \times D_i) / N,$$

where  $x$  is the lower boundary of deepest sample,  $n_i$  is the number of specimen in interval  $i$ ,  $D_i$  the midpoint of sample interval  $i$ , and  $N$  is the total number of individuals for all levels.

For all stations, ALD<sub>10</sub> was calculated for the whole fauna, as well as for individual taxa, on the basis of the numbers of stained individuals found in the successive sediment slices. Isolated individuals separated from the main population by more than 1 cm of “sterile” sediment (without live individuals of the studied taxon) were not inte-

grated in the calculations of the ALD<sub>10</sub>. We suppose that such isolated individuals have been transported downward (outside their normal microhabitat) by bioturbation, or correspond to dead organisms that have been counted erroneously. Overall, weighed ALD<sub>10</sub> values were calculated for each taxon by integrating the results obtained in the 13 cores:

$$\overline{\text{ALD}}_{10} = \sum_{i=1,n} (\text{ALD}_{10}^i \times n_i) / \sum_{i=1,n} n_i,$$

where  $n$  is the number total of cores, ALD<sub>10</sub><sup>*i*</sup> the average living depth for the 10 first cms of the core  $I$ , and  $n_i$  is the number of specimen in the core  $i$ .

In order to evaluate the differences between duplicate and temporally distinct samples, a non-standardised principal component analysis (Davis, 1986) was applied for both size fractions, using the percentages of all taxa with an occurrence of more than 5% in at least one sample.

## 4. Results

### 4.1. Chlorophyll-*a* concentrations from October 1997 to April 2000

In the Bay of Biscay, the spring bloom chlorophyll-*a* increase is the most prominent seasonal event of the year (Fig. 2). This gradual or pulse-like increase of chlorophyll-*a* concentration generally starts with a first pulse at the end of winter, in the second part of March (Fig. 2), which affects the whole Bay of Biscay. A second pulse appears in the second half of April, and lasts through the first weeks of May. This second pulse has a more restricted geographical distribution, and at our sampling station, we observed it only in spring 1999.

In summer, the offshore waters of the Bay of Biscay exhibit rather low chlorophyll-*a* concentrations (0.1–0.5 mg/m<sup>3</sup>); higher values are present over the shelf and shelf break. However, we can imagine that phytoplankton biomass associated with a putative Deep chlorophyll maximum is not represented in satellite images.

In autumn, the scarcity of satellite images allows far reliable estimates of chlorophyll-*a* concentrations. This period is characterised by low surface water phytoplankton biomass for the whole Bay of Biscay. Chlorophyll-*a* concentrations are minimal in late autumn and early winter.

At our station B, the variability of the chlorophyll-*a* concentrations in the overlying surface waters is in close agreement with the basin-wide patterns inferred before. In late autumn and in the winter, chlorophyll-*a* concentrations are low ( $<0.7 \text{ mg/m}^3$ ). They precede high and abrupt chlorophyll-*a* increases (up to  $10 \text{ mg/m}^3$ ), which can be noticed every year in late winter (March 1998, 1999 and 2000). Only in 1999 is a second chlorophyll-*a* increase observed, on the 13th of May. The summer is characterised by intermediate chlorophyll-*a* concentrations (about  $1 \text{ mg/m}^3$ ). In early autumn (October), the chlorophyll-*a* concentrations are comparable to those in summer. Minimal surface water chlorophyll-*a* values, of about  $0.2 \text{ mg/m}^3$ , are reached in October; surface waters remain oligotrophic ( $0.2\text{--}0.5 \text{ mg/m}^3$ ) until January.

#### 4.2. Sedimentary organic deposits

Semi-quantitative observations of the sieve residues of the first half centimetre reveal that cores sampled in October 1997, in June 1999 and in April 2000 (OB1B, OB9B<sup>bis</sup>, OB10B and OB10B<sup>bis</sup>, respectively) contain high amounts of whitish amorphous aggregates composed of diatoms and of radiolarians (Table 1). These organic aggregates were observed after sieving of the samples, and therefore, benthic eventual foraminiferal faunas living within the organic aggregates could not be substantiated.

#### 4.3. Oxygen concentration and redox conditions of interstitial waters

Bottom water oxygen concentrations (Table 1, Fig. 3a) vary from 205 to  $221 \mu\text{mol/l}$ . The zero oxygen boundary varies from 17 mm depth (OB1B, October 1997) to 26 mm depth (OB7B, January 1999). In all cases, there is a steep decrease of

oxygen concentration in the first half cm of the sediment. The nitrate and nitrite concentrations (Fig. 3b) have maximum values (of about  $20 \mu\text{mol/l}$ ) in the first half cm, and show a rapid decrease to background values at about 2.5 cm depth.

#### 4.4. Faunal density and number of taxa

In the  $>150 \mu\text{m}$  fraction, foraminiferal densities vary from 245 to 1346 individuals per core (Fig. 4). Maxima of 1346 and 845 individuals are found in October 1997 (OB1B) and April 2000 (OB10B), respectively. Minimum values are recorded in July 1998 (OB4B, 281 individuals), October 1998 (OB5B, 322 individuals) and April 1999 (245 and 465 individuals in two replicate cores). For the 63– $150 \mu\text{m}$  fraction, foraminiferal densities (only for the topmost 0.5 cm) vary from about 40–2195 individuals (Fig. 4). Maxima of 1505 and 2195 individuals are found in October 1997 (OB1B) and in April 2000 (OB10B<sup>bis</sup>); minimum values are found in December 1998 (OB6B, 37 individuals) and in October 1998 (OB5B, 73 specimens). Part of this variability, however, may be due to the variable sample size of the first 0.5 cm in the case of oblique sediment surfaces.

In the  $>150 \mu\text{m}$  fraction, perforate foraminifera form the main faunal component (60% to 90% of the fauna). Non-fossilising agglutinated taxa account for 10–30%, whereas fossilising agglutinated taxa (maximum 10%) and miliolids (maximum 3.5%) are rare in all cores. Also in the 63– $150 \mu\text{m}$  fraction, the perforate group is largely dominant (50–95%), but non-fossilising agglutinated individuals can represent up to 48% of the total fauna (in April 1999, OB8B). Miliolids and fossilising agglutinated foraminifera account for less than 5% of the foraminiferal faunas.

The number of taxa in the  $>150 \mu\text{m}$  fraction varies from 31 (December 1998, OB6B) to 57 (June 1999, OB9B<sup>bis</sup>), without a clear relation to faunal density. In the 63– $150 \mu\text{m}$  fraction, where only the topmost 0.5 cm was studied, the number of taxa varies from 16 (December 1998, OB6B) to 65 (October 1997, OB1B); there is a clear positive correlation with the faunal density here.






Fig. 3. (a) Dissolved oxygen concentrations in the sediment for the ten cruises; (b) Nitrate + nitrite concentrations in the sediment for the ten cruises.

#### 4.5. Faunal composition and microhabitat

(1) *>150 μm fraction*: For most of the cores, the foraminiferal fauna is strongly concentrated in the oxygenated sediment top layers (Figs. 5a–m). The highest density is normally found in the first half cm. Maximum values were observed in October 1997 (OB1B; ~800 specimens/50 cm<sup>3</sup>) and April 2000 (OB10B; ~500 specimens/50 cm<sup>3</sup>). In most cores, the faunal density drops rapidly in the second cm, where dysoxic conditions prevail. Only low-density foraminiferal faunas are found in the

generally anoxic sediments below 2 cm depth. The cores taken in December 1998 (OB6B) and April 1999 (OB8B<sup>bis</sup>), which contain fair amounts of live individuals below the zero oxygen level, form notable exceptions. Consequently, for most of the cores, the ALD<sub>10</sub> of the total fauna is about 1 cm. Maximum ALD<sub>10</sub> values are found in December 1998 (OB6B; 2.0 cm) and April 1999 (OB8B<sup>bis</sup>; 1.6 cm). The shallowest microhabitat depth was observed in July 1998 (ALD<sub>10</sub>=0.6 cm).

*Uvigerina mediterranea* dominates the faunas in every core (Appendix B); its relative abundance




Fig. 4. Foraminiferal density for both size fractions during the ten cruises. The foraminiferal density for the  $>150\ \mu\text{m}$  fraction is calculated as the total number of live individuals per core of  $72\ \text{cm}^2$ . The foraminiferal density for the  $63\text{--}150\ \mu\text{m}$  fraction is calculated as the total number of live foraminifera in the first half cm of the core. When duplicate cores are available, an average value is plotted, and the values of the individual cores are given (underlined numbers). For all cores, the number of species is given in italics. When duplicate cores are available, the values of the individual cores are also presented (couple of numbers in italics).

fluctuates from 17.8% to 39.9%, with maximum percentages observed in January 1998 (OB2B) and June 1999 (OB9B<sup>bis</sup>). Minimum values are found in July 1998 (OB4B) and April 1999 (OB8B<sup>bis</sup>). *Melonis barleeanus* is the second most abundant species; in most samples it counts for about 10% of the total fauna (4.1–16.7%). *U. peregrina* has a much more variable percentage. It is very frequent in October 1997 (OB1B and OB1B<sup>bis</sup>), when its percentage reaches 21%. In many other samples, the species is rather infrequent (2–5%). *Globobulimina affinis* is well represented in most cores, with relative abundances of about 5%. The highest percentage (12–15%) is found in April 1999 (OB8B and OB8B<sup>bis</sup>). *U. elongatastriata* and *Reophax scorpiurus* account each for about 5% of the faunas. Some other taxa show relatively low background values (1–3%), but can occasionally occur with high percentages. *Cibicidoides pachydermus* shows a 7.2% maximum in January 1998 (OB2B), whereas *Clavulina cylindrica* is well represented (7.7%) in June 1998 (OB3B). *Cribrostomoides subglobosus* shows peak values (9–12%) in June 1998 (OB3B), October 1998 (OB5B), and April and June 1999 (OB8B<sup>bis</sup> and OB9B<sup>bis</sup>). *Pseudoclavulina crustata* shows

one single peak occurrence (9%), in April 1999 (OB8B).

The microhabitat of the majority of the taxa is very shallow (Table 2). The overall, weighed, values are lowest for *C. pachydermus* ( $\text{ALD}_{10} = 0.54\ \text{cm}$ ), *P. crustata* ( $\text{ALD}_{10} = 0.71\ \text{cm}$ ), *R. scorpiurus* ( $\text{ALD}_{10} = 0.75\ \text{cm}$ ), *U. mediterranea* ( $\text{ALD}_{10} = 0.77\ \text{cm}$ ), *U. peregrina* ( $\text{ALD}_{10} = 0.78\ \text{cm}$ ) and *C. subglobosus* ( $\text{ALD}_{10} = 0.82\ \text{cm}$ ); these species are largely restricted to the shallow infaunal microhabitats in the first cm of the sediment (Figs. 5a–m). Their  $\text{ALD}_{10}$  shows very little variability between the 13 cores (Table 2), with the exception of core OB6B, where the microhabitat of all taxa is significantly deeper than normal. The anomalously low numbers in the first 0.5 cm level at this station (in the  $63\text{--}150\ \mu\text{m}$  as well as in the  $>150\ \mu\text{m}$  fraction) suggest that, because of a strongly oblique sediment surface, only a minor part of the  $72\ \text{cm}^2$  surface was sampled. This sampling artefact effects the calculation of the microhabitat, and the density value in the  $63\text{--}150\ \mu\text{m}$  fraction, but not the total density values (down to 10 cm) in the  $>150\ \mu\text{m}$  fraction.

*M. barleeanus* (overall weighed  $\text{ALD}_{10} = 1.30\ \text{cm}$ ), *U. elongatastriata* (overall weighed

ALD<sub>10</sub> = 1.47 cm) and *C. cylindrica* (overall weighed ALD<sub>10</sub> = 1.46 cm) show a rather constant intermediate infaunal microhabitat, with the exception of core OB6B, where all occurrences are deeper. *G. affinis* (overall weighed ALD<sub>10</sub> = 2.84 cm) occupies a deep infaunal microhabitat, in dysoxic or sometimes totally anoxic sediment layers.

(2) 63–150 μm fraction: The densities of the main taxa in the topmost 0.5 cm of the fifteen cores are shown in Fig. 6 and Appendix C. Because of the

extreme variability of foraminiferal density, which may be partially due to a varying sampling volume (see Methods section), the faunal variability may be better represented by percentage data, which are presented in Appendix C. The faunas in this small size fraction are much more variable than those in the > 150 μm fraction. *R. guttiferus* is a frequent taxon in many cores; it accounts for about 27% in June 2000 (OB9B<sup>bis</sup>), and for 10–15% in October 1997 (OB1B), January 1998 (OB2B), December 1998 (OB6B), January 1999




Fig. 5. (a–m) Foraminiferal distribution (number of individuals > 150 μm fraction found in each level, standardised for a 50 cm<sup>3</sup> sediment volume) for 14 available cores. Data for core OB1B in (a) comes from corrected data presented in Fontanier et al. (2002).




Fig. 5 (continued).



### Density of the main 63–150 $\mu\text{m}$ fraction foraminiferal taxa




Fig. 6. Foraminiferal density of the main foraminiferal taxa in the 63–150  $\mu\text{m}$  fraction for the 15 cores. Densities are standardised to a 50  $\text{cm}^3$  sediment volume.

(OB7B), April 1999 (OB8B and OB8B<sup>bis</sup>) (OB9B) and April 2000 (OB10B and OB10B<sup>bis</sup>). *Epistominella exigua* occurs with spectacular peak abundances in April 2000 (OB10Bbis—35.6%), June 1999 (OB9B—32.5%; OB9B<sup>bis</sup>—14%) and October 1997 (9% in both OB1B and OB1B<sup>bis</sup>). *U. peregrina* occurs with about 15% abundance in October 1997 and October 1998 (OB1B and OB5B), whereas *C. carinata* exhibits values from 10–15% in October 1997 (OB1B<sup>bis</sup>), April 2000 (OB10B), January 1998 (OB2B) and December 1998 (OB6B). *G. affinis* shows a spectacular peak occurrence in June 1998 (OB3B—45%), whereas *M. barleeanus* is frequent in July 1998 (OB4B—15%). Many other species, such as *Bolivina*

*spathulata*, *Gyroidina umbonata*, *Nonionella* spp., *Siphogenerina columellaris*, *U. mediterranea*, *R. scorpiurus*, and *Trochammina* spp. show shifts from near-absence to values of about 10%, highlighting the extreme variability of the 63–150  $\mu\text{m}$  fraction.

## 5. Discussion and conclusions

### 5.1. Organic matter deposits

The presence of organic aggregates at the sediment–water interface appears to follow phytoplankton bloom events taking place several weeks

before the cruises (Table 1, Fig. 2). This is especially the case for OB10 (6 weeks after the first 2000 spring bloom) OB9 (6 weeks after the second 1999 spring bloom) and OB1 (4 weeks after a 1997 autumn chlorophyll-*a* maximum). We suppose that the organic aggregates represent the remains of phytoplankton deposits following these surface water bloom periods.

### 5.2. Spatial heterogeneity of the benthic ecosystem at station B

Most of studies dealing with temporal variability of benthic foraminiferal faunas taxa (e.g. Gooday, 1988; Gooday and Lamshead, 1989; Thiel et al., 1990; Gooday and Turley, 1990; Thurston et al., 1994; Kitazato and Ohga, 1995; Silva et al., 1996; Kitazato et al., 2000) avoid the fundamental question what part of the observed differences is due to spatial variability (patchiness) and what part is really due to temporal variability, resulting from the intermittence of export production. Barmawidjaja et al. (1992) moderate their conclusions about seasonal changes of foraminiferal faunas from the Northern Adriatic Sea by underlining the possible importance of a patchiness effect and insist on the necessity to study replicate cores. Thiel et al. (1990) suggest that current activity and microtopography may induce a patchy distribution of phytodetritus at the sea floor. Organic deposits create a mosaic of ephemeral organic rich patches on the ocean floor, which maintain sea floor heterogeneity, and contribute to the high diversity of deep sea benthic communities (Grassle and Morse-Porteous, 1987; Grassle, 1989; Snelgrove et al., 1994, 1996). Hohenegger et al. (1993) demonstrate that foraminiferal faunas have patchy distributions which are interpreted as being food controlled. In a study using duplicate cores, in recently enriched sediments, Silva et al. (1996) show that spatial faunal variability (patchiness) exists but does not obscure studies on the temporal and seasonal changes of the foraminiferal faunas, which are much more prominent.

In order to distinguish between spatial and temporal variability in our cores we performed a non-standardised principal component analysis (Davis, 1986) on the basis of the percentage data

of all taxa which appear with at least 5% in one of the cores. For the >150  $\mu\text{m}$  fraction, this multivariate analysis is based on 15 samples (10 samples and 5 duplicates) and 10 taxa, and yields two significant axes, explaining 68% of the total variability. The eigenvalues for these two axes and the species loadings on the axes are given in Table 3a. The positive side of axis 1 is highly dominated by *U. mediterranea* (0.90), whereas *G. affinis*, *R. scorpiurus*, *U. elongatastriata*, *Cribrostomoides subglobosus*, and *Clavulina cylindrica* all load negatively on the first axis. The positive side of the second axis is strongly dominated by *U. peregrina* (0.92), while *U. mediterranea*, *G. affinis* and *C. subglobosus* load negatively. Fig. 7 shows the position of the 15 cores in the axial plot. The first axis allows separation of all rather poor samples (total density less than 500 individuals, PCA1  $\leq$ 16.0) from all relatively rich samples (total density > 500 specimens, PCA1 >18.0), suggesting that the first axis is related to ecosystem enrichment due to phytodetritus deposits, a phenomenon which is always accompanied by an increase of the percentage of *U. mediterranea*. The second axis tends to separate cores OB1B and OB1B<sup>bis</sup>, which are both strongly enriched in *U. peregrina*, from all other cores with high scores on axis 1. This suggests that there are two possible responses to ecosystem enrichment; in most cases *U. mediterranea* shows the clearest response, but in October 1997 the large increase of *U. peregrina* is the most obvious phenomenon accompanying the faunal density increase. These two possible responses, translated by a different position on axis 2, could be explained by a different quality of the organic matter between the spring and autumn blooms. All five pairs of duplicate cores are fairly close to each other. Although individual cores of these duplicate couples may have a higher similarity to other cores (e.g. core OB9B is much closer to OB7B than to OB9B<sup>bis</sup>), showing the presence of significant spatial variability, the distances between extreme samples are much bigger than the maximum distance between duplicate couples. This suggests that for the >150  $\mu\text{m}$  fraction, temporal variability is more important than spatial variability.

Table 3

(a–c) Results of non-standardised principal component analyses based on the percentages of the main foraminiferal taxa in the > 150 and 63–150  $\mu\text{m}$  fraction (percentage  $\geq 5\%$ ). (a) Eigenvalues and species loadings of the two significant axes for the > 150  $\mu\text{m}$  fraction. (b) Same, for the 63–150  $\mu\text{m}$  fraction. (c) The normative distance between cores in the five-dimensional space (normative distance between the five pairs of replicate cores are given in bold letters)

| (a)                                |       |       |       |       |       |
|------------------------------------|-------|-------|-------|-------|-------|
| > 150 $\mu\text{m}$ fraction       | PCA1  | PCA2  |       |       |       |
| Eigenvalues column                 | 67.8  | 37.3  |       |       |       |
| Percent of trace                   | 44.3  | 24.4  |       |       |       |
| Cumulative percent of trace        | 44.3  | 68.7  |       |       |       |
| Taxa                               |       |       |       |       |       |
| <i>Cibicidoides pachydermus</i>    | 0.11  | 0.12  |       |       |       |
| <i>Globobulimina affinis</i>       | −0.28 | −0.16 |       |       |       |
| <i>Melonis barleeanus</i>          | 0.03  | 0.09  |       |       |       |
| <i>Uvigerina elongatastriata</i>   | −0.16 | −0.11 |       |       |       |
| <i>Uvigerina mediterranea</i>      | 0.90  | −0.19 |       |       |       |
| <i>Uvigerina peregrina</i>         | 0.04  | 0.92  |       |       |       |
| <i>Clavulina cylindrica</i>        | −0.14 | −0.12 |       |       |       |
| <i>Cribrostomoides subglobosus</i> | −0.16 | −0.15 |       |       |       |
| <i>Reophax scorpiurus</i>          | −0.17 | −0.07 |       |       |       |
| <i>Pseudoclavulina crustata</i>    | −0.09 | −0.06 |       |       |       |
| (b)                                |       |       |       |       |       |
| 63–150 $\mu\text{m}$ fraction      | PCA1  | PCA2  | PCA3  | PCA4  | PCA5  |
| Eigenvalues column                 | 172.4 | 123.9 | 51.3  | 36.4  | 23.6  |
| Percent of trace                   | 36.7  | 26.4  | 10.9  | 7.8   | 5.0   |
| Cumulative percent of trace        | 36.7  | 63.1  | 74.1  | 81.8  | 86.8  |
| Taxa                               |       |       |       |       |       |
| <i>Bolivina</i> sp.1               | 0.00  | −0.04 | 0.09  | 0.06  | 0.05  |
| <i>Bolivina spathulata</i>         | −0.04 | 0.04  | 0.16  | 0.40  | 0.10  |
| <i>Bolivina difformis</i>          | −0.06 | 0.06  | 0.04  | 0.05  | −0.06 |
| <i>Bulimina marginata</i>          | −0.03 | −0.05 | 0.08  | −0.06 | −0.02 |
| <i>Cassidulina carinata</i>        | −0.06 | −0.12 | 0.21  | 0.38  | 0.17  |
| <i>Ceratobulimina</i> sp.          | 0.03  | 0.00  | −0.03 | 0.02  | 0.04  |
| <i>Epistominella exigua</i>        | −0.57 | 0.76  | −0.10 | −0.08 | −0.04 |
| <i>Globobulimina affinis</i>       | 0.75  | 0.53  | 0.29  | −0.18 | 0.03  |
| <i>Gyroidina umbonifera</i>        | −0.02 | −0.04 | −0.12 | −0.20 | 0.27  |
| <i>Melonis barleeanus</i>          | 0.14  | −0.05 | −0.38 | 0.04  | −0.50 |
| <i>Nonionella</i> spp.             | 0.09  | 0.00  | −0.13 | 0.10  | −0.22 |
| <i>Nuttalides pusillus</i>         | −0.04 | 0.03  | −0.06 | −0.07 | 0.05  |
| <i>Pullenia</i> spp.               | −0.02 | 0.02  | −0.13 | −0.05 | 0.04  |
| <i>Siphogenerina columellaris</i>  | −0.08 | −0.09 | −0.09 | 0.05  | −0.52 |
| <i>Uvigerina mediterranea</i>      | −0.01 | −0.12 | −0.10 | −0.18 | 0.11  |
| <i>Uvigerina peregrina</i>         | 0.02  | −0.17 | −0.32 | −0.55 | 0.25  |
| <i>Haplophragmoides</i> sp.        | 0.00  | −0.03 | 0.09  | 0.09  | 0.17  |
| <i>Psammosphaera fusca</i>         | −0.01 | −0.07 | 0.01  | 0.18  | −0.06 |
| <i>Reophax guttiferus</i>          | −0.24 | −0.19 | 0.68  | −0.42 | −0.28 |
| <i>Reophax scorpiurus</i>          | 0.01  | −0.11 | 0.04  | 0.02  | 0.17  |
| <i>Reophax</i> sp.1                | −0.02 | −0.09 | 0.09  | 0.07  | 0.10  |
| <i>Trochammina</i> spp.            | −0.09 | −0.01 | 0.13  | −0.17 | −0.31 |



Table 3 (continued)

| (c)                 | OB1B <sup>bis</sup> | OB2B | OB2B <sup>bis</sup> | OB3B | OB4B | OB5B | OB6B | OB7B | O8BB | OB8B <sup>bis</sup> | OB9B | OB9B <sup>bis</sup> | OB10B | OB10B <sup>bis</sup> |
|---------------------|---------------------|------|---------------------|------|------|------|------|------|------|---------------------|------|---------------------|-------|----------------------|
| OB1B                | 22.4                | 20.2 | 19.2                | 50.1 | 21.2 | 13.6 | 14.2 | 14.1 | 10.9 | 15.2                | 28.0 | 19.8                | 17.4  | 31.1                 |
| OB1B <sup>bis</sup> |                     | 16.1 | 15.1                | 49.4 | 27.2 | 24.6 | 24.5 | 16.0 | 18.6 | 11.1                | 26.3 | 26.7                | 11.9  | 31.1                 |
| OB2B                |                     |      | 17.6                | 49.2 | 25.9 | 26.4 | 12.4 | 7.9  | 11.8 | 8.0                 | 36.0 | 21.1                | 10.7  | 38.3                 |
| OB2B <sup>bis</sup> |                     |      |                     | 47.1 | 12.7 | 20.6 | 20.9 | 12.1 | 14.7 | 13.7                | 31.6 | 27.8                | 11.0  | 34.9                 |
| OB3B                |                     |      |                     |      | 46.6 | 48.8 | 52.1 | 48.2 | 48.1 | 48.4                | 56.2 | 55.1                | 48.5  | 59.3                 |
| OB4B                |                     |      |                     |      |      | 21.1 | 23.7 | 18.8 | 19.1 | 23.1                | 38.1 | 32.5                | 20.8  | 40.5                 |
| OB5B                |                     |      |                     |      |      |      | 23.3 | 20.1 | 15.7 | 22.2                | 33.7 | 32.9                | 24.4  | 39.3                 |
| OB6B                |                     |      |                     |      |      |      |      | 9.8  | 9.3  | 13.8                | 37.5 | 18.2                | 16.1  | 39.0                 |
| OB7B                |                     |      |                     |      |      |      |      |      | 5.7  | 6.8                 | 33.3 | 20.7                | 8.5   | 35.9                 |
| OB8B                |                     |      |                     |      |      |      |      |      |      | <b>11.4</b>         | 32.7 | 22.3                | 23.5  | 37.8                 |
| OB8B <sup>bis</sup> |                     |      |                     |      |      |      |      |      |      |                     | 28.4 | 18.2                | 5.5   | 31.2                 |
| OB9B                |                     |      |                     |      |      |      |      |      |      |                     |      | <b>27.6</b>         | 27.9  | 8.9                  |
| OB9B <sup>bis</sup> |                     |      |                     |      |      |      |      |      |      |                     |      |                     | 18.6  | 25.7                 |
| OB10B               |                     |      |                     |      |      |      |      |      |      |                     |      |                     |       | <b>29.9</b>          |

The 63–150  $\mu\text{m}$  data (for the topmost 0.5 cm of the sediment) were subjected to a similar analysis, this time with 22 taxa. Because of extreme variability of the data set, with many different taxa having occasional peak occurrences, this second principal component analyses yields no less than five significant axes, explaining 86.8% of the total variability (Table 3b). In order to evaluate differences between samples, we calculated distances between all samples in the five-dimensional space defined by the significant axes (Table 3c). This table confirms that differences between duplicate samples (11–30 units) are larger in the 63–150 than in the > 150  $\mu\text{m}$  fraction (2–12 units), but that maximum temporal differences (59 units) are still about two times higher than maximum spatial differences.

These multidimensional analyses suggest that small-scale patchiness exists, and cannot be completely ruled out, as a parameter explaining differences between the cores taken at different times. In the > 150  $\mu\text{m}$  fraction, spatial variability seems to be smaller than in the 63–150  $\mu\text{m}$  fraction, and, consequently, temporal variation should be more prominent in this larger size fraction. An important question is why duplicate samples are more different in the 63–150  $\mu\text{m}$  than in the > 150  $\mu\text{m}$  fraction. It is very probable that the high degree of patchiness is related to micro-relief

and current activity, which cause a patchy distribution of organic matter on the ocean floor. In case of abundant input of organic detritus, phytodetritus aggregates will concentrate in depressions on the sea floor. This may explain important small-scale spatial variability in terms of concentration of phytodetrital compounds between duplicate samples (see for example the difference between OB1B and OB1B<sup>bis</sup>). Small opportunistic taxa, such as *E. exigua*, may have a faster response than larger (> 150  $\mu\text{m}$ ) taxa, and will be first in colonising freshly deposited phytoplankton floccules (Gooday and Turley, 1990). This can be seen in October 1997 (OB1B and OB1B<sup>bis</sup>), in June 1999 (OB9B and OB9B<sup>bis</sup>) and in April 2000 (OB10B<sup>bis</sup>), when organic aggregates are observed in the sieve residues, and when *E. exigua* reaches high densities (Table 1, Fig. 6). This taxon could feed on fresh microalgae and reproduce rapidly after phytodetritus deposits following the important surface water bloom periods preceding our sampling (Fig. 2). Furthermore, different small opportunistic taxa may colonise different organic-rich patches, increasing local patchiness. Still another possible explanation for the much higher variability of the 63–150  $\mu\text{m}$  fraction could be that reproductive response to phytodetritus inputs of larger taxa would first be noticed by the presence of juveniles in the small




Fig. 7. Plot of the 15 cores in the 2-D space defined by the two main axes of the non-standardised principal component analysis (PCA1, PCA2), and three distinct faunal assemblages (indicated by the dotted lines).

size fraction. This is clearly the case in October 1997, when juvenile *U. peregrina* peak in the 63–150  $\mu\text{m}$  fraction, and in June 1998, when *G. affinis* exhibits a similar reproductive event. Apparently, the response of the > 150  $\mu\text{m}$  fraction is delayed, and increased numbers of the larger opportunistic taxa are found only after the impact of organic input has already been homogenised for a large surface area. Except for cores OB10B and OB10B<sup>bis</sup>, which were collected in two different multi-tube corer deployments, spatial patchiness recorded for each other couple of duplicate samples suggests strong spatial variability of foraminiferal faunas on a scale of less than a meter. Faunal differences between cores OB10B and OB10B<sup>bis</sup> reflect a larger scale (several hundred metres) foraminiferal variability.

Our results suggest that at least part of the foraminiferal patchiness is due to a spatially heterogeneous distribution of organic matter aggregates at the sediment–water interface. Just as heterotrophic bacterial consortia, which can respond quickly to patchy organic matter deposits (Turley et al., 1988; Thiel et al., 1990), some foraminiferal taxa can also take rapid advantage of freshly deposited organic patches. According to

Gooday and Turley (1990), the patchy distribution of the organic material deposited after bloom events play a major role in structuring deep-sea benthic ecosystems.

### 5.3. Temporal variability of bottom and pore water chemical characteristics

Our data show a relative constancy in the chemistry of the bottom and interstitial waters. Bottom water oxygen concentrations are almost invariable throughout the investigated years; values range from 205 to 221  $\mu\text{mol/l}$ . The vertical profiles of dissolved oxygen and nitrate in the uppermost sediment are rather similar for the 1997–2000 period (Figs. 3a and b). Deeper in the sediment bacterial consortia are able to reduce nitrate in anoxic environments in order to decompose the available organic compounds (Froelich et al., 1979; Fenchel and Finlay, 1995). The ammonia that is created during the mineralisation of organic nitrogen migrates upward to oxygenated layers, where it is transformed back into nitrate and nitrite by nitrifying bacterial consortia (Anschutz et al., 1999). Nitrate and nitrite diffuse downward to the anoxic layer, thus creating a

gradient that overlies the oxygen zero boundary. In our cores, the downward nitrate–nitrite gradients always occur in the 0.25–2.5 cm depth range (Fig. 3b). The depth and amplitude variability is rather limited. No clear sulphate reduction zone was detected at station B; i.e. we detected no decrease with depth of sulphate and no significant increase of sulphide (Anschutz, 2002, pers. com.). Apparently, in this deep open-slope environment, the exported organic matter fluxes are not high enough to sustain a sharp sulphate gradient in the sampled cores. However, sulphate reduction may be a discrete phenomenon in the anoxic part of the core.

It seems that the seasonal variability of the exported organic matter flux is not high enough to induce significant fluctuations of the early diagenetic processes in the upper sediment. Apparently, we are far away from the variable conditions prevailing in the lower part of oxygen minimum zone areas (Jannink et al., 1998) or in some oligotrophic basins where sudden organic matter deposits may severely modify redox patterns at and below the sediment–water interface (Kitazato and Ohga, 1995; Kitazato et al., 2000; Duijnste et al., 2001). At our station B, the phytodetritus deposits following main bloom periods (October 1997, June 1999 and April 2000; Table 1) have at most a minor impact on the redox conditions within the sediment and at the sediment–water interface. This is probably because of the overall eutrophic conditions reigning at our site. Fresh organic matter deposits, which are strongly concentrated at the sediment surface, are preferentially used as food by bacteria (Turley et al., 1988; Thiel et al., 1990; Danovaro et al., 2000a,b), but also by meiofaunal foraminifera, which as a whole appear to be an opportunistic group of organisms. This explains why most foraminiferal taxa are concentrated in the first cm and why the highest foraminiferal densities are most times recorded in the first half cm. The restriction of  $^{234}\text{Th}$  (half-life time 24.1 days) activity to the uppermost 1.5 cm, and the vertical aspect of the  $^{210}\text{Pb}$  profile in the upper cm, strongly suggest that macrofaunal bioturbation is largely limited to the topmost part of the sediment (Jouanneau, 2001, pers. comm.). This rather shallow bioturbation zone will cause

an absence of labile components deeper in the sediment. There, a constant, much slower flux of more refractory organic matter sustains a rather invariable succession of redox zones, where stable dysaerobic and anaerobic bacterial pools develop, which may form a food source for some highly specialised deeper dwelling taxa. Thus, the microhabitat of many foraminiferal taxa appears to be controlled largely by their trophic requirements (Jorissen et al., 1995).

#### 5.4. Temporal variability of foraminiferal faunas at station B

*E. exigua* shows the most important frequency variations in our successive samples. Whereas this taxon is almost absent for most of the winter and summer months, it exhibits strong density increases after the spring and autumn phytoplankton bloom periods; density peaks are recorded in October 1997, in June 1999 and April 2000 (Fig. 6). These strong absolute and relative density variations show a highly opportunistic behaviour, and suggest that this species feeds on fresh organic phytodetritus. These results confirm similar observations in other deep-sea environments (Gooday, 1988, 1993; Gooday and Lamshead, 1989; Gooday and Turley, 1990; Thurston et al., 1994; Loubere, 1998; Jannink et al., 1998). Boltovskoy and Lena (1969) show that in eutrophic shallow water environments, *Epistominella* has a very short reproductive cycle (about one month), and reproduces throughout the year. We think that the intermittent occurrences of *E. exigua* at our station B are a direct response to phytodetritus deposits of a rather ephemeral nature. The species responds to the freshly deposited organic matter by a reproductive event, which is especially clear during the spring bloom of the year 2000 (see OB10B<sup>bis</sup> data). Such a reproductive response has also been shown after food addition in culture experiments (Heinz et al., 2001, 2002).

Other small taxa, which are largely restricted to the 63–150  $\mu\text{m}$  fraction, such as *R. guttiferus*, *N. pusillus*, *C. carinata*, and *B. spathulata*, show a similar density variability, albeit to a lesser degree. The peak periods are essentially the same as those for *E. exigua*, but the weaker response of these

taxa suggests a slightly less opportunistic behaviour (Fig. 6).

*Trochammina* spp. and *Bolivina difformis*, which are also limited to the 63–150  $\mu\text{m}$  fraction, show only a strong density increase after the spring bloom of 2000 (Fig. 6). This differential response suggests that not only the quantity, but also the quality, of the organic matter input could be a controlling factor. The quality of the organic matter may vary in function of the different phytoplankton groups responsible for the surface water bloom conditions.

In the >150  $\mu\text{m}$  fraction, the shallow infaunal species *U. mediterranea* and *U. peregrina* show by far the largest frequency variations and can thus be considered as most opportunistic (Figs. 5a–m). *U. peregrina* shows a major peak (in both size fractions) after the putative 1997 autumn bloom. Apparently it is the most opportunistic species occurring in the >150  $\mu\text{m}$  fraction. *U. mediterranea* is responsible for most of the density variations of the total foraminiferal fauna in this larger size fraction, and shows a relative, as well as absolute, frequency increase in all eutrophic periods. The different behaviour of the two *Uvigerina* species suggests slightly different trophic requirements, where *U. peregrina* and other taxa (e.g. *N. pusillus*), which exclusively show a density increase in the October 1997 sample, may prefer a diet based on microflagellates remains, which are especially abundant in the autumn bloom (Sellmer et al., 1998). Species that peak only after the spring bloom, such as *Trochammina* spp. and *B. difformis*, could prefer to feed on diatoms, coccolithophores, or other microalgae typical of spring blooms (Tréguer et al., 1979; Bender et al., 1992; Fernandez et al., 1995). Finally, *U. mediterranea*, and other species occurring both in autumn and spring, may be less critical with respect to food quality.

*U. mediterranea* and *U. peregrina* have been described in a wide range of mesotrophic to eutrophic settings (e.g. Lutze, 1980; Lutze and Coulbourne, 1984; Corliss, 1985, 1991; Corliss and Emerson, 1990; Jannink et al., 1998; Schmiedl et al., 2000; Morigi et al., 2001). According to De Rijk et al. (2000), they would need an exported labile organic flux of at least 2.5 g C/m<sup>2</sup>/year. The

observations made by Fontanier et al. (2002) at OB1B, where *U. peregrina* is found slightly deeper than *U. mediterranea*, suggesting a higher tolerance to slightly more degraded organic matter, are not confirmed during the other sampling periods, where the microhabitat of both species is very similar. However, in all other samples, *U. peregrina* is much poorer than in OB1B, and the present data may not be representative for its optimum conditions.

The infaunal niches deeper in the sediment are permanently occupied by *M. barleeanus*, *U. elongatastriata* and, still deeper, *G. affinis*. A deeper infauna dominated by *M. barleeanus* and *G. affinis* has been described in numerous mesotrophic-eutrophic oceanic ecosystems (e.g. Corliss, 1985; Harloff and Mackensen, 1997; Jorissen et al., 1995, 1998; Schmiedl et al., 2000). Jorissen (1999) and Fontanier et al. (2002) observe that *M. barleeanus* occurs systematically in the lower part of the oxic zone whereas *G. affinis* is consequently found in the upper part of the totally anoxic zone. Our present results fully confirm this pattern, suggesting that deeper infaunal species may be dependent on anaerobic bacterial stocks degrading more or less refractory organic matter. They could either directly prey on the bacterial stocks (Lee, 1979; Kitazato, 1994), or feed on the organic matter breakdown products (Caralp, 1989; Alve, 1990; Bernhard, 1992; Jorissen, 1999). Although we do not think that *G. affinis* and *M. barleeanus* feed directly on fresh organic matter (concentrated in the upper first cm of the sediment), they do show a significant frequency increase (in the >150  $\mu\text{m}$  fraction) in some of the most productive months (October 1997, December 1998; Figs. 5a–m). A very similar increase of the population density was observed by Heinz et al. (2001) in a culture experiment. This suggests that increased surficial biological activity is rapidly transmitted towards deeper sediment layers, perhaps by an increase of bioturbation and accompanying bacterial activity.

In a previous study, Barmawidjaja et al. (1992) suggested that some infaunal taxa (e.g. *Eggerella scabra*, *Morulaepecta bullosa*, *Textularia agglutinans*) could reproduce close to the sediment–water interface with a time lag with respect to more




Fig. 8. Pelagic–benthic coupling and stepwise benthic ecosystem response to phytodetritus input. See text for further explanation.

opportunistic epifaunal taxa. More recently, Kitazato et al. (2000) speculated that deep infaunal taxa (*Chilostomella ovoidea* and *G. affinis*) are able to profit from fresh organic matter supplies after spring blooms, by moving to the sediment–water interface in order to feed on fresh organic matter and to reproduce at the end of the spring bloom. At our site, *G. affinis* shows a clear reproductive event in June 1998, when an abundant fauna consisting exclusively of juvenile specimens is found in the first half cm (Fig. 6). We think that this reproductive event is a delayed response to the 1998 spring bloom, which took place only when all other species have fallen back to background level (Figs. 5a–m and 6). This taxon does obviously not reproduce at the same time as the more opportunistic shallow infaunal taxa (e.g. *E. exigua*), which are more adapted to profiting quickly from labile phytodetritus input. The *G. affinis* reproductive event could be triggered by increased bacterial activity due to the delayed input (by bioturbation) of large amounts of less labile material in dysoxic

microenvironments, once the more labile components have been consumed by more opportunistic taxa. Such a reproductive behaviour would also characterise intermediate infaunal taxa. An increase in the density of *M. barleeanus* in July 1998 (in both size fractions), accompanied by a shoaling of its microhabitat, could represent a similar reproductive event at one of the least productive moments of our 3-year study. In the eutrophic context of the Bay of Biscay, this migrational and reproductive behaviour is apparently independent of changes of redox gradients but is triggered by changes in the trophic level (presence of bacterial pools; quantity and quality of organic remains).

##### 5.5. Stepwise benthic ecosystem response to phytodetritus input

The results of our study show that it is not easy to detect a straightforward relationship between phytoplankton bloom and benthic foraminiferal faunal characteristics. In case of the important

Table 4

| Species  | References  |
|--|---|
| <i>Adercotryma glomerata</i> (Brady), 1878               | Jones (1994), pl. 34, Figs. 15–18                   |
| <i>Ammoscalaria pseudospiralis</i> (Williamson), 1958    | Jones (1994), pl. 33, Figs. 1–4                     |
| <i>Amphicoryna scalaris</i> (Batsch), 1791               | Jones (1994), pl. 63, Figs. 28–31                   |
| <i>Astacolus crepidulus</i> (Fichtel & Moll), 1798       | Jones (1994), pl. 67, Fig. 20; pl. 68, Fig. 1 and 2 |
| <i>Bigenerina nodosaria</i> d'Orbigny, 1826              | Jones (1994), pl. 44, Figs. 19–24                   |
| <i>Biloculinella irregularis</i> (d'Orbigny) 1839        | d'Orbigny (1839), pl. 8, Fig. 20 and 21             |
| <i>Bolivina alata</i> (Seguenza), 1862                   | Schiebel (1992), pl. 1, Fig. 2                      |
| <i>Bolivina difformis</i> (Williamson), 1958             | Cushman (1937), pl. 15, Fig. 5 and 6                |
| <i>Bolivina spathulata</i> (Williamson), 1858            | Jorissen (1987), pl. 1, Fig. 5                      |
| <i>Bolivina striatula</i> (Cushman), 1922                | Cushman (1937), pl. 18, Fig. 30 and 31              |
| <i>Bolivina subaenariensis</i> Cushman, 1922             | Phleger et al. (1953), pl. 7, Fig. 24 and 25        |
| <i>Bolivina variabilis</i> (Williamson), 1958            | Cushman (1937), pl. 16, Figs. 6, 12–14              |
| <i>Bolivinita quadrilatera</i> (Schwager), 1866          | Jones (1994), pl. 42, Figs. 8–12                    |
| <i>Bulimina aculeata</i> d'Orbigny, 1826                 | Jones (1994), pl. 51, Figs. 7–9                     |
| <i>Bulimina costata</i> d'Orbigny, 1826                  | Van Leeuwen (1989), pl. 8, Fig. 2 and 3             |
| <i>Bulimina inflata</i> Seguenza, 1862                   | Van Leeuwen (1989), pl. 8, Fig. 4                   |
| <i>Cancris auriculus</i> (Fichtel & Moll), 1942          | Jones (1994), pl. 106, Fig. 4                       |
| <i>Cassidulina carinata</i> Silvestri, 1896              | Phleger et al. (1953), pl. 9, Figs. 32–37           |
| <i>Cassidulina crassa</i> d'Orbigny, 1839                | Jones (1994), pl. 54, Fig. 4 and 5                  |
| <i>Chilostomella oolina</i> Schwager, 1878               | Jones (1994), pl. 55, Figs. 12–14                   |
| <i>Cibicides lobatulus</i> Walker & Jacob, 1798          | Jones (1994), pl. 92, Fig. 10                       |
| <i>Cibicides wuellerstorfi</i> (Schwager), 1866          | Van Leeuwen (1989), pl. 10, Figs. 1–9               |
| <i>Cibicoides pachydermus</i> (Rzehac), 1886             | Jones (1994), pl. 94, Fig. 9                        |
| <i>Cibicoides robertsonianus</i> (Brady), 1881           | Van Leeuwen (1989), pl. 9, Figs. 1–3                |
| <i>Cibicoides ungerianus</i> d'Orbigny, 1846             | Jones (1994), pl. 94, Fig. 9                        |
| <i>Clavulina cylindrica</i> d'Orbigny, 1952              | Hofker (1932), Fig. 18 and 19                       |
| <i>Cornuspira involvens</i> (Reuss), 1950                | Jones (1994), pl. 11, Figs. 1–3                     |
| <i>Cribrostomoides subglobosus</i> (Cushman), 1910       | Jones (1994), pl. 34, Figs. 8–10                    |
| <i>Cyclammmina cancellata</i> Brady, 1879                | Jones (1994), pl. 37, Figs. 8–16                    |
| <i>Cystammmina pauciloculata</i> (Brady), 1879           | Jones (1994), pl. 41, Fig. 1                        |
| <i>Dentalina advena</i> (Cushman), 1923                  | Jones (1994), pl. 63, Fig. 1                        |
| <i>Dentalina ariana</i> Patterson & Pettis, 1986         | Jones (1994), pl. 62, Figs. 27–31                   |
| <i>Dentalina bradyensis</i> (Dervieux), 1894             | Jones (1994), pl. 62, Fig. 19 and 20                |
| <i>Dentalina submaciata</i> Parr, 1950                   | Jones (1994), pl. 62, Fig. 25 and 26                |
| <i>Eggerella scabra</i> (Williamson), 1858               | Jones (1994), pl. 47, Figs. 15–17                   |
| <i>Elphidium advenum</i> Cushman, 1922                   | Phleger et al. (1953), pl. 6, Fig. 15               |
| <i>Epistominella exigua</i> (Brady), 1884                | Schiebel (1992), pl. 5, Fig. 9                      |
| <i>Gavelinopsis translucens</i> (Phleger & Parker), 1951 | Schiebel (1992), pl. 4, Fig. 5                      |
| <i>Glandulina ovula</i> d'Orbigny, 1846                  | Jones (1994), pl. 61, Figs. 17–22                   |
| <i>Globbulimina affinis</i> d'Orbigny, 1826              | Phleger et al. (1953), pl. 6, Fig. 32               |
| <i>Globocassidulina subglobosa</i> (Brady), 1881         | Jones (1994), pl. 54, Fig. 17                       |
| <i>Gyroidina altiformis</i> Stewart & Stewart, 1930      | Jorissen (1987), pl. 1, Fig. 11                     |
| <i>Gyroidina umbonata</i> (Silvestri), 1898              | Parker (1958), pl. 3, Fig. 19 and 20                |
| <i>Hanzawaia boueana</i> (d'Orbigny), 1846               | Jorissen (1987), pl. 3, Fig. 10                     |
| <i>Hoeglundina elegans</i> (d'Orbigny), 1826             | Phleger et al. (1953), pl. 9, Fig. 24 and 25        |
| <i>Hyalinea balthica</i> (Schroeter), 1783               | Jones (1994), pl. 112, Fig. 1 and 2                 |
| <i>Karrerella bradyi</i> (Cushman), 1911                 | Jones (1994), pl. 41, Figs. 1–4                     |
| <i>Lagena multilatera</i> McMulloch, 1977                | Jones (1994), pl. 58, Figs. 2–3, 7–8, 22–24         |
| <i>Lenticulina peregrina</i> (Schwager), 1866            | Cushman and McCulloch (1950), pl. 39, Fig. 5        |
| <i>Lenticulina vortex</i> (Fichtel and Moll), 1798       | Jones (1994), pl. 69, Figs. 14–16                   |
| <i>Marginula obesa</i> (Cushman), 1923                   | Jones (1994), pl. 65, Fig. 5 and 6                  |
| <i>Melonis barleanus</i> (Williamson), 1858              | Van Leeuwen (1989), pl. 13, Fig. 1 and 2            |
| <i>Nonion scaphum</i> (Fichtel & Moll), 1798             | Jones (1994), pl. 109, Fig. 12                      |
| <i>Nonionella turgida</i> (Williamson), 1858             | Jones (1994), pl. 109, Figs. 17–19                  |

Table 4 (continued)

| Species  | References   |
|--|--|
| <i>Nouria polymorphinoides</i> Heron-Allen & Earland, 1914 | Loeblich and Tappan (1988), pl. 123, Fig. 11 and 12        |
| <i>Nuttallides pusillus</i> (Parr), 1950                   | Phleger et al. (1953), pl. 9, Fig. 5 and 6                 |
| <i>Nuttallides umboniferus</i> (Cushman), 1933             | Van Leeuwen (1989), pl. 15, Figs. 11–13; pl. 16, Figs. 1–7 |
| <i>Oridorsalis umbonatus</i> Reuss, 1851                   | Van Leeuwen (1989), pl. 17, Figs. 1–13                     |
| <i>Psammospaera fusca</i> Schulze, 1875                    | Jones (1994), Pl. 18, Fig. 1–8                             |
| <i>Pseuclavulina crustata</i> Cushman, 1936                | Jorissen (1987), pl. 1, Fig. 1                             |
| <i>Pseudoeponides falsobeccarii</i> Rouvillois, 1974       | Jorissen (1987), pl. 4, Fig. 3a                            |
| <i>Pullenia bulloides</i> (d'Orbigny), 1826                | Phleger et al. (1953), pl. 10, Fig. 19                     |
| <i>Pullenia quinqueloba</i> (Reuss), 1851                  | Jones (1994), Pl. 84, Fig. 14 and 15                       |
| <i>Pyrgo depressa</i> (d'Orbigny), 1826                    | Jones (1994), Pl. 2, Figs. 12, 16 and 17                   |
| <i>Pyrgo subsphaerica</i> d'Orbigny, 1839                  | Cushman (1929), pl. 18, Fig. 1 and 2                       |
| <i>Pyrgoella sphaera</i> (d'Orbigny), 1839                 | Jones (1994), pl. 2, Fig. 4                                |
| <i>Quinqueloculina seminula</i> (Linné), 1758              | Jones (1994), pl. 5, Fig. 6                                |
| <i>Rectuvigerina phlegeri</i> Le Calvez, 1959              | Schiebel (1992), pl. 3, Fig. 10a–d                         |
| <i>Reophax ampullacea</i> Brady, 1881                      | Jones (1994), pl. 30, Fig. 6                               |
| <i>Reophax dentiliniiformis</i> Brady, 1881                | Jones (1994), pl. 30, Fig. 21 and 22                       |
| <i>Reophax fusiformis</i> (Williamson), 1858               | Jones (1994), pl. 30, Figs. 7–10, ?11                      |
| <i>Reophax guttiferus</i> Brady, 1881                      | Jones (1994), pl. 31, Fig. 10–15                           |
| <i>Reophax scorpiurus</i> Montfort, 1808                   | Loeblich and Tappan (1988), pl. 44, Figs. 1–3              |
| <i>Reophax spiculifer</i> Brady, 1879                      | Jones (1994), pl. 31, Fig. 16 and 17                       |
| <i>Rhabdamina cornuta</i> (Brady), 1879                    | Jones (1994), pl. 22, Fig. 11 and 13                       |
| <i>Robertinoides bradyi</i> (Cushman and Parker), 1936     | Jones (1994), pl. 50, Fig. 18                              |
| <i>Sigmoilopsis schlumbergeri</i> Silvestri, 1904          | Jones (1994), pl. 8, Figs. 1–4                             |
| <i>Siphogenerina columellaris</i> (Brady), 1881            | Jones (1994), pl. 75, Figs. 15–17                          |
| <i>Siphotextularia affinis</i> Fornasini, 1883             | Kohl (1985), pl. 2, Fig. 5                                 |
| <i>Siphotextularia concava</i> (Karrer), 1868              | Jones (1994), pl. 42, Figs. 13–14                          |
| <i>Spiroloculina tenuiseptata</i> Brady, 1884              | Jones (1994), pl. 10, Fig. 5 and 6                         |
| <i>Stainforthia concava</i> (Höglund), 1947                | Timm (1992), pl. 5, Fig. 11                                |
| <i>Technitella melo</i> Norman, 1978                       | Jones (1994), pl. 25, Fig. 7                               |
| <i>Textularia agglutinans</i> d'Orbigny, 1839              | Jones (1994), pl. 43, Figs. 1–3                            |
| <i>Textularia conica</i> d'Orbigny, 1839                   | Le Calvez (1977), p. 18, Fig. 1 and 2                      |
| <i>Textularia sagittula</i> DeFrance, 1824                 | Jorissen (1987), pl. 3, Fig. 12                            |
| <i>Textularia truncata</i> Höglund, 1947                   | Le Calvez (1958), pl. 1, Fig. 5                            |
| <i>Trifarina angulosa</i> (Williamson), 1858               | Jones (1994), pl. 74, Fig. 17 and 18                       |
| <i>Trifarina bradyi</i> Cushman, 1923                      | Jones (1994), pl. 67, Figs. 1–3                            |
| <i>Trochammina inflata</i> (Montagu), 1808                 | Jones (1994), pl. 41, Fig. 4                               |
| <i>Uvigerina elongatastriata</i> (Colom), 1952             | Van der Zwaan et al. (1986), pl. 6, Figs. 1–8              |
| <i>Uvigerina mediterranea</i> Hofker, 1932                 | Van der Zwaan et al. (1986), pl. 5, Figs. 1–7              |
| <i>Uvigerina peregrina</i> Cushman, 1923                   | Van der Zwaan et al. (1986), pl. 1, Figs. 1–6              |
| <i>Uvigerina proboscidea</i> Schawger, 1866                | Van der Zwaan et al. (1986), pl. 12, Figs. 1–4             |
| <i>Valvulineria bradyana</i> (Fornasini), 1900             | Jorissen (1987), pl. 4, Fig. 1 and 2                       |

spring phytoplankton blooms of 1999 and 2000 and the putative autumn bloom of 1997, a significant response of the foraminiferal fauna was only noticed about 6 weeks after surface water chlorophyll maxima. Fig. 8 graphically represents our ideas about the time delays between the surface water phytoplankton bloom and the responses in the benthic ecosystem. A first

important delay will exist between the time of maximum chlorophyll-*a* concentrations and phytodetritus deposits at the ocean floor. This delay is caused by the longevity of the phytoplankton assemblages, the time involved in the formation of organic aggregates, and the time for physical transport to the ocean floor. We do not have a precise idea about the first two parameters

Table 5

| Taxa                                | OB1B  |                  | OB1B <sup>bis</sup> |                  | OB2B  |                  | OB2B <sup>bis</sup> |                  | OB3B  |                  | OB4B  |                  | OB5B  |                  |
|-------------------------------------|-------|------------------|---------------------|------------------|-------|------------------|---------------------|------------------|-------|------------------|-------|------------------|-------|------------------|
|                                     | Total | % of total fauna | Total               | % of total fauna | Total | % of total fauna | Total               | % of total fauna | Total | % of total fauna | Total | % of total fauna | Total | % of total fauna |
| Perforate                           |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| Indet.                              | 1     | 0.07             |                     |                  |       |                  | 1                   | 0.19             |       |                  | 1     | 0.36             |       |                  |
| <i>Amphicoryna scalaris</i>         | 4     | 0.30             | 5                   | 1.04             | 1     | 0.14             | 2                   | 0.38             | 3     | 0.75             | 2     | 0.71             | 1     | 0.31             |
| <i>Astacolus</i> sp.                |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Astacolus crepidulus</i>         |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Bolivina alata</i>               | 6     | 0.44             |                     |                  | 7     | 0.97             | 3                   | 0.57             | 2     | 0.50             | 2     | 0.71             | 8     | 2.48             |
| <i>Bolivina spathulata</i>          |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Bolivina subaenariensis</i>      |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Bolivinita quadrilatera</i>      | 8     | 0.59             |                     |                  |       |                  |                     |                  | 1     | 0.25             |       |                  | 1     | 0.31             |
| <i>Bulimina aculeata</i>            |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Bulimina costata</i>             |       |                  |                     |                  |       |                  |                     |                  | 2     | 0.50             | 1     | 0.36             | 3     | 0.93             |
| <i>Bulimina inflata</i>             |       |                  | 2                   | 0.42             |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Bulimina marginata</i>           | 19    | 1.41             | 11                  | 2.29             | 18    | 2.50             | 9                   | 1.70             | 9     | 2.24             | 4     | 1.42             | 1     | 0.31             |
| <i>Cassidulina carinata</i>         |       |                  | 2                   | 0.42             | 2     | 0.28             |                     |                  |       |                  |       |                  |       |                  |
| <i>Cassidulina crassa</i>           |       |                  |                     |                  |       |                  | 1                   | 0.19             |       |                  |       |                  |       |                  |
| <i>Chilostomella oolina</i>         | 3     | 0.22             |                     |                  | 3     | 0.42             | 13                  | 2.46             | 2     | 0.50             | 1     | 0.36             |       |                  |
| <i>Cibicides lobatulus</i>          | 1     | 0.07             |                     |                  | 1     | 0.14             | 3                   | 0.57             | 2     | 0.50             | 2     | 0.71             |       |                  |
| <i>Cibicides wuellerstorfi</i>      | 1     | 0.07             |                     |                  | 6     | 0.83             | 2                   | 0.38             |       |                  | 1     | 0.36             |       |                  |
| <i>Cibicidoides pachydermus</i>     | 63    | 4.67             | 20                  | 4.17             | 52    | 7.22             | 33                  | 6.24             | 13    | 3.23             | 8     | 2.85             | 10    | 3.11             |
| <i>Cibicidoides ungerianus</i>      |       |                  | 11                  | 2.29             |       |                  |                     |                  | 3     | 0.75             | 6     | 2.14             | 3     | 0.93             |
| <i>Dentalina</i> sp.1               | 1     | 0.07             |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Dentalina advena</i>             |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  | 1     | 0.31             |
| <i>Dentalina ariena</i>             |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Dentalina bradyensis</i>         |       |                  |                     |                  | 1     | 0.14             | 2                   | 0.38             | 3     | 0.75             |       |                  |       |                  |
| <i>Dentalina subemaciata</i>        | 1     | 0.07             |                     |                  |       |                  | 2                   | 0.38             |       |                  |       |                  | 1     | 0.31             |
| <i>Elphidium</i> sp.                | 1     | 0.07             |                     |                  |       |                  |                     |                  | 1     | 0.25             |       |                  |       |                  |
| <i>Epistominella</i> sp.            |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Gavelinopsis translucens</i>     | 1     | 0.07             |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Glandulina ovula</i>             | 3     | 0.22             | 1                   | 0.21             |       |                  | 1                   | 0.19             | 3     | 0.75             |       |                  | 1     | 0.31             |
| <i>Globobulimina affinis</i>        | 81    | 6.00             | 11                  | 2.29             | 19    | 2.64             | 39                  | 7.37             | 33    | 8.21             | 18    | 6.41             | 13    | 4.04             |
| <i>Gyroidina altiformis</i>         | 10    | 0.74             | 4                   | 0.83             | 3     | 0.42             |                     |                  |       |                  | 2     | 0.71             |       |                  |
| <i>Gyroidina orbicularis</i>        | 3     | 0.22             | 5                   | 1.04             |       |                  | 2                   | 0.38             | 1     | 0.25             | 1     | 0.36             | 2     | 0.62             |
| <i>Hanzawaia boueana</i>            |       |                  | 2                   | 0.42             |       |                  |                     |                  | 2     | 0.50             | 1     | 0.36             | 1     | 0.31             |
| <i>Hoeglundina elegans</i>          |       |                  |                     |                  |       |                  | 1                   | 0.19             |       |                  |       |                  |       |                  |
| <i>Hyalinea balthica</i>            | 12    | 0.89             | 3                   | 0.63             | 7     | 0.97             | 2                   | 0.38             | 2     | 0.50             |       |                  | 1     | 0.31             |
| <i>Lagena</i> sp.                   |       |                  |                     |                  | 2     | 0.28             | 1                   | 0.19             |       |                  |       |                  |       |                  |
| <i>Lenticulina</i> sp.1             | 1     | 0.07             |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Lenticulina</i> sp.2             |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Lenticulina peregrina</i>        | 7     | 0.52             |                     |                  | 3     | 0.42             | 3                   | 0.57             | 4     | 1.00             | 2     | 0.71             |       |                  |
| <i>Lenticulina vortex</i>           | 3     | 0.22             |                     |                  | 3     | 0.42             | 1                   | 0.19             | 4     | 1.00             | 1     | 0.36             | 2     | 0.62             |
| <i>Maginula obesa</i>               |       |                  |                     |                  | 3     | 0.42             | 1                   | 0.19             |       |                  |       |                  |       |                  |
| <i>Melonis berleeanus</i>           | 121   | 8.97             | 36                  | 7.50             | 69    | 9.58             | 42                  | 7.94             | 40    | 9.95             | 43    | 15.30            | 34    | 10.56            |
| <i>Nodosaria</i> sp.                |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  | 1     | 0.31             |
| <i>Nonion scaphum</i>               |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Nonionella turgida</i>           |       |                  |                     |                  |       |                  |                     |                  | 1     | 0.25             | 1     | 0.36             |       |                  |
| <i>Nuttallides pusillus</i>         |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Nuttallides umboniferus</i>      | 6     | 0.44             | 4                   | 0.83             | 4     | 0.56             |                     |                  | 3     | 0.75             | 2     | 0.71             | 1     | 0.31             |
| <i>Polymorphinidae</i>              |       |                  | 2                   | 0.42             | 3     | 0.42             |                     |                  |       |                  | 1     | 0.36             |       |                  |
| <i>Pseudoepionides falsobecarii</i> | 1     | 0.07             |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Pullenia quinqueloba</i>         |       |                  |                     |                  | 9     | 1.25             | 4                   | 0.76             | 2     | 0.50             | 1     | 0.36             | 6     | 1.86             |



| OB6B  |                  | OB7B  |                  | OB8B  |                  | OB8B <sup>bis</sup> |                  | OB9B  |                  | OB9B <sup>bis</sup> |                  | OB10B |                  | OB10B <sup>bis</sup> |                  |
|-------|------------------|-------|------------------|-------|------------------|---------------------|------------------|-------|------------------|---------------------|------------------|-------|------------------|----------------------|------------------|
| Total | % of total fauna | Total | % of total fauna | Total | % of total fauna | Total               | % of total fauna | Total | % of total fauna | Total               | % of total fauna | Total | % of total fauna | Total                | % of total fauna |
|       |                  |       |                  |       |                  | 1                   | 0.22             | 4     | 0.68             | 1                   | 0.16             | 2     | 0.24             |                      |                  |
| 1     | 0.16             | 5     | 0.90             |       |                  | 1                   | 0.22             |       |                  | 7                   | 1.11             | 9     | 1.07             | 2                    | 0.36             |
| 1     | 0.16             | 1     | 0.18             |       |                  |                     |                  | 1     | 0.17             |                     |                  |       |                  | 1                    | 0.18             |
| 7     | 1.14             | 20    | 3.61             | 1     | 0.41             | 9                   | 1.94             | 3     | 0.51             | 2                   | 0.32             | 18    | 2.13             | 8                    | 1.43             |
|       |                  |       |                  |       |                  |                     |                  |       |                  | 3                   | 0.48             |       |                  |                      |                  |
|       |                  |       |                  |       |                  | 1                   | 0.22             |       |                  |                     |                  | 1     | 0.12             |                      |                  |
|       |                  | 3     | 0.54             | 1     | 0.41             | 2                   | 0.43             | 3     | 0.51             |                     |                  | 6     | 0.71             | 1                    | 0.18             |
|       |                  |       |                  |       |                  | 1                   | 0.22             |       |                  |                     |                  |       |                  |                      |                  |
| 4     | 0.65             |       |                  |       |                  |                     |                  |       |                  | 1                   | 0.16             |       |                  | 1                    | 0.18             |
| 6     | 0.98             | 2     | 0.36             |       |                  | 10                  | 2.15             | 9     | 1.53             | 4                   | 0.63             | 8     | 0.95             | 5                    | 0.90             |
|       |                  |       |                  |       |                  |                     |                  | 1     | 0.17             | 1                   | 0.16             |       |                  | 2                    | 0.36             |
|       |                  |       |                  |       |                  |                     |                  | 1     | 0.17             |                     |                  |       |                  |                      |                  |
| 4     | 0.65             | 3     | 0.54             | 2     | 0.82             | 19                  | 4.09             | 12    | 2.04             | 2                   | 0.32             | 31    | 3.67             | 18                   | 3.23             |
|       |                  | 2     | 0.36             |       |                  |                     |                  |       |                  | 3                   | 0.48             | 2     | 0.24             | 1                    | 0.18             |
| 15    | 2.44             | 23    | 4.15             | 3     | 1.22             | 1                   | 0.22             | 11    | 1.87             | 11                  | 1.74             | 16    | 1.89             | 10                   | 1.79             |
|       |                  | 9     | 1.62             |       |                  | 3                   | 0.65             | 3     | 0.51             | 5                   | 0.79             | 7     | 0.83             | 8                    | 1.43             |
| 1     | 0.16             | 1     | 0.18             | 1     | 0.41             | 1                   | 0.22             |       |                  |                     |                  |       |                  |                      |                  |
|       |                  |       |                  |       |                  |                     |                  | 1     | 0.17             |                     |                  |       |                  |                      |                  |
|       |                  |       |                  | 1     | 0.41             | 1                   | 0.22             | 2     | 0.34             | 4                   | 0.63             | 1     | 0.12             | 1                    | 0.18             |
|       |                  | 2     | 0.36             | 1     | 0.41             | 1                   | 0.22             | 2     | 0.34             | 1                   | 0.16             | 3     | 0.36             | 1                    | 0.18             |
|       |                  |       |                  |       |                  |                     |                  |       |                  |                     |                  |       |                  |                      |                  |
|       |                  |       |                  |       |                  |                     |                  |       |                  | 1                   | 0.16             |       |                  |                      |                  |
|       |                  |       |                  | 1     | 0.41             |                     |                  | 3     | 0.51             | 1                   | 0.16             | 3     | 0.36             | 21                   | 3.76             |
| 55    | 8.94             | 38    | 6.86             | 31    | 12.65            | 71                  | 15.27            | 51    | 8.67             | 24                  | 3.80             | 41    | 4.85             | 21                   | 3.76             |
| 1     | 0.16             | 3     | 0.54             | 1     | 0.41             | 1                   | 0.22             | 1     | 0.17             | 2                   | 0.32             | 1     | 0.12             | 2                    | 0.36             |
| 3     | 0.49             | 1     | 0.18             | 2     | 0.82             | 1                   | 0.22             | 1     | 0.17             | 1                   | 0.16             | 1     | 0.12             | 2                    | 0.36             |
| 1     | 0.16             |       |                  |       |                  | 1                   | 0.22             | 3     | 0.51             | 1                   | 0.16             | 1     | 0.12             | 1                    | 0.18             |
|       |                  |       |                  |       |                  |                     |                  |       |                  |                     |                  |       |                  |                      |                  |
|       |                  | 14    | 2.53             |       |                  | 5                   | 1.08             | 10    | 1.70             | 6                   | 0.95             | 7     | 0.83             | 2                    | 0.36             |
|       |                  | 2     | 0.36             |       |                  | 1                   | 0.22             |       |                  |                     |                  |       |                  |                      |                  |
|       |                  |       |                  |       |                  |                     |                  | 1     | 0.17             |                     |                  |       |                  |                      |                  |
|       |                  |       |                  | 1     | 0.41             |                     |                  |       |                  |                     |                  |       |                  |                      |                  |
|       |                  | 2     | 0.36             |       |                  | 2                   | 0.43             | 3     | 0.51             | 4                   | 0.63             | 3     | 0.36             | 1                    | 0.18             |
| 1     | 0.16             | 3     | 0.54             | 1     | 0.41             | 1                   | 0.22             | 1     | 0.17             | 4                   | 0.63             | 1     | 0.12             | 3                    | 0.54             |
|       |                  |       |                  |       |                  | 1                   | 0.22             |       |                  |                     |                  |       |                  | 1                    | 0.18             |
| 103   | 16.75            | 40    | 7.22             | 10    | 4.08             | 36                  | 7.74             | 39    | 6.63             | 52                  | 8.24             | 61    | 7.22             | 42                   | 7.53             |
|       |                  |       |                  |       |                  |                     |                  |       |                  |                     |                  |       |                  |                      |                  |
|       |                  |       |                  |       |                  |                     |                  |       |                  |                     |                  |       |                  | 1                    | 0.18             |
| 1     | 0.16             | 1     | 0.18             |       |                  |                     |                  | 5     | 0.85             | 7                   | 1.11             |       |                  |                      |                  |
|       |                  |       |                  |       |                  |                     |                  |       |                  |                     |                  |       |                  |                      |                  |
|       |                  | 1     | 0.18             |       |                  |                     |                  | 7     | 1.19             |                     |                  | 1     | 0.12             | 4                    | 0.72             |

Table 5 (continued)

| Taxa                               | OB1B  |                  | OB1B <sup>bis</sup> |                  | OB2B  |                  | OB2B <sup>bis</sup> |                  | OB3B  |                  | OB4B  |                  | OB5B  |                  |
|------------------------------------|-------|------------------|---------------------|------------------|-------|------------------|---------------------|------------------|-------|------------------|-------|------------------|-------|------------------|
|                                    | Total | % of total fauna | Total               | % of total fauna | Total | % of total fauna | Total               | % of total fauna | Total | % of total fauna | Total | % of total fauna | Total | % of total fauna |
| <i>Rectuvigerina phlegeri</i>      |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  | 1     | 0.31             |
| <i>Robertinoides bradyi</i>        | 3     | 0.22             | 1                   | 0.21             | 1     | 0.14             |                     |                  |       |                  |       |                  |       |                  |
| <i>Rosalina</i> sp.                |       |                  |                     |                  | 3     | 0.42             |                     |                  |       |                  | 2     | 0.71             |       |                  |
| <i>Siphogenerina columellaris</i>  | 30    | 2.22             | 2                   | 0.42             | 17    | 2.36             | 12                  | 2.27             | 3     | 0.75             | 13    | 4.63             | 6     | 1.86             |
| <i>Sphaeroidina bulloides</i>      |       |                  | 1                   | 0.21             |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Trifarina</i> sp.               | 1     | 0.07             |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Trifarina angulosa</i>          | 2     | 0.15             |                     |                  | 1     | 0.14             |                     |                  |       |                  |       |                  |       |                  |
| <i>Uvigerina elongatastriata</i>   | 62    | 4.60             | 14                  | 2.92             | 20    | 2.78             | 39                  | 7.37             | 11    | 2.74             | 20    | 7.12             | 20    | 6.21             |
| <i>Uvigerina mediterranea</i>      | 454   | 33.65            | 139                 | 28.96            | 287   | 39.86            | 163                 | 30.81            | 88    | 21.89            | 50    | 17.79            | 74    | 22.98            |
| <i>Uvigerina peregrina</i>         | 290   | 21.50            | 102                 | 21.25            | 44    | 6.11             | 57                  | 10.78            | 36    | 8.96             | 31    | 11.03            | 27    | 8.39             |
| <i>Uvigerina proboscidea</i>       | 1     | 0.07             |                     |                  |       |                  |                     |                  |       |                  | 1     | 0.36             |       |                  |
| Porcellaneous                      |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Biloculina</i> sp.              |       |                  |                     |                  | 1     | 0.14             |                     |                  |       |                  |       |                  |       |                  |
| <i>Biloculina irregularis</i>      |       |                  |                     |                  | 2     | 0.28             |                     |                  | 1     | 0.25             | 1     | 0.36             |       |                  |
| <i>Cornuspira involvens</i>        | 1     | 0.07             |                     |                  | 5     | 0.69             | 5                   | 0.95             | 2     | 0.50             | 8     | 2.85             | 3     | 0.93             |
| <i>Cruciloculina</i> sp.           | 1     | 0.07             |                     |                  | 1     | 0.14             |                     |                  |       |                  |       |                  |       |                  |
| <i>Miliolinella</i> sp.            |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Pyrgo</i> sp.                   |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Pyrgo depressa</i>              | 1     | 0.07             |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Pyrgo subsphaerica</i>          |       |                  | 15                  | 3.13             | 1     | 0.14             |                     |                  |       |                  | 1     | 0.36             |       |                  |
| <i>Pyrgoella sphaera</i>           |       |                  |                     |                  | 1     | 0.14             |                     |                  |       |                  |       |                  |       |                  |
| <i>Quinqueloculina</i> sp.         |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Quinqueloculina seminula</i>    |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Scutularis</i> sp.              |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Sigmoilina</i> sp.              | 1     | 0.07             |                     |                  | 1     | 0.14             |                     |                  |       |                  |       |                  | 1     | 0.31             |
| <i>Spiroloculina tenuiseptata</i>  |       |                  |                     |                  |       |                  |                     |                  | 1     | 0.25             |       |                  |       |                  |
| <i>Triloculina</i> spp.            |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| Non fossilising agglutinated       |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| Indet.                             |       |                  | 3                   | 0.63             |       |                  |                     |                  |       |                  |       |                  | 1     | 0.31             |
| Agglutinated sp.1                  |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Adercotryma glomerata</i>       |       |                  |                     |                  |       |                  | 1                   | 0.19             |       |                  |       |                  |       |                  |
| <i>Ammolagena</i> sp.              | 5     | 0.37             |                     |                  | 1     | 0.14             |                     |                  |       |                  |       |                  |       |                  |
| <i>Ammoscalaria</i> sp.            | 2     | 0.15             | 4                   | 0.83             | 2     | 0.28             |                     |                  | 2     | 0.50             |       |                  | 1     | 0.31             |
| <i>Ammoscalaria pseudospiralis</i> |       |                  |                     |                  |       |                  | 1                   | 0.19             |       |                  |       |                  |       |                  |
| <i>Clavulina cylindrica</i>        | 10    | 0.74             | 4                   | 0.83             | 17    | 2.36             | 11                  | 2.08             | 31    | 7.71             | 14    | 4.98             | 6     | 1.86             |
| <i>Cribrostomoides subglobosus</i> | 20    | 1.48             | 21                  | 4.38             | 30    | 4.17             | 7                   | 1.32             | 45    | 11.19            | 13    | 4.63             | 41    | 12.73            |
| <i>Cyclammia</i> sp.1              | 1     | 0.07             |                     |                  |       |                  | 1                   | 0.19             |       |                  |       |                  |       |                  |
| <i>Cyclammia</i> sp.2              | 6     | 0.44             | 5                   | 1.04             | 6     | 0.83             | 4                   | 0.76             | 4     | 1.00             | 1     | 0.36             | 8     | 2.48             |
| <i>Cyclammia</i> sp.3              | 7     | 0.52             |                     |                  | 8     | 1.11             | 2                   | 0.38             | 3     | 0.75             |       |                  |       |                  |
| <i>Eggerella scabra</i>            | 6     | 0.44             | 1                   | 0.21             | 15    | 2.08             | 2                   | 0.38             | 6     | 1.49             | 12    | 4.27             | 11    | 3.42             |
| <i>Haplophragmoides</i> sp.        |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Haplophragmoides</i> sp.1       |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Haplophragmoides bradyi</i>     | 2     | 0.15             |                     |                  | 2     | 0.28             |                     |                  | 2     | 0.50             | 1     | 0.36             |       |                  |
| <i>Karrerella</i> sp.              |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Martinotiella</i> sp.           |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Psammosphaera fusca</i>         |       |                  |                     |                  |       |                  | 1                   | 0.19             |       |                  |       |                  |       |                  |
| <i>Reophax</i> sp.                 |       |                  |                     |                  | 8     | 1.11             |                     |                  | 2     | 0.50             |       |                  |       |                  |
| <i>Reophax</i> sp.1                | 3     | 0.22             |                     |                  |       |                  | 1                   | 0.19             |       |                  |       |                  |       |                  |

| OB6B  |            | OB7B  |            | OB8B  |            | OB8B <sup>bis</sup> |            | OB9B  |            | OB9B <sup>bis</sup> |            | OB10B |            | OB10B <sup>bis</sup> |            |      |
|-------|------------|-------|------------|-------|------------|---------------------|------------|-------|------------|---------------------|------------|-------|------------|----------------------|------------|------|
| Total | % of total | Total | % of total | Total | % of total | Total               | % of total | Total | % of total | Total               | % of total | Total | % of total | Total                | % of total |      |
|       |            | 2     | 0.36       |       |            |                     |            | 2     | 0.34       | 2                   | 0.32       |       |            |                      |            |      |
| 9     | 1.46       | 12    | 2.17       |       |            | 4                   | 0.86       | 8     | 1.36       | 14                  | 2.22       | 18    | 2.13       | 14                   | 2.51       |      |
|       |            |       |            |       |            |                     |            |       |            |                     |            | 1     | 0.12       | 2                    | 0.36       |      |
| 21    | 3.41       | 42    | 7.58       | 32    | 13.06      | 22                  | 4.73       | 15    | 2.55       | 37                  | 5.86       | 57    | 6.75       | 24                   | 4.30       |      |
| 225   | 36.59      | 178   | 32.13      | 50    | 20.41      | 90                  | 19.35      | 189   | 32.14      | 251                 | 39.78      | 216   | 25.56      | 206                  | 36.92      |      |
| 45    | 7.32       | 19    | 3.43       | 14    | 5.71       | 33                  | 7.10       | 27    | 4.59       | 11                  | 1.74       | 45    | 5.33       | 50                   | 8.96       |      |
|       |            | 1     | 0.18       |       |            | 1                   | 0.22       | 1     | 0.17       | 1                   | 0.16       | 2     | 0.24       | 4                    | 0.72       |      |
|       |            |       |            |       |            |                     |            |       |            | 1                   | 0.16       |       |            |                      |            |      |
| 3     | 0.49       | 1     | 0.18       | 1     | 0.41       |                     |            | 1     | 0.17       | 1                   | 0.16       | 2     | 0.24       | 1                    | 0.18       |      |
|       |            | 1     | 0.18       |       |            | 2                   | 0.43       | 2     | 0.34       | 3                   | 0.48       | 1     | 0.12       |                      |            |      |
|       |            |       |            |       |            |                     |            |       |            |                     |            | 1     | 0.12       |                      |            |      |
| 3     | 0.49       |       |            |       |            |                     |            |       |            | 1                   | 0.16       | 3     | 0.36       | 1                    | 0.18       |      |
| 2     | 0.33       |       |            |       |            |                     |            |       |            |                     |            | 1     | 0.12       | 1                    | 0.18       |      |
|       |            | 1     | 0.18       |       |            | 3                   | 1.22       | 1     | 0.17       |                     |            |       |            |                      |            |      |
|       |            |       |            |       |            |                     |            |       |            |                     |            | 5     | 0.59       |                      |            |      |
|       |            | 1     | 0.18       |       |            |                     |            |       |            |                     |            | 1     | 0.12       |                      |            |      |
|       |            | 1     | 0.18       |       |            |                     | 1          | 0.22  |            |                     |            |       |            |                      | 1          | 0.18 |
| 3     | 0.49       |       |            |       |            | 4                   | 0.86       |       |            |                     |            | 6     | 0.71       |                      |            |      |
|       |            |       |            |       |            |                     |            |       |            | 8                   | 1.27       |       |            |                      |            |      |
|       |            | 7     | 1.26       |       |            |                     |            | 9     | 1.53       |                     |            |       |            |                      |            |      |
|       |            |       |            | 1     | 0.41       |                     |            |       |            | 1                   | 0.16       | 4     | 0.47       | 5                    | 0.90       |      |
| 3     | 0.49       | 4     | 0.72       | 2     | 0.82       | 18                  | 3.87       | 1     | 0.17       | 3                   | 0.48       | 2     | 0.24       | 11                   | 1.97       |      |
|       |            |       |            | 1     | 0.41       |                     |            | 2     | 0.34       | 2                   | 0.32       |       |            |                      |            |      |
| 32    | 5.20       | 13    | 2.35       | 8     | 3.27       | 30                  | 6.45       | 27    | 4.59       | 16                  | 2.54       | 36    | 4.26       | 6                    | 1.08       |      |
| 25    | 4.07       | 7     | 1.26       | 14    | 5.71       | 41                  | 8.82       | 28    | 4.76       | 61                  | 9.67       | 23    | 2.72       | 27                   | 4.84       |      |
|       |            |       |            |       |            |                     |            | 1     | 0.17       |                     |            |       |            | 5                    | 0.90       |      |
| 2     | 0.33       | 5     | 0.90       | 3     | 1.22       |                     |            | 6     | 1.02       | 4                   | 0.63       | 4     | 0.47       | 1                    | 0.18       |      |
| 4     | 0.65       |       |            | 2     | 0.82       | 1                   | 0.22       | 1     | 0.17       | 1                   | 0.16       | 7     | 0.83       | 4                    | 0.72       |      |
| 6     | 0.98       |       |            |       |            | 5                   | 1.08       | 6     | 1.02       | 10                  | 1.58       | 11    | 1.30       | 7                    | 1.25       |      |
|       |            |       |            |       |            |                     |            |       |            | 1                   | 0.16       |       |            |                      |            |      |
|       |            |       |            |       |            | 1                   | 0.22       |       |            | 4                   | 0.63       |       |            |                      |            |      |
|       |            | 3     | 0.54       | 5     | 2.04       |                     |            | 6     | 1.02       | 1                   | 0.16       | 6     | 0.71       |                      |            |      |
|       |            |       |            |       |            |                     |            |       |            | 1                   | 0.16       |       |            |                      |            |      |
|       |            |       |            |       |            |                     |            |       |            |                     |            |       |            |                      |            |      |
|       |            |       |            |       |            | 4                   | 0.86       |       |            | 1                   | 0.16       |       |            | 1                    | 0.18       |      |
|       |            |       |            |       |            |                     |            |       |            |                     |            |       |            | 6                    | 1.08       |      |
|       |            |       |            |       |            |                     |            | 3     | 0.51       | 1                   | 0.16       | 7     | 0.83       | 2                    | 0.36       |      |
|       |            |       |            | 1     | 0.41       | 1                   | 0.22       |       |            | 1                   | 0.16       | 4     | 0.47       | 3                    | 0.54       |      |

Table 5 (continued)

| Taxa                              | OB1B  |                  | OB1B <sup>bis</sup> |                  | OB2B  |                  | OB2B <sup>bis</sup> |                  | OB3B  |                  | OB4B  |                  | OB5B  |                  |
|-----------------------------------|-------|------------------|---------------------|------------------|-------|------------------|---------------------|------------------|-------|------------------|-------|------------------|-------|------------------|
|                                   | Total | % of total fauna | Total               | % of total fauna | Total | % of total fauna | Total               | % of total fauna | Total | % of total fauna | Total | % of total fauna | Total | % of total fauna |
| <i>Reophax dentaliniformis</i>    | 1     | 0.07             |                     |                  |       |                  |                     |                  | 3     | 0.75             | 1     | 0.36             |       |                  |
| <i>Reophax fustiformis</i>        | 7     | 0.52             |                     |                  |       |                  | 5                   | 0.95             |       |                  |       |                  |       |                  |
| <i>Reophax guttiferus</i>         | 19    | 1.41             | 1                   | 0.21             | 14    | 1.94             | 3                   | 0.57             |       |                  |       |                  |       |                  |
| <i>Reophax scorpiurus</i>         | 25    | 1.85             | 27                  | 5.63             | 4     | 0.56             | 19                  | 3.59             | 14    | 3.48             | 9     | 3.20             | 25    | 7.76             |
| <i>Reophax spiculifer</i>         |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Rhabdammina cornuta</i>        |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Saccammina</i> sp.             |       |                  |                     |                  |       |                  | 3                   | 0.57             |       |                  |       |                  |       |                  |
| <i>Technitella melo</i>           |       |                  |                     |                  | 1     | 0.14             | 2                   | 0.38             |       |                  |       |                  | 1     | 0.31             |
| <i>Trochammina</i> sp.            | 2     | 0.15             | 1                   | 0.21             | 1     | 0.14             |                     |                  | 1     | 0.25             |       |                  | 1     | 0.31             |
| Fossilising agglutinated          |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Bigenenerina nodosaria</i>     | 9     | 0.67             | 4                   | 0.83             | 4     | 0.56             | 1                   | 0.19             | 1     | 0.25             | 1     | 0.36             | 3     | 0.93             |
| <i>Karrerulina</i> sp.            |       |                  |                     |                  |       |                  | 1                   | 0.19             |       |                  |       |                  |       |                  |
| <i>Pseudoclavulina crustata</i>   | 10    | 0.74             | 14                  | 2.92             | 3     | 0.42             | 9                   | 1.70             | 4     | 1.00             |       |                  | 2     | 0.62             |
| <i>Sigmoilopsis schlumbergeri</i> |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Siphotextularia affinis</i>    | 9     | 0.67             |                     |                  | 1     | 0.14             |                     |                  | 3     | 0.75             | 1     | 0.36             |       |                  |
| <i>Textularia</i> sp.             |       |                  |                     |                  | 1     | 0.14             |                     |                  | 1     | 0.25             |       |                  |       |                  |
| <i>Textularia agglutinans</i>     |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Textularia conica</i>          |       |                  |                     |                  | 1     | 0.14             |                     |                  |       |                  |       |                  |       |                  |
| <i>Textularia sagittula</i>       |       |                  | 3                   | 0.63             |       |                  |                     |                  | 2     | 0.50             |       |                  |       |                  |
| Total live foraminifera           | 1349  | 100.00           | 480                 | 100.00           | 720   | 100.00           | 529                 | 100.00           | 402   | 100.00           | 281   | 100.00           | 322   | 100.00           |
| Nbr species                       | 53    |                  | 33                  |                  | 52    |                  | 44                  |                  | 45    |                  | 38    |                  | 36    |                  |
| <i>Glomospira</i> spp.            | 56    |                  | 2                   |                  | 63    |                  | 49                  |                  | 2     |                  | 2     |                  | 1     |                  |
| Arborescent indet.                | 203   |                  | 78                  |                  | 50    |                  |                     |                  | 95    |                  | 52    |                  | 59    |                  |
| Ostracoda                         |       |                  | 8                   |                  | 9     |                  |                     |                  |       |                  |       |                  | 4     |                  |

Table 6

| Taxa                           | OB1B  |                  | OB1B <sup>bis</sup> |                  | OB2B  |                  | OB2B <sup>bis</sup> |                  | OB3B  |                  | OB4B  |                  | OB5B  |                  |
|--------------------------------|-------|------------------|---------------------|------------------|-------|------------------|---------------------|------------------|-------|------------------|-------|------------------|-------|------------------|
|                                | Total | % of total fauna | Total               | % of total fauna | Total | % of total fauna | Total               | % of total fauna | Total | % of total fauna | Total | % of total fauna | Total | % of total fauna |
| Perforate                      |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| Indet.                         | 1     | 0.07             | 2                   | 0.60             |       |                  |                     |                  | 2     | 1.52             |       |                  |       |                  |
| <i>Amphicoryna scalaris</i>    | 3     | 0.20             |                     |                  |       |                  |                     |                  | 1     | 0.76             |       |                  |       |                  |
| <i>Animalinoides</i> sp.       | 1     | 0.07             | 3                   | 0.91             |       |                  |                     |                  |       |                  | 2     | 1.40             | 1     | 1.37             |
| <i>Astacolus</i> sp.           |       |                  |                     |                  |       |                  |                     |                  | 1     | 0.76             |       |                  |       |                  |
| <i>Baggina</i> sp.             |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  | 1     | 1.37             |
| <i>Bolivina</i> sp.            | 8     | 0.53             | 2                   | 0.60             | 3     | 1.20             | 1                   | 0.83             | 4     | 3.03             |       |                  |       |                  |
| <i>Bolivina</i> sp.1           |       |                  |                     |                  | 18    | 7.23             |                     |                  |       |                  |       |                  |       |                  |
| <i>Bolivine alata</i>          |       |                  |                     |                  |       |                  | 3                   | 2.48             | 1     | 0.76             |       |                  |       |                  |
| <i>Bolivina difformis</i>      | 10    | 0.66             | 7                   | 2.11             | 1     | 0.40             | 1                   | 0.83             | 1     | 0.76             | 1     | 0.70             | 1     | 1.37             |
| <i>Bolivina spathulata</i>     | 49    | 3.26             | 44                  | 13.29            | 12    | 4.82             | 8                   | 6.61             | 6     | 4.55             | 2     | 1.40             |       |                  |
| <i>Bolivina striatula</i>      | 2     | 0.13             |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Bolivina subaenariensis</i> | 1     | 0.07             |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Bolivina variabilis</i>     | 5     | 0.33             |                     |                  | 2     | 0.80             |                     |                  | 1     | 0.76             |       |                  | 1     | 1.37             |
| <i>Bolivinita quadrilatera</i> | 15    | 1.00             |                     |                  |       |                  | 1                   | 0.83             |       |                  | 2     | 1.40             | 1     | 1.37             |

| OB6B  |            | OB7B  |            | OB8B  |            | OB8B <sup>bis</sup> |            | OB9B  |            | OB9B <sup>bis</sup> |            | OB10B |            | OB10B <sup>bis</sup> |            |
|-------|------------|-------|------------|-------|------------|---------------------|------------|-------|------------|---------------------|------------|-------|------------|----------------------|------------|
| Total | % of total | Total | % of total | Total | % of total | Total               | % of total | Total | % of total | Total               | % of total | Total | % of total | Total                | % of total |
|       |            |       |            | 1     | 0.41       | 1                   | 0.22       | 6     | 1.02       |                     |            | 3     | 0.36       |                      |            |
|       |            | 2     | 0.36       |       |            | 4                   | 0.86       |       |            | 3                   | 0.48       | 14    | 1.66       |                      |            |
| 20    | 3.25       | 44    | 7.94       | 20    | 8.16       | 20                  | 4.30       | 14    | 2.38       | 21                  | 3.33       | 74    | 8.76       | 15                   | 2.69       |
|       |            |       |            |       |            |                     |            | 3     | 0.51       |                     |            | 9     | 1.07       | 2                    | 0.36       |
|       |            | 8     | 1.44       | 4     | 1.63       | 5                   | 1.08       | 20    | 3.40       | 4                   | 0.63       | 19    | 2.25       | 1                    | 0.18       |
|       |            |       |            |       |            | 3                   | 0.65       |       |            | 2                   | 0.32       | 8     | 0.95       | 1                    | 0.18       |
|       |            | 2     | 0.36       | 1     | 0.41       |                     |            | 1     | 0.17       |                     |            |       |            |                      |            |
| 5     | 0.81       | 8     | 1.44       | 3     | 1.22       |                     |            | 3     | 0.51       | 8                   | 1.27       | 27    | 3.20       | 8                    | 1.43       |
| 2     | 0.33       | 14    | 2.53       | 22    | 8.98       | 3                   | 0.65       | 24    | 4.08       | 5                   | 0.79       | 4     | 0.47       | 5                    | 0.90       |
|       |            |       |            |       |            |                     |            | 1     | 0.17       |                     |            |       |            |                      |            |
|       |            | 2     | 0.36       |       |            |                     |            | 2     | 0.34       |                     |            |       |            | 3                    | 0.54       |
|       |            |       |            |       |            |                     |            |       |            | 2                   | 0.32       |       |            |                      |            |
|       |            | 1     | 0.18       |       |            |                     |            |       |            |                     |            |       |            |                      |            |
| 615   | 100.00     | 554   | 100.00     | 245   | 100.00     | 465                 | 100.00     | 588   | 100.00     | 631                 | 100        | 845   | 100        | 558                  | 100.00     |
| 31    |            | 49    |            | 34    |            | 43                  |            | 53    |            | 57                  |            | 52    |            | 55                   |            |
| 3     |            | 52    |            | 67    |            | 35                  |            | 84    |            | 71                  |            |       |            | 34                   |            |
| 38    |            |       |            |       |            | 2                   |            |       |            |                     |            |       |            | 7                    |            |
| 6     |            |       |            |       |            |                     |            |       |            |                     |            |       |            |                      |            |

| OB6B  |                  | OB7B  |                  | OB8B  |                  | OB8B <sup>bis</sup> |                  | OB9B  |                  | OB9B <sup>bis</sup> |                  | OB10B |                  | OB10B <sup>bis</sup> |                  |
|-------|------------------|-------|------------------|-------|------------------|---------------------|------------------|-------|------------------|---------------------|------------------|-------|------------------|----------------------|------------------|
| Total | % of total fauna | Total | % of total fauna | Total | % of total fauna | Total               | % of total fauna | Total | % of total fauna | Total               | % of total fauna | Total | % of total fauna | Total                | % of total fauna |

|  |  |    |      |   |      |   |      |    |      |    |      |    |      |    |      |
|--|--|----|------|---|------|---|------|----|------|----|------|----|------|----|------|
|  |  |    |      |   |      |   |      | 2  | 0.48 |    |      | 2  | 0.52 | 3  | 0.14 |
|  |  | 1  | 0.32 |   |      |   |      |    |      | 2  | 0.46 | 4  | 1.04 | 1  | 0.05 |
|  |  | 4  | 1.29 |   |      | 5 | 3.40 | 1  | 0.24 |    |      | 2  | 0.52 | 6  | 0.27 |
|  |  |    |      |   |      |   |      |    |      |    |      |    |      |    |      |
|  |  | 1  | 0.32 | 3 | 1.08 | 1 | 0.68 | 3  | 0.71 | 4  | 0.91 | 9  | 2.35 | 4  | 0.18 |
|  |  |    |      |   |      |   |      |    |      |    |      | 1  | 0.26 | 1  | 0.05 |
|  |  | 3  | 0.96 | 2 | 0.72 |   |      | 11 | 2.62 | 15 | 3.42 | 27 | 7.05 | 87 | 3.96 |
|  |  | 30 | 9.65 | 9 | 3.24 | 9 | 6.12 | 23 | 5.48 | 27 | 6.16 | 27 | 7.05 | 96 | 4.37 |
|  |  |    |      |   |      |   |      |    |      |    |      |    |      | 2  | 0.09 |
|  |  |    |      | 1 | 0.36 |   |      |    |      |    |      |    |      |    |      |
|  |  | 4  | 1.29 |   |      |   |      | 2  | 0.48 |    |      | 2  | 0.52 |    |      |
|  |  | 1  | 0.32 | 1 | 0.36 | 1 | 0.68 | 3  | 0.71 | 18 | 4.11 | 17 | 4.44 | 10 | 0.46 |



| 0B6B  |                  | 0B7B  |                  | 0B8B  |                  | 0B8B <sup>bis</sup> |                  | 0B9B  |                  | 0B9B <sup>bis</sup> |                  | 0B10B |                  | 0B10B <sup>bis</sup> |                  |
|-------|------------------|-------|------------------|-------|------------------|---------------------|------------------|-------|------------------|---------------------|------------------|-------|------------------|----------------------|------------------|
| Total | % of total fauna | Total | % of total fauna | Total | % of total fauna | Total               | % of total fauna | Total | % of total fauna | Total               | % of total fauna | Total | % of total fauna | Total                | % of total fauna |
| 1     | 2.70             | 2     | 0.64             | 2     | 0.72             |                     |                  | 7     | 1.67             | 1                   | 0.23             | 5     | 1.31             | 8                    | 0.36             |
|       |                  | 1     | 0.32             |       |                  |                     |                  |       |                  |                     |                  |       |                  |                      |                  |
| 3     | 8.11             | 7     | 2.25             |       |                  | 8                   | 5.44             | 12    | 2.86             | 14                  | 3.20             | 8     | 2.09             | 15                   | 0.68             |
| 4     | 10.81            | 25    | 8.04             | 9     | 3.24             | 9                   | 6.12             | 13    | 3.10             | 18                  | 4.11             | 43    | 11.23            | 135                  | 6.15             |
|       |                  | 5     | 1.61             |       |                  | 1                   | 0.68             | 3     | 0.71             |                     |                  | 7     | 1.83             | 11                   | 0.50             |
|       |                  |       |                  |       |                  |                     |                  |       |                  |                     |                  | 1     | 0.26             |                      |                  |
|       |                  | 2     | 0.64             | 14    | 5.04             | 4                   | 2.72             | 6     | 1.43             | 1                   | 0.23             | 5     | 1.31             | 19                   | 0.87             |
|       |                  |       |                  | 1     | 0.36             |                     |                  |       |                  |                     |                  | 1     | 0.26             | 1                    | 0.05             |
|       |                  |       |                  |       |                  |                     |                  |       |                  |                     |                  |       |                  |                      |                  |
|       |                  | 1     | 0.32             |       |                  |                     |                  |       |                  |                     |                  |       |                  | 1                    | 0.05             |
|       |                  | 1     | 0.32             | 1     | 0.36             |                     |                  |       |                  |                     |                  |       |                  |                      |                  |
|       |                  | 3     | 0.96             | 1     | 0.36             | 2                   | 1.36             | 1     | 0.24             |                     |                  | 6     | 1.57             | 8                    | 0.36             |
|       |                  |       |                  |       |                  |                     |                  |       |                  |                     |                  |       |                  | 1                    | 0.05             |
|       |                  |       |                  |       |                  |                     |                  |       |                  |                     |                  |       |                  |                      |                  |
|       |                  |       |                  |       |                  | 1                   | 0.68             |       |                  |                     |                  |       |                  |                      |                  |
|       |                  | 1     | 0.32             |       |                  | 1                   | 0.68             |       |                  |                     |                  |       |                  | 3                    | 0.14             |
|       |                  | 1     | 0.32             |       |                  | 1                   | 0.68             | 2     | 0.48             |                     |                  | 1     | 0.26             | 1                    | 0.05             |
|       |                  | 2     | 0.64             | 2     | 0.72             | 8                   | 5.44             | 136   | 32.38            | 62                  | 14.16            | 23    | 6.01             | 781                  | 35.58            |
|       |                  | 1     | 0.32             |       |                  | 1                   | 0.68             |       |                  | 3                   | 0.68             | 3     | 0.78             | 11                   | 0.50             |
|       |                  |       |                  | 1     | 0.36             |                     |                  |       |                  |                     |                  |       |                  |                      |                  |
|       |                  | 1     | 0.32             |       |                  |                     |                  | 1     | 0.24             |                     |                  |       |                  |                      |                  |
|       |                  | 1     | 0.32             | 1     | 0.36             |                     |                  | 2     | 0.48             |                     |                  | 1     | 0.26             |                      |                  |
|       |                  |       |                  |       |                  |                     |                  |       |                  |                     |                  |       |                  |                      |                  |
|       |                  |       |                  |       |                  |                     |                  | 1     | 0.24             | 1                   | 0.23             | 2     | 0.52             | 5                    | 0.23             |
| 2     | 5.41             | 2     | 0.64             | 8     | 2.88             | 2                   | 1.36             | 19    | 4.52             | 18                  | 4.11             |       |                  | 2                    |                  |
|       |                  |       |                  |       |                  |                     |                  |       |                  |                     |                  |       |                  | 2                    | 0.09             |
|       |                  | 1     | 0.32             |       |                  |                     |                  | 1     | 0.24             |                     |                  |       |                  | 1                    | 0.05             |
|       |                  | 2     | 0.64             | 3     | 1.08             | 1                   | 0.68             | 3     | 0.71             |                     |                  |       |                  | 12                   | 0.55             |
|       |                  |       |                  | 2     | 0.72             |                     |                  |       |                  |                     |                  | 1     | 0.26             |                      |                  |
|       |                  |       |                  |       |                  |                     |                  | 1     | 0.24             |                     |                  |       |                  |                      |                  |
|       |                  | 8     | 2.57             | 2     | 0.72             |                     |                  | 1     | 0.24             | 1                   | 0.23             | 4     | 1.04             |                      |                  |
|       |                  |       |                  |       |                  |                     |                  | 2     | 0.48             | 1                   | 0.23             |       |                  | 2                    | 0.09             |
| 1     | 2.70             | 1     | 0.32             | 1     | 0.36             | 1                   | 0.68             | 3     | 0.71             | 5                   | 1.14             | 4     | 1.04             | 4                    | 0.18             |
|       |                  |       |                  |       |                  |                     |                  |       |                  |                     |                  |       |                  | 1                    | 0.05             |
|       |                  |       |                  |       |                  |                     |                  |       |                  | 1                   | 0.23             |       |                  |                      |                  |
|       |                  | 5     | 1.61             | 10    | 3.60             | 1                   | 0.68             | 3     | 0.71             | 1                   | 0.23             | 5     | 1.31             | 19                   | 0.87             |
|       |                  |       |                  |       |                  |                     |                  |       |                  |                     |                  |       |                  |                      |                  |
|       |                  | 3     | 0.96             | 12    | 4.32             |                     |                  | 6     | 1.43             | 1                   | 0.23             | 3     | 0.78             | 41                   | 1.87             |
|       |                  | 3     | 0.96             | 2     | 0.72             | 5                   | 3.40             | 14    | 3.33             | 8                   | 1.83             | 12    | 3.13             | 48                   | 2.19             |
|       |                  | 7     | 2.25             | 2     | 0.72             |                     |                  | 1     | 0.24             | 3                   | 0.68             |       |                  | 3                    | 0.14             |
|       |                  |       |                  |       |                  |                     |                  |       |                  |                     |                  |       |                  |                      |                  |
|       |                  |       |                  | 3     | 1.08             |                     |                  |       |                  |                     |                  | 2     | 0.52             | 1                    | 0.05             |
|       |                  |       |                  |       |                  |                     |                  |       |                  |                     |                  |       |                  | 1                    | 0.05             |
|       |                  | 1     | 0.32             | 3     | 1.08             | 3                   | 2.04             | 3     | 0.71             | 1                   | 0.23             | 3     | 0.78             | 84                   | 3.83             |
|       |                  |       |                  |       |                  |                     |                  |       |                  |                     |                  |       |                  |                      |                  |
|       |                  |       |                  |       |                  | 1                   | 0.68             |       |                  |                     |                  | 2     | 0.52             |                      |                  |

Table 6 (continued)

| Taxa                               | 0B1B  |                  | 0B1B <sup>bis</sup> |                  | 0B2B  |                  | 0B2B <sup>bis</sup> |                  | 0B3B  |                  | 0B4B  |                  | 0B5B  |                  |
|------------------------------------|-------|------------------|---------------------|------------------|-------|------------------|---------------------|------------------|-------|------------------|-------|------------------|-------|------------------|
|                                    | Total | % of total fauna | Total               | % of total fauna | Total | % of total fauna | Total               | % of total fauna | Total | % of total fauna | Total | % of total fauna | Total | % of total fauna |
| <i>Siphogenerina columellaris</i>  | 46    | 3.06             | 1                   | 0.30             | 3     | 1.20             | 8                   | 6.61             | 5     | 3.79             | 16    | 11.19            |       |                  |
| <i>Stainforthia</i> sp.            | 1     | 0.07             |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Stainforthia concava</i>        | 1     | 0.07             |                     |                  | 1     | 0.40             | 1                   | 0.83             |       |                  |       |                  |       |                  |
| <i>Trifarina angulosa</i>          |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Trifarina bradyi</i>            | 31    | 2.06             | 5                   | 1.51             |       |                  |                     |                  |       |                  | 1     | 0.70             | 2     | 2.74             |
| <i>Uvigerina elongatastriata</i>   | 7     | 0.47             | 4                   | 1.21             | 1     | 0.40             | 2                   | 1.65             |       |                  |       |                  | 1     | 1.37             |
| <i>Uvigerina mediterranea</i>      | 138   | 9.17             | 8                   | 2.42             | 5     | 2.01             | 10                  | 8.26             | 3     | 2.27             | 3     | 2.10             | 5     | 6.85             |
| <i>Uvigerina peregrina</i>         | 250   | 16.61            | 10                  | 3.02             | 5     | 2.01             | 2                   | 1.65             | 4     | 3.03             | 11    | 7.69             | 11    | 15.07            |
| <i>Uvigerina proboscidea</i>       |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Uvigerina</i> spp.              |       |                  | 1                   | 0.30             |       |                  |                     |                  |       |                  |       |                  | 1     | 1.37             |
| Porcellaneous                      |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Biloculinella</i> sp.           |       |                  |                     |                  |       |                  |                     |                  |       |                  | 1     | 0.70             |       |                  |
| <i>Cornuspira involvens</i>        | 4     | 0.27             |                     |                  | 2     | 0.80             | 3                   | 2.48             | 1     | 0.76             | 7     | 4.90             | 2     | 2.74             |
| <i>Cruciloculina</i> sp.           |       |                  |                     |                  |       |                  |                     |                  |       |                  | 1     | 0.70             |       |                  |
| <i>Pyrgo subsphaerica</i>          |       |                  | 4                   | 1.21             |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Quinqueloculina</i> sp.         | 5     | 0.33             | 3                   | 0.91             | 6     | 2.41             |                     |                  |       |                  |       |                  |       |                  |
| <i>Sigmoilina</i> sp.              | 1     | 0.07             | 1                   | 0.30             | 2     | 0.80             |                     |                  |       |                  |       |                  |       |                  |
| <i>Triloculina</i> sp.             |       |                  | 1                   | 0.30             |       |                  |                     |                  |       |                  |       |                  |       |                  |
| Non fossilising agglutinated       |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| Agglut. sp.A                       | 3     | 0.20             | 1                   | 0.30             | 1     | 0.40             |                     |                  |       |                  |       |                  |       |                  |
| Indet.                             | 11    | 0.73             | 2                   | 0.60             | 1     | 0.40             |                     |                  |       |                  |       |                  |       |                  |
| <i>Adercotryma glomerata</i>       | 18    | 1.20             | 8                   | 2.42             | 10    | 4.02             | 1                   | 0.83             |       |                  | 2     | 1.40             | 1     | 1.37             |
| <i>Ammoscalaria</i> sp.            |       |                  | 2                   | 0.60             |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Clavulina cylindrica</i>        |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Cribrostomoides subglobosus</i> | 5     | 0.33             | 12                  | 3.63             | 6     | 2.41             | 1                   | 0.83             | 5     | 3.79             | 1     | 0.70             |       |                  |
| <i>Eggerella scabra</i>            | 1     | 0.07             |                     |                  | 1     | 0.40             |                     |                  |       |                  | 2     | 1.40             |       |                  |
| <i>Haplophragmoides</i> sp.        | 3     | 0.20             | 9                   | 2.72             | 11    | 4.42             |                     |                  | 2     | 1.52             | 1     | 0.70             |       |                  |
| <i>Psammosphaera fusca</i>         | 8     | 0.53             | 18                  | 5.44             | 6     | 2.41             | 5                   | 4.13             |       |                  | 5     | 3.50             |       |                  |
| <i>Reophax</i> sp.1                | 19    | 1.26             | 12                  | 3.63             | 6     | 2.41             | 4                   | 3.31             |       |                  |       |                  |       |                  |
| <i>Reophax dentaliniformis</i>     | 4     | 0.27             |                     |                  | 1     | 0.40             | 3                   | 2.48             |       |                  | 3     | 2.10             | 1     | 1.37             |
| <i>Reophax fusiformis</i>          |       |                  |                     |                  | 1     | 0.40             |                     |                  |       |                  |       |                  |       |                  |
| <i>Reophax guttiferus</i>          | 179   | 11.89            | 14                  | 4.23             | 41    | 16.47            | 5                   | 4.13             |       |                  | 8     | 5.59             |       |                  |
| <i>Reophax scorpiurus</i>          | 1     | 0.07             | 3                   | 0.91             | 18    | 7.23             |                     |                  |       |                  | 2     | 1.40             | 1     | 1.37             |
| <i>Reophax spiculifer</i>          | 2     | 0.13             | 2                   | 0.60             |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Saccamina</i> spp.              | 50    | 3.32             | 1                   | 0.30             |       |                  |                     |                  |       |                  | 2     | 1.40             | 1     | 1.37             |
| <i>Spiroplectinella</i> sp.        |       |                  | 1                   | 0.30             | 4     | 1.61             |                     |                  |       |                  |       |                  |       |                  |
| <i>Techmitella melo</i>            | 15    | 1.00             | 1                   | 0.30             |       |                  | 3                   | 2.48             |       |                  |       |                  |       |                  |
| <i>Trochammina</i> spp.            | 19    | 1.26             | 1                   | 0.30             | 1     | 0.40             | 4                   | 3.31             | 1     | 0.76             | 5     | 3.50             |       |                  |
| Fossilising agglutinated           |       |                  |                     |                  |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Bigenerina nodosaria</i>        | 5     | 0.33             | 11                  | 3.32             | 1     | 0.40             |                     |                  |       |                  |       |                  | 1     | 1.37             |
| <i>Siphotextularia affinis</i>     | 1     | 0.07             | 1                   | 0.30             |       |                  |                     |                  |       |                  |       |                  |       |                  |
| <i>Siphotextularia concava</i>     |       |                  | 1                   | 0.30             |       |                  |                     |                  |       |                  |       |                  |       |                  |
| Total live foraminifera            | 1505  | 100.00           | 331                 | 100.00           | 249   | 100.00           | 121                 | 100.00           | 132   | 100.00           | 143   | 100.00           | 73    | 100.00           |
| Nbr species                        | 65    |                  | 55                  |                  | 49    |                  | 33                  |                  | 30    |                  | 40    |                  | 34    |                  |
| Ostracoda                          | 32    |                  | 5                   |                  | 9     |                  |                     |                  | 2     |                  | 11    |                  | 3     |                  |
| <i>Glomospira</i> spp.             | 14    |                  |                     |                  | 3     |                  | 9                   |                  | 1     |                  | 4     |                  | 6     |                  |



| 0B6B  |                  | 0B7B  |                  | 0B8B  |                  | 0B8B <sup>bis</sup> |                  | 0B9B  |                  | 0B9B <sup>bis</sup> |                  | 0B10B |                  | 0B10B <sup>bis</sup> |                  |
|-------|------------------|-------|------------------|-------|------------------|---------------------|------------------|-------|------------------|---------------------|------------------|-------|------------------|----------------------|------------------|
| Total | % of total fauna | Total | % of total fauna | Total | % of total fauna | Total               | % of total fauna | Total | % of total fauna | Total               | % of total fauna | Total | % of total fauna | Total                | % of total fauna |
| 2     | 5.41             | 17    | 5.47             | 4     | 1.44             | 3                   | 2.04             |       |                  | 3                   | 0.68             | 24    | 6.27             | 29                   | 1.32             |
|       |                  |       |                  |       |                  |                     |                  |       |                  |                     |                  | 2     | 0.52             | 3                    | 0.14             |
|       |                  |       |                  |       |                  |                     |                  |       |                  |                     |                  |       |                  | 22                   | 1.00             |
|       |                  | 6     | 1.93             | 3     | 1.08             | 1                   | 0.68             | 11    | 2.62             | 2                   | 0.46             | 8     | 2.09             | 13                   | 0.59             |
|       |                  | 4     | 1.29             |       |                  | 1                   | 0.68             | 20    | 4.76             |                     |                  |       |                  | 3                    | 0.14             |
| 5     | 13.51            | 11    | 3.54             | 10    | 3.60             | 6                   | 4.08             | 7     | 1.67             | 8                   | 1.83             | 15    | 3.92             | 22                   | 1.00             |
| 3     | 8.11             | 29    | 9.32             | 26    | 9.35             | 2                   | 1.36             | 1     | 0.24             | 24                  | 5.48             | 5     | 1.31             | 72                   | 3.28             |
|       |                  |       |                  |       |                  |                     |                  |       |                  |                     |                  | 1     | 0.26             | 2                    | 0.09             |
|       |                  | 2     | 0.64             |       |                  |                     |                  | 2     | 0.48             | 1                   | 0.23             |       |                  | 4                    | 0.18             |
|       |                  |       |                  |       |                  |                     |                  |       |                  |                     |                  | 2     | 0.52             |                      |                  |
| 1     | 2.70             | 2     | 0.64             |       |                  | 2                   | 1.36             | 1     | 0.24             | 1                   | 0.23             | 1     | 0.26             | 2                    | 0.09             |
|       |                  | 1     | 0.32             |       |                  |                     |                  |       |                  |                     |                  |       |                  |                      |                  |
|       |                  | 3     | 0.96             | 3     | 1.08             | 1                   | 0.68             | 1     | 0.24             | 5                   | 1.14             |       |                  | 9                    | 0.41             |
|       |                  |       |                  |       |                  | 1                   | 0.68             | 3     | 0.71             |                     |                  |       |                  |                      |                  |
|       |                  |       |                  |       |                  |                     |                  |       |                  | 1                   | 0.23             |       |                  |                      |                  |
|       |                  | 5     | 1.61             |       |                  | 2                   | 1.36             | 2     | 0.48             |                     |                  | 4     | 1.04             |                      |                  |
|       |                  |       |                  |       |                  |                     |                  |       |                  |                     |                  |       |                  | 28                   | 1.28             |
| 1     | 2.70             | 8     | 2.57             | 10    | 3.60             | 5                   | 3.40             | 9     | 2.14             | 9                   | 2.05             | 10    | 2.61             | 38                   | 1.73             |
|       |                  | 1     | 0.32             | 1     | 0.36             | 1                   | 0.68             | 2     | 0.48             | 4                   | 0.91             | 1     | 0.26             | 9                    | 0.41             |
|       |                  | 1     | 0.32             |       |                  |                     |                  |       |                  |                     |                  |       |                  |                      |                  |
|       |                  | 3     | 0.96             | 2     | 0.72             | 4                   | 2.72             | 18    | 4.29             | 13                  | 2.97             | 1     | 0.26             |                      |                  |
|       |                  | 4     | 1.29             | 1     | 0.36             |                     |                  |       |                  | 1                   | 0.23             |       |                  | 11                   | 0.50             |
| 1     | 2.70             | 5     | 1.61             | 13    | 4.68             | 10                  | 6.80             | 15    | 3.57             |                     |                  |       |                  | 5                    | 0.23             |
| 1     | 2.70             | 10    | 3.22             | 8     | 2.88             | 5                   | 3.40             | 2     | 0.48             | 3                   | 0.68             | 4     | 1.04             | 35                   | 1.59             |
| 2     | 5.41             | 4     | 1.29             | 13    | 4.68             | 9                   | 6.12             | 3     | 0.71             | 5                   | 1.14             | 2     | 0.52             | 11                   | 0.50             |
| 1     | 2.70             | 1     | 0.32             | 2     | 0.72             | 2                   | 1.36             | 1     | 0.24             |                     |                  |       |                  | 3                    | 0.14             |
|       |                  |       |                  |       |                  |                     |                  | 2     | 0.48             |                     |                  |       |                  | 2                    | 0.09             |
| 6     | 16.22            | 37    | 11.90            | 32    | 11.51            | 17                  | 11.56            | 22    | 5.24             | 116                 | 26.48            | 41    | 10.70            | 237                  | 10.80            |
|       |                  | 3     | 0.96             | 35    | 12.59            | 2                   | 1.36             | 2     | 0.48             | 1                   | 0.23             | 1     | 0.26             | 2                    | 0.09             |
|       |                  |       |                  | 4     | 1.44             |                     |                  |       |                  | 1                   | 0.23             | 8     | 2.09             |                      |                  |
|       |                  |       |                  |       |                  |                     |                  | 1     | 0.24             | 1                   | 0.23             |       |                  |                      |                  |
|       |                  | 1     | 0.32             | 2     | 0.72             |                     |                  |       |                  |                     |                  | 2     | 0.52             | 3                    | 0.14             |
|       |                  | 3     | 0.96             |       |                  |                     |                  | 1     | 0.24             |                     |                  |       |                  | 2                    | 0.09             |
| 4     | 10.81            | 13    | 4.18             | 10    | 3.60             | 3                   | 2.04             | 6     | 1.43             | 26                  | 5.94             | 17    | 4.44             | 194                  | 8.84             |
|       |                  |       |                  |       |                  |                     |                  |       |                  |                     |                  |       |                  |                      |                  |
|       |                  | 6     | 1.93             | 1     | 0.36             | 2                   | 1.36             | 1     | 0.24             | 7                   | 1.60             | 1     | 0.26             |                      |                  |
|       |                  |       |                  | 2     | 0.72             |                     |                  |       |                  |                     |                  |       |                  | 1                    | 0.05             |
|       |                  |       |                  |       |                  |                     |                  | 1     | 0.24             |                     |                  |       |                  | 1                    | 0.05             |
| 37    | 100.00           | 311   | 100.00           | 278   | 100.00           | 147                 | 100.00           | 420   | 100.00           | 438                 | 100.00           | 383   | 100.00           | 2195                 | 100.00           |
| 16    |                  | 56    |                  | 45    |                  | 40                  |                  | 52    |                  | 42                  |                  | 49    |                  | 64                   |                  |
|       |                  | 5     |                  | 11    |                  |                     |                  | 8     |                  | 8                   |                  |       |                  | 13                   |                  |
| 1     |                  | 21    |                  | 16    |                  |                     |                  | 3     |                  | 3                   |                  |       |                  | 12                   |                  |

(Lampitt, 1985); transport to 550 m depth will take about one week (McCave, 1975; Lampitt, 1985; Deuser, 1986). Many publications suggest that the first response in the benthic environment will be by heterotrophic bacteria (Fenchel and Jørgensen, 1977; Lochte and Turley, 1988; Jumars et al., 1989; Rowe, 1991; Della Groce et al., 1996; Danovaro et al., 2000a, b). Meiofaunal organisms, such as ostracods, nematode, and annelids, are thought to be secondary deposit feeders, which will have a delayed response. Larger meiofauna and macrofauna are extremely scarce at our 550 m station (Sorbe, 2001, pers. com.). The first foraminiferal response to phytodetritus supply, about 4–6 weeks after maximum surface water chlorophyll-*a* values, can be seen in the most opportunistic, small surface dwelling taxa like *E. exigua* and *R. guttiferus*. This delay may be shorter, but the temporal resolution of our study does not allow us to precisely determine the exact timing of the response of the foraminiferal opportunistic taxa. However, this estimate corresponds very well with the observations of Heinz et al. (2002), who describe a clear increase of the foraminiferal density about three weeks after food addition in a laboratory experiment. Slightly less opportunistic taxa such as *B. spathulata*, *C. carinata*, *N. pusillus*, *U. peregrina* and *U. mediterranea* could respond somewhat later. In the first phase of benthic foraminiferal response, the degree of patchiness appears to be very high, probably due to the spatial heterogeneity of phytodetritus deposits at the sea floor. Because of the rather superficial depth of bioturbation, the transmission of benthic ecosystem enrichment towards deeper sediment layers is a slow process. This explains why a reproductive event of intermediate and deep infaunal taxa is only noticed 2–3 months after maximum chlorophyll-*a* values (e.g. *G. affinis* in June 1998, and *M. barleeanus* in July 1998).

### Acknowledgements

We would like to thank the French national program PROOF (INSU-CNRS) for sponsoring the OXYBENT program. We have special and kind thoughts for the crews and the captains of the

Côte de la Manche, our scientific ship during all campaigns. We would like to thank the Space Applications Institute/Marine Environment Unit (SAI/ME) from the Joint Research Center (European Commission) (JRC/EC) and more precisely N. Hoepffner and G. Zibordi, for the easy access to the on-line archives of chlorophyll-*a* concentration (SeaWIFS data). We are grateful for the SeaWIFS project (NASA/ GSFC DAAC), which provides very useful satellite images. We also thank Cecilia Laprida, Sandra Langenzaal, Gerhard Schmiedl, Pierre Carbonel and Jean-Claude Sorbe for the very interesting and helpful discussions we had about macro- and meiofaunal ecology in the Bay of Biscay and elsewhere. We have particular thanks for Ralf Schiebel and Luis Lampert for insights and pieces of advice about phytoplankton changes in the northeastern Atlantic Ocean. We are grateful to Jean-Marie Jouanneau for providing us access to his data on radioactive elements in our cores. We thank Jean-Jacques Pichon for his help in identifying the observed phytodetritus components. We highly appreciate the very constructive criticism of two anonymous reviewers.

### Appendix A

Species of benthic foraminifera recognised at station B from the Bay of Biscay, with references to plates and figures in the literature on Atlantic and Mediterranean foraminifera (Table 4).

### Appendix B

Census data for benthic foraminifera in the > 150 µm size fractions for all 15 cores. Data for core OB1B comes from corrected data presented in Fontanier et al. (2002).

N.B. Numbers are not standardised for sediment volume (Table 5).

### Appendix C

Census data for benthic foraminifera in the 63–150 µm size fractions for all 15 cores. N.B.

Numbers are not standardised for sediment volume (Table 6).

## References

- Altenbach, A.V., 1985. Die Biomasse der benthischen Foraminiferen. Auswertungen von "Meteor"-Expedition im östlichen Nordatlantik. Doctorat Thesis, University of Kiel, Germany.
- Altenbach, A.V., 1988. Deep sea benthic foraminifera and flux rate of organic carbon. *Revue de Paléobiologie* (special vol.) 2, 719–720.
- Altenbach, A.V., Sarnthein, M., 1989. Productivity record in benthic foraminifera. In: Berger, W.H., Smetacek, V.S., Wefer, G. (Eds.), *Productivity of the Ocean: Present and Past*. Wiley, Chichester, pp. 255–269.
- Alve, E., 1990. Variations in estuarine foraminiferal biofacies with diminishing oxygen conditions in Dramsfjord, SE Norway. In: Hemleben, C., Kaminski, M.A., Kuhnt, W., Scott, D.B. (Eds.), *Paleoecology, Biostratigraphy, Paleocceanography and Taxonomy of Agglutinated Foraminifera*. Kluwer, Dordrecht, pp. 661–694.
- Anderson, L., 1979. Simultaneous spectrophotometric determination of nitrite and nitrate by flow injection analysis. *Analytica Chimica Acta* 110, 123–128.
- Anschutz, P., Hyacinthe, C., Carbonel, P., Jouanneau, J.M., Jorissen, F.J., 1999. La distribution du phosphore inorganique dans les sédiments modernes du Golfe de Gascogne. *Comptes Rendus de l'Académie des Sciences*, Paris, 328, 765–771.
- Antoine, D., Andre, J.M., Morel, A., 1996. Ocean primary production, 2, estimation at global scale from satellite (coastal zone color scanner) chlorophyll. *Global Biogeochemical Cycles* 10, 57–70.
- Barmawidjaja, D.M., Jorissen, F.J., Puskaric, S., Van der Zwaan, G.J., 1992. Microhabitat selection by benthic foraminifera in the northern adriatic sea. *Journal of Foraminiferal Research* 22, 297–317.
- Barnett, P.R.O., Watson, J., Connely, D., 1984. A multiple corer for taking virtually undisturbed sample from shelf, bathyal and abyssal sediments. *Oceanologica Acta* 7, 399–408.
- Beaufort, L., Heussner, S., 1999. Coccolithophorids on the continental slope of the Bay of Biscay—production, transport and contribution to mass fluxes. *Deep-Sea Research II* 46, 2146–2174.
- Bender, M., Ducklow, H., Kiddon, J., Marra, J., Martin, J., 1992. The carbon balance during the 1989 spring bloom in the North Atlantic Ocean, 47°N, 20°W. *Deep-Sea Research* 39, 1707–1725.
- Berger, W.H., Wefer, G., 1990. Export productivity: seasonality and intermittency, and paleoceanographic implications. *Palaeogeography, Palaeoclimatology, Palaeoecology* 89, 245–254.
- Bernhard, J.M., 1988. Postmortem vital staining in benthic foraminifera: duration and importance in population and distributional studies. *Journal of Foraminiferal Research* 18, 143–146.
- Bernhard, J.M., 1992. Benthic foraminiferal distribution and biomass related to porewater oxygen content: central California continental slope and rise. *Deep-Sea Research* 39, 585–605.
- Billet, D.S.M., Lampitt, R.S., Rice, A.L., Mantoura, R.F.C., 1983. Seasonal sedimentation of phytodetritus to the deep sea benthos. *Nature* 302, 520–522.
- Boltovskoy, E., Lena, H., 1969. Seasonal occurrences, standing crop and production in benthic foraminifera of Puerto Deseado. *Contributions from the Cushman Foundation for Foraminiferal Research XX* (3), 87–95.
- Boucher, J., 1985. Caractéristiques physiques et biologiques. In: Laubier, L., Monniot, C., (Eds.), *Peuplements Profonds du Golfe de Gascogne: Campagne BIOGAS. IFREMER*, pp. 25–40.
- Brunland, K.W., Beinfang, P.K., Bishop, J.K., Eglinton, G., Ittekkot, V.A.W., Lampitt, R., Sarnthein, M., Theide, J., Walsh, J.J., Wefer, G., 1989. Flux to the sea floor. In: Berger, W.H., Smetacek, V.S., Wefer, G. (Eds.), *Productivity of the Ocean: Present and Past*. Wiley, Chichester, pp. 193–215.
- Buzas, M., Culver, S.J., Jorissen, F.J., 1993. A statistical evaluation of the microhabitats of living (stained) infaunal benthic foraminifera. *Marine Micropaleontology* 20, 3–4.
- Caralp, H.M., 1989. Abundance of *Bulimina exilis* and *Melonis barleeaanum*: relationship to the quality of marina organic matter. *Geo-Marine Letters* 9, 37–43.
- Corliss, B.H., 1985. Microhabitats of benthic foraminifera within deep-sea sediments. *Nature* 314, 435–438.
- Corliss, B.H., 1991. Morphology and microhabitat preferences of benthic foraminifera from the northwest Atlantic Ocean. *Marine Micropaleontology* 17, 195–236.
- Corliss, B.H., Chen, C., 1988. Morphotype patterns of Norwegian Sea deep-sea benthic foraminifera and ecological implications. *Geology* 16, 716–719.
- Corliss, B.H., Emerson, S., 1990. Distribution of Rose Bengal stained deep-sea benthic foraminifera from the Nova Scotia continental margin and gulf of maine. *Deep-sea Research* 37, 381–400.
- Cushman, J.A., 1929. *The Foraminifera of the Atlantic Ocean*. Part 6. Miliolidae, Ophthalmitidae and Fischerinidae. Smithsonian Institution, United States National Museum, Bulletin 104.
- Cushman, J.A., 1937. A monograph of the subfamily Virguliniinae of the foraminiferal family Buliminidae. *Spec. Publ. Cushman Lab. Foramin. Res.*, Vol. 9, 1–228, pls. 1–124, Sharon, MA.
- Cushman, J.A., McCulloch, I., 1950. Some Virguliniinae in the collections of the Allan Hancock Foundation. *Alan Hancock Pacific Expedition* 6, 295–364.
- Danovaro, R., Tselepidis, A., Otegui, A., Della Croce, N., 2000a. Dynamics of meiofaunal assemblages on the continental shelf and deep-sea sediments of the Cretan Sea

- (Ne Mediterranean): relationships with seasonal changes in food supply. *Progress in Oceanography* 46, 367–400.
- Danovaro, R., Marralle, D., Dell'Anno, A., Della Groce, N., Tselepides, A., Fabiano, M., 2000b. Bacterial response to seasonal changes in labile organic matter composition on the continental shelf and bathyal sediments of the Cretan Sea. *Progress in Oceanography* 46, 345–366.
- Davis, J.C., 1986. *Statistics and Data Analysis in Geology*, 2nd Edition. Wiley, New York, 656pp.
- Della Groce, N., Danovaro, R., Fabiano, M., Albertelli, G., Tselepides, A., 1996. Benthic bacteria and seasonal changes of organic input in the deep-sea sediments of the Cretan Sea preliminary results. *Journal de Recherche Océanographique* 21, 1 and 2, 21–28.
- De Rijk, S., Jorissen, F.J., Rohling, E.J., Troelstra, S.R., 2000. Organic flux control on bathymetric zonation of Mediterranean benthic foraminifera. *Marine Micropaleontology* 40, 151–166.
- Deuser, W.G., 1986. Seasonal and interannual variations in deep water particles fluxes in the Sargasso Sea and their relation to surface hydrography. *Deep-Sea Research* 33A, 225–246.
- Duijnste, I., de Lugt, I., Vonk Noordegraaf, H., van der Zwaan, B., 2001. Dynamics of benthic foraminifera from northern Adriatic Sea. Doctorat Thesis, University of Utrecht, pp. 39–58 (Chapter 3).
- Fenchel, T.M., Jørgensen, B.B., 1977. Detritus food chains of aquatic ecosystems: the role of bacteria. *Advances in Microbial Ecology* 1, 1–58.
- Fenchel, T.M., Finlay, B.J., 1995. *Ecology and Evolution in Anoxic Worlds*. Oxford University Press, Oxford, 276pp.
- Fernandez, E., 1990. Composicion, distribucion y produccion del fitoplancton en el Cantabrico Central. Thesis doctoral, Universidad de Oviedo, 388pp.
- Fernandez, E., Boyd, P., Holligan, P.M., Harbour, D.S., 1993. Production of organic and inorganic carbon within a large-scale coccolithophore bloom in the northeast Atlantic Ocean. *Marine Ecology Progress Series* 9, 271–285.
- Fernandez, E., Maranon, E., Cabal, J., Alvarez, F., Anadon, R., 1995. Vertical particle flux in outer shelf waters of the southern Bay of Biscay in summer 1993. *Oceanologica Acta* 18, 3, 379–384.
- Fontanier, C., Jorissen, F.J., Licari, L., Alexandre, A., Anschutz, P., Carbonel, P., 2002. Live benthic foraminiferal faunas from the Bay of Biscay: faunal density, composition, and microhabitats, *Deep-Sea Research I*, in press.
- Froelich, P.N., Klinkhammer, G.P., Bender, M.L., Luedke, N.A., Heath, G.R., Cullen, D., Dauphin, P., Hammond, D., Hartman, B., Maynard, V., 1979. Early oxidation of organic matter in pelagic sediments of the Eastern Equatorial Atlantic: suboxic diagenesis. *Geochemica et Cosmochimica Acta* 43, 1075–1090.
- Froidefond, J.M., Castaing, P., Jouanneau, J.M., 1996. Distribution of suspended matter in a coastal upwelling area. Satellite data and in situ measurements. *Journal of Marine System* 8, 91–105.
- Gooday, A.J., 1988. A response by benthic foraminifera to the deposition of phytodetritus in the deep-sea. *Nature London* 332, 70–73.
- Gooday, A., 1993. Deep-sea benthic foraminifera species which exploit phytodetritus: characteristic features and controls on distribution. *Marine Micropaleontology* 22, 187–205.
- Gooday, A.J., Lamshead, P.J.D., 1989. Influence of seasonally deposited phytodetritus on benthic foraminiferal populations in the bathyal northeast Atlantic: the species response. *Marine Ecology Progress Series* 5, 53–67.
- Gooday, A.J., Turley, C.M., 1990. Responses by benthic organisms to inputs of organic material to the ocean floor: A review. *Philosophical Transactions of the Royal Society of London, Series A* 331, 119–138.
- Grassle, J.F., 1989. Species diversity in deep-sea communities. *Trends in Ecology and Evolution* 4, 12–15.
- Grassle, J.F., Morse-Porteous, L.S., 1987. Macrofaunal colonization of disturbed deep-sea environments and the structure of deep-sea benthic communities. *Deep-sea Research* 34, 1911–1950.
- Harloff, J., Mackensen, A., 1997. Recent benthic foraminiferal associations and ecology of the Scotia Sea and Argentine Basin. *Marine Micropaleontology* 31, 1–29.
- Harris, G.P., 1986. *Phytoplankton Ecology: Structure, Function and Fluctuation*. University Press, Cambridge, Cambridge, 384pp.
- Heinz, P., Schmiel, G., Kitazato, H., Hemleben, C., 2001. Response of deep-sea benthic foraminifera from the Mediterranean Sea to simulated phytoplankton pulses under laboratory conditions. *Journal of Foraminiferal Research* 31 (3), 210–227.
- Heinz, P., Hemleben, C., Kitazato, J., 2002. Time-response of cultured deep-sea benthic foraminifera to different algal diets. *Deep-Sea Research I* 49, 517–537.
- Helder, W., Bakker, J.F., 1985. Shipboard comparison of micro- and mini electrodes for measuring oxygen in marine sediments. *Limnology and Oceanography* 30, 1106–1109.
- Herguera, J.C., Berger, W.H., 1991. Paleoproductivity from benthic foraminifera abundance; glacial and postglacial change in the west-equatorial Pacific. *Abstracts with Program Geological Society of America* 23 (5), 107.
- Heussner, S., Durrieu de Madron, X., Radakovitch, O., Beaufort, L., Biscaye, P.E., Carbonne, J., Delsaut, N., Etcheber, H., Monaco, A., 1999. Spatial and temporal patterns of downward particle fluxes on the continental slope of the Bay of Biscay (Northeastern Atlantic). *Deep-sea Research II* 46, 2101–2146.
- Hofker, J., 1932. Notizen uber die Foraminiferen des Golfes von Neapel. III. Die foraminiferen Fauna der Ammontatura. *Publicazioni della Stazione Zoologica di Napoli*, Vol. 12, Fasc. 1.
- Hohenegger, J., Piller, W., Baal, C., 1993. Horizontal and vertical spatial microdistribution of foraminifera in the shallow subtidal Gulf of Trieste, Northern Adriatic Sea. *Journal of foraminiferal Research* 23, 79–101.

- Holligan, P.M., Viollier, M., Harbour, D.S., Camus, P., Champagne-Philippe, M., 1983. Satellite and ship studies of coccolithophore production along a continental shelf-edge. *Nature* 304, 339–342.
- Hyacinthe, C., Anschutz, P., Carbonel, P., Jouanneau, J.M., Jorissen, F.J., 2001. Early diagenetic processes in the muddy sediments of the Bay of Biscay. *Marine Geology* 177, 111–128.
- Jannink, N.T., Zachariasse, W.J., Van der Zwaan, G.J., 1998. Living (Rose Bengal stained) benthic foraminifera from the Pakistan continental margin (northern Arabian Sea). *Deep-Sea Research I* 45, 1483–1513.
- Jones, R.W., 1994. *The Challenger Foraminifera*. Oxford Science Publications—The Natural History Museum, 149pp.
- Jorissen, F.J., 1987. The distribution of benthic foraminifera in the Adriatic Sea. *Marine Micropaleontology* 12, 21–48.
- Jorissen, F.J., 1999. Benthic foraminiferal microhabitats. In: Sen Gupta, B.K. (Ed.), *Foraminifera*. Kluwer Academic Publishers, Dordrecht, pp. 161–179.
- Jorissen, F.J., de Stigter, H.C., Widmark, J.G.V., 1995. A conceptual model explaining benthic foraminiferal microhabitats. *Marine Micropaleontology* 22, 3–15.
- Jorissen, F.J., Wittling, I., Peypouquet, J.P., Rabouille, C., Relexans, J.C., 1998. Live benthic foraminiferal faunas off Cap Blanc, NW Africa: community structure and microhabitats. *Deep-Sea Research I* 45, 2157–2188.
- Jumars, P.A., Altenbach, A.V., De Lange, G.J., Emerson, S.R., Hargrave, B.T., Prahl, F.G., Reimers, C.E., Steiger, T., Suess, E., 1989. Transformation of Seafloor-arriving fluxes into the sedimentary record. In: Berger, W.H., Smetacek, V.S., Wefer, G. (Eds.), *Productivity of the Ocean: Present and Past*. Wiley, Chichester, pp. 291–311.
- Kitazato, H., 1994. Foraminiferal microhabitats in four marine environments around Japan. *Marine Micropaleontology* 24, 29–41.
- Kitazato, H., Ohga, T., 1995. Seasonal changes in deep-sea benthic foraminiferal populations: results of long-term observations at Sagami Bay, Japan. In: Sakai, H., Nozaki, Y. (Eds.), *Biogeochemical Processes and Ocean Flux Studies in the Western Pacific*. Terra Scientific, Tokyo, pp. 331–342.
- Kitazato, H., Shirayama, Y., Nakatsuka, T., Fujiwara, S., Shimanaga, M., Kato, Y., Okada, Y., Kanda, J., Yamaoka, A., Masukawa, T., Suzuki, K., 2000. Seasonal phytodetritus deposition and responses of bathyal benthic foraminiferal populations in Sagami Bay, Japan: preliminary results from “Project Sagami 1996–1999”. *Marine Micropaleontology* 40, 135–149.
- Kohl, B., 1985. Early Pliocene Benthic Foraminifera from the Salina Basin, Southeastern Mexico. *Bulletins of American Paleontology* 88/322, 1–173.
- Laborde, P., Urrutia, J., Valencia, V., 1999. Seasonal variability of primary production in the Cap-Ferret Canyon area (Bay of Biscay) during the ECOFER cruises. *Deep-Sea Research II* 46, 2057–2079.
- Lambhead, P.J.D., Gooday, A.J., 1990. The impact of seasonally deposited phytodetritus on epifaunal and shallow infaunal benthic foraminiferal populations in the bathyal northeast Atlantic: the assemblage response. *Deep-Sea Research* 37, 1263–1283.
- Lampitt, R.S., 1985. Evidence for seasonal deposition of detritus to deep sea floor and its subsequent resuspension. *Deep-Sea Research* 32, 885–897.
- Le Calvez, Y., 1958. Les Foraminifères de La Mer Celtique. *Revue des Travaux del Institut des Pêches Maritimes* 22 (2), 147–209.
- Le Calvez, Y., 1977. Révision des foraminifères de la collection d’Orbigny. II. Foraminifères de l’île de Cuba. *Cahiers de Micropaléontologie* 1, 1–128.
- Le Corre, P., Treguer, P., 1976. Caractéristiques chimiques et planctoniques du Golfe de Gascogne et du Proche Atlantique. Campagne POLYGAS A (20.10 au 04.11.1972) et PHYGAS 32 (24.04 au 08.05.1973). Résultats  $\{\text{tf} = \text{"MacAccR"}\}$   $\{\text{tf} = \text{"Times"}\}$ -sultatsdescampagnesàlamer, *CNEXO9*, 306pp.
- Lee, J.J., 1979. Nutrition and physiology of the foraminifera. In: Levandowsky, M., Hutner, S.H. (Eds.), *Biochemistry and Physiology of Foraminifera*. Academic Press, New York, pp. 42–66.
- Lochte, K., Turley, C.M., 1988. Bacteria and cyanobacteria associated with phytodetritus in the deep sea. *Nature* 333, 67–69.
- Loeblich, A.R., Tappan, H., 1988. *Foraminifera Genera and their Classification*. Van Nostrand Reinhold, New York, 970pp.
- Loeblich, A.R., Tappan, H., 1988. *Foraminifera Genera and their Classification—Plates*. Van Nostrand Reinhold, New York, 212pp.
- Lohrenz, S.E., Knauer, G.A., Asper, V.L., Tuel, M., Michaels, A.F., Knap, A.H., 1992. Seasonal variability in primary production and particle flux in the north-western Sargasso Sea: US JGOFS Bermuda Atlantic Time-series Study. *Deep-sea Research* 39, 1373–1992.
- Loubere, P., 1998. The impact of seasonality on the benthos as reflected in the assemblages of deep-sea foraminifera. *Deep-Sea Research I* 45, 409–432.
- Loubere, P., Fariduddin, M., 1999. Quantitative estimation of global patterns of surface ocean biological productivity and its seasonal variations on time scales from centuries to millennia. *Global Biogeochemical Cycles* 13, 115–133.
- Lutze, G.F., 1980. Depth distribution of benthic foraminifera on the continental margin off NW Africa. “Meteor” Forschungsergebnisse, Reihe C 32, 31–80.
- Lutze, G., Coulbourn, W., 1984. Recent benthic foraminifera from the continental margin off northwest Africa: community structure and distribution. *Marine Micropaleontology* 8, 361–401.
- McCave, I.N., 1975. Vertical flux of particles in the ocean. *Deep-sea Research* 22, 491–502.
- Morigi, C., Jorissen, F.J., Gervais, A., Guichard, S., Borsetti, A.M., 2001. Benthic foraminiferal faunas in surface sediments off NW Africa: Relationship with the organic flux

- to the ocean floor. *Journal of Foraminiferal Research* 31, 350–368.
- Ogawa, N., Tazuin, P., 1973. Contribution à l'étude hydrologique et géochimique du Golfe de Cap-Breton. *Bulletin de l'Institut Géologique du Bassin d'Aquitaine, Bordeaux* 14, 19–46.
- Ohga, T., Kitazato, H., 1997. Seasonal changes in bathyal foraminiferal populations in response to the flux of organic matter (Sagami Bay, Japan). *Terra Nova* 9, 33–37.
- d'Orbigny, A., 1839. Voyage dans l'Amérique méridionale—Foraminifères. 5, 86pp.
- Parker, F.L., 1958. Eastern Mediterranean Foraminifera. Reports of the Swedish deep-sea expedition 1947–1948, vol. VIII: Sediment cores from the Mediterranean and the Red Sea, Vol. 4, 283pp.
- Parsons, T.R., Lalli, C.M., 1988. Comparative oceanic ecology of the plankton communities of the subarctic Atlantic and Pacific Oceans. *Marine Biology Annual Review* 26, 317–359.
- Pfannkuche, O., Thiel, H., 1987. Meiobenthic stocks and benthic activity on the NE-Svalbard Shelf and the Nansen Basin. *Polar Biology* 7, 253–266.
- Phleger, F.B., Parker, F.L., Peirson, J.F., 1953. North Atlantic Foraminifera. Reports of the Swedish deep-sea expedition 1947–1948, Vol. VII: Sediment cores from the North Atlantic, Vol. 1, 122pp.
- Revsbech, N.P., 1983. In-situ measurements of oxygen profiles of sediments by use of oxygen microelectrodes. In: Ganuger, E., Forstner, H. (Eds.), *Polarographic Oxygen Sensors*. Springer, Berlin, pp. 265–273.
- Revsbech, N.P., Jørgensen, B.B., 1986. Microelectrodes: their use in microbial ecology. *Advances in Microbial Ecology* 9, 293–352.
- Rowe, G.T., Sibuet, M., Deming, J., Khripounoff, A., Tietjen, J., Macko, S., Theroux, R., 1991. 'Total' sediment biomass and preliminary estimates of organic carbon residence time in deep-sea benthos. *Marine Ecology Progress Series* 79, 99–114.
- Schiebel, R., 1992. Rezente benthische Foraminiferen in Sedimenten des Schelfes und oberen Kontinentalhanges im Golf von Guinea (Westafrika). *Berichte—Reports. Geologisch-Paläontologisches Institut und Museum Christian-Albrechts-Universität Kiel, Deutschland* 51, 126pp.
- Schmiedl, G., de Bovée, F., Buscail, R., Charrière, B., Hemleben, C., Medernach, L., Picon, P., 2000. Trophic control of benthic foraminiferal abundance and microhabitat in the bathyal Gulf of Lions, western Mediterranean Sea. *Marine Micropaleontology* 40, 167–188.
- Sellmer, C., Fehner, U., Nachtigall, K., Reineke, C., Fritsche, P., Lisok, K., Obermüller, B., Adam, D., 1998. Planktological Studies. In: Mienert, J., Graf, G., Hemleben, C., Kremling, K., Pfannkuche, O., Schulz-Bull, D. (Eds.), *Nordatlantik 1996. Meteor-Berichte* 98–2, pp. 197–200.
- Silva, K.A., Corliss, B.C., Rathburn, A.E., Thunnell, R.C., 1996. Seasonality of living benthic foraminifera from the San Pedro Basin, California Borderland. *Journal of Foraminiferal Research* 26, 71–93.
- Snelgrove, P.V.R., Grassle, J.F., Petrecca, R.F., 1994. Macrofaunal response to artificial enrichments and depressions in the deep-sea habitat. *Journal of Marine Research* 52, 345–369.
- Snelgrove, P.V.R., Grassle, J.F., Petrecca, R.F., 1996. Experimental evidence for aging food patches as a factor contributing to high deep-sea macrofaunal diversity. *Limnology and Oceanography* 41, 605–614.
- Strickland, J.D.H., Parsons, T.R., 1972. A practical handbook of seawater analysis. *Bulletin of Fisheries Resource B Canada* 167, 311.
- Thiel, H., Pfannkuche, O., Schrieber, G., Lochte, K., Gooday, A.J., Hemleben, C., Montoura, R.F.C., Turley, C.M., Patching, J.W., Rieman, F., 1990. Phytodetritus on the deep-sea floor in a central oceanic region of the north-east Atlantic. *Biological Oceanography* 6, 203–239.
- Thomsen, L., Altenbach, A.V., 1993. Vertical and areal distribution of foraminiferal abundance and biomass in microhabitats around inhabited tubes of marine echinurids. *Marine Micropaleontology* 20, 303–309.
- Thurston, M., Bett, B., Rice, A., Jackson, P., 1994. Variations in the invertebrate abyssal megafauna in the North Atlantic Ocean. *Deep-Sea Research I* 41, 1321–1348.
- Timm, S., 1992. Rezente Tiefsee-Benthosforaminiferen aus Oberflächensedimenten des Golfes von Guinea (Westafrika)—Taxonomie, Verbreitung, Ökologie und Korngrößenfraktionen—Reports, Geologisch-Paläontologisches Institut und Museum Christian-Albrechts-Universität Kiel, Deutschland 59, 155pp.
- Tréguer, P., Le Corre, P., Grall, J.R., 1979. The seasonal variations of nutrients in the upper waters of the Bay of Biscay region and their relation to phytoplanktonic growth. *Deep-Sea Research* 26 (10A), 1121–1152.
- Turley, C.M., Lochte, K., Patterson, D.J., 1988. A barophilic flagellates isolated from 4500 m in the mid-North Atlantic. *Deep-Sea Research* 35, 1079–1092.
- Turley, C.M., Gooday, A.J., Green, J.C., 1993. Maintenance of abyssal benthic foraminifera under high pressure and low temperature: some preliminary results. *Deep-Sea Research I* 40, 643–652.
- Van der Zwaan, G.J., Jorissen, F.J., Verhallen, P.J.J.M., Von Daniels, C.H., 1986. Atlantic-European Oligocene to recent *Uvigerina*: taxonomy, paleoecology and paleobiogeography. *Utrecht Micropaleontological Bulletins* 35, 240pp.
- Van Leeuwen, R.J.W., 1989. Sea-floor distribution and Late Quaternary faunal patterns of planktonic and benthic foraminifera in the Angola Basin. *Utrecht Micropaleontological Bulletins* 38, 288pp.
- Wroblewski, J.S., 1989. A model of the spring bloom in the North Atlantic and its impact on ocean optics. *Limnology and Oceanography* 34, 1563–1571.
- Walton, W.R., 1952. Techniques for recognition of living foraminifera. *Contributions from the Cushman Foundation for Foraminiferal Research* 3, 56–60.