

Seasonal dynamics and spatial distribution of lepidopterans in selected locations in Mordovia, Russia

ALEXANDER B. RUCHIN

Joint Directorate of the Mordovia State Nature Reserve and National Park «Smolny», Russia.
e-mail: ruchin.alexander@gmail.com

Manuscript received: 8 January 2021. Revision accepted: 16 April 2021.

Abstract. Ruchin AB. 2021. *Seasonal dynamics and spatial distribution of lepidopterans in selected locations in Mordovia, Russia. Biodiversitas 22: 2569-2575.* The research was conducted in 2019-2020 in the Republic of Mordovia (European part of Russia). In order to collect Lepidoptera, beer traps were used where beer served as bait. The field surveys were carried out from April to October in various forest habitats. The air temperature was recorded. To elucidate the spatial distribution of Lepidoptera, various habitats were examined in two study plots. In total, more than 23 thousand Lepidoptera specimens were examined. The largest number of Lepidoptera was collected in oak forests in contrast to the four other forest habitats. In all habitats, population dynamics were similar and characterized by the same trends of decrease and increase. During the season, there were three peaks of abundance for this group. A moderate first peak was recorded in the second half of May, while the maximum peak occurred in the first half of July. Autumn population peaks depend on temperature, while spring and summer peaks are associated with the flight of imagos and an increase in the lepidopteran abundance in habitats. In open habitats, the abundance of Lepidoptera was lower. On forest edges, the number of Lepidoptera considerably exceeded one in open habitats (river sandbanks, willow thickets, forest glades, and floodplain meadows) and closed habitats (inside the forests). Forest edges are a hotspot of Lepidoptera abundance and biodiversity in forest habitats. According to the vertical gradient, the number of lepidopterans was higher at the height of 7.5 m than at the height of 1.5 m. At the lower height, the number of Lepidoptera did not vary as considerably as at higher heights.

Keywords: Butterflies, dynamics, insects, Mordovia, State Nature Reserve

INTRODUCTION

The spatial distribution of species and individual taxonomic groups among habitats has been studied for quite a long time (Chursina and Ruchin 2018; Cicort-Lucaciu 2020; Proshchalykin and Sergeev 2020; Sergeev and Makarkin 2021). For example, many studies were devoted to the vertical distribution of various insect groups (Sutton and Hudson 1980; DeVries 1988; Rubik 1993; Kirstova et al. 2017). In temperate forests, parasitic Hymenoptera is more abundant in the herb layer of the forests than at higher heights (Preisser et al. 1998). In other experiments, a considerable proportion of Vespidae specimens were caught by traps established in the forest crowns (Ulyshen et al. 2011). Giovanni et al. (2017) showed that in the undergrowth layer, most amount of the Sphecidae community consists of species preying on dipterans and spiders. The sex ratio of syrphids differed considerably between the two layers, as females were caught mainly on the ground, while males were caught mainly in the canopy (Birtele and Hardersen 2012). Some differences in the height preferences of Diptera were found during surveys using beer traps (Dvořák et al. 2020). Other studies indicated an increase in the Diptera abundance in the forest canopy (Maguire et al. 2014; Gossner et al. 2016; Krivosheina and Krivosheina 2019).

The largest number of Neuroptera species was caught in the tree crowns in five various forest areas (Gruppe and Schubert 2001). Saure and Klelhorn (1993) found 22 and

24 species in the crowns of pine and oak, respectively. According to Duelli et al. (2002), Neuroptera showed their highest species abundance in the glacier belt and in the crowns. In the forest depth, the species number peaked in the canopy. High level of Chrysopidae biodiversity has been identified in the tree canopy (Makarkin and Ruchin 2019). Some species of Scolytinae (Coleoptera) were associated with traps exposed at heights of 7-21 m, while other species were associated with heights of 1.2 m (Procházka et al. 2018). The vertical stratification of Chrysomelidae was more pronounced in wet habitats than in dry ones (Charles and Basset 2005; Sergeev 2020). The distribution of Cerambycidae beetles was dependent on the height (Graham et al. 2012). The abundance and biomass of Cetoniinae were higher in trap set at a height of 10.5 m. The abundance of Rutelinae is higher at a height of 4.5, 7.5 and 10.5 m (Puker et al. 2020).

There were clear differences in diversity and abundance of lepidopterans between canopy and understory in the rainforests of Brazil (Santos et al. 2017). Schulze et al. (2001) showed that under forest canopy there is a unique butterfly fauna that differs from the Lepidoptera fauna of the surface layers. In addition, the Lepidoptera biodiversity varies in space not only vertically, but also horizontally. Thus, the variety of Lepidoptera in naturally occurring lake edge is very different from the pasture-forest edge. The comparison showed that in natural and disturbed forest areas, the distribution of species abundance has considerable differences (Devries et al. 1999). For our

studies of Lepidoptera, we used beer traps with baits, which have simple construction.

MATERIALS AND METHODS

Study area and field survey design

All studies were conducted in the Republic of Mordovia in 2019-2020 (Temnikov district, Mordovia State Nature Reserve and its immediate surroundings) (Figure 1). Mordovia State Nature Reserve is located on the right bank of the Moksha River and covers an area of 321.62 km². According to the natural zoning, the forest area of the reserve is included in the zone of coniferous-broad-leaved forests on the border with the forest steppe. Forest covers 89.3% of the total area. Pine (*Pinus sylvestris* L.) is the main forest-forming species in the Mordovia State Nature Reserve. It forms pure or mixed plant communities in the southern, central and western parts of the Protected Area. Birch (*Betula pendula* Roth) forests occupy the second

rank of the forest-forming tree species in the Mordovia State Nature Reserve. These are mainly secondary communities in the areas of cuttings and burnt pine forests. Oak (*Quercus robur* L.) forests occupy a relatively small area of the Mordovia State Nature Reserve. They are common in the Moksha River floodplain in the western part. Spruce (*Picea abies* L.) forests and alder (*Alnus glutinosa* (L.) Gaertn.) forests are located mainly in floodplains of rivers and streams by occupying small areas. The main areas of floodplain meadows are located along the Moksha River in the southwest of Mordovia State Nature Reserve. The protected area borders the Nizhny Novgorod region in the north.

Collection of Lepidoptera specimens was carried out by crown traps of our own design. A five-liter plastic container with a window cut out on one side at a distance of 10 cm from the bottom was used as a trap (Ruchin et al. 2020). Beer or wine was used as bait. For fermentation, sugar, jam, and honey were added in each specific case.




Figure 1. Research locations in the Mordovia State Nature Reserve, Russian Federation and its surroundings. Note: A. Surroundings of the cordons Taratinsky; B. Surroundings of the Sosnovka village

Seasonal experiments

To study seasonal dynamics, traps were placed in five various habitats (a detailed description of the habitats is given in Ruchin et al. (2021). The habitat was named by the prevailing tree species in them: 1 - the forest area with a predominance of *Pinus sylvestris* (pine forest); 2 - the forest area with a predominance of *Tilia cordata* (lime forest); 3 - the forest area with a predominance of *Populus tremula* (aspen forest); 4 - the forest area with a predominance of *Betula pendula* (birch forest); 5 - the forest area with a predominance of *Quercus robur* (oak forest).

In each habitat, two traps were installed under the forest canopy at a distance of 5 m from each other. Traps were hung on tree trunks in a crown 7m to 8m in height. The sampling period ranged from six to 17 days. The studies were conducted from April to October 2019. The air temperature was measured daily during the day and night (maximum and minimum values). During the experiment, 9012 *Lepidoptera* individuals were captured and examined.

Horizontal and vertical distribution

In these experiments, both the horizontal distribution of *Lepidoptera* over individual habitats and the vertical distribution within each habitat were studied (Table 1).

Pairs of traps at two heights (below-above) within the same habitat were located side by side (4-5 m from each other). The studies were conducted from April to August 2020. In total, 10 collections (expositions) were carried out. During the experiment, 6266 *Lepidoptera* individuals were captured and examined.

Horizontal distribution

In these experiments, we studied the horizontal distribution of *Lepidoptera* over individual habitats. All traps were located at a height of 1.5 m (Table 2).

The studies were conducted from May to August 2020. In total, 12 collections (expositions) were carried out. During the experiment, 8296 *Lepidoptera* individuals were captured and examined.

Data analyses

When analyzing the results, we used only data on the quantitative parameter (number) of all *Lepidoptera* specimens in traps during the exposure time. Determination of the *Lepidoptera* species was difficult due to poor quality of specimens and the inability to determine the collected material. Exposure time is the period between hanging a trap and taking samples for analysis (expressed in days). All data from individual collections were averaged for the entire duration of the experiments (Experiments 2 and 3).

Table 1. Brief description of the habitat

Name of the habitat	Description of the habitat	Height of the trap location, m
River sandbanks	Open habitat, 5 m from the edge of the Moksha River; herbaceous vegetation is weak	1.5
Willow thickets	Near the floodplain deciduous forest, transitional habitat	1.5
Glade in the forest	Glade in the floodplain deciduous forest, overgrown with nettle	1.5 below
Glade in the forest	Glade in the floodplain deciduous forest, overgrown with nettle	7.5 above
Edge of the forest	Edge of the floodplain deciduous forest, western side, maximum sun illumination	1.5 below
Edge of the forest	Edge of the floodplain deciduous forest, western side, maximum sun illumination	7.5 above
Floodplain meadow	Center of the floodplain meadow, grassy cover is well-developed, herb layer height is up to 1.2 m; herb species diversity is considerable	1.5
Edge of the forest	Edge of the floodplain deciduous forest, eastern side	1.5 below
Edge of the forest	Edge of the floodplain deciduous forest, eastern side	7.5 above
In the depth of the forest	In the depth of the floodplain deciduous forest	1.5 below
In the depth of the forest	In the depth of the floodplain deciduous forest	7.5 above
Edge of the forest	Edge of the floodplain deciduous forest, north side, no direct sunlight	1.5 below
Edge of the forest	Edge of the floodplain deciduous forest, north side, no direct sunlight	7.5 above

Table 2. Brief description of the habitat

Name of the habitat	Description of the habitat
Open habitat	Dry meadow; herbaceous layer is represented by various cereals
The edge of a mixed forest	A young forest appeared as a result of self-seeding in a meadow
Open habitat	Dry meadow; herbaceous layer is represented by various cereals and perennial herbs
Open habitat	Moistened meadow, in a hollow which has water in spring; herbaceous layer is represented by various perennial herbs
The edge of a young pine forest	The edge of a young pine forest formed as a result of self-seeding in a meadow
In the depth of a medium-aged birch forest	The herbaceous layer is poorly expressed, high shading occurs due to the high crown density

RESULTS AND DISCUSSION

Experiments demonstrated that the highest number of Lepidoptera (more than twice) was found in oak forests. The number of specimens differed little from other habitats. Floodplain deciduous forests are very abundant in the species diversity of plants in both herbaceous and understory layers. Apparently, this is the reason for the high number of Lepidoptera there. Gilbert and Singer (1975) and Shapiro (1975) suggested that the relationships of lepidopterans with host plants and climate may explain much of the distribution patterns of the insects. It is also known that there is a direct relationship between the abundance of Lepidoptera and the species diversity of plants (Root et al. 2017).

Similar to majority of other insects, the seasonal cycles of lepidopterans are strictly related to seasonal changes in temperature, day duration, humidity, and other factors. Abundance is influenced by climatic factors that determine the conditions of reproduction and survival (Owen 1971). Seasonality more often depends on the change of rainy and dry seasons in the tropics (Owen et al. 1972; Ribeiro et al. 2010; Grøtan et al. 2012). Photoperiod and temperature indicate the main impact on seasonal phenomena in temperate forests (Roy and Sparks 2000; Altermatt 2012; Zografou et al. 2014; Brooks et al. 2017; Ruchin et al. 2018; Colom et al. 2021).

The seasonal dynamics of Lepidoptera abundance were expressed quite well and it was quite natural (Figure 2). The abundance dynamics were similar in all habitats. In late April, the number of Lepidoptera specimens was very low. Then there was a gradual increase in this parameter. During the season, there were three peaks in the number of lepidopterans. A small first peak in numbers was recorded in the second half of May. Apparently, it was associated

with the appearance of imago of the first spring species from various families.

The most considerable peak of Lepidoptera abundance was found in the first half of July. It was clearly associated with the flight and activity of imago in late spring, and summer species after the May high temperatures, which contributed to the accelerated development of larvae. This peak was preceded by a decline in the Lepidoptera number in late June - early July, which is associated with certain diapauses before the appearance of adult insects. The second peak was recorded in early October. In our opinion, this peak was not associated with the Lepidoptera reproduction, but with the activity of these insects. This is indicated by the correlation (ratio of decreases and increases in number) of the trends of the temperature dynamics in September and October and trends of the number of Lepidoptera. The October peaks (most likely all autumn peaks) of increase in air temperatures contribute to the increase in the activity of Lepidoptera in all habitats.

Figure 3 shows the results of experiments on the spatial distribution of Lepidoptera in individual habitats and within each habitat.

In open areas (meadows, river sandbanks), the number of Lepidoptera was low. In the forest depth (habitats 10 and 11), the number of specimens was less than at the nearest edge of the forest (habitats 8 and 9). In low-level traps (h=1.5) (habitats 8 and 10), the number of Lepidoptera was almost the same (26.3 and 26.6 specimens/day, respectively). This means that the activity of Lepidoptera species differs little at low heights.

The highest values of Lepidoptera abundance were obtained in the fringe habitats (Figure 3). In total, all three habitats showed high and slightly different results on the Lepidoptera abundance (Figure 4).




Figure 2. Seasonal abundance of Lepidoptera specimens collected in various habitats from April to October. The graph shows the dynamics of day and night temperatures. Y-axis indicates a numerical expression (number in specimen per day, temperature in °C)




Figure 3. The average number of *Lepidoptera* (in specimen/day) in all habitats in experiment No. 2




Figure 4. The number of *Lepidoptera* (in specimens/day) in three habitats on the forest edges. There are data about different heights and total values




Figure 5. The ratio of the number of *Lepidoptera* individuals in beer traps located 1.5 m from the ground (below) and in the forest canopy at a height of 7-8 m (above)

However, we noted certain differences in the number of lepidopterans caught at different heights. Thus, it was observed that at a height of 1.5 m, the number of *Lepidoptera* on the western edge significantly exceeded the number of *Lepidoptera* at a similar height in two other habitats. On the contrary, at a height of 7.5 m, the number of *Lepidoptera* on the northern edge was higher than in edges of other expositions.

We assume that in this case, this ratio was influenced by microclimatic conditions, such as illumination,

humidity, wind, etc. Especially the first two factors could have an impact on this ratio. *Lepidoptera* is heliothermic organisms, their flight depends on sunlight intensity (Shapiro 1975). Therefore, open sunny edges are the most preferred for butterflies (Kuussaari et al. 2007). In floodplain meadows in the summer months, there is a frequent drop of dew on the herbaceous layer. Usually, the dew disappears after the sun rises and the surface layer of air warms up. However, at the northern edge, in contrast to the western and eastern edges, the surface layer of air warms up more slowly. In such conditions, in this habitat near the herbaceous cover, the temperature is lower than in similar conditions on other edges. At the same time, temperature differences are less pronounced in the tree crowns. It is possible that it is precise because of such microclimatic conditions that the ratio of *Lepidoptera* abundance changes towards an increase in the number of individuals in the upper traps.

Figure 5 shows data on the ratio of *Lepidoptera* abundance at two heights in various locations. In all the habitats, this ratio was shifted towards an increase in the number of *Lepidoptera* at a height of 7.5 m. In the forest depths, in a clearing in the forest, and in the eastern edge, the ratio of *Lepidoptera* was twice higher in traps at two heights. However, on the western and northern edges, this ratio changed in the direction of an increase in the number in the upper traps (Figure 4).

The study of the vertical *Lepidoptera* distribution in temperate deciduous forests demonstrated that there are insect communities unevenly distributed vertically. These patterns are determined by multiple factors acting simultaneously (Ulyshen 2011). Many studies demonstrated that there are also certain patterns in the vertical *Lepidoptera* distribution (DeVries et al. 1997; Walla et al. 2004; Santos et al. 2017).

At low height, the number of *Lepidoptera* in various habitats did not vary as much as at higher heights (Figure 6). As we pointed out above, especially in this case, the western edge was distinguished, where the highest number of individuals was observed. On the northern edge, the number of *Lepidoptera* was lower at low heights than on the river sandbanks, where the herbaceous layer and plant species diversity serving for feeding *Lepidoptera* are much lower. Apparently, the microclimatic conditions described above are crucial in the *Lepidoptera* activity and the attraction of this group to bait.

A separate experiment on the horizontal *Lepidoptera* distribution in individual habitats showed that forest edges adjacent to meadow habitats play a crucial role in the number of *Lepidoptera* specimens (Figure 7).

At the edge of the young mixed forest, the number of *Lepidoptera* individuals exceeded the one in the meadow habitat, located 30 m away, by almost three times. The number of *Lepidoptera* was 2.5 times higher at the edge of a young pine forest than in a nearby meadow habitat (No. 4). At the same time, the differences between meadow habitats with similar conditions (No. 1 and No. 3) were minimal. However, in a more humid meadow habitat, the number of *Lepidoptera* differed from them.




Figure 6. The average number of Lepidoptera (specimens/day) in all habitats at a height of 1.5 m




Figure 7. The average number of Lepidoptera (specimens/day) in all habitats in experiment No. 3

Despite certain differences in the population ratio at various heights (Figure 5), forest edges are a hotspot of Lepidoptera abundance and biodiversity in forest habitats (Figures 3 and 7). This is confirmed by other researchers. For example, Lepidoptera species abundance was highest in semi-natural meadows and forest edges (Kuussaari et al. 2007). In the eucalyptus forest and at its edge, the diversity of butterflies was higher than in the adjacent habitats (Bragança et al. 1998). In natural parks of Brazil, similar regularities were found expressed in the increase in the Lepidoptera species richness on the forest edges (Melo et al. 2019).

In conclusion, the largest number of Lepidoptera was found in oak forests in contrast to other four forest habitats. In all habitats, population dynamics were similar, being characterized by the same rates of decrease and increase. In late April, the number of lepidopterans was low. Then there was a gradual increase in the number of traps. During the season, there were three peaks in the number of lepidopterans. A small first peak was recorded in the second half of May, and the maximum peak occurred in the first half of July. This peak was preceded by a decrease in numbers in late June - early July, associated with certain diapauses before the appearance of adult butterflies. The second peak was recorded in early October. Autumn population peaks depend on temperature, while spring and summer peaks are associated with the flight of adults and an increase in the number of butterflies in habitats.

In open habitats, the number of Lepidoptera was usually lower. The highest values of Lepidoptera abundance were obtained in forest edge habitats. Thus, forest edges are a hotspot of abundance and biodiversity in forest habitats. The ratio of the number of butterflies in all habitats was shifted towards an increase at a height of 7.5 m. At low heights, the number of Lepidoptera did not vary as much as at higher heights.

REFERENCES

- Altermatt F. 2012. Temperature-related shifts in butterfly phenology depend on the habitat. *Glob Chang Biol* 18: 2429-2438. DOI: 10.1111/j.1365-2486.2012.02727.x.
- Birtel D, Hardersen S. 2012. Analysis of vertical stratification of Syrphidae (Diptera) in an oak-hornbeam forest in Northern Italy. *Ecol Res* 27: 755-763. DOI: 10.1007/s11284-012-0948-2.
- Bragança MAL, Zanon JC, Picanço M, Laranjeiro AJ. 1998. Effects of environmental heterogeneity on Lepidoptera and Hymenoptera populations in *Eucalyptus* plantations in Brazil. *Forest Ecol Manag* 103 (2-3): 287-292. DOI: 10.1016/S0378-1127(97)00226-0.
- Brooks SJ, Self A, Powney GD, Pearse WD, Penn M, Paterson GLJ. 2017. The influence of life-history traits on the phenological response of British butterflies to climate variability since the late-19th century. *Ecography* 40: 1152-1165. DOI: 10.1111/ecog.02658.
- Charles E, Basset Y. 2005. Vertical stratification of leaf-beetle assemblages (Coleoptera: Chrysomelidae) in two forest types in Panama. *J Trop Ecol* 21: 329-336. DOI: 10.1017/S0266467405002300.
- Chursina MA, Ruchin AB. 2018. A checklist of Syrphidae (Diptera) from Mordovia, Russia. *Halteres* 9: 57-73. DOI: 10.5281/zenodo.1255874.
- Cicort-Lucaciu AŞ. 2020. Road-killed ground beetles prove the presence of *Carabus hungaricus* (Coleoptera: Carabidae) in North-Western Romania. *Nat Conser Res* 5 (3): 134-138. DOI: 10.24189/ncr.2020.035.
- Colom P, Traveset A, Carreras D, Stefanescu C. 2021. Spatio-temporal responses of butterflies to global warming on a Mediterranean island over two decades. *Ecolog Entomol* 46. DOI: 10.1111/een.12958.
- DeVries PJ. 1988. Stratification of fruit-feeding nymphalid butterflies in a Costa Rican rainforest. *J Res Lepidoptera* 26: 98-108.
- DeVries PJ, Murray D, Lande R. 1997. Species diversity in vertical, horizontal, and temporal dimensions of a fruit-feeding butterfly community in an Ecuadorian rainforest. *Biol J Linnean Soc* 62: 343-364. DOI: 10.1111/j.1095-8312.1997.tb01630.x.
- DeVries PJ, Walla TR, Greeney HF. 1999. Species diversity in spatial and temporal dimensions of fruit-feeding butterflies from two Ecuadorian rainforests. *Biol J Linnean Soc* 68: 333-353. DOI: 10.1111/j.1095-8312.1999.tb01175.x.
- Duelli P, Obrist MK, Flückiger PF. 2002. Forest edges are biodiversity. *Acta Zool Acad Sci Hungaricae* 48(2): 75-87.
- Dvořák L, Dvořáková K, Oboňa J, Ruchin AB. 2020. Selected Diptera families caught with beer traps in the Republic of Mordovia (Russia). *Nat Conser Res* 5(4): 65-77. DOI: 10.24189/ncr.2020.057.

- Gilbert LE, Singer MC. 1975. Butterfly ecology. *Ann Rev Ecol Syst* 6: 365-397. DOI: 10.1146/annurev.es.06.110175.002053.
- Giovanni F, Mei M, Cerretti P. 2017. Vertical stratification of selected Hymenoptera in a remnant forest of the Po Plain (Italy, Lombardy) (Hymenoptera: Ampulicidae, Crabronidae, Sphecidae). *Fragmenta Entomol* 49(1): 71-77. DOI: 10.4081/fe.2017.233.
- Gossner MM, Struwe J-F, Sturm S, Max S, McCutcheon M, Weisser WW, et al. 2016. Searching for the optimal sampling solution: Variation in invertebrate communities, sample condition and DNA quality. *PLoS ONE* 11(2): e0148247. DOI: 10.1371/journal.pone.0148247.
- Graham EE, Poland TM, McCullough DG, Millar JG. 2012. A comparison of trap type and height for capturing cerambycid beetles (Coleoptera). *J Econ Entomol* 105: 837-846. DOI: 10.1603/EC12053.
- Grotan V, Lande R, Engen S, Sæther BE, DeVries PJ. 2012. Seasonal cycles of species diversity and similarity in a tropical butterfly community. *J Animal Ecol* 81: 714-723. DOI: 10.1111/j.1365-2656.2011.01950.x.
- Gruppe A, Schubert H. 2001. The spatial distribution and plant specificity of Neuropterida in different forest sites in Southern Germany (Raphidioptera and Neuroptera). *Beiträge zur Entomol* 51: 517-527. DOI: 10.21248/contrib.entomol.51.2.517-527.
- Kirstová M, Pyszko P, Šipoš J, Drozd P, Kočárek P. 2017. Vertical distribution of earwigs (Dermaptera: Forficulidae) in a temperate lowland forest, based on sampling with a mobile aerial lift platform. *Entomol Sci* 20(1): 57-64. DOI: 10.1111/ens.12229.
- Krivosheina NP, Krivosheina MG. 2019. Saproxylic Diptera (Insecta) of the Lazovsky State Nature Reserve (Russia). *Nat Conser Res* 4(3): 78-92. DOI: 10.24189/ncr.2019.052.
- Kuussaari M, Heliölä J, Luoto M, Pöyry J. 2007. Determinants of local species richness of diurnal Lepidoptera in boreal agricultural landscapes. *Agric Ecosyst Environ* 122(3): 366-376. DOI: 10.1016/j.agee.2007.02.008.
- Maguire DY, Robert K, Brochu K, Larrivé M, Buddle CM, Wheeler TA. 2014. Vertical stratification of beetles (Coleoptera) and flies (Diptera) in temperate forest canopies. *Environ Entomol* 43: 9-17. DOI: 10.1603/EN13056.
- Makarkin VN, Ruchin AB. 2019. New data on Neuroptera and Raphidioptera of Mordovia (Russia). *Kavkaz Entomol Bull* 15(1): 147-157. DOI: 10.23885/181433262019151-147157.
- Melo DHA, Duarte M, Mielke OHH, Robbins RK, Freitas AVL. 2019. Butterflies (Lepidoptera: Papilionoidea) of an urban park in Northeastern Brazil. *Biota Neotr* 19 (1): e20180614. DOI: 10.1590/1676-0611-bn-2018-0614.
- Owen DF. 1971. *Tropical Butterflies*. Oxford University Press, London.
- Owen DF, Owen J, Chanter DO. 1972. Seasonal changes in relative abundance and estimates of species diversity in a family of tropical butterflies. *Oikos* 23: 200-205. DOI: 10.2307/3543406.
- Preisser E, Smith DC, Lowman MD. 1998. Canopy and ground-level insect distribution in a temperate forest. *Selbyana* 19(2): 141-146.
- Proshchalykin MYu, Sergeev ME. 2020. New distribution data of *Apis cerana ussuriensis* (Hymenoptera, Apidae) from Primorsky Krai, Russia. *Nat Conser Res* 5(4): 111-112. DOI: 10.24189/ncr.2020.049.
- Procházka J, Cizek L, Schlaghamerský J. 2018. Vertical stratification of scolytine beetles in temperate forests. *Insect Conserv Divers* 11: 534-544. DOI: 10.1111/icad.12301.
- Puker A, Correa CMA, Silva AS, Silva JVO, Korasaki V, Grossi PC. 2020. Effects of fruit-baited trap height on flower and leaf chafer scarab beetles sampling in Amazon rainforest. *Entomol Sci* 23(3): 245-255. DOI: 10.1111/ens.12418.
- Ribeiro DB, Prado PI, Brown KS, Freitas AVL. 2010. Temporal diversity patterns and phenology in fruit-feeding butterflies in the Atlantic forest. *Biotropica* 42: 710-716. DOI: 10.1111/j.1744-7429.2010.00648.x.
- Root HT, Verschuyt J, Stokely T, Hammond P, Scherr MA, Betts MG. 2017. Plant diversity enhances moth diversity in an intensive forest management experiment. *Ecol Appl* 27: 134-142. DOI: 10.1002/eap.1426.
- Roy DB, Sparks TH. 2000. Phenology of British butterflies and climate change. *Glob Chang Biol* 6: 407-416. DOI: 10.1046/j.1365-2486.2000.00322.x.
- Rubik DW. 1993. Tropical pollinators in the canopy and understory: Field data and theory for stratum preferences. *J Insect Behav* 6: 659-673. DOI: 10.1007/BF01201668.
- Ruchin AB, Egorov LV, Khapugin AA. 2021. Seasonal activity of Coleoptera attracted by fermental crown traps in forest ecosystems of Central Russia. *Ecolog Quest* 32(1): 37-53. DOI: 10.12775/EQ.2021.004.
- Ruchin AB, Egorov LV, Khapugin AA, Vikhrev NE, Esin MN. 2020. The use of simple crown traps for the Insect collection. *Nat Conser Res* 5(1): 87-108. DOI: 10.24189/ncr.2020.008.
- Ruchin AB, Egorov LV, Semishin GB. 2018. Fauna of click beetles (Coleoptera: Elateridae) in the interfluvium of Rivers Moksha and Sura, Republic of Mordovia, Russia. *Biodiversitas* 19 (4): 1352-1365. DOI: 10.13057/biodiv/d190423.
- Santos J, Iserhard C, Carreira J, Freitas A. 2017. Monitoring fruit-feeding butterfly assemblages in two vertical strata in seasonal Atlantic forest: temporal species turnover is lower in the canopy. *J Trop Ecol* 33(5): 345-355. DOI: 10.1017/S0266467417000323.
- Saure C, Kielhorn KH. 1993. Lacewings in the tree canopy of oak and Scots pine (Neuroptera: Coniopterygidae, Hemerobiidae, Chrysopidae). *Faun-Ökol Mitt* 6: 391-402.
- Schulze CH, Linsenmair KE, Fiedler K. 2001. Understorey versus canopy: Patterns of vertical stratification and diversity among Lepidoptera in a Bornean rain forest. In: Linsenmair KE, Davis AJ, Fiala B, Speight MR (eds). *Tropical Forest Canopies: Ecology and Management*. Forestry Sciences. Vol. 69. Springer, Dordrecht. DOI: 10.1007/978-94-017-3606-0_11.
- Sergeev ME. 2020. Species composition and biotopic distribution of leaf beetles (Coleoptera: Megalopodidae, Chrysomelidae) in the Sikhote-Alin State Nature Reserve (Russia). *Nat Conser Res* 5(2): 80-88. DOI: 10.24189/ncr.2020.020.
- Sergeev ME, Makarkin VN. 2021. The first record of *Euroleon polyspilus* from the Sikhote-Alin State Nature Reserve, Russia, with remarks on its biology. *Nat Conser Res* 6(1): 100-101. DOI: 10.24189/ncr.2021.011.
- Shapiro AM. 1975. The temporal component of butterflies species diversity. In: Cody ML, Diamond JM (eds). *Ecology and Evolution of Communities*. The Belknap Press of Harvard University, London. 545 pp.
- Sutton SL, Hudson PJ. 1980. The vertical distribution of small flying insects in the lowland rain forest of Zaire. *Zool J Linnean Soc* 68: 111-124. DOI: 10.1111/j.1096-3642.1980.tb01921.x.
- Ulyshen MD. 2011. Arthropod vertical stratification in temperate deciduous forests: Implications for conservation-oriented management. *Forest Ecol Manag* 261: 1479-1489. DOI: 10.1016/j.foreco.2011.01.033.
- Ulyshen MD, Soon V, Hanula JL. 2011. Vertical distribution and seasonality of predatory wasps (Hymenoptera: Vespidae) in a temperate deciduous forest. *Florida Entomol* 94(4): 1068-1070. DOI: 10.1653/024.094.0450.
- Walla TR, Engen S, DeVries PJ, Lande R. 2004. Modeling vertical beta-diversity in tropical butterfly communities. *Oikos* 107: 610-618. DOI: 10.1111/j.0030-1299.2004.13371.x.
- Zografou K, Kati V, Grill A, Wilson RJ, Tzirkalli E, Pamperis LN et al. 2014. Signals of climate change in butterfly communities in a Mediterranean Protected Area. *PLoS ONE* 9(1): e87245. DOI: 10.1371/journal.pone.0087245.