

1 **Type of paper: Primary research article**

2

3 **Title:**

4 Seasonal variability of foliar photosynthetic and morphological
5 traits and drought impacts in a Mediterranean mixed forest

6 Keywords (10): Seasonality, leaf position, $V_{c,max}$, J_{max} , leaf traits, abiotic stress, *Pinus halepensis*,
7 *Quercus ilex*, *Quercus pubescens*, *Arbutus unedo*

8

9 Running head: SEASONAL PHOTOSYNTHESIS MORPHOLOGY MIXED FOREST

10

11 D. Sperlich^{1,2*}, C.T. Chang^{1,2}, J. Peñuelas^{2,3}, C. Gracia^{1,2}, S. Sabaté^{1,2}

12

13 ¹ Departament d'Ecologia, Facultat de Biologia, Universitat de Barcelona, Diagonal 645, 08028 Barcelona, Spain,

14 ² CREAM, Cerdanyola del Vallès, 08193 Barcelona, Catalonia, Spain

15 ³ CSIC, Global Ecology Unit CREAM-CSIC-UAB, Cerdanyola del Vallès, 08193 Barcelona, Catalonia, Spain

16 * Author for correspondence

17 *Dominik Sperlich*

18 *Tel.: +34 9358 14850*

19 *E-mail: Dominik@creaf.uab.es*

20

21 **Word counts:**

22 Abstract	195
23 Introduction	1177
24 Material & Methods	1744
25 Results	1682
26 Discussion	2521
27 Acknowledgments	75
28 Total (excl. abstract)	7199

29 **Number of:**

Figures Tables Supp.information

8 (colour) 5 12 (Fig. S1-S3, Tables S1-S6, Notes S1-S2)

30

31 **Abstract**

32 The Mediterranean region is a hot spot of climate change vulnerable to increased
33 droughts and heat waves. Scaling carbon fluxes from leaf to landscape levels is
34 particularly challenging under drought conditions. We aimed to improve the
35 mechanistic understanding of the seasonal acclimation of photosynthesis and
36 morphology in sunlit and shaded leaves of four Mediterranean trees (*Quercus ilex* L.,

37 *Pinus halepensis* Mill., *Arbutus unedo* L., and *Q. pubescens* Willd.) under natural
38 conditions. $V_{c,max}$ and J_{max} were not constant, and mesophyll conductance was not
39 infinite, as assumed in most terrestrial biosphere models, but varied significantly
40 between seasons, tree species, and leaf position. Favourable conditions in winter led to
41 photosynthetic recovery and growth in the evergreens. Under moderate drought,
42 adjustments in the photo/biochemistry and stomatal/mesophyllic diffusion behaviour
43 effectively protected the photosynthetic machineries. Severe drought, however, induced
44 early leaf senescence mostly in *A. unedo*, *Q. pubescens*, and significantly increased leaf
45 mass per area in *Q. ilex* and *P. halepensis*. Shaded leaves had lower photosynthetic
46 potentials but cushioned negative effects during stress periods. Species-specificity,
47 seasonal variations, and leaf position are key factors to explain vegetation responses to
48 abiotic stress and hold great potential to reduce uncertainties in terrestrial biosphere
49 models especially under drought conditions.

50

51 ***Introduction***

52 The Mediterranean region is dominated by arid or semi-arid ecosystems where high evaporative
53 demand and low soil-water content during the summer dry period are the main ecological
54 limitations to plant growth (Specht 1969, Di Castri 1973). The resilience of plants to drought
55 and heat waves is determined by their frequency and duration, which are projected to become
56 much more severe under current climate change scenarios - particularly in the Mediterranean
57 region (Somot et al. 2008, Friend 2010, IPCC 2013). Increased drought-induced defoliation
58 (Poyatos et al. 2013) associated with the depletion of carbon reserves (Galiano et al. 2012) can
59 ultimately lead to catastrophic hydraulic failure and tree mortality (Urli et al. 2013, Choat
60 2013). Drought-induced forest impacts and diebacks in the Mediterranean region have been
61 reported in numerous studies (Peñuelas et al. 2001, Martínez-Vilalta and Piñol 2002,
62 Raftoyannis et al. 2008, Allen et al. 2010, Carnicer et al. 2011, Matusick et al. 2013) and can
63 lead to shifts in vegetation composition (Jump and Penuelas 2005, Anderegg et al. 2013) and to
64 a higher risk of forest fires (Piñol et al. 1998, Pausas et al. 2008). The challenge in the
65 Mediterranean region in the coming years will be to learn how carbon uptake and growth in
66 species and communities will respond to these changes, and how forest management strategies
67 can be adapted to cushion the negative impacts of climate change on forests (Sabaté 2002,
68 Bugmann et al. 2010).

69 In past decades, ecosystem models on regional or global level contributed substantially
70 to our understanding of the implications of climate change on a coarse scale where field
71 experiments are limited (Luo 2007). Much uncertainty, however, remains in the modelled
72 feedback of the global carbon cycle to climatic warming (Friedlingstein et al. 2014) and in the

73 understanding and modelling of species responses to climate change (Luo 2007, McDowell et
74 al. 2008, Beaumont et al. 2008). Photosynthesis is generally overestimated in the main Earth
75 system models, with significant regional variations (Anav et al. 2013). Two critical parameters,
76 the maximum rate of carboxylation ($V_{c,max}$) and the maximum rate of electron transport (J_{max}),
77 are a prerequisite for scaling foliar photosynthesis to the canopy level at which global dynamic
78 models operate (Friedlingstein et al. 2006, Friedlingstein and Prentice 2010). These two
79 parameters describe the biochemical limitations to carbon assimilation, but are not easily
80 measured. So relatively little data of their variability between species or seasons are available.
81 $V_{c,max}$ and J_{max} are thus often used as constants for various plant functional types and seasons or,
82 in some cases, are derived from other parameters such as leaf nitrogen content (Grassi and
83 Magnani 2005, Walker et al. 2014). Moreover, extreme climatic conditions and inter-annual
84 variability in arid and semi-arid regions are challenging for scaling carbon assimilation patterns
85 from one year to another (Reynolds et al. 1996, Morales et al. 2005, Gulías et al. 2009).
86 Simulations of ecosystem carbon fluxes are consequently limited, first, by underrepresented
87 temporal variability of photosynthetic parameters and soil-water patterns, and second by our
88 limited understanding of the effects of water stress on both carbon uptake and release (Hickler
89 et al. 2009, Niinemets and Keenan 2014). The modelling performance in Mediterranean-type
90 ecosystems is thus particularly poor and stresses the need for a better mechanistic description of
91 photosynthetic processes under water stress (Morales et al. 2005, Keenan et al. 2011, Zheng et
92 al. 2012, Vargas et al. 2013). Mesophyll conductance, g_m , might play a future key role in
93 improving model performance of photosynthesis under drought conditions (Keenan et al. 2010).

94 The photosynthetic limitations of Mediterranean vegetation, especially under drought,
95 have been extensively studied (for a review see Flexas *et al.*, 2014), but fewer studies have
96 thoroughly assessed the seasonal behaviour of photosynthesis and morphology under natural
97 conditions in a mixed mature forest. The information gained from seedlings under controlled
98 conditions can only poorly represent the physiological mechanisms of the long-term acclimation
99 to variable environmental conditions in mature trees (Flexas et al., 2006; Mittler, 2006;
100 Niinemets, 2010). Seedlings or saplings are characterised by higher metabolism and enzymatic
101 function, lower leaf dry mass per unit area (LMA), and higher photosynthetic potential relative
102 to mature trees (Johnson & Ball, 1996; Bond, 2000; Niinemets, 2014). Responses to short-term
103 stress are related to the mechanisms of prompt reactions (Flexas et al., 2006). Under natural
104 conditions, however, mature trees acclimate to gradually developing water stress through the
105 photosynthetic pathway (biochemical, stomatal or mesophyllic) (e.g. Martin-StPaul et al. 2013),
106 but also through foliar traits such as nitrogen, LMA etc. (Poorter et al. 2009). Less work has
107 evaluated simultaneously the variations of photosynthetic and morphological traits in response
108 to abiotic stress conditions. The variation of these traits is largely species specific (Orshan 1983,
109 Chaves et al. 2002, Gratani and Varone 2004, Krasteva et al. 2013), although within-canopy

110 gradients can play an additional overriding role (Valladares and Niinemets 2008, Sperlich et al.
111 2014). Mixed forests provide ideal test conditions where we can observe distinct species-
112 specific strategies coping equally with the yearly variability of environmental conditions.

113 The aim of this study was to investigate the impact of seasonal environmental changes
114 (above all drought) on foliar photosynthetic and morphological traits of the winter-deciduous
115 sub-Mediterranean *Quercus pubescens*, two evergreen sclerophyllous species (*Quercus ilex* and
116 *Arbutus unedo*) and an early-successional drought-adapted conifer, *Pinus halepensis*. *P.*
117 *halepensis* is characterised as isohydric following a water saving and photoinhibition-tolerant
118 strategy (Martínez-Ferri et al. 2004, Baquedano and Castillo 2006, Sperlich et al. 2014). *Q. ilex*
119 L. is a late-successional, slow growing, water-spending, photoinhibition-avoiding, anisohydric
120 tree species with a plastic hydraulic and morphological behaviour (Villar-Salvador et al. 1997,
121 Fotelli et al. 2000, Corcuera et al. 2004, Ogaya and Peñuelas 2006, Limousin et al. 2009). The
122 winter-deciduous anisohydric *Q. pubescens* follows a similar drought-avoiding strategy as *Q.*
123 *ilex*, but maximizes gas exchange during a shorter growing season (Baldochi et al. 2009),
124 resulting in high transpiration rates throughout the summer (Poyatos et al. 2008). Over extensive
125 areas of the Mediterranean region *Q. ilex* and *Q. pubescens* form the terminal point of secondary
126 succession (Lookingbill and Zavala 2000). *A. unedo* - relict of the humid-subtropical Tertiary
127 tree flora (Gratani and Ghia, 2002a and references therein) – is typically occurring as shrub or
128 small tree in the macchia ecosystems and holding an intermediate position concerning stomatal-
129 sensitivity (Beyschlag et al. 1986, Vitale and Manes 2005, Barbeta et al. 2012) and photoinhibition-
130 sensitivity (Sperlich et al. 2014). Prolonged climate stress might disadvantage *A. unedo* being
131 more drought sensitive than the companion species (Ogaya and Peñuelas 2004, Barbeta et al.
132 2012).

133 Our particular interests were to distinguish the species-specific strategies and to explore
134 the eco-physiological mechanism behind drought responses by examining the fine tuning of
135 foliar photosynthetic potentials/rates and foliar morphological traits. We hypothesized that i)
136 seasonal environmental changes (above all drought) affect the photosynthetic and ii)
137 morphological traits, iii) mesophyll conductance (g_m) strongly constrains
138 photosynthesis under drought conditions, iv) the seasonal acclimation varies qualitatively and
139 quantitatively with species and v) light environment (leaf canopy position). We thus created a
140 matrix of photosynthetic parameters that could be incorporated into process-based ecosystem
141 models to improve estimates of carbon flux in the Mediterranean region.

142

143 **Materials and methods**

144 ***Field site***

145 The experimental site Can Balasc is located in the coastal massif of the Collserola Natural Park
146 (8500 ha), in the province of Barcelona, northeastern Spain (41° 25' N, 2° 04' E, 270 m a.s.l.).
147 Seasonal summer droughts, warm temperatures and mild winters characterise the typical
148 Mediterranean climate with a mean August temperature of 22.8 °C and a mean January
149 temperature of 7.9 °C. Mean annual precipitation and temperature are 723 mm and 15.1 °C
150 (1951-2010), respectively (Ninyerola et al. 2007a, 2007b). Sensors for measuring air
151 temperature (HMP45C, Vaisala Oyj, Finland) and solar radiation (SP1110 Skye Instruments
152 Ltd., Powys, UK) were installed at a height of 3 m, in a clearing *ca.* 1 km from the plot.

153 ***Stand structure***

154 Our study site is characterised by a dense forest stand (1429 stems ha⁻¹) with a two-layered
155 canopy consisting of a dense layer of *Quercus* species surmounted by shelter trees of the early-
156 successional and fast growing Aleppo Pine (*P. halepensis* Mill.). The mean heights of each
157 layer are 9.9 m and 17.1 m, respectively. The *Quercus* species are the late-successional
158 evergreen Holm Oak (*Q. ilex* L.) and the deciduous Pubescent Oak (*Q. pubescens* Willd.). The
159 Strawberry tree (*A. unedo* L.) grows usually as a shrub being widely abundant in the macchia
160 ecosystems of the Iberian peninsula (Beyschlag et al. 1986, Reichstein et al. 2002). In our study
161 site, however, *A. unedo* occurs scattered in the tree canopy (mean height 8.1 m) enriching the
162 forest diversity with its flowering and fruiting habit. The trees with the biggest dimensions are
163 the pines followed by the two *Quercus* species and at last by *A. unedo* (mean DBH of 33.7,
164 12.9, 9.6 cm, respectively). The forest succession has reached the final stage: The dense
165 *Quercus* canopy is out-competing the early-successional *P. halepensis* by suppressing the
166 growth of the light demanding pine seedlings and saplings. More details of stand history and
167 field site are described in Sperlich *et al.* (2014).

168 ***Sampling method***

169 We conducted eight field campaigns from June 2011 to February 2013. The sampling periods
170 are presented in Table 2 and Figure 1. We avoided difficulties encountered during field
171 measurements such as deviations from the standard temperature (25 °C) or unpredictable plant
172 responses (patchy stomatal conductance) (Mott and Buckley 1998, 2000) by analysing sampled
173 twigs in the laboratory. We cut twigs with a pruning pull from sunlit and shaded leaf positions,
174 optimally at similar heights. The twigs were immediately re-cut under water in the field,
175 wrapped in plastic bags to minimise transpiration, stored in water buckets, and transported to
176 the laboratory. Five replicates of each leaf position and tree species were collected for the
177 analysis of gas exchange. The twigs were pre-conditioned in the laboratory at room temperature

178 (24-28 °C) in dim light for 1-3 d and were freshly cut every morning. More details and
179 references can be found in Sperlich *et al.* (2014).

180 ***Analyses of gas exchange and chlorophyll fluorescence***

181 Gas exchange and chlorophyll fluorescence were measured with a Li-Cor LI-6400XT Portable
182 Photosynthesis System equipped with a LI-6400-40 Leaf Chamber Fluorometer (Li-Cor, Inc.,
183 Lincoln, USA). Response curves for foliar net assimilation versus CO₂ concentration were
184 recorded in parallel with the chlorophyll fluorescence measurements. In some cases the sunlit
185 leaves of *Q. ilex* were too small to fill the leaf cuvette (2 cm²) and so the measured parameters
186 were adjusted after the measurements. For *P. halepensis*, we positioned a layer of needles (ca.
187 10-15) on the leaf cuvette, avoiding gaps and overlays, and sealed the gaskets with Blu-tack
188 (Bostik SA, La Plaine St Denis, France) to keep the needles in position. The preparation and
189 acclimation of the leaves prior to recording the response curves were conducted as in Sperlich *et*
190 *al.* (2014).

191 ***CO₂ experiments***

192 The CO₂-response curves were recorded at a leaf temperature (T_{Leaf}) of 25 °C and a quantum
193 flux density of 1000 μmol photons m⁻² s⁻¹. The CO₂ concentrations in the leaf chamber (C_a) used
194 to generate the response curves were 400→300→200→150→100→50→400→400→600→
195 800→1200→2000 μmol CO₂ mol air⁻¹. The minimum and maximum times for stabilising net
196 assimilation rate (A_{net} in μmol CO₂ m⁻² s⁻¹), stomatal conductance (g_s in mol H₂O m⁻² s⁻¹), and
197 stomatal internal CO₂ concentrations (C_i in μmol CO₂ mol air⁻¹) for each log were set to 4 and 6
198 min, respectively.

199 ***Calculation of chlorophyll fluorescence parameters***

200 F_m' and F_s were used to estimate the effective quantum yield of photosystem II (Φ_{PSII}, unitless)
201 as:

$$202 \quad \Phi_{\text{PSII}} = \frac{(F_{\text{m}}' - F_{\text{s}})}{F_{\text{m}}'} \quad (1)$$

203 where F_s is the steady-state fluorescence of a fully light-adapted sample, and F_m' is the maximal
204 fluorescence yield reached after a pulse of intense light. The effective quantum yield of PSII
205 represents the fraction of photochemically absorbed photons for a light-adapted leaf. The
206 electron-transport rate based on the effective quantum yield of PSII (J_{CF} in μmol electron m⁻² s⁻¹)
207 was calculated as

$$208 \quad J_{\text{CF}} = \varepsilon * \Phi_{\text{PSII}} * \alpha_{\text{L}} \quad (2)$$

209 ε is a scaling factor accounting for the partitioning of intercepted light between photosystem I
210 (PSI) and PSII. We assumed that light was equally distributed between both photosystems (ε =
211 0.5) (Bernacchi *et al.* 2002, Niinemets *et al.* 2005). α_L (unitless) is the foliar absorbance; we

212 used the following values: 0.932 for *Q. ilex* and 0.912 for *P. halepensis* for both sunlit and
 213 shaded leaves, 0.935 for sunlit leaves of *A. unedo*, 0.917 for shaded leaves of *A. unedo*, 0.939
 214 for sunlit leaves of *Q. pubescens*, and 0.900 for shaded leaves of *Q. pubescens*. For the
 215 determination of α_L , foliar reflectance and transmittance were measured at midday in August
 216 2012 using a UniSpec Spectral Analysis System spectroradiometer (PP Systems, Haverhill,
 217 USA). The ambient photosynthetic electron transport (J_{amb}) was defined as the value of J_{CF} at a
 218 CO_2 concentration of $400 \mu\text{mol } CO_2 \text{ mol air}^{-1}$ and a PPFD of $1000 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$. The
 219 relationship between J_{amb} and the net assimilation rate (J_{amb}/A_{net}) was used for the analyses of
 220 alternative electron sinks other than carbon metabolism. Calculations of F_v/F_m and NPQ can be
 221 found in the supplementary material (Note S1).

222 ***Estimation of mesophyll conductance***

223 We estimated g_m (in $\text{mol m}^{-2} \text{ s}^{-1} \text{ bar}^{-1}$) using the variable- J method by Harley et al. (1992):

$$224 \quad g_m = \frac{A_{net}}{C_i - \frac{[\Gamma^* J_{CF} + 8(A_{net} + R_d)]}{J_{CF} - 4(A_{net} + R_d)}} \quad (6)$$

225 where Γ^* is the CO_2 concentration at which the photorespiratory efflux of CO_2 equals the rate of
 226 photosynthetic CO_2 uptake, and R_d is the mitochondrial respiration of a leaf in light conditions
 227 and was estimated from the light-response curves combining gas exchange and measurements
 228 with the CF- method proposed by Yin et al. (2009). See supplementary material for details
 229 (Note S2). The chloroplastic CO_2 concentration (C_c in $\mu\text{mol } CO_2 \text{ mol air}^{-1}$) was determined as:

$$230 \quad C_c = C_i - \frac{A_{net}}{g_m} \quad (7)$$

231 ***Photosynthesis model***

232 The photosynthesis model of Farquhar et al. (1980) considers photosynthesis as minimum of the
 233 potential rates of Rubisco activity (A_c) and ribulose-1,5-bisphosphate (RuBP) regeneration (A_j).
 234 The model was further complemented with a third limitation (A_p) that considers the limitation
 235 by triose-phosphate use (TPU) at high CO_2 concentrations when the CO_2 response shows a
 236 plateau or decrease (Sharkey 1985). However, we rarely detected A_p limitations and TPU was
 237 therefore discarded in our analyses. A_{net} was then determined by the minimum of these two
 238 potential rates from an A/C_c curve:

$$239 \quad A_{net} = \min\{A_c, A_j\} \quad (8)$$

240 where

$$241 \quad A_c = V_{cmax} * \left[\frac{C_c - \Gamma^*}{C_c + K_c \left(1 + \frac{o}{K_o}\right)} \right] - R_d \quad (9)$$

242 where $V_{c,max}$ (in $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$) is the maximum rate of Rubisco carboxylation, K_c is the
243 Michaelis-Menten constant of Rubisco for CO_2 , O is the partial pressure of O_2 at Rubisco, and
244 K_o is the Michaelis-Menten constant of Rubisco for O_2 , taken from Bernacchi et al. (2002). The
245 equation representing photosynthesis limited by RuBP regeneration is:

$$246 \quad A_j = J * \left[\frac{C_c - \Gamma^*}{4C_c + 8\Gamma^*} \right] - R_d \quad (10)$$

247 where J (in $\mu\text{mol electron m}^{-2} \text{ s}^{-1}$) is the rate of electron transport. We assumed that J becomes
248 J_{max} under light and CO_2 saturation when the maximum possible rate of electron transport is
249 theoretically achieved, although we may have underestimated the true J_{max} (for further details
250 see Buckley & Diaz-Espejo, 2014). $V_{c,max}$ and J_{max} define the biochemical potential to drive
251 photosynthesis and are summarised in the term “photosynthetic potential” (Niinemets et al.
252 2006). Curves were fit, and diffusion leakage was corrected, as in Sperlich *et al.* (2014).

253 ***Foliar morphology, chemical analyses, and assessment of crown*** 254 ***condition***

255 Foliar morphological traits were measured on fully expanded leaves ($n = 60$ per leaf position
256 and species) from the excised twigs in five sampling campaigns in spring and autumn 2011^a
257 (2011^a indicates sampling during a drought), and winter, spring, and summer 2012. Immediately
258 after the gas exchange analyses, we measured fresh weight (FW, mg) and projected leaf surface
259 area (LA, cm^2) (including petioles) with Photoshop from scanned leaves at 300 dpi. We oven-
260 dried the leaves at 70°C for 48 h and weighed the leaves for dry weight (DW, mg) and
261 measured leaf thickness (LT, mm) with a portable dial thickness gauge (Baxlo Precisión,
262 Barcelona, Spain). We then calculated the percentage of the leaf water content (WC) as $[1 -$
263 $(\text{DW}/\text{FW})] * 100$. Leaf mass per area (LMA) (mg cm^{-2}) was calculated as the ratio of DW to LA
264 and leaf tissue density (D, mg cm^{-3}) as the ratio of LMA to LT. Foliar Succulence (S) was
265 calculated as $(\text{FW} - \text{DW})/\text{LA}$. We ground the leaves to a fine powder using a MM400 mixer mill
266 (Retsch, Hahn, Germany), encapsulated a sample of 0.7 mg in tin foil and determined carbon
267 and nitrogen contents by EA/IRMS (Elemental Analyzer/Isotope Ratio Mass Spectrometry) and
268 GC/C/IRMS (Gas Chromatography/Combustion/IRMS). The crown condition was assessed
269 using ‘International Co-operative Programme on Assessment and Monitoring of Air
270 Pollution Effects on Forests’ (ICP Forests) standards (Eichhorn et al. 2010).

271 ***Statistical analyses***

272 We performed the statistical analyses with the R version 3.0.2 (<http://www.r-project.org/>). The
273 matrix of photosynthetic and morphological traits was subjected to principal component
274 analyses (PCAs) to summarise the principal factors explaining the variation in these parameters.
275 Differences in the parameters between sunlit and shaded leaves were determined with Student’s

276 *t*-tests ($P \leq 0.05$). The normality of the data was tested with Shapiro-Wilk tests. If the data were
277 not normally distributed, they were normalised. One-factorial analyses of variance (ANOVAs)
278 with season as the main factor were used to test for differences in the parameters in each species
279 and leaf position. Significant differences were determined at $P \leq 0.05$ with Fisher's least
280 significance difference (LSD) tests. Bonferroni correction was used for familywise error rate.
281 Linear regression analyses were conducted to study the relationships among various leaf traits
282 such as A_{net}/g_s , A_{net}/g_m , $J_{\text{max}}/V_{c,\text{max}}$, g_m/g_s , $J_{\text{amb}}/A_{\text{net}}$. With analyses of co-variance (ANCOVAs),
283 we tested for differences in regression slopes and intercepts. We applied a non-linear regression
284 analysis using the nls function in R to study the relationship of g_m/LMA .
285

286 **Results**

287 ***Environmental and crown conditions***

288 The year 2011 was characterised by 30% more precipitation than the climatic average of 723
289 mm (1951-2010) (Ninyerola et al. 2007a, 2007b) (Tab. 3, Fig. 1), and no drought-induced leaf
290 shedding was observed. The winter from 1 December 2011 to 31 January 2012 was relatively
291 mild with average maximum and minimum temperatures of 11.8 and 4.2 °C, respectively,
292 coinciding with high photosynthetic potentials and shoot growth. The precipitation in 2012 was
293 20% lower than the climatic average (Table 3). *A. unedo* and *Q. pubescens* were strongly
294 defoliated during summer 2012; *Q. ilex* and *P. halepensis* to a lesser extent (Table 4). *Q. ilex*
295 showed some discoloration in the more exposed sites. Only one individual of *P. halepensis*
296 showed discoloration. The defoliated *Q. pubescens* trees recovered completely in 2013. In
297 contrast, heavily affected individuals of *A. unedo* showed an irreversible dieback of the main
298 leading branches but also vigorous re-sprouting in 2013.

299 ***Effect of season, tree species and leaf position on photosynthetic*** 300 ***parameters***

301 In Fig. 2a, we present the PCA for the morphological and photosynthetic parameters. No
302 rotation was applied to the space of the PC's. $V_{c,max}$, J_{max} , and g_s were negatively correlated with
303 N_{mass} , C_{mass} , NPQ, and g_m . F_v/F_m , g_s , and water content (WC) were negatively correlated with
304 nitrogen and carbon per unit leaf area (N_{area} , C_{area}), LMA, and density (D). Nitrogen per unit leaf
305 mass (N_{mass}) and g_m correlated well with LT (Fig. 2). A_{net} was correlated negatively with
306 succulence (S) and positively with g_m . PC1 and PC2 explained 37.2 and 20.4% of the variation,
307 respectively. The datapoints within the cluster circles in Fig. 1b-d exhibited similar behaviours
308 in photosynthetic and morphological traits. Leaf positions, seasons, and species could be
309 separated. Sunlit leaves were characterised by higher values on the orthogonal axis. The
310 horizontal axes separated *A. unedo* and *Q. pubescens* from *Q. ilex* and *P. halepensis*. The
311 orthogonal axes separated *Q. ilex* from *P. halepensis* with generally positive values. The
312 seasonality was further investigated for each species and leaf position with ANOVAs for each
313 photosynthetic and morphological parameter.

314 *Q. ilex*

315 *Q. ilex* had the most plastic response to the environmental conditions. The sunlit leaves of *Q.*
316 *ilex* exhibited strong declines in several photosynthetic parameters from summer 2011 to
317 autumn 2011^a. $V_{c,max}$, A_{net} and g_s were significantly ($P < 0.05$), and J_{max} and g_m were marginally
318 significantly lower ($P < 0.10$) (Fig. 3 a1-b1). The means of the majority of the photosynthetic
319 parameters recovered after the first rains in autumn 2011^b (2011^b indicates sampling after the
320 drought), reaching pre-drought values, but accompanied by a high standard error. This recovery

321 was thus only significant for J_{\max} and g_m . Surprisingly, $V_{c,\max}$ and J_{\max} peaked in winter and not,
322 as expected, in spring. From that peak we observed significant declines from winter to spring to
323 summer 2012. In contrast to the pattern of $V_{c,\max}$ and J_{\max} , F_v/F_m , A_{net} , and g_s peaked in spring
324 2012 (Figs. 3c1, 4a1-b1). These parameters then also declined significantly in summer 2012.
325 Interestingly, g_m peaked in summer 2012 in parallel with a reduction in g_s (Fig. 4c1). The
326 photosynthetic parameters of shaded leaves in *Q. ilex* showed a similar trend, declining after the
327 drought in 2011 and recovering after the autumn rains (Figs. 3, 4). The parameter means of
328 shaded leaves remained relatively stable throughout the season, in contrast to the pattern in
329 sunlit leaves, except for a peak of $V_{c,\max}$ and J_{\max} in spring 2012. The photosynthetic parameters
330 in *Q. ilex* were significantly lower in shaded leaves. During periods of stress, however, the
331 photosynthetic parameters of sunlit leaves declined and had values similar to those of shaded
332 leaves (Table 6, Figs. 3, 4).

333 *P. halepensis*

334 Mean $V_{c,\max}$, J_{\max} , and F_v/F_m were generally higher in sunlit leaves of *P. halepensis* than the
335 other species (Figs. 3, 4). The seasonal variation of the photosynthetic potential was not as
336 strongly pronounced as in *Q. ilex*, and mean $V_{c,\max}$ and J_{\max} remained relatively high and stable
337 in 2011 (Fig. 3a1-b1). The 2012 drought had comparatively stronger effects on $V_{c,\max}$ and J_{\max}
338 than the 2011 drought. Mean A_{net} , g_s , and g_m , however, were significantly lower in autumn 2011^a
339 (Fig. 4a1-c1). These values recovered quickly and significantly after the first autumn rains. The
340 relatively high $V_{c,\max}$, J_{\max} , and F_v/F_m during this period reflected a stronger limitation of g_s and
341 g_m than of the biochemistry imposed on A_{net} . A_{net} recovered in winter 2012 due to the mild
342 conditions (Fig. 4a1). The 2012 summer drought significantly reduced the high values of A_{net}
343 observed in winter 2012, but not as much as after the 2011 drought (Fig. 4a1). Both g_s and g_m
344 remained relatively stable during this period, so the reductions in A_{net} were due to biochemical
345 limitations ($V_{c,\max}$ and J_{\max}) (Figs. 3, 4). Sunlit and shaded leaves differed the least in *P.*
346 *halepensis*; only $V_{c,\max}$ and J_{\max} were significantly different (Table 6). The sunlit and shaded
347 leaves of *P. halepensis* had similar patterns of seasonal variation, but changes between seasonal
348 campaigns were not significant (Fig. 3a1-b1).

349 *A. unedo*

350 Similar to *Q. ilex*, the photosynthetic parameters in *A. unedo* varied strongly seasonally but had
351 high standard errors (Figs. 3, 4). A_{net} decreased significantly in winter 2012, in contrast to J_{\max}
352 and $V_{c,\max}$ that peaked in the same campaign (Figs. 3a1-b1 and 4a1). A decline in g_s and g_m in
353 this campaign suggested that they more strongly regulated A_{net} (Fig. 4b1-c1). A_{net} , g_s , and g_m
354 peaked in spring 2012. These increases were significant for A_{net} and g_s and marginally
355 significant for g_m relative to the other field campaigns (Fig. 4a1-4c1). The photosynthetic
356 parameters were generally lower in the shaded leaves of *A. unedo*, but with no clear pattern and
357 high variability (Table 5).

358 *Q. pubescens*

359 The photosynthetic potentials were much higher in *Q. pubescens* than in the other species but
360 also had high standard errors (Fig. 3a1-b1). The 2012 summer drought led to a decline of the
361 photosynthetic potentials by approximately one third. These decreases were only significant for
362 the average of spring 2011 and spring 2012 relative to the average of summer 2011 and summer
363 2012. A_{net} showed a similar trend, with a peak in spring 2012 being reduced significantly by the
364 2012 summer drought (Fig. 4a1). Stomatal control was more strongly pronounced than
365 mesophyllic control (Fig. 4b1-c1). Shaded leaves had higher A_{net} , g_m , and g_s means throughout
366 the campaigns, in contrast to lower means of $V_{c,\text{max}}$ and J_{max} (Figs. 3a1-c2, 4a1-b2). Shaded
367 leaves generally showed lower values than sunlit leaves and were less affected by the droughts
368 (Figs. 3a1-c2, 4a1-c2).

369 ***Morphological parameters***

370 The foliar traits of *P. halepensis* and *Q. ilex* acclimated most strongly to drought. LMA was
371 significantly higher in *P. halepensis* and *Q. ilex* in both shaded and sunlit leaves in summer
372 2012 compared to the previous field campaigns (Fig. 5a1-a2). This was similar in *A. unedo* but
373 less pronounced. LMA had no clear pattern in *Q. pubescens*. Elevated LMA was accompanied
374 by higher values of leaf density (D), succulence (S), and carbon content, indicating a more
375 sclerophyllic and succulent structure as response to the drier conditions in 2012 (Figs. S2, S3).
376 N_{mass} was significantly higher in spring and summer 2012 for *Q. ilex* and *P. halepensis* (shaded
377 and sunlit leaves) and for shaded leaves of *A. unedo*, but not for *Q. pubescens* (Fig. 5b1-b2).

378 ***Relationships of photosynthetic and morphological parameters***

379 In order to analyse the general pattern of several relationships of the photosynthetic parameters
380 and foliar traits, we used ANCOVAs to test for differences in the slopes between seasons across
381 all species.

382 The slope the A_{net}/g_s relationship was significantly steeper in summer and autumn 2011^a in all
383 species compared to the other field campaigns (Fig. 6a1, Table S1), suggesting an increased
384 intrinsic water-use efficiency during the dry period in 2011. Shaded leaves had a similar
385 conservative water-use strategy in autumn 2011^a (Fig. 6a2). Shallower slopes in autumn 2011^b
386 in both leaf positions represent rapid responses (less than one week) to the post-drought rains
387 easing the strict stomatal control.

388 The A_{net}/g_m relationship in autumn 2011^a also had a significantly steeper slope in both sunlit and
389 shaded leaves recovering after the first rains in autumn 2011^b (Fig. 6b1-2, Table S2). In the drier
390 year 2012, g_m imposed less resistance on photosynthetic assimilation compared to the wet year
391 2011. The slope of the A_{net}/g_m relationship was significantly higher for winter 2012 than spring
392 and summer 2012, suggesting a stronger control of g_m on photosynthesis in winter. The autumn
393 2011^a and summer 2012 droughts had strong effects on the slope of A_{net}/g_m in shaded leaves.

394 With the ANCOVA of the relationship of g_m and g_s , we investigated the proportional diffusion
395 limitation on photosynthesis. We observed seasonal differences across all species (Fig. 6c,
396 Table S3). Mesophyllic control was stronger in the dry autumn 2011^a and the two winter
397 periods. In contrast, stomatal control was higher than mesophyllic control in the mild 2011
398 summer drought. This was most strongly pronounced in *P. halepensis* and *Q. ilex* (data not
399 shown).

400 The slope in the relationship of $V_{c,max}$ and J_{max} was significantly steeper in autumn 2011^a for
401 both sunlit and shaded (Fig. 7a, Table S4) leaves due to a stronger reduction in $V_{c,max}$ compared
402 to J_{max} . The overall $J_{max}/V_{c,max}$ ratios were 1.09 for sunlit and 1.24 for shaded leaves. The slope
403 of the J_{amb}/A_{net} relationship in sunlit and shaded leaves was significantly lower in the more
404 humid periods (autumn 2011^b, winter 2012, and winter 2013), indicating lower protective
405 energy dissipation and alternative electron pathways under favourable conditions (Fig. 7b, Table
406 S5).

407 Increased foliar sclerophylly led to higher LMAs and thus to higher diffusion resistances in the
408 mesophyll, as shown by the relationship between g_m and LMA (Fig. 8, Table S6). In spring
409 2012 and summer 2012, we detected a less negative exponent (hence a gentler curve) (-0.953
410 and -0.800, respectively) compared to winter 2012 and autumn 2011^a) (-1.486 and -1.533,
411 respectively). This shows that, regardless of the drier conditions and higher LMA in 2012, g_m
412 was higher in this period reflecting a regulatory mechanism of g_m in the CO₂ diffusion pathway
413 (in line with the results of the g_m/g_s analyses).

414

415 **Discussion**

416 **Photosynthetic seasonality and effects of drought**

417 We found that $V_{c,max}$ and J_{max} acclimated strongly to the seasonal changes in temperature and
418 water availability in agreement with previous studies (Vitale and Manes 2005, Corcuera et al.
419 2005, Misson et al. 2006, Ribeiro et al. 2009, Limousin et al. 2010). High radiation and water
420 stress can have a combinatory negative effect on the photosynthetic apparatus, especially in
421 sunlit leaves. Stomata close to avoid transpiration loss and hydraulic failure, but stomatal
422 closure impairs the diffusion of the CO_2 needed in the chloroplasts, the site of carboxylation.
423 $V_{c,max}$ is a proxy for the maximum potential rate of carboxylation, which is carried out by
424 Rubisco, a costly nitrogen-rich protein. The temporary unemployment of Rubisco due to limited
425 substrate (CO_2) availability leads to its de-activation and, during chronic water stress, to its
426 decomposition (Parry 2002, Chaves and Oliveira 2004, Lawlor and Tezara 2009). High
427 incoming radiation that cannot efficiently be dissipated in the Calvin cycle over-excites the
428 photoreaction centres (photoinhibition) and produces reactive oxygen species (ROS) that
429 damage the photosystems and the ATP synthase- needed for the carbon reactions (Epron et al.
430 1993). Leaves prevent harmful excess energy with protective actions such as the reorganisation
431 of the thylakoid membrane, closure of reaction centres, and reduced antennal size (Huner et al.
432 1998, Maxwell and Johnson 2000, Ensminger et al. 2012, Verhoeven 2014). These actions
433 reduce PSII efficiency and J_{max} , and enhance alternative energy pathways to prevent damage on
434 the molecular level on the cost of a lower carbon assimilation.

435 The trees in our study site maintained considerable rates of A_{net} during moderate drought
436 through improved water relations via g_s and g_m control. The relatively stable F_v/F_m values
437 indicate that the protective actions against photoinhibitory stress were effective. The trees
438 showed trunk rehydration after the first autumn rain (Sánchez-Costa *et al.*, unpublished results)
439 and quickly recovered their photosynthetic potential, suggesting that the Rubisco content
440 remained unaffected by moderate drought. The drought impacts were much more severe in the
441 dry year 2012, illustrating the vulnerability of tree physiology to the depletion of soil-water
442 reserves during the early growing season. The severity of drought strongly determined the
443 relative limitations of g_s and g_m on photosynthesis, especially in *Q. ilex* and *P. halepensis*.
444 Stomatal closure regulated photosynthesis during both the moderate and severe droughts; g_m , in
445 contrast, decreased under moderate, but increased under severe drought. We postulate that
446 altered g_m can ease the leaf internal CO_2 diffusion needed for photosynthesis, especially under
447 chronic water stress when depleted non-structural carbohydrates (NSCs) make plants
448 particularly reliant on photosynthetic products for refinement, repair, and protective actions
449 (Niinemets et al. 2009). Major changes of Φ_{PSII} , F_v/F_m , and photosynthetic potentials across all

450 species reflected these refinements of the photosynthetic apparatus as responses to chronic water
451 stress in summer 2012.

452 These acclimatisations occurred not only under dry and hot conditions, but also in winter at high
453 radiation and low temperature. Nevertheless, favourable winter conditions in 2012 resulted in
454 biochemical recovery (peak of $V_{c,max}$ and J_{max}), new shoot growth, and moderate transpiration
455 across species (often exceeding summer values) (Sánchez-Costa *et al.*, unpublished results).
456 Year-round growth patterns with several flushes during the year have also been reported in other
457 studies (Alonso *et al.* 2003). Under novel climatic conditions, favourable conditions in winter
458 may be crucial in the competition between evergreen and deciduous tree species.

459 We observed a highly species-specific pattern. *Q. ilex* and *A. unedo* followed a water-spending,
460 anisohydric strategy that maintained A_{net} and g_s in parallel with lower $V_{c,max}$ and J_{max} . In contrast,
461 *P. halepensis* had significantly decreased g_s , consistent with the conservative water-use strategy
462 and strict stomatal control of isohydric species (Borghetti *et al.* 1998, Martínez-Ferri *et al.*
463 2000). *Q. ilex* generally responded most plastically by rapidly adjusting the photosynthetic
464 machinery to the prevailing conditions (García-Plazaola *et al.* 1997, 1999, Martínez-Ferri *et al.*
465 2004). *P. halepensis* was the most tolerant to photoinhibition and had the most robust
466 photosynthetic machinery to combat abiotic stress (Baquedano and Castillo 2006, Sperlich *et al.*
467 2014). The mesophyll diffusion limitation was lowest in *Q. pubescens* and *A. unedo*, as we
468 claim, due to their deciduous/semi-evergreen foliar habits and lower LMAs (see also Tomás *et al.*
469 *et al.*, 2014). *Q. pubescens* must maximise gas exchange during a shorter growing season, leading
470 to high photosynthetic potentials, A_{net} (Baldocchi *et al.* 2009) and transpiration rates throughout
471 the summer (Sánchez-Costa *et al.*, unpublished results; Poyatos *et al.*, 2008).

472 **Responses specific to leaf position**

473 The seasonality of photosynthetic parameters was qualitatively different between leaf positions
474 (Niinemets *et al.* 2006, Vaz *et al.* 2011) and was mostly pronounced in sunlit leaves. Shaded
475 leaves cushioned the negative climatic effects, maintaining their functionality compared to
476 sunlit leaves. Foliar anatomy, morphology, and biochemistry were highly specialised and
477 dependent on the light regime, leading to smaller but also thicker sunlit leaves and broader and
478 thinner shaded leaves (Kull and Niinemets 1993, Terashima and Hikosaka 1995, Niinemets
479 2001). Shaded leaves had lower N, photosynthetic potentials, carbon metabolisms and higher
480 $J_{max}/V_{c,max}$ ratio (see also Le Roux *et al.* 2001). Shaded leaves invest in higher J_{max} relative to
481 $V_{c,max}$ in order to increase the light-use efficiency. Responses specific to leaf position, however,
482 differed among tree species due to distinct foliar morphologies and crown architectures. The
483 sun-exposed crown position of *P. halepensis*, surmounting the forest canopy resulted in high
484 photosynthetic potentials and a low $J_{max}/V_{c,max}$ ratio throughout the crown. Pine needles attain
485 nearly saturated photosynthetic rates over a wide range of diurnal and seasonal variation in
486 radiation due to their cylindrical shape and steep angles (Jordan and Smith 1993, Lusk *et al.*

487 2003). Similarly, *Q. pubescens* showed a low differentiation between sunlit and shaded leaves.
488 A low $J_{\max}/V_{c,\max}$ ratio throughout the crown suggests a higher proportion of sunlit leaves. In
489 contrast, the comparatively higher $J_{\max}/V_{c,\max}$ ratio of sunlit leaves in *A. unedo* reflects a more
490 shaded growth environment explained by its subordinated position in the forest canopy. The *Q.*
491 *ilex* canopy was dense with a high proportion of shaded leaves, in line with its shade tolerance.
492 Hence, leaf position specific responses were highest in *Q. ilex*. The comparatively higher
493 photosynthetic values in sunlit leaves decreased partly below the level of shaded leaves under
494 stress conditions (see also Sperlich *et al.*, 2014). Shaded leaves are less exposed to the dramatic
495 changes in radiation and temperature in the outer canopy and can be of particular importance for
496 *Q. ilex* to attain a positive net carbon ratio during stress periods (Valladares *et al.* 2008). We
497 stress that the solar environment of the leaves is a crucial factor for assessing tree performance,
498 especially in a competitive environment.

499 **Acclimation of foliar morphology**

500 Mediterranean trees acclimate to water deficits with higher investments in structural
501 compounds, thereby increasing leaf density and succulence (Niinemets 2001, Ogaya and
502 Peñuelas 2006, Poorter *et al.* 2009). Foliar traits are known to be good indicators for the ability
503 of Maquis-species to respond to decreases in rainfall under climate change (Gratani and Varone
504 2006, Ogaya and Peñuelas 2007). We confirm that severe water deficit resulted in increased LT
505 and reduced LA and consequently in higher LMA. It was reported that the plasticity of leaf
506 morphology is generally higher than the plasticity of foliar chemistry and assimilation rates over
507 a wide range of woody species (Niinemets 2001). Under moderate drought, however, foliar
508 morphology was less plastic than foliar chemistry and assimilation rates (Quero *et al.* 2006);
509 severe water stress affected both to a similar extent. Leaf trait acclimation strongly constrained
510 mesophyll conductance under severe drought, especially in *Q. ilex* and *P. halepensis* (see also
511 Tomás *et al.* 2013). We postulate that foliar traits served best as proxies for drought acclimation
512 in *Q. ilex* (Grossoni *et al.* 1998, Bussotti *et al.* 2000) and *P. halepensis* (Alonso *et al.* 2003),
513 both characterised by high leaf-longevities. These changes may be accompanied by increased
514 leaf vein density that may helped to increase the tolerance to foliar hydraulic dysfunction in
515 Mediterranean plants (Nardini *et al.* 2014). The foliar traits of *A. unedo* and *Q. pubescens*
516 acclimated the least, so leaves were susceptible to foliar hydraulic dysfunction and drought-
517 deciduousness. We attribute this species-specificity in leaf trait acclimation to functional
518 differences of leaf investment costs and distinct leaf shedding strategies between deciduous /
519 semi-deciduous (*Q. pubescens* and *A. unedo*) to evergreen sclerophyllic species (*Q. ilex* and *P.*
520 *halepensis*) which we will elaborate further in the following chapter.

521 **Crown defoliation in summer 2012**

522 The lack of rain in early 2012 predisposed the vegetation to leaf senescence observed in summer
523 2012, with high variability across and within species. Leaf senescence was highest in *A. unedo*

524 and *Q. pubescens* – showing partly completely defoliated crowns. *Q. ilex* and mostly *P.*
525 *halepensis* overcame this period with marginal leaf shedding. Stored NSCs strongly determine
526 the recovery of xylem hydraulic conductivity by vessel refilling and the resistance of water
527 transport to drought under prolonged evaporative demand (Ogasa et al. 2013). Depleted NSCs
528 may limit the ability to recover from embolisms (Galiano et al. 2012). *A. unedo* is susceptible to
529 hydraulic dysfunction induced by depleted NSC (e.g. Rosas et al., 2013) which might explain
530 the severe branch dieback of *A. unedo* in our study. As shrubby species characteristic of
531 Maquis-biomes (Beyschlag et al. 1986, Harley et al. 1986), *A. unedo* likely faced a trade-off
532 between growing tall and risking hydraulic dysfunction due to high xylem tension under severe
533 soil-water deficits (Choat et al. 2012). Though, *A. unedo* might contend with severe climatic
534 stress through its strong capacity to resprout (see also Ogaya & Peñuelas, 2004).

535 Pines follow a strategy of water conservation and embolism avoidance, because they have a low
536 capacity to store carbohydrates (Meinzer et al. 2009). *P. halepensis* had a high growth-based
537 water-use efficiency ($WUE_{BAI} = \text{Basal area increment}/\text{Tree transpiration}$) during severe drought
538 (Sánchez-Costa et al., unpublished results), through the combinatory effect of photosynthetic
539 downregulation, foliar-trait acclimation, and improved gas exchange. Thus, this tree species is
540 comparatively the most productive one, especially under drought, confirming its high
541 competitiveness in dry habitats (Zavala and Zea 2004, Maseyk et al. 2008, de Luis et al. 2011).
542 Sánchez-Costa et al. (unpublished results) observed a higher WUE_{BAI} in *Q. pubescens* compared
543 to *Q. ilex* during the soil-moisture deficit in 2012. The “low-cost” leaves of the deciduous *Q.*
544 *pubescens* facilitate drought senescence, so that the reduced transpiratory surface area can
545 effectively avoid damage from hydraulic cavitation and xylem embolism (Ogaya and Peñuelas
546 2006, Barbeta et al. 2013). Fully refoliated crowns in the following growing season was
547 evidence of its success relative to *A. unedo*. The extraordinarily high photosynthetic potentials
548 in the remaining leaves were probably due to a mechanism to compensate for the reduced total
549 leaf area, as indicated by the higher translocation of leaf nitrogen before leaf shedding.

550 *Q. ilex* can effectively tolerate the effects of drought by reducing its LMA and by allowing low
551 water potentials (anisohydric behaviour) (Villar-Salvador et al. 1997, Ogaya and Peñuelas 2006,
552 Limousin et al. 2009). Its hydraulic features are highly plastic, because yearly vessel diameter
553 and recovery are well coupled with annual rainfall (Fotelli et al. 2000, Corcuera et al. 2004). *Q.*
554 *ilex*, however, was also severely effected in 2012, shedding leaves (Tognetti et al. 1998),
555 reducing radial growth and WUE_{BAI} (Sánchez-Costa et al., unpublished results). The positive
556 A_{net} , despite the reduced WUE_{BAI} , suggests that photosynthetic products were used for the
557 maintenance and recovery of xylem hydraulic conductivity instead of growth (Castell et al.
558 1994). In fact, *Quercus* species show generally a good ability in vessel refilling after xylem
559 embolism (Carnicer et al. 2013).

560 **Implications for the global carbon cycle and modelling**

561 There is evidence that the use of seasonally variable photosynthetic potentials reduces
562 uncertainties in modelled ecosystem carbon fluxes relative to the use of constant values (Wilson
563 et al. 2001, Tanaka et al. 2002, Kosugi et al. 2003, 2006, Medvigy et al. 2013). The significant
564 seasonal acclimation of $V_{c,max}$ and J_{max} observed in our study demonstrates that prognostic
565 models should account for seasonal variation, especially in drought-prone areas. Also, the
566 significant role of g_m under abiotic stress periods highlights its importance for estimating the
567 whole-carbon gain. It is now widely accepted that the apparent values of $V_{c,max}$ and J_{max} derived
568 from A/C_i curves are, from a physiological point of view, incorrect. A recent study by Sun *et al.*
569 (2014a) for nearly 130 C_3 species showed that the assumption of infinite g_m in the
570 parameterization of CO_2 -response curves underestimates $V_{c,max}$ and J_{max} by up to 75 and 60%,
571 respectively. Terrestrial biosphere models on regional or global scales are most commonly
572 calibrated on A/C_i -based parameters and therefore use apparent values of $V_{c,max}$ and J_{max} .
573 Incorporating values of $V_{c,max}$ and J_{max} parameterised on A/C_c curves would clearly lead to
574 erroneous results, because their use requires the incorporation of g_m and different Rubisco
575 kinetic parameters into the sub-models of photosynthesis. Therefore, the use of consistent
576 equations and parameters when incorporating parameters from experimental studies into
577 vegetation models is inevitable to correctly estimate photosynthesis (Rogers et al. 2014). From a
578 modelling point of view, it might seem questionable why including g_m and A/C_c - based
579 parameters would improve simulation results and not just increase model complexity.
580 Terrestrial biosphere models are currently well calibrated against observational data despite
581 their use of apparent $V_{c,max}$ and J_{max} . Another criticism often raised is that there are still potential
582 errors in various methods to estimate g_m (and subsequently $V_{c,max}$ and J_{max}) including the
583 variable J - method (used in this study) (Pons et al. 2009, Tholen et al. 2012, Gu and Sun 2014).
584 Nonetheless, large uncertainties remain in the simulations of the future CO_2 fluxes of the global
585 carbon cycle (Anav et al. 2013, Friedlingstein et al. 2014). Patterns of temperature and
586 precipitation are highly uncertain in these models due to both a lack of scientific understanding
587 and model representation (Booth et al. 2012).

588 These uncertainties could partly explain the poor modelling performance for Mediterranean-
589 type ecosystems, because the mechanistic description of the photosynthetic processes under
590 water stress is not very well developed (Morales et al. 2005, Keenan et al. 2011, Zheng et al.
591 2012, Vargas et al. 2013). As we have shown, the limitations imposed by g_m on photosynthetic
592 assimilation can decrease relatively more than the limitations imposed by g_s or biochemistry
593 ($V_{c,max}$ and J_{max}) under drought or winter stress. This distinction has important consequences for
594 the control of water-use efficiency and holds great potential for improving the estimation of
595 ecosystem carbon fluxes under drought conditions (Niinemets et al. 2009a). As already
596 mentioned above, the issue of whether (and how) to include g_m in models is actively debated by
597 physiologists and modellers (see also Rogers et al. 2014). Keenan *et al.* (2010a) showed that g_m

598 was the missing constraint for accurately capturing the response of terrestrial vegetation
599 productivity to drought. Yet relatively little information is available from modelling exercises
600 that have included g_m in their algorithms, and more research in this field is needed.

601 Concluding the above, we underline that we need to consider the seasonality of photosynthetic
602 potentials and mesophyll conductance to explain eco-physiological responses to abiotic stress.
603 These two factors should deserves much more attention in terrestrial biosphere modelling
604 because they hold great potential to reduce model uncertainties, especially under Mediterranean
605 climatic conditions.

606

607 *Acknowledgments*

608 We thank Elisenda Sánchez-Costa and Sílvia Poblador for her assistances in the field and lab
609 work. The research was funded by the European Community's Seventh Framework Programme
610 GREENCYCLESII (FP7 2007-2013) under grant agreement n° 238366 and by the Ministerio de
611 Economica y Competividad under grant agreement n° CGL2011-30590-C02-01
612 (MED_FORESTREAM project) and n° CSD2008-00040 (Consolider-Ingenio MONTES
613 project). JP acknowledges funding from the Spanish Government grant CGL2013-48074-P, the
614 Catalan Government project SGR 2014-274, and the European Research Council Synergy grant
615 ERC-SyG-610028 IMBALANCE-P. M. Ninyerola and M. Batalla (Unitat de Botànica, UAB)
616 provided the climatic database (CGL 2006-01293, MICINN).

617

618 Conflict of Interest: none declared

619

620 **References:**

- 621 Allen CD, Macalady AK, Chenchouni H, Bachelet D, McDowell N, Vennetier M, Kitzberger T, Rigling A,
622 Breshears DD, Hogg EH (Ted) (2010) A global overview of drought and heat-induced tree mortality
623 reveals emerging climate change risks for forests. For Ecol Manage 259:660–684.
624 <http://linkinghub.elsevier.com/retrieve/pii/S037811270900615X> (18 July 2011, date last accessed
625).
- 626 Alonso R, Elvira S, Inclán R, Bermejo V, Castillo FJ, Gimeno BS (2003) Responses of Aleppo pine to ozone.
627 In: Karnosky DF, Percy KE, Chappelka AH, Simpson C, Pikkariainen J (eds) Air Pollution, Global
628 Change and Forests in the New Millenium. Elsevier Ltd., pp 211–230.
- 629 Anav A, Friedlingstein P, Kidston M, Bopp L, Ciais P, Cox P, Jones C, Jung M, Myrneni R, Zhu Z (2013)
630 Evaluating the Land and Ocean Components of the Global Carbon Cycle in the CMIP5 Earth System
631 Models. J Clim 26:6801–6843. <http://journals.ametsoc.org/doi/abs/10.1175/JCLI-D-12-00417.1>
632 (27 August 2014, date last accessed).
- 633 Anderegg WRL, Kane JM, Anderegg LDL (2013) Consequences of widespread tree mortality triggered by
634 drought and temperature stress. Nat Clim Chang 3:30–36.
- 635 Baldocchi DD, Ma S, Rambal S, Misson L, Ourcival J-M, Limousin J-M, Pereira J, Papale D (2009) On the
636 differential advantages of evergreenness and deciduousness in mediterranean oak woodlands: a
637 flux perspective. Ecol Appl 20:1583–1597.
- 638 Baquedano FJ, Castillo FJ (2006) Comparative ecophysiological effects of drought on seedlings of the
639 Mediterranean water-saver *Pinus halepensis* and water-spenders *Quercus coccifera* and *Quercus*
640 *ilex*. Trees 20:689–700. <http://link.springer.com/10.1007/s00468-006-0084-0> (17 June 2013, date
641 last accessed).
- 642 Barbeta A, Ogaya R, Peñuelas J (2012) Comparative study of diurnal and nocturnal sap flow of *Quercus*
643 *ilex* and *Phillyrea latifolia* in a Mediterranean holm oak forest in Prades (Catalonia, NE Spain).
644 Trees 26:1651–1659. <http://link.springer.com/10.1007/s00468-012-0741-4> (8 August 2013, date
645 last accessed).
- 646 Barbeta A, Ogaya R, Peñuelas J (2013) Dampening effects of long-term experimental drought on growth
647 and mortality rates of a Holm oak forest. Glob Chang Biol 19:1–12.
648 <http://www.ncbi.nlm.nih.gov/pubmed/23712619> (10 August 2013, date last accessed).
- 649 Beaumont LJ, Hughes L, Pitman a J (2008) Why is the choice of future climate scenarios for species
650 distribution modelling important? Ecol Lett 11:1135–46.
651 <http://www.ncbi.nlm.nih.gov/pubmed/18713269> (23 July 2011, date last accessed).
- 652 Bernacchi CJ, Portis AR, Nakano H, Caemmerer S Von, Long SP (2002) Temperature Response of
653 Mesophyll Conductance . Implications for the Determination of Rubisco Enzyme Kinetics and for
654 Limitations to Photosynthesis in Vivo. Plant Physiol 130:1992–1998.
- 655 Beyschlag W, Lange OL, Tenhunen JD (1986) Photosynthesis und Wasserhaushalt der immergrünen
656 mediterranen Hartlaubpflanze *Arbutus unedo* L. im Jahresverlauf am Freilandstandort in Portugal
657 I. Tagesläufe von CO₂-Gaswechsel und Transpiration unter natürlichen Bedingungen. Flora
658 178:409–444.
- 659 Booth BBB, Jones CD, Collins M, Totterdell IJ, Cox PM, Sitch S, Huntingford C, Betts R a, Harris GR, Lloyd J
660 (2012) High sensitivity of future global warming to land carbon cycle processes. Environ Res Lett
661 7:024002. [20](http://stacks.iop.org/1748-</p></div><div data-bbox=)

- 662 9326/7/i=2/a=024002?key=crossref.0608852a9b4c87899a93d976c7428fa8 (3 November 2014,
663 date last accessed).
- 664 Borghetti M, Cinnirella S, Magnani F, Saracino A (1998) Impact of long-term drought on xylem embolism
665 and growth in *Pinus halepensis* Mill. *Trees* 12:187–195.
- 666 Buckley TN, Diaz-Espejo A (2015) Reporting estimates of maximum potential electron transport rate.
667 *New Phytol* 205:14–17. <http://onlinelibrary.wiley.com/doi/10.1111/nph.13018/abstract> (15
668 September 2014, date last accessed).
- 669 Bugmann H, Palahí M, Bontemps J, Tomé M (2010) Trends in modeling to address forest management.
670 *For Syst* 19:3–7.
- 671 Bussotti F, Borghini F, Celesti C, Leonzio C, Bruschi P (2000) Leaf morphology and macronutrients in
672 broadleaved trees in central Italy. *Trees* 14:361–368.
673 <http://link.springer.com/10.1007/s004680000056> (18 September 2014, date last accessed).
- 674 Carnicer J, Barbeta A, Sperlich D, Coll M, Peñuelas J (2013) Contrasting trait syndromes in angiosperms
675 and conifers are associated with different responses of tree growth to temperature on a large
676 scale. *Front Plant Sci* 4:409.
677 [http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3797994&tool=pmcentrez&renderty
678 pe=abstract](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3797994&tool=pmcentrez&rendertype=abstract) (12 November 2013, date last accessed).
- 679 Carnicer J, Coll M, Ninyerola M, Pons X, Sánchez G, Peñuelas J (2011) Widespread crown condition
680 decline, food web disruption, and amplified tree mortality with increased climate change-type
681 drought. *Proc Natl Acad Sci U S A* 108:1474–8.
682 [http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3029725&tool=pmcentrez&renderty
683 pe=abstract](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3029725&tool=pmcentrez&rendertype=abstract) (13 August 2013, date last accessed).
- 684 Castell C, Terradas J, Tenhunen JD (1994) Water relations, gas exchange, and growth of resprouts and
685 mature plant shoots of *Arbutus unedo* L. and *Quercus ilex* L. *Oecologia* 98:201–211.
- 686 Di Castri F (1973) Climatographical comparison between Chile and the western coast of North America.
687 In: Di Castri F, Mooney HA (eds) *Mediterranean-type ecosystems: origin and structure*. Springer,
688 Berlin, pp 21–36.
- 689 Chaves MM, Oliveira MM (2004) Mechanisms underlying plant resilience to water deficits: prospects for
690 water-saving agriculture. *J Exp Bot* 55:2365–84. <http://www.ncbi.nlm.nih.gov/pubmed/15475377>
691 (14 August 2013, date last accessed).
- 692 Chaves MM, Pereira JS, Maroco J, Rodrigues ML, Ricardo CPP, Osorio ML, Carvalho I, Faria T, Pinheiro C
693 (2002) How Plants Cope with Water Stress in the Field . *Photosynthesis and Growth*. *Ann Bot*
694 89:907–916.
- 695 Choat B (2013) Predicting thresholds of drought-induced mortality in woody plant species. *Tree Physiol*
696 33:669–671. <http://www.ncbi.nlm.nih.gov/pubmed/23878170> (7 August 2013, date last accessed
697).
- 698 Choat B, Jansen S, Brodribb TJ, Cochard H, Delzon S, Bhaskar R, Bucci SJ, Feild TS, Gleason SM, Hacke UG,
699 Jacobsen AL, Lens F, Maherali H, Martínez-Vilalta J, Mayr S, Mencuccini M, Mitchell PJ, Nardini A,
700 Pittermann J, Pratt RB, Sperry JS, Westoby M, Wright IJ, Zanne AE (2012) Global convergence in
701 the vulnerability of forests to drought. *Nature* 491:752–5.
- 702 Corcuera L, Morales F, Abadia a., Gil-Pelegrin E (2004) The effect of low temperatures on the
703 photosynthetic apparatus of *Quercus ilex* subsp. *ballota* at its lower and upper altitudinal limits in

- 704 the Iberian peninsula and during a single freezing-thawing cycle. *Trees* 19:99–108.
705 <http://www.springerlink.com/index/10.1007/s00468-004-0368-1> (31 October 2012, date last
706 accessed).
- 707 Corcuera L, Morales F, Abadía a, Gil-Pelegrín E (2005) Seasonal changes in photosynthesis and
708 photoprotection in a *Quercus ilex* subsp. *ballota* woodland located in its upper altitudinal extreme
709 in the Iberian Peninsula. *Tree Physiol* 25:599–608.
710 <http://www.ncbi.nlm.nih.gov/pubmed/15741152>
- 711 Eichhorn J, Roskams P, Ferretti M, Mues V, Szepesi A, Durrant D (2010) Visual assessment of crown
712 condition and damaging agents. 49 pp. Manual Part IV. Man methods criteria Harmon sampling,
713 assessment, Monit Anal Eff air Pollut For UNECE ICP For Program Co-ord Centre,
714 Hamburg[<http://www.icp-forests.org/Manual.htm>]. <http://www.icp-forests.org/Manual.htm>
- 715 Ensminger I, Berninger F, Streb P (2012) Response of photosynthesis to low temperature. In: Flexas J,
716 Loreto F, Medrano H (eds) *Terrestrial Photosynthesis in a Changing Environment - A Molecular,
717 Physiological and Ecological Approach*. Cambridge University Press, Cambridge, pp 272–289.
- 718 Epron D, Dreyer E, Aussenac G (1993) A comparison of photosynthetic responses to water stress in
719 seedlings from 3 oak species: *Quercus petraea* (Matt) Liebl, *Q. rubra* L and *Q. cerris* L. *Ann Des Sci
720 For* 50:48–60. <http://www.afs-journal.org/10.1051/forest:19930704>
- 721 Flexas J, Diaz-Espejo A, Gago J, Gallé A, Galmés J, Gulías J, Medrano H (2014) Photosynthetic limitations
722 in Mediterranean plants: A review. *Environ Exp Bot* 103:12–23.
723 <http://linkinghub.elsevier.com/retrieve/pii/S0098847213001238> (6 March 2014, date last
724 accessed).
- 725 Fotelli MN, Radoglou KM, Constantinidou HI (2000) Water stress responses of seedlings of four
726 Mediterranean oak species. *Tree Physiol* 20:1065–75.
727 <http://www.ncbi.nlm.nih.gov/pubmed/11269958>
- 728 Friedlingstein P, Betts R, Bopp L, Bloh W Von, Brovkin V, Doney S, Eby M, Fung I, Govindasamy B, John J,
729 Jones C, Joos F, Kato T, Kawamiya M, Knorr W, Lindsay K, Matthews HD, Raddatz T, Rayner P, Reick
730 C, Roeckner E, Schnitzler KG, Schnurr R, Strassmann K, Thompson S, J Weaver A, Yoshikawa C,
731 Zeng N (2006) Climate –carbon cycle feedback analysis, results from the C4MIP model
732 intercomparison. *J Clim In review*
- 733 Friedlingstein P, Meinshausen M, Arora VK, Jones CD, Anav A, Liddicoat SK, Knutti R (2014) Uncertainties
734 in CMIP5 Climate Projections due to Carbon Cycle Feedbacks. *J Clim* 27:511–526.
735 <http://journals.ametsoc.org/doi/abs/10.1175/JCLI-D-12-00579.1> (8 September 2014, date last
736 accessed).
- 737 Friedlingstein P, Prentice I (2010) Carbon–climate feedbacks: a review of model and observation based
738 estimates. *Curr Opin Environ Sustain* 2:251–257.
739 <http://linkinghub.elsevier.com/retrieve/pii/S1877343510000473>
- 740 Friend AD (2010) Terrestrial plant production and climate change. *J Exp Bot* 61:1293–309.
741 <http://www.ncbi.nlm.nih.gov/pubmed/20202998> (3 March 2012, date last accessed).
- 742 Galiano L, Martínez-Vilalta J, Sabatés S, Lloret F (2012) Determinants of drought effects on crown
743 condition and their relationship with depletion of carbon reserves in a Mediterranean holm oak
744 forest. *Tree Physiol* 32:478–89. <http://www.ncbi.nlm.nih.gov/pubmed/22499595> (1 February
745 2013, date last accessed).

- 746 García-Plazaola JI, Artetxe U, Beccheril, María J (1999) Diurnal changes in antioxidant and carotenoid
747 composition in the Mediterranean sclerophyll tree *Quercus ilex* (L.) during winter. *Plant Sci*
748 143:125–133.
- 749 García-Plazaola JI, Faria T, Abadia J, Abadia A, Chaves MM, Pereira JS (1997) Seasonal changes in
750 xanthophyll composition and photosynthesis of cork oak (*Quercus suber* L.) leaves under
751 mediterranean climate. *J Experimental Bot* 48:1667–1674.
- 752 Grassi G, Magnani F (2005) Stomatal, mesophyll conductance and biochemical limitations to
753 photosynthesis as affected by drought and leaf ontogeny in ash and oak trees. *Plant, Cell Environ*
754 28:834–849. <http://www.blackwell-synergy.com/doi/abs/10.1111/j.1365-3040.2005.01333.x>
- 755 Gratani L, Ghia E (2002) Adaptive strategy at the leaf level of *Arbutus unedo* L. to cope with
756 Mediterranean climate. *Flora* 197:275–284.
- 757 Gratani L, Varone L (2004) Adaptive photosynthetic strategies of the Mediterranean maquis species
758 according to their origin. *Photosynthetica* 42:551–558.
- 759 Gratani L, Varone L (2006) Long-time variations in leaf mass and area of Mediterranean evergreen
760 broad-leaf and narrow-leaf maquis species. *Photosynthetica* 44:161–168.
- 761 Grossoni P, Bussotti F, Tani C, Gravano E, Antarelli S, Bottacci A (1998) Morpho-anatomical alterations in
762 leaves of *Gagus vatica* L. and *Quercus ilex* L. in different environmental stress condition.
763 *Chemosphere* 36:919–924.
764 <http://www.sciencedirect.com/science/article/pii/S0045653597101485> (22 September 2014, date
765 last accessed).
- 766 Gu L, Sun Y (2014) Artificial responses of mesophyll conductance to CO₂ and irradiance estimated with
767 the variable J and online isotope discrimination methods. *Plant Cell Environ* 37:1231–1249.
768 <http://doi.wiley.com/10.1111/pce.12232> (15 November 2013, date last accessed).
- 769 Gulías J, Cifre J, Jonasson S, Medrano H, Flexas J (2009) Seasonal and inter-annual variations of gas
770 exchange in thirteen woody species along a climatic gradient in the Mediterranean island of
771 Mallorca. *Flora - Morphol Distrib Funct Ecol Plants* 204:169–181.
772 <http://linkinghub.elsevier.com/retrieve/pii/S036725300800114X> (2 August 2013, date last
773 accessed).
- 774 Harley PC, Loreto F, Di Marco G, Sharkey TD (1992) Theoretical Considerations when Estimating the
775 Mesophyll Conductance to CO₂ Flux by Analysis of the Response of Photosynthesis to CO₂. *Plant*
776 *Physiol* 98:1429–1436.
777 [http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1080368&tool=pmcentrez&renderty
778 pe=abstract](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1080368&tool=pmcentrez&rendertype=abstract)
- 779 Harley PC, Tenhunen JD, Lange OL (1986) Use of an analytical model to study limitations on net
780 photosynthesis in *Arbutus unedo* under field conditions. *Oecologia* 70:393–401.
- 781 Hickler T, Fronzek S, Araújo MB, Schweiger O, Thuiller W, Sykes MT (2009) An ecosystem model-based
782 estimate of changes in water availability differs from water proxies that are commonly used in
783 species distribution models. *Glob Ecol Biogeogr* 18:304–313. [http://doi.wiley.com/10.1111/j.1466-
784 8238.2009.00455.x](http://doi.wiley.com/10.1111/j.1466-8238.2009.00455.x) (9 August 2013, date last accessed).
- 785 Huner NPA, Öquist G, Sarhan F (1998) Energy balance and acclimation to light and cold. *Trends Plant Sci*
786 3:224–230. <http://linkinghub.elsevier.com/retrieve/pii/S1360138598012485>

- 787 IPCC (2013) Summary for Policymakers. In: T.F. S, Qin D, Plattner G-K, Tignor M, Allen SK, Boschung J,
788 Nauels A, Xia Y, Bex V, Midgley PM (eds) Climate change 2013: the physical science basis.
789 Cambridge University Press, Cambridge (UK) and New York (USA)
- 790 Jordan DN, Smith WK (1993) Simulated influence of leaf geometry on sunlight interception and
791 photosynthesis in conifer needles. *Tree Physiol* 13:29–39.
792 <http://www.ncbi.nlm.nih.gov/pubmed/14969899>
- 793 Jump AS, Penuelas J (2005) Running to stand still: adaptation and the response of plants to rapid climate
794 change. *Ecol Lett* 8:1010–1020. <http://doi.wiley.com/10.1111/j.1461-0248.2005.00796.x> (3 June
795 2013, date last accessed).
- 796 Keenan T, Maria Serra J, Lloret F, Ninyerola M, Sabate S (2011) Predicting the future of forests in the
797 Mediterranean under climate change, with niche- and process-based models: CO2 matters! *Glob
798 Chang Biol* 17:565–579. <http://doi.wiley.com/10.1111/j.1365-2486.2010.02254.x> (23 July 2011,
799 date last accessed).
- 800 Keenan T, Sabate S, Gracia C (2010) The importance of mesophyll conductance in regulating forest
801 ecosystem productivity during drought periods. *Glob Chang Biol* 16:1019–1034.
802 <http://doi.wiley.com/10.1111/j.1365-2486.2009.02017.x> (19 June 2011, date last accessed).
- 803 Kosugi Y, Shibata S, Kobashi S (2003) Parameterization of the CO₂ and H₂O gas exchange of several
804 temperate deciduous broad-leaved trees at the leaf scale considering seasonal changes. *Plant Cell
805 Environ* 26:285–301.
- 806 Kosugi Y, Takanashi S, Matsuo N, Tanaka K, Tanaka H (2006) Impact of leaf physiology on gas exchange
807 in a Japanese evergreen broad-leaved forest. *Agric For Meteorol* 139:182–199.
808 <http://linkinghub.elsevier.com/retrieve/pii/S0168192306001602> (28 August 2013, date last
809 accessed).
- 810 Krasteva V, Alexandrov V, Chepishcheva M, Dambov S, Stefanov D, Yordanov I, Goltsev V (2013) Drought
811 induced damages of photosynthesis in bean and plantain plants analyzed in vivo by chlorophyll a
812 fluorescence. *Bulg J Plant Physiol* 19:39–44.
- 813 Kull O, Niinemets U (1993) Variations in leaf morphometry and nitrogen concentration in *Betula pendula*
814 Roth., *Corylus avellana* L. and *Lonicera xylosteum* L. *Tree Physiol* 12:311–8.
815 <http://www.ncbi.nlm.nih.gov/pubmed/14969921>
- 816 Lawlor DW, Tezara W (2009) Causes of decreased photosynthetic rate and metabolic capacity in water-
817 deficient leaf cells: a critical evaluation of mechanisms and integration of processes. *Ann Bot*
818 103:561–79.
819 [http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2707350&tool=pmcentrez&renderty
820 pe=abstract](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2707350&tool=pmcentrez&rendertype=abstract) (14 August 2013, date last accessed).
- 821 Limousin J-M, Misson L, Lavoie A-V, Martin NK, Rambal S (2010) Do photosynthetic limitations of
822 evergreen *Quercus ilex* leaves change with long-term increased drought severity? *Plant Cell
823 Environ* 33:863–75. <http://www.ncbi.nlm.nih.gov/pubmed/20051039> (23 July 2011, date last
824 accessed).
- 825 Limousin JM, Rambal S, Ourcival JM, Rocheteau a., Joffre R, Rodriguez-Cortina R (2009) Long-term
826 transpiration change with rainfall decline in a Mediterranean *Quercus ilex* forest. *Glob Chang Biol*
827 15:2163–2175. <http://doi.wiley.com/10.1111/j.1365-2486.2009.01852.x> (13 August 2013, date
828 last accessed).

- 829 Lookingbill TR, Zavala MA (2000) Spatial pattern of *Quercus ilex* and *Quercus pubescens* recruitment in
830 *Pinus halepensis* dominated woodlands. *J Veg Sci* 11:607–612.
- 831 De Luis M, Novak K, Raventós J, Gričar J, Prislán P, Čufar K (2011) Cambial activity, wood formation and
832 sapling survival of *Pinus halepensis* exposed to different irrigation regimes. *For Ecol Manage*
833 262:1630–1638. <http://linkinghub.elsevier.com/retrieve/pii/S0378112711004415> (31 March 2014,
834 date last accessed).
- 835 Luo Y (2007) Terrestrial Carbon–Cycle Feedback to Climate Warming. *Annu Rev Ecol Evol Syst* 38:683–
836 712. <http://www.annualreviews.org/doi/abs/10.1146/annurev.ecolsys.38.091206.095808>
- 837 Lusk CH, Wright I, Reich PB (2003) Photosynthetic differences contribute to competitive advantage of
838 evergreen angiosperm trees over evergreen conifers in productive habitats. *New Phytol* 160:329–
839 336.
- 840 Martínez-Ferri E, Balaguer L, Valladares F, Chico JM, Manrique E (2000) Energy dissipation in drought-
841 avoiding and drought-tolerant tree species at midday during the Mediterranean summer. *Tree*
842 *Physiol* 20:131–138. <http://treephys.oxfordjournals.org/cgi/doi/10.1093/treephys/20.2.131>
- 843 Martínez-Ferri E, Manrique E, Valladares F, Balaguer L (2004) Winter photoinhibition in the field involves
844 different processes in four co-occurring Mediterranean tree species. *Tree Physiol* 24:981–90.
845 <http://www.ncbi.nlm.nih.gov/pubmed/15234895>
- 846 Martínez-Vilalta J, Piñol J (2002) Drought-induced mortality and hydraulic architecture in pine
847 populations of the NE Iberian Peninsula. *For Ecol Manage* 161:247–256.
- 848 Martin-StPaul NK, Limousin J-M, Vogt-Schilb H, Rodríguez-Calcerrada J, Rambal S, Longepierre D, Misson
849 L (2013) The temporal response to drought in a Mediterranean evergreen tree: comparing a
850 regional precipitation gradient and a throughfall exclusion experiment. *Glob Chang Biol* 19:2413–
851 26. <http://www.ncbi.nlm.nih.gov/pubmed/23553916> (6 August 2013, date last accessed).
- 852 Maseyk KS, Lin T, Rotenberg E, Grünzweig JM, Schwartz A, Yakir D (2008) Physiology–phenology
853 interactions in a productive semi-arid pine forest. *New Phytol* 178:603–616.
- 854 Matusick G, Ruthrof KX, Brouwers NC, Dell B, Hardy GSJ (2013) Sudden forest canopy collapse
855 corresponding with extreme drought and heat in a mediterranean-type eucalypt forest in
856 southwestern Australia. *Eur J For Res* 132:497–510.
- 857 Maxwell K, Johnson GN (2000) Chlorophyll fluorescence — a practical guide. *J Exp Bot* 51:659–668.
- 858 McDowell N, Pockman WT, Allen CD, Breshears DD, Cobb N, Kolb T, Plaut J, Sperry J, West A, Williams
859 DG, Yezzer E a (2008) Mechanisms of plant survival and mortality during drought: why do some
860 plants survive while others succumb to drought? *New Phytol* 178:719–39.
861 <http://www.ncbi.nlm.nih.gov/pubmed/18422905>
- 862 Medvigy D, Jeong S-J, Clark KL, Skowronski NS, Schäfer KVR (2013) Effects of seasonal variation of
863 photosynthetic capacity on the carbon fluxes of a temperate deciduous forest. *J Geophys Res*
864 *Biogeosciences* 118:1703–1714. <http://doi.wiley.com/10.1002/2013JG002421> (26 March 2014,
865 date last accessed).
- 866 Meinzer FC, Johnson DM, Lachenbruch B, McCulloh K a., Woodruff DR (2009) Xylem hydraulic safety
867 margins in woody plants: coordination of stomatal control of xylem tension with hydraulic
868 capacitance. *Funct Ecol* 23:922–930. <http://doi.wiley.com/10.1111/j.1365-2435.2009.01577.x> (25
869 May 2013, date last accessed).

- 870 Misson L, Tu KP, Boniello R a, Goldstein AH (2006) Seasonality of photosynthetic parameters in a multi-
871 specific and vertically complex forest ecosystem in the Sierra Nevada of California. *Tree Physiol*
872 26:729–41. <http://www.ncbi.nlm.nih.gov/pubmed/16510388>
- 873 Morales P, Sykes MT, Prentice IC, Smith P, Smith B, Bugmann H, Zierl B, Friedlingstein P, Viovy N, Sabate
874 S, Sanchez A, Pla E, Gracia C a., Sitch S, Arneth A, Ogee J (2005) Comparing and evaluating process-
875 based ecosystem model predictions of carbon and water fluxes in major European forest biomes.
876 *Glob Chang Biol* 11:2211–2233. <http://doi.wiley.com/10.1111/j.1365-2486.2005.01036.x> (18 July
877 2011, date last accessed).
- 878 Mott KA, Buckley TN (1998) Stomatal heterogeneity. *J Experimental Bot* 49:407–417.
- 879 Mott KA, Buckley TN (2000) Patchy stomatal conductance: emergent collective behaviour of stomata.
880 *Trends Plant Sci* 1385:258–262.
- 881 Nardini A, Lo Gullo MA, Trifilò P, Salleo S (2014) The challenge of the Mediterranean climate to plant
882 hydraulics: Responses and adaptations. *Environ Exp Bot* 103:68–79.
883 <http://www.sciencedirect.com/science/article/pii/S0098847213001548> (5 September 2014, date
884 last accessed).
- 885 Niinemets Ü (2001) Global-scale climatic controls of leaf dry mass per area, density, and thickness in
886 trees and shrubs. *Ecology* 82:453–469.
- 887 Niinemets Ü, Cescatti A, Rodeghiero M, Tosens T (2005) Leaf internal diffusion conductance limits
888 photosynthesis more strongly in older leaves of Mediterranean evergreen broad-leaved species.
889 *Plant, Cell Environ* 28:1552–1566. <http://doi.wiley.com/10.1111/j.1365-3040.2005.01392.x>
- 890 Niinemets Ü, Cescatti A, Rodeghiero M, Tosens T (2006) Complex adjustments of photosynthetic
891 potentials and internal diffusion conductance to current and previous light availabilities and leaf
892 age in Mediterranean evergreen species *Quercus ilex*. *Plant, Cell Environ* 29:1159–1178.
893 <http://doi.wiley.com/10.1111/j.1365-3040.2006.01499.x> (23 July 2011, date last accessed).
- 894 Niinemets U, Díaz-Espejo A, Flexas J, Galmés J, Warren CR (2009) Role of mesophyll diffusion
895 conductance in constraining potential photosynthetic productivity in the field. *J Exp Bot* 60:2249–
896 70. <http://www.ncbi.nlm.nih.gov/pubmed/19395391> (20 July 2011, date last accessed).
- 897 Niinemets Ü, Keenan T (2014) Photosynthetic responses to stress in Mediterranean evergreens:
898 mechanisms and models. *Environ Exp Bot* 103:24–41.
899 <http://linkinghub.elsevier.com/retrieve/pii/S0098847213001925> (26 November 2013, date last
900 accessed).
- 901 Ninyerola M, Pons X, Roure JM (2007a) Objective air temperature mapping for the Iberian Peninsula
902 using spatial interpolation and GIS. *Int J Climatol* 27:1231–1242.
903 <http://dx.doi.org/10.1002/joc.1462>
- 904 Ninyerola M, Pons X, Roure JM (2007b) Monthly precipitation mapping of the Iberian Peninsula using
905 spatial interpolation tools implemented in a Geographic Information System. *Theor Appl Climatol*
906 89:195–209. <http://link.springer.com/10.1007/s00704-006-0264-2> (14 December 2014, date last
907 accessed).
- 908 Ogasa M, Miki NH, Murakami Y, Yoshikawa K (2013) Recovery performance in xylem hydraulic
909 conductivity is correlated with cavitation resistance for temperate deciduous tree species. *Tree*
910 *Physiol* 33:335–44. <http://www.ncbi.nlm.nih.gov/pubmed/23492871> (22 May 2013, date last
911 accessed).

- 912 Ogaya R, Peñuelas J (2004) Phenological patterns of *Quercus ilex*, *Phillyrea latifolia*, and *Arbutus unedo*
913 growing under a field experimental drought. *Écoscience* 11:263–270.
- 914 Ogaya R, Peñuelas J (2006) Contrasting foliar responses to drought in *Quercus ilex* and *Phillyrea latifolia*.
915 *Biol Plant* 50:373–382.
- 916 Ogaya R, Peñuelas J (2007) Leaf mass per area ratio in *Quercus ilex* leaves under a wide range of climatic
917 conditions. The importance of low temperatures. *Acta Oecologica* 31:168–173.
918 <http://linkinghub.elsevier.com/retrieve/pii/S1146609X06001482> (18 February 2014, date last
919 accessed).
- 920 Orshan G (1983) Approaches to the Definition of Mediterranean Growth Forms. In: Kruger FJ, Mitchell
921 DT, Jarvis JUM (eds) *Mediterranean type-ecosystems - The role of nutrients*. Springer-Verlag,
922 Berlin - Heidelberg, pp pp. 86–100.
- 923 Parry M a. J (2002) Rubisco Activity: Effects of Drought Stress. *Ann Bot* 89:833–839.
- 924 Pausas JG, Llovet J, Rodrigo A, Ramon Vallejo (2008) Are wildfires a disaster in the Mediterranean
925 basin ? – A review. *Int J Wildl Fire* 17:713–723.
- 926 Peñuelas J, Lloret F, Montoya R (2001) Severe Drought Effects on Mediterranean Woody Flora in Spain.
927 *For Sci* 47:214–218.
- 928 Piñol J, Terradas J, Lloret F (1998) Climate warming, wildfire hazard, and wildfire occurrence in coastal
929 eastern Spain. *Clim Change* 38:345–357.
- 930 Pons TL, Flexas J, von Caemmerer S, Evans JR, Genty B, Ribas-Carbo M, Brugnoli E (2009) Estimating
931 mesophyll conductance to CO₂: methodology, potential errors, and recommendations. *J Exp Bot*
932 60:2217–34. <http://www.ncbi.nlm.nih.gov/pubmed/19357431> (12 March 2012, date last accessed
933).
- 934 Poorter H, Niinemets Ü, Poorter L, Wright IJ, Villar R (2009) Causes and consequences of variation in leaf
935 mass per area (LMA): a meta-analysis. *New Phytol* 182:565–588.
936 <http://doi.wiley.com/10.1111/j.1469-8137.2009.02830.x>
- 937 Poyatos R, Aguadé D, Galiano L, Mencuccini M, Martínez-Vilalta J (2013) Drought-induced defoliation
938 and long periods of near-zero gas exchange play a key role in accentuating metabolic decline of
939 Scots pine. *New Phytol*. <http://www.ncbi.nlm.nih.gov/pubmed/23594415> (31 May 2013, date last
940 accessed).
- 941 Poyatos R, Llorens P, Piñol J, Rubio C (2008) Response of Scots pine (*Pinus sylvestris* L.) and pubescent
942 oak (*Quercus pubescens* Willd.) to soil and atmospheric water deficits under Mediterranean
943 mountain climate. *Ann For Sci* 65:306–306.
- 944 Quero JL, Villar R, Marañón T, Zamora R (2006) Interactions of drought and shade effects on seedlings of
945 four *Quercus* species: physiological and structural leaf responses. *New Phytol* 170:819–33.
946 <http://www.ncbi.nlm.nih.gov/pubmed/16684241> (16 September 2014, date last accessed).
- 947 Raftoyannis Y, Spanos I, Radoglou K (2008) The decline of Greek fir (*Abies cephalonica* Loudon):
948 Relationships with root condition. *Plant Biosyst* 142:386–390.
949 <http://www.tandfonline.com/doi/abs/10.1080/11263500802151017> (6 August 2013, date last
950 accessed).

- 951 Reichstein M, Tenhunen JD, Roupsard O, Ourcival J-M, Rambal S, Dore S, Valentini R (2002) Ecosystem
952 respiration in two Mediterranean evergreen Holm Oak forests: drought effects and decomposition
953 dynamics. *Funct Ecol* 16:27–39. <http://doi.wiley.com/10.1046/j.0269-8463.2001.00597.x>
- 954 Reynolds JF, Kemp PR, Acock B, Chen J, Moorhead DL (1996) Progress, Limitations and Challenges in
955 Modeling the Effects of Elevated CO₂ on Plants and Ecosystems. In: Koch GW, Mooney HA (eds)
956 Carbon Dioxide and Terrestrial Ecosystems. Academic Press, San Diego - New York - Boston -
957 London - Sydney - Tokio - Toronto, pp 347–380.
- 958 Ribeiro RV, Machado EC, Santos MG, Oliveira RF (2009) Seasonal and diurnal changes in photosynthetic
959 limitation of young sweet orange trees. *Environ Exp Bot* 66:203–211.
960 <http://linkinghub.elsevier.com/retrieve/pii/S0098847209000653> (23 July 2011, date last accessed
961).
- 962 Rogers A, Medlyn BE, Dukes JS (2014) 9th New Phytologist Workshop, Montauk, New York, USA, April
963 2014. In: Improving representation of photosynthesis in Earth System Models. pp 12–14.
964 <http://onlinelibrary.wiley.com/doi/10.1111/nph.12972/abstract>
- 965 Rosas T, Galiano L, Ogaya R, Peñuelas J, Martínez-Vilalta J (2013) Dynamics of non-structural
966 carbohydrates in three Mediterranean woody species following long-term experimental drought.
967 *Front Plant Sci* 4:1–16.
968 [http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3795346&tool=pmcentrez&renderty
969 pe=abstract](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3795346&tool=pmcentrez&rendertype=abstract) (6 August 2014, date last accessed).
- 970 Le Roux X, Walcroft a S, Sinoquet H, Chaves MM, Rodrigues a, Osorio L (2001) Photosynthetic light
971 acclimation in peach leaves: importance of changes in mass:area ratio, nitrogen concentration,
972 and leaf nitrogen partitioning. *Tree Physiol* 21:377–86.
973 <http://www.ncbi.nlm.nih.gov/pubmed/11282577>
- 974 Sabaté S (2002) Likely effects of climate change on growth of *Quercus ilex*, *Pinus halepensis*, *Pinus*
975 *pinaster*, *Pinus sylvestris* and *Fagus sylvatica* forests in the Mediterranean region. *For Ecol Manage*
976 162:23–37. <http://linkinghub.elsevier.com/retrieve/pii/S0378112702000488>
- 977 Sharkey TD (1985) Photosynthesis in intact leaves of C₃ plants: Physics, physiology and rate limitations.
978 *Bot Rev* 51:53–105.
- 979 Somot S, Sevault F, Déqué M, Crépon M (2008) 21st century climate change scenario for the
980 Mediterranean using a coupled atmosphere–ocean regional climate model. *Glob Planet Change*
981 63:112–126. <http://linkinghub.elsevier.com/retrieve/pii/S0921818107001774> (13 August 2013,
982 date last accessed).
- 983 Specht RL (1969) A comparison of the sclerophyllous vegetation characteristics of Mediterranean type
984 climates in France, California, and southern Australia. I. Structure, morphology, and succession.
985 *Aust J Bot* 17:277–292.
- 986 Sperlich D, Chang CT, Peñuelas J, Gracia C, Sabaté S (2014) Foliar photochemical processes and carbon
987 metabolism under favourable and adverse winter conditions in a Mediterranean mixed forest,
988 Catalonia (Spain). *Biogeosciences* 11:5657–5674.
989 <http://www.biogeosciences.net/11/5657/2014/bg-11-5657-2014.html> (16 October 2014, date last
990 accessed).
- 991 Sun Y, Gu L, Dickinson RE, Pallardy SG, Baker J, Cao Y, Damatta FM, Dong X, Ellsworth D, van Goethem D,
992 Jensen AM, Law BE, Loos R, Martins SCV, Norby RJ, Warren J, Weston D, Winter K (2014)
993 Asymmetrical effects of mesophyll conductance on fundamental photosynthetic parameters and

- 994 their relationships estimated from leaf gas exchange measurements. *Plant Cell Environ* 37:978–
995 994. <http://www.ncbi.nlm.nih.gov/pubmed/24117476> (18 November 2013, date last accessed).
- 996 Tanaka K, Kosugi Y, Nakamura A (2002) Impact of leaf physiological characteristics on seasonal variation
997 in CO₂, latent and sensible heat exchanges over a tree plantation. *Agric For Meteorol* 114:103–
998 122. <http://linkinghub.elsevier.com/retrieve/pii/S0168192302001284>
- 999 Terashima I, Hikosaka K (1995) Comparative ecophysiology of leaf and canopy photosynthesis. *Plant Cell*
1000 *Environ* 18:1111–1128.
- 1001 Tholen D, Ethier G, Genty B, Pepin S, Zhu X-G (2012) Variable mesophyll conductance revisited:
1002 theoretical background and experimental implications. *Plant Cell Environ* 35:2087–103.
1003 <http://www.ncbi.nlm.nih.gov/pubmed/22590996> (6 June 2014, date last accessed).
- 1004 Tognetti R, Longobucco A, Raschi A (1998) Vulnerability of xylem to embolism in relation to plant
1005 hydraulic resistance in *Quercus pubescens* and *Quercus ilex* co-occurring in a Mediterranean
1006 coppice stand in central Italy. *New Phytol* 139:437–447. <http://doi.wiley.com/10.1046/j.1469-8137.1998.00207.x> (17 September 2014, date last accessed).
- 1008 Tomás M, Flexas J, Copolovici L, Galmés J, Hallik L, Medrano H, Ribas-Carbó M, Tosens T, Vislap V,
1009 Niinemets Ü (2013) Importance of leaf anatomy in determining mesophyll diffusion conductance
1010 to CO₂ across species: quantitative limitations and scaling up by models. *J Exp Bot* 64:2269–81.
1011 <http://www.ncbi.nlm.nih.gov/pubmed/23564954> (19 March 2014, date last accessed).
- 1012 Tomás M, Medrano H, Brugnoli E, Escalona JM, Martorell S, Pou A, Ribas-Carbó M, Flexas J (2014)
1013 Variability of mesophyll conductance in grapevine cultivars under water stress conditions in
1014 relation to leaf anatomy and water use efficiency. *Aust J Grape Wine Res*:n/a–n/a.
1015 <http://doi.wiley.com/10.1111/ajgw.12069> (10 April 2014, date last accessed).
- 1016 Urli M, Porté AJ, Cochard H, Guengant Y, Burlett R, Delzon S (2013) Xylem embolism threshold for
1017 catastrophic hydraulic failure in angiosperm trees. *Tree Physiol* 33:672–683.
1018 <http://www.ncbi.nlm.nih.gov/pubmed/23658197> (6 August 2013, date last accessed).
- 1019 Valladares F, Niinemets Ü (2008) Shade Tolerance, a Key Plant Feature of Complex Nature and
1020 Consequences. *Annu Rev Ecol Evol Syst* 39:237–257.
1021 <http://www.annualreviews.org/doi/abs/10.1146/annurev.ecolsys.39.110707.173506> (21 March
1022 2014, date last accessed).
- 1023 Valladares F, Zaragoza-Castells J, Sánchez-Gómez D, Matesanz S, Alonso B, Portsmouth A, Delgado A,
1024 Atkin OK (2008) Is shade beneficial for mediterranean shrubs experiencing periods of extreme
1025 drought and late-winter frosts? *Ann Bot* 102:923–33.
1026 [http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2712400&tool=pmcentrez&renderty
1027 pe=abstract](http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2712400&tool=pmcentrez&rendertype=abstract) (28 March 2014, date last accessed).
- 1028 Vargas R, Sonnentag O, Abramowitz G, Carrara A, Chen JM, Ciais P, Correia A, Keenan TF, Kobayashi H,
1029 Ourcival J, Papale D, Pearson D, Pereira JS, Piao S, Rambal S, Baldocchi DD (2013) Drought
1030 Influences the Accuracy of Simulated Ecosystem Fluxes : A Model-Data Meta-analysis for
1031 Mediterranean Oak Woodlands. *Ecosystems* 16:749–764.
- 1032 Vaz M, Maroco J, Ribeiro N, Gazarini LC, Pereira JS, Chaves MM (2011) Leaf-level responses to light in
1033 two co-occurring *Quercus* (*Quercus ilex* and *Quercus suber*): leaf structure, chemical composition
1034 and photosynthesis. *Agrofor Syst* 82:173–181. <http://link.springer.com/10.1007/s10457-010-9343-6>
1035 (28 March 2014, date last accessed).

- 1036 Verhoeven A (2014) Sustained energy dissipation in winter evergreens. *New Phytol* 201:57–65.
1037 <http://doi.wiley.com/10.1111/nph.12466>
- 1038 Villar-Salvador P, Castro-Díez P, Pérez-Rontomé C, Montserrat-Martí G (1997) Stem xylem features in
1039 three *Quercus* (Fagaceae) species along a climatic gradient in NE Spain. *Trees* 12:90–96.
1040 <http://link.springer.com/10.1007/PL00009701> (30 August 2014, date last accessed).
- 1041 Vitale M, Manes F (2005) Role of changing environmental parameters in leaf gas exchange of *Arbutus*
1042 *unedo* L. assessed by field and laboratory measurements. *Photosynthetica* 43:99–106.
- 1043 Walker AP, Beckerman AP, Gu L, Kattge J, Cernusak L a., Domingues TF, Scales JC, Wohlfahrt G,
1044 Wullschlegel SD, Woodward FI (2014) The relationship of leaf photosynthetic traits - V_{cmax} and
1045 J_{max} - to leaf nitrogen, leaf phosphorus, and specific leaf area: a meta-analysis and modeling
1046 study. *Ecol Evol* 4:3218–3235. <http://doi.wiley.com/10.1002/ece3.1173> (28 July 2014, date last
1047 accessed).
- 1048 Wilson KB, Baldocchi DD, Hanson PJ, Ridge O (2001) Leaf age affects the seasonal pattern of
1049 photosynthetic capacity and net ecosystem exchange of carbon in a deciduous forest. *Plant, Cell*
1050 *Environ* 24:571–583.
- 1051 Yin X, Struik PC, Romero P, Harbinson J, Evers JB, Van Der Putten PEL, Vos J (2009) Using combined
1052 measurements of gas exchange and chlorophyll fluorescence to estimate parameters of a
1053 biochemical C photosynthesis model: a critical appraisal and a new integrated approach applied to
1054 leaves in a wheat (*Triticum aestivum*) canopy. *Plant Cell Environ* 32:448–64.
1055 <http://www.ncbi.nlm.nih.gov/pubmed/19183300> (9 November 2012, date last accessed).
- 1056 Zavala M a., Zea E (2004) Mechanisms maintaining biodiversity in Mediterranean pine-oak forests:
1057 insights from a spatial simulation model. *Plant Ecol* (formerly *Veg* 171:197–207.
1058 <http://link.springer.com/10.1023/B:VEGE.0000029387.15947.b7>
- 1059 Zheng Y, Zhong Z, Zhou JJ, Zhou H (2012) Evaluations of different leaf and canopy photosynthesis
1060 models: a case study with black locust (*Robinia pseudoacacia*) plantations on a loess plateau.
1061 *Pakistan J Bot* 44:531–539.
- 1062

1063 **Figure captions**

1064 **Fig. 1.** Environmental variables are presented for the day of the year (DOY) from January 2011 until
1065 February 2013; a) atmospheric vapour pressure deficit (VPD), b) rainfall in mm c) soil water content in
1066 $\text{cm}^3 \text{cm}^{-3}$ (gap in data is due to power cut), d) maximum and minimum temperatures in $^{\circ}\text{C}$ on the primary
1067 y-axes (in dark circles) and radiation in W m^{-2} (in light crosses, foreground) on the secondary y-axes.
1068 Field campaigns are indicated (acronyms of seasons are detailed in Tab. 2).

1069
1070 **Fig. 2.** Principal component analyses (PCA) for a) all trees species, leaf positions, and seasons, b) with
1071 differentiation between sunlit and shaded leaves, c) with differentiation between seasonal campaigns, and
1072 d) with differentiation between species. We used a subset of all data where both morphological and
1073 photosynthetic information was available. Fifteen parameters were used in the PCA: net assimilation rate
1074 (A_{net}), stomatal conductance (g_s), mesophyll conductance (g_m), maximum carboxylation rate ($V_{c,\text{max}}$),
1075 maximum electron transport rate (J_{max}), nonphotochemical quenching (NPQ), maximum quantum
1076 efficiency of PSII (F_v/F_m), leaf thickness (LT), leaf mass per area (LMA), leaf density (D), water content
1077 (WC), nitrogen content per leaf unit area (N_{area}), nitrogen content per leaf unit mass (N_{mass}), carbon
1078 content per leaf unit area (C_{area}), and carbon content per leaf unit mass (C_{mass}). The directions of the
1079 arrows indicate the higher levels of the parameters. Principal component (PC) 1 explains 37.2% of the
1080 variation, and PC 2 explained 20.4%. The ellipses are normal probability contour lines of 68% for the
1081 factors in b) leaf positions, c) seasons, and d) species.

1082
1083 **Fig. 3.** Line graphs depicting seasonal changes of a) maximum carboxylation rate ($V_{c,\text{max}}$), b) maximum
1084 electron-transport rate (J_{max}), and c) maximum quantum efficiency of PSII (F_v/F_m) for *Q. ilex*, *P.*
1085 *halepensis*, *A. unedo*, and *Q. pubescens* in sunlit (1) and shaded (2) leaves. Seasonal campaigns were
1086 conducted in spring 2011 (sp11), summer 2011 (su11), autumn 2011^a (au11 a), autumn 2011^b (au11b),
1087 winter 2012 (wi12), spring 2012 (sp12), summer 2012 (su12), and winter 2013 (wi13). Missing data
1088 points were due to limitations of labour and equipment. Vertical bars indicate standard errors of the
1089 means ($n = 3-5$).

1090
1091 **Fig. 4.** Line graphs depicting seasonal changes of a) net assimilation (A_{net}), b) stomatal conductance (g_s),
1092 and c) mesophyll conductance (g_m) for *Q. ilex*, *P. halepensis*, *A. unedo*, and *Q. pubescens* in sunlit (1) and
1093 shaded (2) leaves. Seasonal campaigns were conducted in spring 2011 (sp11), summer 2011 (su11),
1094 autumn 2011^a (au11 a), autumn 2011^b (au11b), winter 2012 (wi12), spring 2012 (sp12), summer 2012
1095 (su12), and winter 2013 (wi13). Missing data points were due to limitations of labour and equipment.
1096 Vertical bars indicate standard errors of the means ($n = 3-5$).

1097
1098 **Fig 5.** Seasonal changes of the relationships between a) net assimilation (A_{net}) and stomatal conductance
1099 (g_s), b) A_{net} and mesophyll conductance (g_m), and c) g_m and g_s in sunlit (1) and shaded (2) leaves. The
1100 regression lines represent the seasonal changes across species. For regression equations see Table S1-3.
1101 The relationships are shown as a thin solid line for spring 2011, short dashes for summer 2011, dots-

1102 dashes for autumn 2011^a), small dots for autumn 2011^b), dashes for winter 2012, large dots for spring
1103 2012, large dots-dashes for summer 2012, and a thick solid line for winter 2013. Statistical differences in
1104 the slopes between seasonal campaigns were tested by ANCOVAs.

1105

1106 **Fig 6.** Seasonal changes of the relationships between a) the maximum electron-transport rate (J_{\max}) and
1107 the maximum carboxylation rate ($V_{c,\max}$) and b) the electron-transport rate from chlorophyllic
1108 fluorescence (J_{amb}) and net assimilation (A_{net}) at ambient CO_2 concentrations and saturating light in sunlit
1109 (a) and shaded (b) leaves. The regression lines represent the seasonal changes across species. For
1110 regression equations see Table S4-5. The relationships are shown as a thin solid line for spring 2011,
1111 short dashes for summer 2011, dots-dashes for autumn 2011^a), small dots for autumn 2011^b), dashes for
1112 winter 2012, large dots for spring 2012, large dots-dashes for summer 2012, and a thick solid line for
1113 winter 2013.

1114

1115 **Fig 7.** Seasonal changes of the relationship for all species and leaf positions between a) mesophyll
1116 conductance (gm) and leaf mass per area (LMA). We used a subset of morphological and photosynthetic
1117 data. Non-linear regression lines of the form $y = x^{-b}$ were fitted to the data. The upper curve is for summer
1118 2012 ($b = 0.800$), the middle curve is for spring 2012 ($b = 0.953$) and the lower two overlaying curves are
1119 for autumn 2011a) ($b = 1.533$) and winter 2012 ($b = 1.486$).

1120

1121 **Supporting information**

1122 Additional supporting information may be found in the online version of this article.

1123 **Fig. S1.** Line graphs depicting seasonal changes of a) effective quantum efficiency of PSII
1124 (Φ_{PSII}), and b) nonphotochemical quenching (NPQ) for *Q. ilex*, *P. halepensis*, *A. unedo*, and *Q.*
1125 *pubescens* in sunlit (1) and shaded (2) leaves. Missing data points were due to limitations of
1126 labour and equipment. Vertical bars indicate standard errors of the means ($n = 3-5$).

1127 **Fig. S2.** Bar charts depicting seasonal changes of a) succulence (S), b) leaf density (D), c) water
1128 content (WC), and d) leaf thickness (LT) for *Q. ilex*, *P. halepensis*, *A. unedo*, and *Q. pubescens*
1129 in sunlit (1) and shaded (2) leaves. Error bars indicate standard errors of the means ($n = 3-5$).

1130 **Fig. S3.** Bar charts depicting seasonal changes of a) nitrogen per unit leaf area (N_{area}) and b)
1131 carbon per unit leaf mass (C_{mass}) for *Q. ilex*, *P. halepensis*, *A. unedo*, and *Q. pubescens* in sunlit
1132 (1) and shaded (2) leaves. Error bars indicate standard errors of the means ($n = 3-5$).

1133 **Table S1.** Regression equations and coefficients of determination (R^2) for A_{net}/g_s for sunlit and
1134 shaded leaves of *Q. ilex*, *P. halepensis*, *A. unedo*, and *Q. pubescens* in eight sampling
1135 campaigns

1136 **Table S2.** Regression equations and coefficients of determination (R^2) for A_{net}/g_m for sunlit and
1137 shaded leaves of *Q. ilex*, *P. halepensis*, *A. unedo*, and *Q. pubescens* in eight sampling
1138 campaigns.

1139 **Table S3.** Regression equations and coefficients of determination (R^2) for $J_{\text{max}}/V_{c,\text{max}}$ for sunlit
1140 and shaded leaves of *Q. ilex*, *P. halepensis*, *A. unedo*, and *Q. pubescens* in eight sampling
1141 campaigns.

1142 **Table S4.** Regression equations and coefficients of determination (R^2) for $J_{\text{amb}}/A_{\text{net}}$ for sunlit and
1143 shaded leaves of *Q. ilex*, *P. halepensis*, *A. unedo*, and *Q. pubescens* in eight sampling
1144 campaigns.

1145 **Table S5.** Regression equations and coefficients of determination (R^2) for g_m/g_s for sunlit and
1146 shaded leaves of *Q. ilex*, *P. halepensis*, *A. unedo*, and *Q. pubescens* in eight sampling
1147 campaigns.

1148 **Table S6.** Regression equations and coefficients of determination (R^2) for a) g_m/LMA and b)
1149 $V_{c,\text{max}}/N_{\text{area}}$ in four seasonal campaigns and for sunlit and shaded leaf positions for *Q. ilex*, *P.*
1150 *halepensis*, *A. unedo*, and *Q. pubescens*.

1151 **Note S1** Calculation of maximum quantum yield of PSII and nonphotochemical quenching

1152 **Note S2** Light experiments and estimation of day respiration

1153 *This is a pre-copyedited, author-produced PDF of an article accepted for publication in Tree*
1154 *physiology* following peer review. The version of record Sperlich, D., et al. "Seasonal variability of foliar
1155 photosynthetic and morphological traits and drought impacts in a Mediterranean mixed forest" in *Tree*
physiology, Vol. 35 issue 5 (May 2015), p. 501-520 is available online at: DOI 10.1093/treephys/tpv017