

1 Title: Seasonal Variations in Physical Fitness among Elementary School Children

2

3 Authors:

4 Dr. Claudia Augste

5 Institute of Sport Science, University of Augsburg, Germany

6 Universitätsstr. 3

7 86159 Augsburg

8 Germany

9 Phone: ++49 821 598 – 2814

10 Fax: ++49 821 598 – 2828

11 E-mail: claudia.augste@sport.uni-augsburg.de

12

13 Prof. Dr. Stefan Künzell

14 Institute of Sport Science, University of Augsburg, Germany

15 Universitätsstr. 3

16 86159 Augsburg

17 Germany

18 Phone: ++49 821 598 – 2824

19 Fax: ++49 821 598 – 2828

20 E-mail: stefan.kuenzell@sport.uni-augsburg.de

21

22 **Abstract**

23 There is empirical evidence that children's physical activity is dependent on climatic conditions. In
24 addition, a correlation between physical activity level and physical fitness has been identified. In this
25 longitudinal study, we investigate whether seasons have an influence on physical fitness. 145 German
26 elementary school children were tested every six months over a 2-year period. We used the German
27 Motor Test 6-18 to assess physical fitness. Performance in the 6-minute endurance run ($p < .001$),
28 bidirectional jumping ($p < .001$), the standing long jump ($p = .026$), the 20 m sprint ($p = .006$), and
29 the stand-and-reach task ($p = .017$) was significantly better in the summer than in the winter. There
30 were no differences in ability to balance backwards ($p = .120$); in the winter, the results for push-ups
31 ($p < .001$) and sit-ups ($p < .001$) were better than in the summer. We have shown that physical fitness
32 is significantly influenced by the season. Consequently, when children's fitness tests are used (e.g.,
33 as the basis for intervention programs, for classifying health risk groups or for recognising talent),
34 the season in which testing occurred should be reported and accounted for in future studies.

35 *Keywords:* physical fitness, primary school children, German Motor Test, gender, longitudinal
36 study

37

38 **Introduction**

39 Physical fitness is an important aspect of human development. Specific attention is paid to
40 the physical fitness of children. Physical fitness enables a healthy lifestyle and allows children to
41 participate in sports. Health care organisations are concerned about the physical fitness of children
42 in each new cohort, sport organisations want to maximise the abilities of youth, and school
43 administrators want to evaluate the effectiveness of sports lessons. Because of the importance of
44 physical fitness, many different tests have been developed over the last half-century (e.g., Morrow,
45 Zhu, Franks, Meredith, & Spain, 2009). To assess an individual's physical fitness level, the
46 individual's scores must be compared with scores from a norming sample. In this article, we
47 investigate whether the scores on physical fitness tests are influenced by the season in which they
48 are measured. We argue that the amount of physical activity (PA) that children perform depends on
49 the season; moreover, some parameters of physical fitness co-vary with PA. Thus, it might be
50 possible that physical fitness test scores – both for individuals and for the norming sample – are
51 dependent on the time of year in which they were assessed.

52 Several studies have shown that PA has an influence on physical fitness. Children with
53 higher PA levels have better total scores in physical fitness (Castelli & Valley, 2007; Fogelholm,
54 Stigman, Huisman, & Metsämuuronen, 2008; Hikiyara et al., 2007; Hume et al., 2008; Sasayama,
55 Okishima, Mizuuchi, & Adachi, 2009; Williams et al., 2008; Wrotniak, Epstein, Dorn, Jones, &
56 Kondilis, 2006). However, most authors do not only analyse total physical fitness scores; they also
57 distinguish between subdomains. Depending on the focus of the study, different aspects of physical
58 fitness are assessed. To focus on abilities, it is common to address the subdomains of aerobic fitness
59 or endurance, strength, speed, flexibility, and coordination. To focus on fundamental motor skills,
60 the common subdomains are locomotion and object-control. The most frequently studied
61 subdomain is aerobic fitness because it often serves as an indicator for children's health. In a
62 review, Boreham and Riddoch (2001) revealed that more active children display healthier

63 cardiovascular profiles. Several other recent studies consistently reported positive correlations
64 between PA and aerobic fitness (Castelli & Valley, 2007; Dencker et al., 2006; Fogelholm et al.,
65 2008; Hands, Larkin, Parker, Straker, & Perry, 2009; Hussey, Bell, Bennett, O'Dwyer, & Gormley,
66 2007; Kristensen et al., 2010; Magnusson, Sveinsson, Arngrimsson, & Johannsson, 2008; Sasayama
67 et al., 2009). Measuring locomotor skills as a subdomain also reveals consistent advantages for
68 more active children (Hume et al., 2008; Williams et al., 2008). Several major studies have reported
69 correlations between a higher PA level and jumping (standing long jump, 5-jump, jumping side-to-
70 side, jumping back-and-forth) or sprinting (Fogelholm et al., 2008; Hikahara et al., 2007; Sacchetti
71 et al., 2012; Sasayama et al., 2009).

72 The correlations between other subdomains and PA, however, are not always so clear. Some
73 researchers have found correlations between PA and object-control skills (Fogelholm et al., 2008;
74 Sasayama et al., 2009); others have not (Williams et al., 2008). Some have found correlations only
75 among boys (Cliff, Okely, Smith, & McKeen, 2009; Hume et al., 2008), or stronger correlations for
76 boys than for girls (Hikihara et al., 2007). In addition, for strength and muscle endurance, the
77 findings are ambiguous. Whereas some studies reveal strong correlations between PA levels and
78 tasks such as curl-ups, sit-ups, push-ups, or grip strength (Castelli & Valley, 2007; Fogelholm et al.,
79 2008), others found no correlations (Hands et al., 2009; Hikahara et al., 2007) or correlations among
80 boys but not among girls (Sasayama et al., 2009). One study even found a negative correlation with
81 upper body strength (Hands et al., 2009). Again, no clear correlation could be found between
82 flexibility and children's PA level; the results differed depending on age, gender, and assessment
83 method (Castelli & Valley, 2007; Fogelholm et al., 2008; Hands et al., 2009; Hikahara et al., 2007;
84 Sacchetti et al., 2012; Sasayama et al., 2009). Potential causes for the ambiguous results might be
85 the differing measurements in PA and physical fitness assessment across the studies. PA assessment
86 methods ranged from self-reports like 7-day physical activity recall (Castelli & Valley, 2007),
87 physical activity questionnaires (Sacchetti et al., 2012) or reports on frequency and duration of

88 sweating during organized and non-organized activity (Fogelholm et al., 2008), to objective
89 measures like pedometers (Castelli & Valley, 2007; Hands et al., 2009) or accelerometers (Hikahara
90 et al., 2007; Sasayama et al., 2009). Likewise, physical fitness parameters were measured with
91 different tests in almost every study. Nevertheless, we summarise that certain subdomains of
92 physical fitness, such as aerobic fitness and locomotor skills, are evidently correlated with PA.

93 PA is subject to seasonal fluctuations. Recently, research has focused on the influence of the
94 season and weather conditions on PA. In a systematic review, Tucker and Gilliland (2007) analysed
95 37 studies conducted between 1980 and 2006, including more than a quarter of a million
96 participants in total. According to their analysis, seasonality had an effect on PA level. For adults,
97 men and women had higher levels of PA in the summer than in the winter (Matthews et al., 2001).
98 In some countries, pre-schoolers and elementary school children participate in more PA in warmer
99 months than in colder months (North Canada: Carson, Spence, Cutumisu, Boule, & Edwards, 2010;
100 UK: Rowlands, Pilgrim, & Eston, 2009; Denmark: Kristensen et al., 2008; Norway: Kolle, Steene-
101 Johannessen, Andersen, & Anderssen, 2009). However, for adolescents, there was no clear
102 relationship between the seasons and PA (Denmark: Kristensen et al., 2008; Norway: Kolle et al.,
103 2009; Cyprus: Loucaides, Chedzoy, & Bennett, 2003; Canada: Bélanger, Gray-Donald, O'Loughlin,
104 Paradis, & Hanley, 2009).

105 Analysing PA by gender reveals that boys are usually more physically active than girls
106 (Baranowski, Thompson, DuRant, Baranowski, & Puhl, 1993; Beighle, Alderman, Morgan, & Le
107 Masurier, 2008; Dencker et al., 2006; Hume et al., 2008; Hussey et al., 2007; Kristensen et al.,
108 2008; Loucaides et al., 2003; Magnusson et al., 2008; Peiró-Velert, Devís-Devís, Beltrán-Carrillo,
109 & Fox, 2008; Rowlands & Eston, 2005; Sasayama et al., 2009). Interestingly, interactions between
110 gender and season have been found among pre-schoolers (Baranowski et al., 1993) and elementary
111 school children (Beighle et al. 2008), indicating that there is less of a difference between the PA
112 levels of boys and girls in the winter than in the spring and summer. In summary, Rowlands and

113 Hughes' (2006) suggestion to measure PA at multiple time points to account for seasonal variation
114 should certainly be followed.

115 Given the influence of seasons on PA and the correlations between PA and some
116 subdomains of physical fitness, it seems plausible that seasons have an effect on physical fitness
117 (see Figure 1). The objective of the present study is to test a hypothesis about the influence of the
118 seasons on tests of physical fitness.

119 **Methods**

120 To assess the influence of seasons on physical fitness, we conducted a panel study in which
121 we tested the participants' physical fitness over a two-year time period.

122 *Participants*

123 A total of 145 children participated in the tests. The sample size varied between 114 and
124 145, with an approximately equal number of boys and girls, over the five data collection points (see
125 Table 1). The average age of all participating children at the time of the first measurement in grade
126 2 was 7.9 years ($SD = \pm 0.59$); at the end of grade 4, the average age was 9.9 years ($SD = \pm 0.60$).
127 The children were recruited from three elementary schools in Augsburg, Germany. Two classes
128 from each school were randomly selected. The directors of the schools agreed to participate in the
129 study. Except for the parents of one child who was excluded from the study, informed consent was
130 obtained from the parents of all the selected children. All procedures were approved by the local
131 ethical commission for research on human participants.

132 *Measures*

133 **Physical fitness.** To test the physical fitness of the children, we used the German Motor Test
134 6-18, a standardised test battery (Bös, 2009), which is performed indoors. It consists of eight items
135 that test different subdomains of physical fitness. To assess children's flexibility, the stand-and-

136 reach test was performed. The children were asked to stand on a bench, bend forward with straight
137 legs, and reach down as far as possible. To measure speed, a 20-meter sprint was performed from an
138 upright start position. In German literature, coordination is differentiated into two subcategories,
139 coordination under time pressure and coordination in a task requiring precision (Bös, 2009). The
140 task used for the latter required the children to balance backwards on bars that were 6 cm, 4.5 cm,
141 and 3 cm in width. The children had two attempts to take a maximum of 8 steps on each bar without
142 touching the floor. Coordination under time pressure was measured with a bidirectional jumping
143 task. The children had to jump sideways within two marked fields (50 cm x 50 cm) as many times
144 as possible in 15 sec. To assess strength endurance, two tests were performed: sit-ups for 40 sec and
145 push-ups for 40 sec. The sit-ups were performed in a very common way, but the push-up task had
146 two special characteristics. First, the participants began each push-up lying flat on the floor with
147 both hands touching their back. Second, in the “up” position, the participants had to lift one hand
148 off the floor and touch the other hand. Using this method ensured that every push-up was completed
149 and that no shortcuts were taken. However, it also added a coordination aspect to the task. Power
150 was assessed with the standing long jump. The test for aerobic fitness in the German Motor Test is a
151 6-minute endurance run in which children run steadily around a volleyball court. Details about the
152 test battery have been described by Bös (2009). The objectivity (average over the items: .95) and
153 retest reliability (average over the items: .82) of the test battery were good, and the battery has been
154 validated for assessing endurance, strength, flexibility, coordination, and speed (Bös, 2009).

155 **Seasons.** Augsburg is a city in southern Germany located at 48° 22' N latitude and 10° 54' E
156 longitude. The climate is a hybrid of the humid Atlantic climate and the dry continental climate.
157 The region experiences four seasons with average temperatures of 8.0° C in spring (March to May),
158 16.7° C in summer (June to August), 8.5° C in fall (September to November), and -0.4° C in winter
159 (December to February) (Stadt Augsburg, 2010). For the study, seasonal variations between
160 summer and winter were tested. The measurements in the summer followed the months May/June

161 with mean temperatures of about 14 to 17° C, the measurements in the winter followed the months
162 November/December with mean temperatures of 4 to 0° C.

163 *Study Design and Procedures*

164 The data were collected every six months from June 2008 to June 2010. Of the five
165 measurements three were conducted in the summer (16 – 23 June 2008; 16 June – 07 July 2009; 14
166 – 17 June 2010) and two in the winter (02 – 18 December 2008; 26 November 2009 – 21 January
167 2010). The children from the six classes were tested during their regular physical education classes
168 in school, so they wore their usual gym clothing and shoes. Body weight and height were measured
169 without shoes. After five minutes of a general warm-up, the children were divided into groups of
170 three, and they completed seven tests in random order. As defined in the test manual (Bös, 2009),
171 the 6-minute endurance run was always performed as the last test in groups of approximately 12
172 children. All of the data were recorded by trained university staff members.

173 *Data Analysis*

174 Test results were recorded for every test item (see Figure 2). To analyse the differences
175 between the mean test scores at the different points in time and the interaction with gender, a 2-
176 factorial analysis of variance (ANOVA) was calculated with a 5-level within-subject factor (time)
177 and a 2-level between-subjects factor (gender: male, female). To compare physical fitness between
178 different seasons, the data were Z-transformed using the norming sample with analogous age and
179 gender (see Figure 3). This procedure eliminates the improvements in fitness associated with gender
180 and increasing age. To determine the difference in physical fitness between summer and winter, the
181 Z-values of the test items for each of the three measures in summer and for the two measures in
182 winter were summed, and then the averages of the two seasons were calculated. This procedure was
183 chosen, because the aim of the study was not to find differences between single time points of
184 measurement, but across the summer and winter season in general. A two-factorial ANOVA with a

185 two-level within-subject factor (season: summer, winter) and a two-level between-subjects factor
186 (gender: male, female) was conducted to assess whether there were significant differences between
187 physical fitness in summer and winter and to assess the interaction between the season and gender.
188 A significance level of 0.05 was applied for all statistical tests.

189 **Results**

190 Descriptive data for the sample are provided in Table 1. In the investigation period, the
191 children's average height increased by 11 cm, and their weight increased by 8 kg. Between 19% and
192 25% of the children were classified as overweight during the time of measurement, according to the
193 German norm (Kromeyer-Hauschild et al., 2001). More than half of the children regularly practiced
194 with sports teams or clubs.

195 *Development of physical fitness*

196 The absolute values of the 8 test items over the 5 measurements are shown in Figures 2a-h
197 for boys, girls, and both. The distribution of all 8 test items was Gaussian at every time of
198 measurement (Kolmogorov-Smirnov Statistic). The stand-and-reach test was the only test item
199 without a significant increase in performance over the 2 years (see Figure 2a), ($F(4, 74) = 2.02, p =$
200 $.100, \eta^2 = .10$), indicating that flexibility was quite stable throughout elementary school. Except in
201 the first measurement, the girls had significantly higher flexibility scores than the boys. For push-
202 ups and sit-ups, performance improved significantly over the two years ($F(4, 74) = 56.09, p < .001,$
203 $\eta^2 = .75$ for push-ups and $F(4, 73) = 20.49, p < .001, \eta^2 = .53$ for sit-ups). Although there were
204 almost no differences between the boys and the girls for push-ups (see Figure 2b), the boys
205 outperformed the girls in sit-ups in winter 2008, $t(105) = 2.143, p = .034, CI [0.24, 6.03]$, and in
206 winter 2009, $t(110) = 2.635, p = .010, CI [0.82, 5.81]$ (see Figure 2c). For the balancing task, we
207 found significant positive development, $F(4, 74) = 14.26, p < .001, \eta^2 = .44$, with an interaction
208 between time and gender, $F(4, 74) = 2.51, p = .049, \eta^2 = .12$. The girls performed significantly

209 better than the boys in grade 2 (summer 2008: $t(114) = -2.93, p = .004, CI [-8.31, -1.60]$) and grade
210 3 (winter 2008: $t(105) = -2.02, p = .046, CI [-6.74, -0.67]$; summer 2009: $t(112) = -3.56, p = .001,$
211 $CI [-8.90, -2.53]$), but there was no significant difference between the genders in grade 4 (see Figure
212 2d). The children had the most improvement in bidirectional jumping (see Figure 2e, $F(4, 73) =$
213 $107.234, p < .001, \eta^2 = .855$), over the two years. On this task, the performance of the boys and the
214 girls reached almost identical values over the years in elementary school. Significant improvements
215 also occurred in the standing long jump (see Figure 2f, $F(4, 72) = 7.162, p < .001, \eta^2 = .285$), with
216 the boys outperforming the girls in grade 3 (winter 2008: $t(105) = 2.06; p = .042, CI [0.00, 0.15]$;
217 summer 2009: $t(112) = 2.08; p = .040, CI [0.00, 0.14]$). Sprint time to cover 20 m decreased
218 significantly over the two years (see Figure 2g, $F(4, 73) = 10.825, p < .001, \eta^2 = .372$). The
219 development of speed was very similar between the boys and the girls, but by the end of grade 4,
220 the boys were significantly ahead of the girls, $t(107) = 2.31; p = .023, CI [-0.35, -0.03]$). The
221 distance covered in the 6-minute endurance run increased significantly (see Figure 2h, $F(4, 70) =$
222 $10.426, p < .001, \eta^2 = .373$), and there were no interactions with gender.

223 *Seasonal differences*

224 For 5 of the 8 test items, the average summer results were significantly higher than the
225 average winter results. The test with the largest summer advantage was the 6-minute endurance run,
226 ($F(1, 80) = 18.27, p < .001, \eta^2 = .19$). In this test, we found a significant interaction between the
227 season and gender, $F(1, 80) = 5.22, p = .025, \eta^2 = .06$. Furthermore, in the summer, the children
228 performed significantly better on the stand-and-reach test ($F(1, 85) = 5.91, p = .017, \eta^2 = .07$),
229 bidirectional jumping, ($F(1, 84) = 16.33, p < .001, \eta^2 = .16$), the standing long jump, ($F(1, 83) =$
230 $5.14, p = .026, \eta^2 = .06$), and the 20 m sprint, ($F(1, 85) = 7.91, p = .006, \eta^2 = .09$). The two tasks
231 requiring strength endurance, push-ups, and sit-ups, were the only tests that were performed
232 significantly better in winter (push-ups: $F(1, 85) = 49.96, p < .001, \eta^2 = .37$; sit-ups: $F(1, 85) =$

233 26.14, $p < .001$, $\eta^2 = .24$). Season had no significant influence on balancing performance, $F(1, 86)$
234 $= 2.47$, $p = .120$, $\eta^2 = .03$. To summarise, seasonal variations in motor test performance were
235 identified in seven out of eight test items. There were better results for push-ups and sit-ups in the
236 winter and better results for the stand-and-reach, bidirectional jumping, the standing long jump, the
237 20 m sprint, and the 6-min endurance run in the summer (see Figure 3).

238 **Discussion**

239 *Fitness subdomains*

240 For some tasks, the Z-scores that the participants obtained were far better than the scores by
241 the norming sample. This difference may have occurred because the participants became more
242 proficient in the coordination tasks over the five measurements. For the strength, endurance, and
243 speed tasks, the performances were similar to the German average. Regarding the fitness test results
244 for the summer and winter measures, we found different effects for the different subdomains of
245 fitness.

246 **Aerobic fitness.** A review of the literature suggests that there is a positive correlation
247 between PA and aerobic fitness (Castelli & Valley, 2007; Dencker et al., 2006; Fogelholm et al.,
248 2008; Hands, et al., 2009; Kristensen, et al., 2010; Sasayama, et al., 2009). Therefore, we
249 hypothesised that the children's performances on aerobic fitness tasks would be better in the
250 summer, when children's PA is higher. In accordance with our expectations, the participants in this
251 study had significantly better test scores on the 6-min endurance run in the summer compared to the
252 winter. Concerning gender differences, our results showed that the seasonal effect for this task was
253 even clearer for girls than for boys, although Hussey et al. (2007) did not find a correlation between
254 the endurance test and PA in girls.

255 **Speed.** A summer advantage was also apparent in the speed task. Again, this finding was
256 consistent with our expectation, even though the correlation between PA and speed has not been

257 thoroughly explored. Whereas one comprehensive study on Australian 9-12-year-olds revealed
258 correlations between vigorous PA and the sprint run for boys and girls (Hume et al., 2008), other
259 studies have demonstrated positive correlations between greater PA and sprint performance among
260 boys but not among girls (Sacchetti et al., 2012; Sasayama et al., 2009). In our study, the negative
261 influence of winter was very similar for boys and girls.

262 **Power.** Most authors found positive correlations between PA and tasks requiring power,
263 such as the standing long jump (Hikihara et al., 2007), the vertical jump (Hume et al., 2008) or the
264 5-jump (Fogelholm et al., 2008). Our finding that the season significantly influenced performance
265 on the standing long jump is consistent with the existing literature. Again, both boys and girls
266 performed better in the summer than in the winter, although some authors have found that the
267 positive influence of PA only affects standing long jump performance for boys (Sacchetti et al.,
268 2012).

269 **Coordination under time-pressure.** Bidirectional jumping is a task that requires
270 coordination under time pressure. Similar tasks, such as the side-step task or back-and-forth
271 jumping, were significantly correlated with PA for both boys and girls (Fogelholm et al., 2008;
272 Hikihara et al., 2007). In our study, the results for bidirectional jumping were significantly better in
273 the summer than in the winter. The girls lost their initial advantage over the boys between grade 2
274 and grade 4. For this task, the negative influence of winter seemed to be more relevant for the girls.

275 **Coordination requiring precision.** Balancing backwards is a task that tests children's
276 ability to achieve coordination with precision (Bös, 2009). In some German studies (Schmidt,
277 2008), correlations have been shown between children's participation in sports clubs and their
278 balancing performance. To date, however, no international studies have addressed balancing tasks
279 and PA. From a logical point of view, it could be assumed that longer outdoor play times would
280 improve performance in balancing. However, the season did not have a significant influence on
281 performing this task.

282 **Strength endurance.** The two tasks testing strength endurance in the present study were
283 push-ups and sit-ups performed for 40 seconds each. As described above, push-ups were performed
284 with special characteristics, adding a coordination aspect. Thus, the sample in this study had very
285 high Z-values for this task compared to the norming sample because a learning effect was evident
286 for this special push-up task over the five points of measurement. Nevertheless, it is surprising that
287 in the comparison of the seasons, children performed significantly better in winter on both strength
288 endurance tasks (push-ups and sit-ups). Previous studies investigating on the effects of PA on these
289 types of tasks have revealed conflicting results. Among children, Castelli and Valley (2007) showed
290 that there were correlations between PA and curl-ups and push-ups, whereas Hikiyara et al. (2007)
291 found no correlation with sit-ups. Sasayama et al. (2009) showed a significant correlation between
292 PA and sit-ups for boys but not for girls. In adolescents, Fogelholm et al. (2008) also found a
293 correlation with sit-ups. Hands et al. (2009), however, revealed contrary results. PA was not
294 correlated with abdominal muscle endurance (curl ups), and high levels of PA were negatively
295 correlated with upper body strength (chest pass). Although the findings are inconsistent, the
296 considerable improvements in the winter measures in our study can hardly be explained. A closer
297 look at the data showed that it was mainly the sports club members who improved in the winter,
298 which may be one plausible explanation for this phenomenon.

299 **Flexibility.** The flexibility of the elementary school children on the stand-and-reach task
300 was higher in the summer compared to the winter. This result is in alignment with the findings of
301 other authors. For age groups that were similar to the group in our study, PA level has been found to
302 be related to flexibility on the sit-and-reach task (Castelli & Valley, 2007; Hikiyara et al., 2007).
303 Some authors, however, have found a correlation only for boys but not for girls (Sasayama et al.,
304 2009), no correlation (Sacchetti et al., 2012) or even a negative correlation (Hands et al., 2009). The
305 latter was contributed to the lower muscle tone of inactive people.

306 We expected bigger differences between the boys and the girls in summer because of some
307 findings among elementary school children that have shown that there were fewer differences in PA
308 levels of boys and girls in the winter compared to spring and summer (Beighle et al., 2008). This
309 did not occur in our study. The only task with an interaction between season and gender was the 6-
310 min endurance run, but the difference between the boys and the girls was lower in the summer than
311 in the winter.

312 *Caveats*

313 Except for strength endurance and balancing, the physical fitness of the tested children in the
314 summer was better than their physical fitness in the winter. Most of the results were in accordance
315 with our hypotheses. However, the scope of the study still remains to be defined. It seems plausible
316 that the seasonal differences in physical fitness are related to climatic conditions. Presumably, this
317 constrains the scope of our study to areas of latitude in which a seasonal difference between warm
318 summers and cold winters can be observed. Moreover, the observed interactions between gender
319 and season in some of the subdomains of fitness are presumably dependent on cultural concepts of
320 boys and girls playing outdoors.

321 **Conclusions**

322 We have shown that physical fitness depends on the season. However, seasonal influences
323 do not affect all subdomains of physical fitness in the same way. Previous research has shown that
324 higher levels of PA did not result in higher performances in all physical fitness subdomains. Thus,
325 when comparing seasonal influences, researchers must always take into account which physical
326 fitness subdomain is being examined.

327 When child fitness is compared to norming standards, the season in which the children have
328 been tested should be taken into account. For example, because aerobic fitness is an indicator for
329 health risks, the children would be classified into the risk group because of their conspicuous test

330 result in the endurance run in the winter. Researchers developing test standard values should either
331 consider measures of all seasons or, at least, document when the norm values were collected. The
332 limits used for declaring the needs of intervention programs, for classifying children into health risk
333 groups and for selecting talent must be adapted. When different cohorts are involved in
334 comparisons, the season when the measures were taken should be considered. Otherwise, the
335 differences may be due to seasonal effects rather than to secular trends.

336 The season seems to have an influence, not only on PA of elementary school children but
337 also on their physical fitness. This fact should be the focus of future studies.

338

339

References

- 340 Baranowski, T., Thompson, W. O., DuRant, R. H., Baranowski, J., & Puhl, J. (1993). Observations
341 on physical activity in physical locations: age, gender, ethnicity, and month effects.
342 *Research Quarterly for Exercise and Sport*, *64*, 127–133.
- 343 Beighle, A., Alderman, B., Morgan, C. F., & Le Masurier, G. (2008). Seasonality in children's
344 pedometer-measured PA levels. *Research Quarterly for Exercise and Sport*, *79*, 256–260.
- 345 Bélanger, M., Gray-Donald, K., O'Loughlin, J., Paradis, G., & Hanley, J. (2009). Influence of
346 weather conditions and season on physical activity in adolescents. *Annals of epidemiology*,
347 *19*, 180–186. doi: 10.1016/j.annepidem.2008.12.008
- 348 Bös, K. (2009). *Deutscher Motorik Test 6-18 (DMT 6-18)* [German Motor Test 6-18]. Hamburg:
349 Czwalina.
- 350 Boreham, C., & Riddoch, C. (2001). The physical activity, fitness and health of children. *Journal of*
351 *Sports Sciences*, *19*, 915–929. doi: 10.1080/026404101317108426
- 352 Carson, V., Spence, J. C., Cutumisu, N., Boule, N., & Edwards, J. (2010). Seasonal variation in
353 physical activity among preschool children in a northern Canadian city. *Research Quarterly*
354 *for Exercise and Sport*, *81*, 392–399.
- 355 Castelli, D. M., & Valley, J. A. (2007). Chapter 3: The relationship of physical fitness and motor
356 competence to physical activity. *Journal of Teaching in Physical Education*, *26*, 358–374.
- 357 Cliff, D. P., Okely, A. D., Smith, L. M., & McKeen, K. (2009). Relationships between fundamental
358 movement skills and objectively measured physical activity in preschool children. *Pediatric*
359 *Exercise Science*, *21*, 436-449.
- 360 Dencker, M., Thorsson, O., Karlsson, M., Lindén, C., Svensson, J., & Wollmer, P., Anderson, L. B.
361 (2006). Daily physical activity and its relation to aerobic fitness in children aged 8-11 years.
362 *European Journal of Applied Physiology*, *96*, 587–592. doi: 10.1007/s00421-005-0117-1

- 363 Fogelholm, M., Stigman, S., Huisman, T., & Metsämuuronen, J. (2008). Physical fitness in
364 adolescents with normal weight and overweight. *Scandinavian Journal of Medicine &*
365 *Science in Sports, 18*, 162–170. doi: 10.1111/j.1600-0838.2007.00685.x
- 366 Hands, B., Larkin, D., Parker, H., Straker, L., & Perry, M. (2009). The relationship among physical
367 activity, motor competence and health-related fitness in 14-year-old adolescents.
368 *Scandinavian Journal of Medicine & Science in Sports, 19*, 655–663. doi: 10.1111/j.1600-
369 0838.2008.00847.x
- 370 Hikiyama, Y., Sasayama, K., Okishima, K., Mizuuchi, H., Yoshitake, Y., & Adachi, M., &
371 Takamatsu, K. (2007). The difference of relationships between physical activity variables
372 and physical fitness in children and adolescents: With special reference to amount and
373 intensity of physical activity. *Japanese Journal of Physical Fitness and Sport, 56*, 327–338.
- 374 Hume, C., Okely, A., Bagley, S., Telford A., Booth, M., Crawford, D., & Salmon, J. (2008). Does
375 weight status influence associations between children's fundamental movement skills and
376 physical activity? *Research Quarterly for Exercise and Sport, 79*, 158–165.
- 377 Hussey, J., Bell, C., Bennett, K., O'Dwyer, J., & Gormley, J. (2007). Relationship between the
378 intensity of physical activity, inactivity, cardiorespiratory fitness and body composition in 7-
379 10-year-old Dublin children. *British Journal of Sports Medicine, 41*, 311–316. doi:
380 10.1136/bjism.2006.032045
- 381 Kolle, E., Steene-Johannessen, J., Andersen, L. B., & Anderssen, S. A. (2009). Seasonal variation
382 in objectively assessed physical activity among children and adolescents in Norway: a cross-
383 sectional study. *The International Journal of Behavioral Nutrition and Physical Activity,*
384 *6*(36). doi:10.1186/1479-5868-6-36
- 385 Kristensen, P., Møller, N. C., Korsholm, L., Kolle, E., Wedderkopp, N., Froberg K., & Andersen, L.
386 B. (2010). The association between aerobic fitness and physical activity in children and

- 387 adolescents: the European youth heart study. *European Journal of Applied Physiology*, *110*,
388 267–275. doi: 10.1007/s00421-010-1491-x
- 389 Kristensen, P., Korsholm, L., Møller, N. C., Wedderkopp, N., Andersen, L. B., & Froberg K.
390 (2008). Sources of variation in habitual physical activity of children and adolescents: the
391 European youth heart study. *Scandinavian Journal of Medicine & Science in Sports*, *18*,
392 298–308. doi: 10.1111/j.1600-0838.2007.00668.x
- 393 Kromeyer-Hauschild, K., Wabitsch, M., Kunze, D., Geller, F., Geiß, H., Hesse, V., ...Hebebrand, J.
394 (2001). Perzentile für den Body-Mass-Index für das Kindes- und Jugendalter unter
395 Heranziehung verschiedener deutscher Stichproben [Percentiles for the body-mass-index for
396 children and adolescents consulting different German samples]. *Monatsschrift*
397 *Kinderheilkunde*, *149*, 807-818. doi:10.1007/s001120170107
- 398 Loucaides, C. A., Chedzoy, S. M., & Bennett, N. (2003). Pedometer-assessed physical (ambulatory)
399 activity in Cypriot children. *European Physical Education Review*, *9*, 43–55. doi:
400 10.1177/1356336X03009001179
- 401 Magnusson, K., Sveinsson, T., Arngrimsson, S. A., & Johannsson, E. (2008). Predictors of fatness
402 and physical fitness in nine-year-old Icelandic school children. *International Journal of*
403 *Pediatric Obesity*, *3*, 217–225. doi: 10.1080/17477160802169482
- 404 Matthews, C. E., Freedson, P. S., Hebert, J. R., Stanek, E. J., Merriam, P. A., & Rosal, M. C., et al.
405 (2001). Seasonal variation in household, occupational, and leisure time physical activity:
406 longitudinal analyses from the seasonal variation of Blood Cholesterol Study 108. *American*
407 *Journal of Epidemiology*, *153*, 172–183. doi: 10.1093/aje/153.2.172
- 408 Morrow, J. R., Zhu, W., Franks, B. D., Meredith, M. D., & Spain, C. (2009). 1958-2008: 50 Years
409 of Youth Fitness Tests in the United States. *Research Quarterly for Exercise and Sport*, *80*,
410 1–11.

- 411 Peiró-Velert, C., Devís-Devís, J., Beltrán-Carrillo, V., & Fox, K. (2008). Variability of Spanish
412 adolescents' physical activity patterns by seasonality, day of the week and demographic
413 factors. *European Journal of Sport Science*, 8, 163–171. doi: 10.1080/17461390802020868
- 414 Rowlands, A. V., & Eston, R. G. (2005). Comparison of accelerometer and pedometer measures of
415 physical activity in boys and girls, ages 8–10 years. *Research Quarterly for Exercise and*
416 *Sport*, 76, 251–257.
- 417 Rowlands, A. V., & Hughes, D. R. (2006). Variability of physical activity patterns by type of day
418 and season in 8-10-year-old boys. *Research Quarterly for Exercise and Sport*, 77, 391–395.
- 419 Rowlands, A. V., Pilgrim, E., & Eston, R. (2009). Seasonal changes in children's physical activity:
420 An examination of group changes, intra-individual variability and consistency in activity
421 pattern across season. *Annals of Human Biology*, 36, 363–378. doi:
422 10.1080/03014460902824220
- 423 Sacchetti, R., Cecilian, A., Garulli, A., Masotti, A., Poletti, G., Beltrami, P., & Leoni, E. (2012).
424 Physical fitness of primary school children in relation to overweight prevalence and physical
425 activity habits. *Journal of Sports Sciences*, 30, 633-640. doi:
426 10.1080/02640414.2012.661070
- 427 Sasayama, K., Okishima, K., Mizuuchi, H., & Adachi, M. (2009). Relationship of daily physical
428 activity and fitness in elementary school children. *Japanese Journal of Physical Fitness and*
429 *Sport*, 58, 259–304.
- 430 Schmidt, W. (2008). Zur Bedeutung des Sportvereins im Kindesalter [The importance of sports
431 clubs for children]. In W. Schmidt (Ed.), *Zweiter Deutscher Kinder- und Jugendsportbericht*
432 (pp. 373-390). Schorndorf: Hofmann.
- 433 Stadt Augsburg (2010). *Statistisches Jahrbuch der Stadt Augsburg 2009* [Statistical year book of
434 the city of Augsburg 2009]. Retrieved from [http://www.augsburg.de/fileadmin/www/dat/](http://www.augsburg.de/fileadmin/www/dat/01au/statistik/Jahrbuch/2009_Gesamt/Jahrbuch2009_Gesamtausgabe.pdf)
435 [01au/statistik/Jahrbuch/2009_Gesamt/Jahrbuch2009_Gesamtausgabe.pdf](http://www.augsburg.de/fileadmin/www/dat/01au/statistik/Jahrbuch/2009_Gesamt/Jahrbuch2009_Gesamtausgabe.pdf)

- 436 Tucker, P., & Gilliland, J. (2007). The effect of season and weather on physical activity: a
437 systematic review. *Public Health, 121*, 909–922. doi: 10.1016/j.puhe.2007.04.009
- 438 Williams, H. G., Pfeiffer K. A., O'Neill, J. R., Dowda, M., McIver, K. L., Brown, W. H., & Pate, R.
439 P. (2008). Motor skill performance and physical activity in preschool children. *Obesity, 16*,
440 1421–1426. doi:10.1038/oby.2008.214
- 441 Wrotniak, B. H., Epstein, L. H., Dorn, J. M., Jones, K. E., & Kondilis V. A. (2006). The
442 relationship between motor proficiency and physical activity in children. *Pediatrics, 118*,
443 1758–1765. doi:10.1542/peds.2006-0742
- 444

445 Table 1. *Demographic Information.*

Variable	T1		T2		T3		T4		T5	
Total # of participants	125		130		134		145		114	
Boys	63		66		69		76		58	
Girls	62		64		65		69		56	
	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>
Age years	7.9	.59	8.3	.56	9.0	.59	9.4	.53	9.9	.60
Height cm	131	5.9	134	6.2	137	6.4	140	6.7	142	7.0
Weight kg	29.5	6.7	31.5	7.6	33.6	8.0	36.2	9.3	37.9	9.4
BMI kg/m ²	17.1	3.0	17.3	3.1	17.7	3.2	18.3	3.7	18.6	3.8
underweight %	5.2		8.5		5.2		5.3		4.4	
normal weight %	71.6		70.8		75.7		73.5		70.2	
overweight %	23.3		20.8		19.1		21.2		25.4	
Sports club members %	62		59		57		63		65	

446

447





Figure 1. Correlations (continuous lines) and supposed correlations (dashed line) between season, physical activity and physical fitness.

448

449



450 Figure 2a)-h). Scores on the test items of boys and girls over the 5 times of measurement. $n_{\text{girls}} =$

451 42; a), b), d) $n_{\text{boys}} = 37$; c), e), g) $n_{\text{boys}} = 36$; f) $n_{\text{boys}} = 35$; h) $n_{\text{boys}} = 33$




Figure 3. Z-values of the single test items averaged over the season and compared across season. *** $p < .001$, ** $p < .01$, * $p < .05$.