


SCIENTIFIC REPORTS

OPEN

Sedum alfredii SaNramp6 Metal Transporter Contributes to Cadmium Accumulation in Transgenic *Arabidopsis thaliana*

Shuangshuang Chen^{1,2}, Xiaojiao Han^{1,2}, Jie Fang^{1,2,3}, Zhuchou Lu⁴, Wenmin Qiu^{1,2}, Mingying Liu^{1,2}, Jian Sang^{1,2}, Jing Jiang^{1,2} & Renying Zhuo^{1,2} 

The plant natural resistance-associated macrophage protein (Nramp) family plays an important role in tolerance to heavy metal stress. However, few *Nramps* have been functionally characterized in the heavy metal-accumulating plant *Sedum alfredii*. Here, *Nramp6* was cloned and identified from *S. alfredii* and its function analyzed in transgenic *Arabidopsis thaliana*. *SaNramp6* cDNA contains an open reading frame of 1, 638 bp encoding 545 amino acids. *SaNramp6*'s expression can be induced by cadmium (Cd) stress, and, after treatment, it peaked at one week and 12 h in the roots and leaves, respectively. *SaNramp6* localized to the plasma membrane in protoplasts isolated from *A. thaliana*, *Nicotiana benthamiana* lower leaf and onion (*Allium cepa*) epidermal cells. The heterologous expression of *SaNramp6* in the $\Delta ycf1$ yeast mutant increased the Cd content in yeast cells. *SaNramp6* also rescued the low Cd accumulation of the *A. thaliana nramp1* mutant. Transgenic *A. thaliana* expressing *SaNramp6* exhibited high Cd accumulation levels, as determined by a statistical analysis of the Cd concentration, translocation factors and net Cd²⁺ fluxes under Cd stress. Thus, *SaNramp6* may play a significant role in improving Cd accumulation, and the gene may be useful for the biotechnological development of transgenic plants for phytoremediation.

A well-balanced cellular concentration of essential metals such as iron (Fe), copper (Cu) and manganese (Mn), plays a fundamental role in the normal growth and development of plants¹. However, the absorption of heavy metals such as lead (Pb), cadmium (Cd) and arsenic (As), can upset the normal metabolism within plant cells and also cause damage to human and animal health through the cumulative effects of the food chain. For example, Cd is a carcinogenic factor closely related to the generation of breast and kidney cancer^{2,3}, and high levels of Pb toxicity can lead to irreversible damage to the central nervous system⁴. The ever-increasing worldwide contamination of soil and water by heavy metals is a problem that demands a prompt solution⁵. Phytoremediation is presently regarded as an eco-friendly and cost-effective strategy to clean heavy metal-polluted soils with the help of hyperaccumulating plants⁶. Among the more than 400 naturally hyperaccumulating plants, an ecotype of *Sedum alfredii* that co-hyperaccumulates Cd, Zn and Pb was first found in China^{7,8}. Previous physiological studies suggested that this ecotype is a promising hyper-accumulator for the decontamination of polluted soils, because it can accumulate up to nine g of Cd per kg of leaf dry weight (DW)^{9–11}. However, the detailed molecular mechanism underlying its hyperaccumulation and tolerance of heavy metals is still unclear. Taking advantage of this genetic resource for the breeding of future phytoremediation-associated plants requires a functional analysis of potential heavy metal-responsive genes in the hyperaccumulating ecotype of *S. alfredii*.

Metal transporters are essential for the maintenance of appropriate metal ions concentrations within different cellular compartments^{12,13}. Among the identified metal transporters, natural resistance-associated macrophage protein genes (*Nramps*) are considered to play potentially important roles mediating metal ion homeostasis at

¹State Key Laboratory of Forest Genetics and Breeding, Xiangshan Road, Beijing, 100091, P.R. China. ²Key Laboratory of Tree Breeding of Zhejiang Province, The Research Institute of Subtropical of Forestry, Chinese Academy of Forestry, Hangzhou, 311400, China. ³Chemical Biology Center, Lishui Institute of Agricultural Sciences, Lishui, Zhejiang Province, 323000, China. ⁴Biotechnology Research Center of China, Three Gorges University, Yichang, Hubei, 443002, China. Shuangshuang Chen, Xiaojiao Han and Jie Fang contributed equally to this work. Correspondence and requests for materials should be addressed to R.Z. (email: zhuory@gmail.com)

multiple cellular levels in plants. First cloned in mouse, *Nramp* gene family members are relatively evolutionarily conserved throughout organisms, including plants, animals, yeast and bacteria¹⁴. The *Nramp* genes comprise a small family represented by six members in *Arabidopsis thaliana*¹⁵, 12 members in rice (*Oryza sativa*; <http://www.ncbi.nlm.nih.gov/gene/?term=Nramp+Oryza+sativa>), eight members in soybean (*Glycine max*; <http://www.phytozome.net/soybean>) and six members in poplar (*Populus trichocarpa*)¹⁶. Several NRAMP members have been experimentally characterized in *A. thaliana* and are involved in the uptake, intracellular transport, translocation and detoxification of metals^{14,17,18}. They are all membrane spanning proteins, with the 10–12 hydrophobic transmembrane domains characteristic of metal transporters¹⁹. When overexpressed in yeast, *AtNramp1*, *AtNramp3* and *AtNramp4* show high affinities for Fe, Mn and Cd, whereas *AtNramp6* can transport Cd, but not Fe or Mn^{20–23}. In rice, *OsNramp1* shows transport activity for Cd and Fe, but not Mn. *OsNramp4* is the first transporter identified for the trivalent aluminium ion, and the knockout of *OsNramp5* results in a significantly reduced Cd uptake^{18,24,25}. *Nramp* genes have also been cloned and characterized from other plants, such as tomato (*Solanum lycopersicum*)²⁶, soybean²⁷ and some metal-hyperaccumulating species. A better understanding of the mechanisms used by metal transporters will provide insights into the detoxification and accumulation of toxic heavy metals in plants.

Although *Nramp* genes have been cloned and analyzed in other plant species, few studies have been reported regarding *Nramps* in the hyperaccumulating ecotype of *S. alfredii*. The transcriptome of *S. alfredii* under Cd stress indicated that an *Nramp* gene was greatly up-regulated after CdCl₂ treatment²⁸. The gene has an 80% homology with *AtNramp6*. Here, we described the isolation and characterization of the *Nramp* gene *SaNramp6* from *S. alfredii*. A subcellular localization analysis indicated that *SaNramp6* is a plasma membrane transporter. Moreover, the overexpression of *SaNramp6* in *A. thaliana* increased the uptake and accumulation of Cd. Thus, *SaNramp6* may be a potentially important heavy metal-responsive gene that could be useful for phytoremediation. This work will aid in understanding heavy metal hyperaccumulation and tolerance in *S. alfredii*.

Results

Isolation and sequence analysis of *SaNramp6*. To identify the function of *SaNramp6* from *S. alfredii*, a full-length cDNA sequence of 2,055 nucleotides was isolated, comprising a 1,638-bp open reading frame, and 95-bp 5′ and 322-bp 3′-untranslated regions. The specific primers *SaNramp6*-F and *SaNramp6*-R were used to amplify the sequence of the *SaNramp6* from genomic DNA to investigate the genomic structure of *SaNramp6*. The genomic sequence spanned 3,587 bp including 10 introns and 11 exons (Fig. 1a). A sequence comparison revealed that *SaNramp6* is similar to members of group I from *A. thaliana* (Fig. 1b).

The predicted protein encoded by *SaNramp6* contained 545 amino acid residues with a putative molecular weight of 58.44 kD and an isoelectric point of 7.97. The deduced amino acid sequence was not predicted to have a signal peptide by SignalP software (ExPASy). Based on analyses using the CELLO and SOSUI programs, we hypothesized that this protein is located at the plasma membrane and has 11 transmembrane domains (Fig. 1c).

Multiple sequence alignments with *SaNramp6* revealed high levels of similarity to the *Nramps* of other species (Fig. 2a). To investigate the evolutionary relationships among *Nramps* from different species, a phylogenetic analysis was performed based on the amino acid sequences. As shown in Fig. 2b, *SaNramp6* shows 80% sequence similarity to *Nramp6* from *A. thaliana*, 79% to *Nramp6* from *Theobroma cacao*, 78% to *Nramp1* from *Populus trichocarpa*, and 71% to *Nramp1* from *Nicotiana tabacum*. A phylogenetic analysis revealed that the *SaNramp6* was most closely related to *AtNramp6* (Fig. 2b). Based on this, we designated this gene as *SaNramp6* (GenBank accession no. KF887490).

Expression profiles of *SaNramp6* under CdCl₂ stress. To examine transcriptional changes under CdCl₂ stress, the expression of *SaNramp6* was monitored at different Cd-stress treatment times in leaves, stems, and roots. Without heavy metal treatment, *SaNramp6* was highly expressed in roots and leaves (Fig. 3a). However, the relative expression levels of *SaNramp6* varied greatly in different tissues under Cd treatment period progressed. Despite starting at a higher level, *SaNramp6*'s expression was not induced in leaves and in fact was reduced during treatment, reaching only a maximum of less than one-fold of the initial level at 12 h (Fig. 3d). In stems, *SaNramp6*'s expression increased gradually before 12 h, and then declined (Fig. 3c). By contrast, *SaNramp6*'s transcript accumulation was highly induced in roots (Fig. 3b). It began to increase within 12 h of treatment and peaked at around one week (14-fold).

***SaNramp6*'s expression enhances Cd²⁺ sensitivity and increases Cd²⁺ content in yeast.** To investigate the cellular function of *SaNramp6*, the protein was expressed in *Saccharomyces cerevisiae* yeast mutant (*Δycf1*) susceptible to Cd excess. *SaNramp6* and empty vector-complemented *Δycf1* cells were grown in SG-U medium overnight. Cells grown overnight were used for spotting on SG-U agar plates supplemented with 0, 15 and 20 μM CdCl₂ at indicated dilutions. The Cd supplementation of the medium caused more considerable growth inhibition in yeast cells expressing *SaNramp6* than in the control (Fig. 4a). We also analyzed the relative growth in liquid media in the presence of Cd in yeast cells. The growth of *Δycf1* cells expressing *SaNramp6* were lower than cells transformed with the empty vector (Fig. 4b). The growth inhibition due to the functional *SaNramp6* in *Δycf1* suggested that *SaNramp6* may facilitate the import of Cd inside the yeast.

To test our hypothesis that *SaNramp6* may mediate the Cd uptake, the metal content was measured in yeast cells expressing *SaNramp6* or the vector that were grown in the presence of Cd. A significantly enhanced accumulation of Cd was observed in yeast cells expressing *SaNramp6* compared with the control (Fig. 4c)

Subcellular localization of *SaNramp6*. Bioinformatics analysis using the CELLO v2.5 program software predicted that *SaNramp6* is localized in plasma membrane.




Figure 1. *SaNramp6* gene structure. **(a)** Genomic organization of *SaNramp6*. Black boxes and lines denote exons and introns, respectively. The numbers refer to the position between the exons and introns, **(b)** Comparison of the genomic DNA structure of *SaNramp6* and several *Nramp* genes of *Arabidopsis* available in GenBank. The white boxes represent the introns, and the grey boxes represent the exons. The numbers indicate the length of the sequence. I and II indicate the groups of *Nramps* in *A. thaliana*, **(c)** Transmembrane domains predicted by the SOSUI program. (*AtNramp1*: AT1G80830; *AtNramp2*: AT1G47240; *AtNramp3*: AT2G23150; *AtNramp4*: AT5G67330; *AtNramp5*: AT4G18790; *AtNramp6*: AT1G15960).

To test the prediction, the subcellular localization of SaNramp6 was analyzed by transiently expressing the SaNramp6-GFP fusion protein in protoplasts isolated from *A. thaliana*, onion epidermal cells and *N. benthamiana* epidermal cells, respectively. As shown in Fig. 5, visualized fluorescence indicated that the SaNramp6-GFP signal was localized at the plasma membrane, whereas the green fluorescent signal in the GFP control vector was distributed throughout the cytosol taken chlorophyll as control in protoplasts of *A. thaliana*.

The plasma membrane localization of *SaNramp6* was further confirmed by the transient expression of SaNramp6-GFP in onion epidermal cells and *N. benthamiana* epidermal cells. The fusion protein was found to be targeted to the plasma membrane by colocalization with FM4-64 within 5 min of the onset of staining (Supplementary Figure S1). These results indicate that SaNramp6 is localized at the plasma membrane, consistent with the prediction by the CELLO software.

***SaNramp6* participates in oxidative damage in transgenic *Arabidopsis*.** The production of reactive oxygen species (ROS) in the different lines was analyzed using H₂O₂ and O₂⁻ accumulation. As shown in Fig. 6, the contents of H₂O₂ and O₂⁻ in the transgenic lines (OE 2 and OE 3) were markedly increased and both were nearly 30% higher than those in WT line. However, they were decreased or slightly increased in the mutant *nramp1* (*Atnr*) and rescue of *nramp1* lines (*Atnr*-N24 and *Atnr*-N28), respectively (Fig. 6a,b,e,f). Thus, upon Cd stress, the overexpression of *SaNramp6* could result in a high level of H₂O₂ and O₂⁻ accumulation.

We next examined the scavenging ability of ROS by determining CAT and POD activities. The concentration of CAT and POD in the different lines had no difference in the control. Nevertheless, the CAT activity was dramatically increased in transgenic lines as was the POD activity under Cd treatment (Fig. 6c,d,g,h). Thus, the overexpression of *SaNramp6* caused more damage and enhanced CAT and POD activities under Cd treatment.

In addition, the four lines (WT, overexpression lines, mutant line, rescue lines) had no obvious differences in roots without Cd treatment (Supplementary Figure S2). However, the root length of transgenic lines (OE 2 and OE 3) was longer than that in the other lines in two weeks after the Cd treatment (Supplementary Figure S2).

Overexpression of *SaNramp6* resulted in an increased Cd concentration. Time-dependent Cd-uptake experiments using aerial parts and roots were conducted to evaluate the differences in Cd-uptake abilities by the different organs of the four lines. The time-dependent experiment on the four lines (WT, overexpression lines, mutant line, rescue lines) showed that Cd concentration increased as the treatment period progressed




Figure 2. Comparison of *SaNramp6* to *Nramps* of other species based on *Nramp* amino acid sequences from different species. Accession numbers for sequences used are listed in Table 2.

Primers	Sequence (5'-3')	Description
<i>SaN1</i> -F	ATGGCATCAACTGTCGAAACGC	qRT-PCR
<i>SaN1</i> -R	ACATGCCAATTCCACAGCGA	qRT-PCR
<i>SaNramp6</i> -F	ATGGCATCAACTGTCGAAACGC	Gene specific amplification
<i>SaNramp6</i> -R	CTACTCTAAGACAGCTCTGCGTTGCGG	Gene specific amplification
<i>SaNramp6</i> -GF	CACCATGGCATCAACTGTCGAAACGC	Gene specific amplification
<i>SaNramp6</i> -RT-F	TGTTTGGCGATTGTGCCAAG	qRT-PCR
<i>SaNramp6</i> -RT-R	ACATGCCAATTCCACAGCGA	qRT-PCR
<i>UBC9</i> -F	TGGCGTCGAAAAGGATTCTGA	qRT-PCR
<i>UBC9</i> -R	CCTTCGGTGGCTTGAATGGATA	qRT-PCR
<i>AtActin</i> -F	GCACCCTGTCTCTTACCG	qRT-PCR
<i>AtActin</i> -R	AACCCTCGTAGATTGGCACA	qRT-PCR
APL	AAGCAGTGGTATCAACGCAGAGTACGC(G) ₁₀	5'-RACE adapter primer
APS	AAGCAGTGGTATCAACGCAGAGT	5'-RACE universal primer
B 26	GACTCTAGACGACATCGA(T) ₁₈	3'-RACE adapter primer
B 25	GACTCTAGACGACATCGA	3'-RACE universal primer
5P1	ATGCAATTGAAACAAGAAAACGAGG	Reverse primer for 5'-RACE
5P2	ATGCCATCTATCGATCAAAGTGT	Reverse primer for 5'-RACE
3P1	AGGCTGGCGTGGTTGATACATGTG	Forward primer for 3'-RACE
3P2	TGTGCGAATCGGATCAAGTTT	Forward primer for 3'-RACE

Table 1. Degenerate and specific primers used in this work.

and the pattern of Cd uptake by roots displayed an initial slower stage during the first eight hours, followed by a second, rapid stage over the subsequent two weeks and it was significant lower in OE 3 than in WT, *Atnr* and *Atnr*-N24 (Fig. 7a). The Cd concentrations in the roots of all of the lines increased remarkably under Cd-stress conditions for two weeks. However, compared with the other lines, the OE 3 had significant higher concentration of Cd in its aerial parts. Cd was transferred to aboveground parts began in 8 h and increased steadily in the following two weeks. The translocation factor of the transgenic lines was markedly higher than that of the other




Figure 3. Expression patterns of *SaNramp6* in *S. alfredii* under Cd stress. (a) Different tissue without any heavy metal treatments, (b) Root, (c) Stem, (d) Leaf. The normalized mRNA levels without treatment (y-axis “Relative mRNA expression”) were set arbitrarily to 1. Bars indicate means \pm standard deviations (SDs) of at least three independent biological experiments. Different letters on the bars indicate significant difference between the treatments. P-values of the two-way ANOVAs of Time, Cd (Cd treatment) and their interaction (Time \times Cd) are indicated. * $P < 0.05$; ** $P < 0.01$.




Figure 4. *SaNramp6* expression increases Cd^{2+} sensitivity and Cd^{2+} content in yeast. (a) Growth of $\Delta ycf1$ yeast cells expressing *SaNramp6* on plates containing SG-U without CdCl_2 (Left) or supplemented with $15\mu\text{M}$ CdCl_2 (Middle) and $20\mu\text{M}$ CdCl_2 (Right), (b) Time-dependent growth of yeast strains in SG-U liquid medium supplemented with $5\mu\text{M}$ CdCl_2 , (c) Cd content of $\Delta ycf1$ yeast cells expressing *SaNramp6* grown for 48 h in liquid SG-U supplemented with $5\mu\text{M}$ CdCl_2 . Bars indicate means \pm standard deviations (SDs) of at least three independent biological experiments. One or two asterisks indicate a significant difference at $P < 0.05$ or $P < 0.01$ from the $\Delta ycf1 + EV$.




Figure 5. Subcellular localization of SaNramp6. Control, Non-transformed protoplast; GFP-vector, protoplast transformed with p35S-GFP vector; SaNramp6-GFP, protoplast transformed with SaNramp6-GFP fusion. Scale bar = 7.5 μ m.

Species	Name	Accession numbers
<i>Arabidopsis thaliana</i>	AtNramp1	NP_178198.1
<i>Arabidopsis thaliana</i>	AtNramp6	NP_173048.3
<i>Populus trichocarpa</i>	PtNramp1	XP_006368514.1
<i>Morus notabilis</i>	MnNramp6	EXB50420.1
<i>Triticum urartu</i>	TuNramp3	EMS65084.1
<i>Nicotiana tabacum</i>	NtNramp1	BAH66919.1
<i>Selaginella moellendorffii</i>	SmNramp	XP_002966634.1
<i>Solanum melongena</i>	SmNramp1	BAM34953.1
<i>Solanum torvum</i>	StNramp1	BAM34952.1
<i>Brassica juncea</i>	BjNramp	ACR16683.1
<i>Medicago truncatula</i>	MtNramp	XP_003602053.1
<i>Arachis hypogaea</i>	AhNramp1	AFQ37304.1
<i>Theobroma cacao</i>	TcNramp6	XP_007023419.1
<i>Oryza sativa Japonica</i>	OsNramp5	NP_001059312.1

Table 2. Names and accession numbers of the Nramp protein family members.

lines after Cd exposure for two weeks (Fig. 7b). These results indicated that the transgenic lines may have a better absorption capacity for Cd. Thus, *SaNramp6* may influence the accumulation ability of Cd in *S. alfredii*.

To decipher the phenomenon of Cd accumulation in the four different lines, Cd²⁺ was measured in 30 d after 30 μ M Cd treatment. Significantly higher Cd²⁺-influx rates were identified in overexpression lines (OE 2 and OE 3) compared with WT line (Fig. 8a,b) and markedly lower in *Atnr* line (Fig. 8c,d); however, there were no differences between rescue lines (*Atnr*-N24 and *Atnr*-N28) and WT line (Fig. 8c,d).

Discussion

Here, we cloned an *Nramp* family member from a heavy metal-accumulating ecotype of *S. alfredii* and the results showed that it conferred the ability to accumulate Cd in overexpression transgenic *A. thaliana*. Cd is a strongly toxic heavy metal transported across plant membranes by physiological metal transporters²⁹. To date, various gene families related to the transport of Cd have been reported, such as the P-type ATPase superfamily³⁰, ABC transporters³¹ and the CE family³². Among these metal transporters, the *Nramp* family is widely distributed in mammals, fungi and bacteria.




Figure 6. ROS accumulation responses to Cd stress and physiological indicators in four different lines - WT (wild type); OE 2 and OE 3 (overexpression lines); *Atnr* (mutant line); *Atnr*-N24 and *Atnr*-N28 (rescue lines). (a,e) H_2O_2 , (b,f) O_2^- , (c,g) CAT activity, (d,h) POD activity. Control, without Cd treatment; Cd treatment, 30 μM Cd treatment for two weeks. Bars indicate means \pm standard deviations (SDs) of at least three independent biological experiments. One or two asterisks indicate a significant difference at $P < 0.05$ or $P < 0.01$ from wild type.

The determination of *SaNramp6*'s subcellular localization is important for understanding its potential roles in the process of accumulating metals. *AtNramp6* is located in a vesicular-shaped endomembrane compartment and works as an intracellular Cd transporter²³. Similarly, *OsNramp1* was localized to the plasma membrane in onion epidermal cells, and the overexpression of *OsNramp1* results in a Cd accumulation in the leaves³³. The soybean Nramp homologue, *GmDmt*, is located on the peribacteroid membrane of root nodules and mediates ferrous iron uptake in yeast²⁷. In *Thlaspi japonicum* H., a nickel (Ni) hyperaccumulator, *TjNramp4* could specifically transport Ni and increase Ni concentrations³⁴. The deduced amino acid sequence of *SaNramp6* shares an 80% identity with *AtNramp6*, and the phylogenetic tree also indicated that *SaNramp6* was most similar to *AtNramp6*. In addition, our subcellular localization analysis showed that *SaNramp6* was located in the plasma membrane. Thus, *SaNramp6* could function as a metal transporter in the plasma membrane (Fig. 5).

An plasma membrane localization is consistent with *SaNramp6* conferring Cd uptake by increasing the Cd content in plant tissues. *AtNramp3* is involved in increased metal tolerance or accumulation. However, *AtNramp6* leads to Cd hypersensitivity when overexpressed in *Arabidopsis*, and *Arabidopsis* plants lacking *AtNramp6* are more resistant to Cd than WT lines²³. In rice, *OsNramp1* participates in cellular Cd uptake or transport and the overexpression of *OsNramp1* enhances tolerance to Cd and increases Cd accumulation in shoots²⁴. *OsNramp5* is a major transporter for Cd uptake, influencing Cd absorption in both solution and soil cultures^{25,33}. Hence, compared with the control, *SaNramp6*'s overexpression in *A. thaliana* could accumulate more ROS of roots when exposed to CdCl_2 . We believe that understanding its functions in plants will facilitate the development of Cd-accumulating plants.

The growth of the Cd sensitive yeast strain ($\Delta ycf1$) transformed with the empty vector was inhibited by Cd, and yeast cells harbouring the *SaNramp6*-expression vector exhibited weaker growth activities. The results indicated that *SaNramp6* cannot complement the Cd sensitivity or rescue the Cd-sensitivity phenotype in the mutant yeast strain. However, the Cd concentration of the *SaNramp6*-expression strain was 10% higher than that of the empty vector strain (Fig. 4). The induction of *SaNramp6* expression by CdCl_2 suggested that the gene might be involved in responding to heavy metal stress and is a transporter for Cd uptake in *S. alfredii*. Consistent results have been reported using a similar approach. Thomine *et al.*²² found that the growth of transgenic yeast expressing *AtNramp1*, *AtNramp3*, *AtNramp4* was strongly reduced in liquid cultures supplemented with 3 μM CdCl_2 compared with the control, and these genes increased the Cd content in yeast. However, *AtNramp3*-OE in *Arabidopsis* were found to be hypersensitive to Cd. *TcNRAMP3*'s-expression increased Cd sensitivity and the Cd content in yeast, and *TcNRAMP3*-OE in tobacco resulted in a slight sensitivity of root growth to Cd³⁵. The growth of yeast strain $\Delta ycf1$ was affected by *OsNRAMP5*, which is involved in Cd transport²⁵. Therefore, these data showed that some NRAMP members could increase Cd sensitivity and Cd concentration. Here, our results from *SaNramp6* in transgenic yeast with Cd-sensitivity phenotype and Cd concentration (Fig. 4) and that in transgenic *A. thaliana* were consistent with the previous results (Fig. 7).

The capacity to reduce Cd-associated oxidation may be an important mechanism contributing to Cd uptake and transport. To test the role of *SaNramp6* in heavy metal-stress tolerance, a functional analysis was carried out by overexpressing *SaNramp6* in *A. thaliana* and rescuing the *Arabidopsis* mutant *nramp1*. In the physiological assays (Fig. 6), the root lengths of overexpression transgenic plants were markedly longer and the contents of H_2O_2 , O_2^- , CAT and POD were also higher than those in WT lines, which suggested that the overexpression of *SaNramp6* enhanced the Cd uptake and accumulation in transgenic plants.

Taken together, we have functioned *SaNramp6* in transgenic yeast and *A. thaliana* and the data presented in this study suggested that *SaNramp6* may be a critical Cd transporter responsible for Cd accumulation in *S. alfredii*. It was hard to place the functions of *SaNramp6* into specific categories such as uptake or translocation.




Figure 7. Time-dependent Cd-uptake experiments (a) and Cd-translocation factors (Tf) (b) in four lines. WT (wild type); OE 3 (overexpression lines); *Atnr* (mutant line); *Atnr*-N24 (rescue lines). Bars indicate means \pm standard deviations (SDs) of at least three independent biological experiments. Different letters on the bars indicate significant difference at same time within treatments (WT, overexpression lines, mutant line, rescue lines). P-values of the two-way ANOVAs of Time, Cd (Cd treatment) and their interaction (Time \times Cd) are indicated. * $P < 0.05$; ** $P < 0.01$.

The improved Cd uptake caused by *SaNramp6* may be due to the exertion of direct effects on several major pathways or may work in cooperation with other genes participating heavy metal uptake, transport, sequestration and detoxification. A similar case was reported recently, the uptake of Fe in roots by *NRAMP1* requires the partnership of another transporter, *IRT1*, in *A. thaliana*³⁶. Although the function of *SaNramp6* is still unclear, the gene appears to be related to the hyperaccumulator characteristic of *S. alfredii*. These findings will contribute to understanding the function of *Nramp* genes and provide experimental evidence and theoretical guidance for further studies.

Methods

Plant materials and growth conditions. A hyperaccumulating ecotype of *S. alfredii* was collected from the area of an old Pb/Zn mine in Quzhou City, Zhejiang Province, P. R. China. The plants were water-cultivated in an artificial climate chamber at 25°C with a 16 h light/8 h dark cycle. The *S. alfredii* seedlings used for the stress treatment were asexual propagated to ensure consistency and grown in half-strength Hoagland-Arnon solution for about two weeks until relatively vigorous roots grew. For the expression analyses of target genes, plants were treated with 400 μM CdCl₂ for 0 h, 0.5 h, 6 h, 12 h, 24 h, 48 h, 72 h, 96 h, 1 week and 2 week. Each treatment was replicated three times. All samples were quickly frozen in liquid nitrogen followed by storage at -80°C until use.




Figure 8. Comparison of Cd concentrations and net Cd²⁺-influx rates in four lines - WT (wild type); OE 2 and OE 3 (overexpression lines); *Atnr* (mutant line); *Atnr*-N24 and *Atnr*-N28 (rescue lines). **(a,c)** Cd²⁺ flux rates with Cd treatment for 24 h. **(b,d)** Mean flow rates of Cd²⁺. Bars indicate means \pm standard deviations (SDs) of at least three independent biological experiments. One or two asterisks indicate a significant difference at $P < 0.05$ or $P < 0.01$ from wild type.

A. thaliana (ecotype Columbia) was grown in a controlled environmental chamber at 22 °C under a long-day cycle (16 h light, 8 h dark), with a white light intensity of approximately 125 mmol·m⁻²·s⁻¹ and 70% relative humidity. Overexpression lines, mutant and rescue lines, were selected for physiological assays. The seeds were surface sterilized and germinated on 1/2 Murashige and Skoog (MS) agar plates containing 25 mg·L⁻¹ hygromycin. Whereafter, 30 d-old *Arabidopsis* seedlings were soaked in Hoagland-Arnon solution with or without 50 μ M CdCl₂ for 24 h and then used to measure the Cd²⁺ flux. For physiological assays, 30 d-old homozygous transgenic seedlings were soaked in Hoagland-Arnon solution with or without 30 μ M CdCl₂ for two weeks, and treatments for 0 h, 10 min, 20 min, 30 min, 1 h, 2 h, 4 h, 8 h, 12 h, 24 h, 48 h, 96 h, 1 week and 2 week used for Cd-uptake assay.

RNA preparation, cDNA synthesis and DNA extraction. Leaves, stems and roots were harvested after each treatment, and all of the samples were frozen in liquid nitrogen and stored at -80 °C for analysis. Total RNA was isolated from the tissues using the Total RNA Purification Kit (NORGEN, Thorold, Canada). First-strand cDNA was then synthesized from 2 μ g of total RNA by the Superscript RT III first-strand cDNA synthesis kit followed by RNase H (Invitrogen, Carlsbad, USA) treatment. Genomic DNA was isolated from seedling leaves using cetyltrimethyl ammonium bromide (CTAB) method as described by Murray and Thompson³⁷.

Cloning of *SaNramp6* gene. The full-length *SaNramp6* cDNA was amplified by reverse transcription-PCR (RT-PCR) and rapid amplification of cDNA ends-PCR (RACE-PCR).

The internal fragment of *SaNramp6* was isolated from *S. alfredii* using the specific primers *SaNramp6*-F and *SaNramp6*-R, which were designed according to transcriptome data²⁸. To obtain the 3'-end cDNA and 5'-ready cDNA, four gene specific primers 3P1, 3P2, 5P1 and 5P2, were designed and synthesized based on the sequence of the cloned internal fragment. The cloning was performed as described by Wang *et al.*³⁸. Additionally, the genomic sequence of *SaNramp6* was amplified by PCR using genomic DNA as the template with primers *SaNramp6*-F and *SaNramp6*-R. All of the primers are listed in Table 1.

Bioinformatics analysis of *SaNramp6*. To compare the genomic structure, the genomic sequences of *A. thaliana Nramps* from GenBank were searched and the intron-exon structure was analyzed.

Translation and protein analyses of *SaNramp6* were initially performed using ExPASy tools (<http://www.expasy.org/tools/>). CELLO v.2.5: subCELLular LOCALization predictor (<http://cello.life.nctu.edu.tw/>) and SOSUI version 1.11 (<http://bp.nuap.nagoya-u.ac.jp/sosui/>) were used to predict subcellular localization and transmembrane domains, respectively. For the multiple sequence alignment, Clustal Omega (<http://www.ebi.ac.uk/Tools/msa/>) was performed to align amino acid sequences first, and subsequently, the results were edited by GeneDoc. Additionally, a phylogenetic tree was constructed by MEGA 5.2 software using the Neighbour-joining method with 1,000 replicates based on amino acid sequences of the NRAMP proteins. The known NRAMP protein sequences from NCBI GenBank are shown in Table 2.

Expression pattern analysis. SYBR-based quantitative real-time PCR (qRT-PCR) reactions (SYBR premix EX Tag reagent, TaKaRa, Da Lian, China) were carried out in triplicate on a 7300 Real-Time PCR System (Applied Biosystems, CA, USA) according to the manufacturer's instructions. Relative gene expression was estimated based on the $2^{-\Delta\Delta Ct}$ method, applying the geometric mean of two reference genes: *UBC9* and *TUB39*⁴⁰. All of the primers for RT-qPCR are listed in Table 1.

Expression vector construction. The open reading frame of *SaNramp6* was amplified by PCR using High Fidelity KOD-Plus DNA Polymerase (Toyobo, Japan) from the cDNA of *S. alfredii* using the specific primers *SaNramp6*-GF and *SaNramp6*-R (Table 1). The yeast expression vector pYES2.1-*SaNramp6* was generated using pYES2.1 TOPO[®] TA Expression Kit (Invitrogen, Carlsbad, USA). For subcellular location and plant expression vector, the purified PCR products were then cloned into the Gateway entry vector pENTR/D-Topo (Invitrogen, Carlsbad, USA) and positive clones were further sequenced to verify the direction and sequence accuracy. The sequence-verified plasmid was then recombined in pK7WGF2.0 and pH2GW7.0⁴¹ to generate pK7W-GF2.0-*SaNramp6* and pH2GW7.0-*SaNramp6*, respectively.

Subcellular localization of *SaNramp6*. The correct plasmid pK7WGF2.0-*SaNramp6* fused to the green fluorescent protein (GFP) was extracted by Plasmid Maxprep Kit (Vigorous, Beijing, China). Free vector p35S-GFP was used as control. *A. thaliana* protoplast isolation and transfection were performed as previously described⁴². The subcellular location of *SaNramp6* was further investigated by transient expression in onion epidermal cells and *Nicotiana benthamiana* lower leaf epidermal cells as described by Liu *et al.*⁴³ and Zheng *et al.*⁴⁴, respectively. A LSM510 confocal laser scanning microscope (Carl Zeiss, Oberkochen, Germany) was used to observe the signals.

Heterologous expression of *SaNramp6* in yeast. The *Saccharomyces cerevisiae* strain BY4742 $\Delta ycf1$ (*MAT α* ; *his3 Δ 1*; *leu2 Δ 0*; *met15 Δ 0*; *ura3 Δ 0*; *YDR135c::kanMX4*) was a Cd-sensitive mutant, which lacked the ability to compartmentalize Cd into vacuoles⁴⁵, was used to assess the Cd tolerance of *SaNramp6*. The yeast transformation was performed using the lithium acetate method⁴⁶. Yeast $\Delta ycf1$ cells transformed with the empty pYES2.0 vector were used as controls. The transformed yeast cells were selected on synthetic defined medium lacking uracil. For complementation assays, a series of three 1:10 dilutions from each culture was spotted onto synthetic-galactose-uracil (SG-U) agar plates supplemented with 0, 15 and 20 μ M CdCl₂ and incubated at 30 °C for three days. The relative growth of transformants was determined by measuring the OD₆₀₀ at 6 h intervals. For the Cd-uptake assay, yeast cells transformed with the empty or *SaNramp6* vector were grown for 48 h at 30 °C on SG-U supplemented with 5 μ M CdCl₂, then measured the Cd content.

Detection of the *Arabidopsis Atnr* mutant by *Atnramp1*. To understand the functions of *SaNramp6*, we obtained mutant alleles from the SALK collection of sequence-indexed T-DNA insertions⁴⁷. However, the mutant alleles of *AtNramp6* were not found. Therefore, a single insertion line (SALK_053236; *nramp1-1*) was confirmed for *SaNramp6* because *AtNramp6* and *AtNramp1* have similar genomic structure. A homozygous mutant was detected by PCR using the primers (LP/RP and universal primers BP) designed based on the T-DNA website (<http://signal.salk.edu/tdnaprimers.2.html>) (data not shown). The collected homozygous mutant seeds were air-dried and stored at 4 °C.

Generation of transgenic *A. thaliana*. The recombinant plasmid pH2GW7.0-*SaNramp6* was introduced into *Agrobacterium tumefaciens* strain EHA105. *A. thaliana* ecotype Columbia plants were transfected by the floral dip method⁴⁸. Positive transformants were selected based on hygromycin (Hyg, 20 μ g·mL⁻¹) resistance and confirmed by PCR and RT-PCR using the primers described above, *AtActin* (Table 1) was the internal control. Homozygous lines were identified by screening for non-segregation from each independent transformant (T₃ generation).

Physiological analysis of *SaNramp6* transgenic, mutant and rescue of mutant lines. Six overexpression lines (designated OE) with high transcriptional levels of *SaNramp6* and 26 rescue of *Arabidopsis* mutant lines were obtained in this study. Among them, the OE 2, OE 3, *Atnr*-N24 and *Atnr*-N28 lines were selected in the following study owing to their phenotypes.

To investigate the potential effects of *SaNramp6* in *A. thaliana*, *SaNramp6*-OE *A. thaliana* (OE 2 and OE 3), homozygous mutant *A. thaliana* (*Atnr*) and the rescue of the *Arabidopsis* mutants (*Atnr*-N24 and *Atnr*-N28), as well as wild type, were used for abiotic stress-related physiological analyses, including root length, peroxidase

(POD) activity, catalase (CAT) content, H₂O₂ and superoxide anion accumulations, and measurements of the Cd²⁺ flux. All of the experiments were independently carried out three times.

As for analyzing peroxidase (POD) activity, catalase (CAT) content, H₂O₂ and superoxide anion accumulations, approximately 0.1 g of root tissue was ground in liquid nitrogen and placed in 2-mL tubes. The extraction of these physiological indices used the appropriate kits according to the instruction manual (Comin, Suzhou, China).

The net Cd²⁺ fluxes in the roots of *Arabidopsis* were measured noninvasively by the Younger USA NMT Service Centre (Xuyue, Beijing) using the NMT system (NMT100 Series, Younger, USA LLC, Amherst, MA, USA). Prior to the flux measurement, the roots were equilibrated for 15 min in testing liquid (0.05 mM CdCl₂, 0.1 mM KCl, 0.02 mM CaCl₂, 0.02 mM MgCl₂, 0.5 mM NaCl, 0.1 mM Na₂SO₄ and 0.3 mM MES, pH 5.7). Then, the transmembrane Cd²⁺ flux in roots was measured of different lines (120 μm to root apex) for 15 min by a Cd²⁺-selective microelectrode. All of the measurements were repeated at least six times independently.

Cd concentration assay. To test the characteristic of *SaNramp6*'s Cd accumulation in *A. thaliana*, *SaNramp6*-OE *A. thaliana* (OE 3), homozygous mutant *A. thaliana* (*Atnr*), the rescue of the *Arabidopsis* mutant (*Atnr*-N28) and wild type were used in this experiment.

Roots and aerial parts were harvested individually for the Cd concentration analysis, and roots were resorbed by dipping in 1 mM EDTA for 30 min, and then washed three times with distilled water. All of the samples containing yeast cells for Cd determination were dried at 105 °C for 30 min, and then placed at 70 °C until they reached a constant weight. The dried samples were digested with a concentrated acid mixture of HNO₃, HClO₄, and H₂SO₄ (volume ratio = 4:1:0.5) at 250 °C for 8 h. The metal concentration in the digested solution was determined by atomic absorption spectrometry (M6; SOLLAR) and an inductively coupled plasma-mass spectrometer (ICP-MS; NexION 300; PerkinElmer) after dilution.

Data processing. Data were exhibited as the means ± standard deviations (SDs) of at least three independent biological experiments. Statistical analysis was performed using SPSS 17.0 statistics software. To test significant changes in mRNA relative expression and Cd concentration, time and Cd treatment were regarded as the main factors. Tukey-HSD method was used to correct all P-values of these multiple comparisons. In addition, one asterisk (*) or two asterisk (**), significantly different from control at P = 0.05, 0.01, respectively.

The translocation factor for Cd within a plant was expressed by the concentration in the aerial parts (μg·g⁻¹DW)/the concentration in the roots (μg·g⁻¹DW), which showed the Cd-translocation properties from roots to aerial parts⁴⁹.

References

1. Fox, T. C. & Guerinot, M. L. Molecular biology of cation transport in plants. *Plant Biology* **49**, 669–696 (1998).
2. Satarug, S., Garrett, S. H., Sens, M. A. & Sens, D. A. Cadmium, environmental exposure, and health outcomes. *Ciência & Saúde Coletiva* **16**, 2587–2602 (2010).
3. Nawrot, T. *et al.* Environmental exposure to cadmium and risk of cancer: a prospective population-based study. *Lancet Oncology* **7**, 119–126 (2006).
4. Mason, L. H., Harp, J. P. & Han, D. Y. Pb neurotoxicity: neuropsychological effects of lead toxicity. *Biomed Research International* **2014** (2014).
5. Mcgrath, S. P., Zhao, F. J. & Lombi, E. Plant and rhizosphere processes involved in phytoremediation of metal-contaminated soils. *Plant and Soil* **232**, 207–214 (2001).
6. Mcgrath, S. P. & Zhao, F. J. Phytoextraction of metals and metalloids from contaminated soils. *Current Opinion in Biotechnology* **14**, 277–282 (2003).
7. Yang, X. E., Long, X. X., Ni, W. Z. & Fu, C. X. *Sedum alfredii* H: A new Zn hyperaccumulating plant first found in China. *Science Bulletin* **47**, 1634–1637 (2002).
8. Yang, X. E. & Stoffella, P. J. Cadmium tolerance and hyperaccumulation in a new Zn-hyperaccumulating plant species (*Sedum alfredii* Hance). *Plant and Soil* **259**, 181–189 (2004).
9. Lu, L. *et al.* Enhanced root-to-shoot translocation of cadmium in the hyperaccumulating ecotype of *Sedum alfredii*. *Journal of Experimental Botany* **59**, 3203–3213 (2008).
10. Yang, X. *et al.* Zinc compartmentation in root, transport into xylem, and absorption into leaf cells in the hyperaccumulating species of *Sedum alfredii* Hance. *Planta* **224**, 185–195 (2006).
11. Tian, S. & Brown, P. Cellular sequestration of cadmium in the hyperaccumulator plant species *Sedum alfredii*. *Plant Physiology* **157**, 1914–1925 (2011).
12. Page, M. D. & Merchant, S. S. Two chlamydomonas CTR copper transporters with a novel cys-met motif are localized to the plasma membrane and function in copper assimilation. *Plant Cell* **21**, 928–943 (2009).
13. Vert, G., Briat, J. F. & Curie, C. *Arabidopsis* *IRT2* gene encodes a root-periphery iron transporter. *Plant Journal* **26**, 181–189 (2001).
14. Nevo, Y. & Nelson, N. The NRAMP family of metal-ion transporters. *Biochimica Et Biophysica Acta Molecular Cell Research* **1763**, 609–620 (2006).
15. Segond, D. *et al.* NRAMP genes function in *Arabidopsis thaliana* resistance to *Erwinia chrysanthemi* infection. *Plant Journal for Cell & Molecular Biology* **58**, 195–207 (2009).
16. Migeon, A. *et al.* Genome-wide analysis of plant metal transporters, with an emphasis on poplar. *Cellular & Molecular Life Sciences* **67**, 3763–3784 (2010).
17. Colangelo, E. P. & Guerinot, M. L. Put the metal to the petal: metal uptake and transport throughout plants. *Current Opinion in Plant Biology* **9**, 322–330 (2006).
18. Xia, J., Yamaji, N., Kasai, T. & Ma, J. F. Plasma membrane-localized transporter for aluminum in rice. *Proceedings of the National Academy of Sciences of the United States of America* **107**, 18381–18385 (2010).
19. Cellier, M. *et al.* Nramp defines a family of membrane proteins. *Proceedings of the National Academy of Sciences* **92**, 10089–10093 (1995).
20. Curie, C., Alonso, J. M., Le, J. M., Ecker, J. R. & Briat, J. F. Involvement of NRAMP1 from *Arabidopsis thaliana* in iron transport. *Biochemical Journal* **347**(Pt 3), S34–S36 (2000).
21. Lanquar, V. *et al.* Mobilization of vacuolar iron by AtNRAMP3 and AtNRAMP4 is essential for seed germination on low iron. *Embo Journal* **24**, 4041–4051 (2006).

22. Thomine, S. & Schroeder, J. I. Cadmium and iron transport by members of a plant metal transporter family in *Arabidopsis* with homology to *Nramp* genes. *Proceedings of the National Academy of Sciences of the United States of America* **97**, 4991–4996 (2000).
23. Cailliatte, R., Lapeyre, B., Briat, J. F., Mari, S. & Curie, C. The *NRAMP6* metal transporter contributes to cadmium toxicity. *Biochemical Journal* **422**, 217–228 (2009).
24. Takahashi, R. *et al.* The *OsNRAMP1* iron transporter is involved in Cd accumulation in rice. *Journal of Experimental Botany* **62**, 4843–4850 (2011).
25. Sasaki, A., Yamaji, N., Yokosho, K. & Ma, J. F. *Nramp5* is a major transporter responsible for manganese and cadmium uptake in rice. *Plant Cell* **24**, 2155–2167 (2012).
26. Berczky, Z., Wang, H. Y., Schubert, V., Ganai, M. & Bauer, P. Differential regulation of *Nramp* and *Irt* metal transporter genes in wild type and iron uptake mutants of tomato. *Journal of Biological Chemistry* **278**, 24697–24704 (2003).
27. Kaiser, B. N. *et al.* The soybean *NRAMP* homologue, *GmDMT1*, is a symbiotic divalent metal transporter capable of ferrous iron transport. *Plant Journal for Cell & Molecular Biology* **35**, 295–304 (2003).
28. Han, X. *et al.* Integration of small RNAs, degradome and transcriptome sequencing in hyperaccumulator *Sedum alfredii* uncovers a complex regulatory network and provides insights into cadmium phytoremediation. *Plant Biotechnology Journal* **33**, 552–575 (2016).
29. Clemens, S., Antosiewicz, D. M., Ward, J. M., Schachtman, D. P. & Schroeder, J. I. The plant cDNA *LCT1* mediates the uptake of calcium and cadmium in yeast. *Proceedings of the National Academy of Sciences of the United States of America* **95**, 12043–12048 (1998).
30. Alvarez, M. E. *et al.* Reactive Oxygen intermediates mediate a systemic signal network in the establishment of plant immunity. *Cell* **92**, 773–784 (1998).
31. Rea, P. A., Li, Z. S., Lu, Y. P., Drozdowicz, Y. M. & Martinoia, E. From vacuolar gs-x pumps to multispecific ABC transporters. *Plant Biology* **49**, 727–760 (1998).
32. Paulsen, I. T. & Saier, M. H. A novel family of ubiquitous heavy metal ion transport proteins. *The Journal of Membrane Biology* **156**, 99–103 (1997).
33. Ishimaru, Y. *et al.* Characterizing the role of rice *NRAMP5* in manganese, iron and cadmium transport. *Scientific Reports* **2**, 989–993 (2012).
34. Mizuno, T. *et al.* Cloning of three *ZIP/Nramp* transporter genes from a Ni hyperaccumulator plant *Thlaspi japonicum* and their Ni²⁺-transport abilities. *Plant Physiology & Biochemistry Ppb* **43**, 793–801 (2005).
35. Wei, W. *et al.* The *Thlaspi caerulescens* *NRAMP* homologue *TcNRAMP3* is capable of divalent cation transport. *Molecular Biotechnology* **41**, 15–21 (2009).
36. Castaings, L., Caquot, A., Loubet, S. & Curie, C. The high-affinity metal transporters *NRAMP1* and *IRT1* team up to take up iron under sufficient metal provision. *Scientific Reports* **6** (2016).
37. Murray, M. G. & Thompson, W. F. Rapid isolation of high molecular weight plant DNA. *Nucleic Acids Research* **8**, 4321–4325 (1980).
38. Wang, M., Zhang, Y., Wang, J., Wu, X. & Guo, X. A novel *MAP* kinase gene in cotton (*Gossypium hirsutum* L.), *GhMAPK*, is involved in response to diverse environmental stresses. *Journal of Biochemistry & Molecular Biology* **40**, 325–332 (2007).
39. Vandesompele, J. *et al.* Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biology* **3**, 1–11 (2002).
40. Sang, J. *et al.* Selection and validation of reference genes for real-time quantitative PCR in hyperaccumulating ecotype of *Sedum alfredii* under different heavy metals stresses. *Plos One* **8**, e82927–e82927 (2013).
41. Karimi, M., Inzé, D. & Depicker, A. GATEWAY vectors for *Agrobacterium*-mediated plant transformation. *Trends in Plant Science* **7**, 193–195 (2002).
42. Yoo, S. D., Cho, Y. H. & Sheen, J. *Arabidopsis* mesophyll protoplasts: a versatile cell system for transient gene expression analysis. *Nature Protocols* **2**, 1565–1572 (2007).
43. Liu, M. *et al.* Functional characterization of a gene in *sedum alfredii* hance resembling rubber elongation factor endowed with functions associated with cadmium tolerance. *Frontiers in Plant Science* **7** (2016).
44. Zheng, H. *et al.* A Rab-E GTPase mutant acts downstream of the Rab-D subclass in biosynthetic membrane traffic to the plasma membrane in tobacco leaf epidermis. *Plant Cell* **17**, 2020–2036 (2005).
45. Li, Z. S. & Rea, P. A. A new pathway for vacuolar cadmium sequestration in *Saccharomyces cerevisiae*: YCF1-catalyzed transport of bis (glutathionato) cadmium. *Proceedings of the National Academy of Sciences of the United States of America* **94**, 42–47 (1997).
46. Szczyńska, M. S., Wemmie, J. A., Moye-Rowley, W. S. & Thiele, D. J. A yeast metal resistance protein similar to human cystic fibrosis transmembrane conductance regulator (CFTR) and multidrug resistance-associated protein. *Journal of Biological Chemistry* **269**, 22853–22857 (1994).
47. Alonso, J. M. & Ecker, J. R. Genome-wide insertional mutagenesis of *Arabidopsis thaliana*. *Science* **301**, 653–657 (2003).
48. Clough, S. J. & Bent, A. F. Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *The Plant Journal* **16**, 735–743 (1998).
49. Stoltz, E. & Greger, M. Accumulation properties of As, Cd, Cu, Pb and Zn by four wetland plant species growing on submerged mine tailings. *Environmental & Experimental Botany* **47**, 271–280 (2002).

Acknowledgements

We thank the anonymous referees and the editor for the comments and suggestions that helped improve the manuscript. This work was supported by Basic Scientific Research Project of Nonprofit Central Research Institutions (No. CAFYBB2012040 and TGB2013008), National High-tech R&D Program of China No. 2013AA102701-3, the National Natural Science Foundation of China (No. 31200465), the National Key R&D Program of China (No. 2016YFD0800801) and the National Nonprofit Institute Research Grant of CAF (No. CAFYBB2012040, No. RISF2014010).

Author Contributions

S.C., X.H., J.F., R.Z. planned and designed the research. S.C., X.H., and J.F. performed the experiments. Z.L., W.M., M.L., J.S. and J.J. contributed analytical tools, and S.C., X.H. wrote the manuscript and coordinated its revision. R.Z. contributed reagents/materials/funds support. All authors read and provided helpful discussions, and approved the final version.

Additional Information

Supplementary information accompanies this paper at <https://doi.org/10.1038/s41598-017-13463-4>.

Competing Interests: The authors declare that they have no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2017