

Self-Adjoint Extensions by Additive Perturbations

ANDREA POSILICANO

Abstract. Let $A_{\mathcal{N}}$ be the symmetric operator given by the restriction of A to \mathcal{N} , where A is a self-adjoint operator on the Hilbert space \mathcal{H} and \mathcal{N} is a linear dense set which is closed with respect to the graph norm on $D(A)$, the operator domain of A . We show that any self-adjoint extension A_{Θ} of $A_{\mathcal{N}}$ such that $D(A_{\Theta}) \cap D(A) = \mathcal{N}$ can be additively decomposed by the sum $A_{\Theta} = \bar{A} + T_{\Theta}$, where both the operators \bar{A} and T_{Θ} take values in the strong dual of $D(A)$. The operator \bar{A} is the closed extension of A to the whole \mathcal{H} whereas T_{Θ} is explicitly written in terms of a (abstract) boundary condition depending on \mathcal{N} and on the extension parameter Θ , a self-adjoint operator on an auxiliary Hilbert space isomorphic (as a set) to the deficiency spaces of $A_{\mathcal{N}}$. The explicit connection with both Kreĭn's resolvent formula and von Neumann's theory of self-adjoint extensions is given.

Mathematics Subject Classification (2000): 47B25 (primary), 47B38, 47F05 (secondary).

1. – Introduction

Given a self-adjoint operator $A : D(A) \subseteq \mathcal{H} \rightarrow \mathcal{H}$, let $A_{\mathcal{N}}$ be the restriction of A to \mathcal{N} , where $\mathcal{N} \subsetneq D(A)$ is a dense linear subspace which is closed with respect to the graph norm. Then $A_{\mathcal{N}}$ is a closed, densely defined, symmetric operator. Since $\mathcal{N} \neq D(A)$, $A_{\mathcal{N}}$ is not essentially self-adjoint, as A is a non-trivial extension of $A_{\mathcal{N}}$, and, by the famed von Neumann's formulae [15], we know that $A_{\mathcal{N}}$ has an infinite family of self-adjoint extensions A_U parametrized by the unitary maps U from \mathcal{K}_+ onto \mathcal{K}_- , where $\mathcal{K}_{\pm} := \text{Kernel}(-A_{\mathcal{N}}^* \pm i)$ denotes the deficiency spaces.

In Section 2 we define a family A_{Θ} of extensions of $A_{\mathcal{N}}$ by means of a Kreĭn-like formula i.e. by explicitly giving its resolvent $(-A_{\Theta} + z)^{-1}$ (see Theorem 2.1). By using the approach developed in [16], we describe the domain of A_{Θ} in terms of the boundary condition $\tau\phi_{\star} = \Theta Q_{\phi}$, where $\tau : D(A) \rightarrow \mathfrak{h}$ is a surjective continuous linear mapping with $\text{Kernel } \tau = \mathcal{N}$, $\Theta : D(\Theta) \subseteq \mathfrak{h} \rightarrow \mathfrak{h}$ is self-adjoint and \mathfrak{h} is a Hilbert space isomorphic (as a set) to \mathcal{K}_{\pm} .

In Section 3 we use the resolvent $(-A_\Theta + z)^{-1}$ given in Theorem 2.1 to re-write A_Θ in a more appealing way as a sum $\bar{A} + T_\Theta$ where both \bar{A} and T_Θ take values in the strong dual (with respect to the graph norm) of $D(A)$ (see Theorem 3.1); \bar{A} is nothing else that the closed extension of A to the whole Hilbert space \mathcal{H} and T_Θ is explicitly given in terms of the maps τ and Θ giving the boundary conditions. This result gives an extension, and a rephrasing in terms of boundary conditions, of the results obtained in [10] (and references therein, in particular [13]), where A is strictly positive and \mathcal{N} is closed in $D(A^{1/2})$ (see Remark 3.5). As regards boundary conditions the reader is also referred to [9], where $A = -\Delta + \lambda$, $\lambda > 0$, \mathcal{N} the kernel of the evaluation map along a regular submanifold, and to [17], where A is an arbitrary injective self-adjoint operator.

Successively, in Section 4, we study the connection of the self-adjoint extensions defined in the previous sections with the ones given by von Neumann's theory [15]. We prove (see Theorem 4.1) that the operator $\tilde{A} = \bar{A} + T$ defined in Theorem 3.4, of which the self-adjoint $A_\Theta = \bar{A} + T_\Theta$ is a restriction, coincides with $A_{\mathcal{N}}^*$; moreover we explicitly define a map on self-adjoint operators $\Theta : D(\Theta) \subseteq \mathfrak{h} \rightarrow \mathfrak{h}$ to unitary operators $U : \mathcal{K}_+ \rightarrow \mathcal{K}_-$ such that $A_\Theta = A_U$, where A_U denotes the von Neumann's extension corresponding to U . Such correspondence is then explicitly inverted (see Theorem 4.3). This shows (see Corollary 4.4) that $\tilde{A} = \bar{A} + T$ coincides with a self-adjoint extension \hat{A} of $A_{\mathcal{N}}$ such that $D(\hat{A}) \cap D(A) = \mathcal{N}$ if and only if the boundary condition $\tau\phi_* = \Theta Q_\phi$ holds for some self-adjoint operator Θ .

In Section 5 we conclude with some examples both in the case of finite and infinite deficiency indices. Example 5.1 (also see Remark 4.2) shows that, in the case $\dim \mathcal{K}_\pm < +\infty$, our results reproduce the theory of finite rank perturbations as given in [3], Section 3.1, and thus they can be viewed as an extension of such a theory to the infinite rank case. In Example 5.2 we give two examples in the infinite rank case: infinitely many point interaction in three dimensions and singular perturbations, supported on d -sets with $0 < n - d < 2s$, of translation invariant pseudo-differential operators with domain the Sobolev space $H^s(\mathbb{R}^n)$.

Notations and definitions

- Given a Banach space \mathcal{X} we denote by \mathcal{X}' its strong dual.
- $L(\mathcal{X}, \mathcal{Y})$ denotes the space of linear operators from the Banach space \mathcal{X} to the Banach space \mathcal{Y} ; $L(\mathcal{X}) \equiv L(\mathcal{X}, \mathcal{X})$.
- $B(\mathcal{X}, \mathcal{Y})$ denotes the Banach space of bounded, everywhere defined, linear operators on the Banach space \mathcal{X} to the Banach space \mathcal{Y} ; $B(\mathcal{X}) \equiv B(\mathcal{X}, \mathcal{X})$.
- Given $A \in L(\mathcal{X}, \mathcal{Y})$ densely defined, the closed operator $A' \in L(\mathcal{Y}', \mathcal{X}')$ is the adjoint of A i.e.

$$\forall \phi \in D(A) \subseteq \mathcal{X}, \quad \forall \lambda \in D(A') \subseteq \mathcal{Y}', \quad (A'\lambda)(\phi) = \lambda(A\phi).$$

- If \mathcal{H} is a complex Hilbert space with scalar product (conjugate-linear with respect to the first variable) $\langle \cdot, \cdot \rangle$, then $C_{\mathcal{H}} : \mathcal{H} \rightarrow \mathcal{H}'$ denotes the conjugate-linear isomorphism defined by

$$(C_{\mathcal{H}} \psi)(\phi) := \langle \psi, \phi \rangle.$$

- The Hilbert adjoint $A^* \in \mathcal{L}(\mathcal{H}_2, \mathcal{H}_1)$ of the densely defined linear operator $A \in \mathcal{L}(\mathcal{H}_1, \mathcal{H}_2)$ is defined as

$$A^* := C_{\mathcal{H}_1}^{-1} \cdot A' \cdot C_{\mathcal{H}_2}.$$

- F and $*$ denote Fourier transform and convolution respectively.
- $H^s(\mathbb{R}^n)$, $s \in \mathbb{R}$, is the usual scale of Sobolev-Hilbert spaces, i.e. $H^s(\mathbb{R}^n)$ is the space of tempered distributions with a Fourier transform which is square integrable with respect to the measure with density $(1 + |x|^2)^s$.

2. – Extensions by a Kreĭn-like formula

Given the Hilbert space \mathcal{H} with scalar product $\langle \cdot, \cdot \rangle$ (we denote by $\|\cdot\|$ the corresponding norm and put $C \equiv C_{\mathcal{H}}$), let $A : D(A) \subseteq \mathcal{H} \rightarrow \mathcal{H}$ be a self-adjoint operator and let $\mathcal{N} \subsetneq D(A)$ be a linear dense set which is closed with respect to the graph norm on $D(A)$. We denote by \mathcal{H}_+ the Hilbert space given by the set $D(A)$ equipped with the scalar product $\langle \cdot, \cdot \rangle_+$ leading to the graph norm, i.e.

$$\langle \phi_1, \phi_2 \rangle_+ := \langle (A^2 + 1)^{1/2} \phi_1, (A^2 + 1)^{1/2} \phi_2 \rangle.$$

We remark that in the sequel we will avoid to identify \mathcal{H}_+ with its dual. Indeed we will use the duality map induced by the scalar product on \mathcal{H} (see the next section for the details).

Being \mathcal{N} closed we have $\mathcal{H}_+ = \mathcal{N} \oplus \mathcal{N}^\perp$ and we can then consider the orthogonal projection $\pi : \mathcal{H}_+ \rightarrow \mathcal{N}^\perp$. From now on, since this gives advantages in concrete applications where usually a variant of π is what is known in advance, more generally we will consider a linear map

$$\tau : \mathcal{H}_+ \rightarrow \mathfrak{h}, \quad \tau \in \mathcal{B}(\mathcal{H}_+, \mathfrak{h}),$$

where \mathfrak{h} is a Hilbert space with scalar product $\langle \cdot, \cdot \rangle_{\mathfrak{h}}$, such that

$$(2.1) \quad \text{Range } \tau = \mathfrak{h}$$

and

$$(2.2) \quad \overline{\text{Kernel } \tau} = \mathcal{H},$$

the bar denoting here the closure in \mathcal{H} .

We put

$$\mathcal{N} := \text{Kernel } \tau.$$

By (2.1) one has $\mathfrak{h} \simeq \mathcal{H}_+ / \text{Kernel } \tau \simeq \mathcal{N}^\perp$ so that

$$\mathcal{H}_+ \simeq \mathcal{N} \oplus \mathfrak{h}.$$

Regarding (2.2) we have the following

LEMMA 2.1. *Hypothesis (2.2) is equivalent to*

$$\text{Range } \tau' \cap \mathcal{H}' = \{0\},$$

when one uses the embedding of \mathcal{H}' into $\mathcal{H}'_+ \supseteq \text{Range } \tau'$ given by the map $\phi \mapsto \langle C^{-1}\phi, \cdot \rangle$.

PROOF. Defining as usual the annihilator of \mathcal{N} by

$$\mathcal{N}^0 := \{\lambda \in \mathcal{H}'_+ : \forall \phi \in \mathcal{N}, \lambda(\phi) = 0\}$$

one has that denseness of \mathcal{N} is equivalent to

$$\mathcal{N}^0 \cap \mathcal{H}' = \{0\}.$$

Since $\overline{\text{Range } \tau'} = \mathcal{N}^0$ the proof is concluded if the range of τ' is closed. This follows from the closed range theorem since the range of τ is closed by the surjectivity hypothesis. \square

Being $\rho(A)$ the resolvent set of A , we define $R(z) \in \mathbb{B}(\mathcal{H}, \mathcal{H}_+)$, $z \in \rho(A)$, by

$$R(z) := (-A + z)^{-1}$$

and we then introduce, for any $z \in \rho(A)$, the two linear operators $\check{G}(z) \in \mathbb{B}(\mathcal{H}, \mathfrak{h})$ and $G(z) \in \mathbb{B}(\mathfrak{h}, \mathcal{H})$ by

$$\check{G}(z) := \tau \cdot R(z), \quad G(z) := \check{G}(\bar{z})^*.$$

By (2.2) one has

$$(2.3) \quad \text{Range } G(z) \cap D(A) = \{0\},$$

and, as an immediate consequence of the first resolvent identity for $R(z)$ (see [16], Lemma 2.1)

$$(2.4) \quad (z - w) R(w) \cdot G(z) = G(w) - G(z).$$

These relations imply

$$(2.5) \quad \text{Range } (G(w) - G(z)) \subseteq D(A)$$

and

$$\text{Range } (G(w) + G(z)) \cap D(A) = \{0\}.$$

By [16] (combining Theorem 2.1, Proposition 2.1, Lemma 2.2, Remarks 2.10, 2.12 and 2.13) one then obtains the following

THEOREM 2.2. *Given $z_0 \in \mathbb{C} \setminus \mathbb{R}$ define*

$$G_\star := \frac{1}{2} (G(z_0) + G(\bar{z}_0)) \quad G_\diamond := \frac{1}{2} (G(z_0) - G(\bar{z}_0))$$

and, given then any self-adjoint operator $\Theta : D(\Theta) \subseteq \mathfrak{h} \rightarrow \mathfrak{h}$, define

$$R_\Theta(z) := R(z) + G(z) \cdot (\Theta + \Gamma(z))^{-1} \cdot \check{G}(z), \quad z \in W_\Theta \cup \mathbb{C} \setminus \mathbb{R},$$

where

$$\Gamma(z) := \tau \cdot (G_\star - G(z))$$

and

$$W_\Theta := \{\lambda \in \mathbb{R} \cap \rho(A) : 0 \in \rho(\Theta + \Gamma(\lambda))\}.$$

Then R_Θ is the resolvent of the self-adjoint extension of $A_{\mathcal{N}}$ defined by

$$\begin{aligned} D(A_\Theta) &:= \{\phi \in \mathcal{H} : \phi = \phi_\star + G_\star Q_\phi, \phi_\star \in D(A), Q_\phi \in D(\Theta), \tau \phi_\star = \Theta Q_\phi\}, \\ A_\Theta \phi &:= A \phi_\star + \operatorname{Re}(z_0) G_\star Q_\phi + i \operatorname{Im}(z_0) G_\diamond Q_\phi. \end{aligned}$$

PROOF. Here we just give the main steps of the proof referring to [16], Section 2, for the details. One starts writing the presumed resolvent of an extension \tilde{A} of $A_{\mathcal{N}}$ as

$$\tilde{R}(z) = R(z) + B(z) \cdot \tau \cdot R(z) \equiv R(z) + B(z) \cdot \check{G}(z),$$

where $B(z) \in \mathcal{B}(\mathfrak{h}, \mathcal{H})$ has to be determined. Self-adjointness requires $\tilde{R}(z)^* = \tilde{R}(\bar{z})$ or, equivalently,

$$(2.6) \quad G(\bar{z}) \cdot B(z)^* = B(\bar{z}) \cdot \check{G}(\bar{z}).$$

Therefore posing $B(z) = G(z) \cdot \Lambda(z)$, where $\Lambda(z) \in \mathcal{B}(\mathfrak{h})$, (2.6) is equivalent to

$$(2.7) \quad \Lambda(z)^* = \Lambda(\bar{z}).$$

The resolvent identity

$$(2.8) \quad (z - w) \tilde{R}(w) \tilde{R}(z) = \tilde{R}(w) - \tilde{R}(z)$$

is then equivalent to

$$(2.9) \quad \Lambda(w) - \Lambda(z) = (z - w) \Lambda(w) \cdot \check{G}(w) \cdot G(z) \cdot \Lambda(z).$$

Suppose now that there exist a (necessarily closed) operator

$$\Gamma(z) : D \subseteq \mathfrak{h} \rightarrow \mathfrak{h}$$

and an open set $Z \subseteq \rho(A)$, invariant with respect to complex conjugation, such that

$$\forall z \in Z, \quad \Gamma(z)^{-1} = \Lambda(z).$$

Then (2.9) forces $\Gamma(z)$ to satisfy the relation

$$(2.10) \quad \Gamma(z) - \Gamma(w) = (z - w) \check{G}(w) \cdot G(z),$$

whereas (2.7), at least in the case $\Gamma(z)$ is densely defined, and has a bounded inverse given by $\Lambda(z)$ as we are pretending, is equivalent to

$$(2.11) \quad \Gamma(z)^* = \Gamma(\bar{z}).$$

By [16], Lemma 2.2, for any self-adjoint Θ , the linear operator

$$\Theta + \tau \cdot (G_\star - G(z))$$

satisfies (2.10), (2.11) and, by [16], Proposition 2.1, has a bounded inverse for any $z \in W_\Theta \cup \mathbb{C} \setminus \mathbb{R}$ (at this point hypothesis (2.1) is used). Therefore (see the proof of Theorem 2.1 in [16])

$$R_\Theta(z) := R(z) + G(z) \cdot (\Theta + \Gamma(z))^{-1} \cdot \check{G}(z)$$

is the resolvent of a self-adjoint operator A_Θ (here hypotheses (2.2) is needed). For any $z \in W_\Theta \cup \mathbb{C} \setminus \mathbb{R}$ one has

$$(2.12) \quad D(A_\Theta) = \{\phi \in \mathcal{H} : \phi = \phi_z + G(z) \cdot (\Gamma + \Theta(z))^{-1} \cdot \tau \phi_z, \phi_z \in D(A)\},$$

$$(2.13) \quad (-A_\Theta + z)\phi = (-A + z)\phi_z,$$

the definition of A_Θ being z -independent thanks to resolvent identity (2.8). Being $G(z)$ injective, (2.3) and (2.5) imply

$$\phi_w + G(w)Q_1 = \phi_z + G(z)Q_2 \quad \Rightarrow \quad Q_1 = Q_2$$

and so the definition

$$Q_\phi := (\Theta + \Gamma(z))^{-1} \cdot \tau \phi_z$$

is z -independent. Therefore any $\phi \in D(A_\Theta)$ can be equivalently re-written as

$$\phi = \phi_z + G(z)Q_\phi,$$

where $Q_\phi \in D(\Theta)$ and

$$\tau \phi_z = \Theta Q_\phi + \Gamma(z)Q_\phi.$$

This implies, for any $\phi \in D(A_\Theta)$,

$$\begin{aligned}\phi &= \frac{1}{2}(\phi_{z_0} + G(z_0)Q_\phi + \phi_{\bar{z}_0} + G(\bar{z}_0)Q_\phi) \equiv \phi_\star + G_\star Q_\phi, \\ \tau\phi_\star &\equiv \frac{1}{2}\tau(\phi_{z_0} + \phi_{\bar{z}_0}) = \Theta Q_\phi + \frac{1}{2}(\Gamma(z_0)Q_\phi + \Gamma(\bar{z}_0)Q_\phi) = \Theta Q_\phi, \\ A_\Theta\phi &= \frac{1}{2}(A\phi_{z_0} + z_0G(z_0)Q_\phi + A\phi_{\bar{z}_0} + \bar{z}_0G(\bar{z}_0)Q_\phi) \\ &\equiv A\phi_\star + \operatorname{Re}(z_0)G_\star Q_\phi + i\operatorname{Im}(z_0)G_\diamond Q_\phi.\end{aligned}$$

Conversely any $\phi = \phi_\star + G_\star Q_\phi$, $\phi_\star \in D(A)$, $\Theta Q = \tau\phi_\star$, admits the decomposition $\phi = \phi_z + G(z) \cdot (\Theta + \Gamma(z))^{-1} \cdot \tau\phi_z$, where

$$\phi_z := \phi_\star + (G_\star - G(z))Q_\phi.$$

Note that $\phi_z \in D(A)$ by (2.5) and $\tau\phi_z = (\Theta + \Gamma(z))Q_\phi$. \square

REMARK 2.3. The results quoted in the previous theorem are consequences of an alternative version of Kreĭn's resolvent formula. The original one was obtained in [11], [12], [18] for the cases where $\dim \mathcal{K}_\pm = 1$, $\dim \mathcal{K}_\pm < +\infty$, $\dim \mathcal{K}_\pm = +\infty$ respectively; also see [4], [6], [14] for more recent formulations. In standard Kreĭn's formula (usually written with $z_0 = i \equiv \sqrt{-1}$) the main ingredient is the orthogonal projection $P : \mathcal{H} \rightarrow \mathcal{K}_+$ whereas we used, exploiting the a priori knowledge of the self-adjoint operator A , the map τ , which plays the role of the orthogonal projection $\pi : \mathcal{H}_+ \rightarrow \mathcal{N}^\perp$. Thus the knowledge of $A_\mathcal{N}^*$ is not needed. The version given in [16] allows τ to be not surjective and \mathfrak{h} can be a Banach space; the use of the map τ simplifies the exposition and makes easier to work out concrete applications. Indeed, as we already said, frequently what is explicitly known is the map τ and \mathcal{N} is then simply defined as its kernel: see the many examples in [16] where τ is the trace (restriction) map along some null subset of \mathbb{R}^n and A is a (pseudo-)differential operator. Moreover this approach allows a natural formulation in terms of the boundary condition $\tau\phi_\star = \Theta Q_\phi$. Note that, since $G_\star Q_\phi \in D(A)$ if and only if $Q_\phi = 0$, once the reference point z_0 has been chosen, the decomposition $\phi = \phi_\star + G_\star Q_\phi$ of a generic element ϕ of $D(A_\Theta)$ by a regular part $\phi_\star \in D(A)$ and a singular one $G_\star Q_\phi \in \mathcal{H} \setminus D(A)$ is univocal.

REMARK 2.4. As regards the definition of $R_\Theta(z)$, the one given in the theorem above is not the only possible definition of the operator $\Gamma(z)$. Any other not necessarily bounded, densely defined operator satisfying

$$\begin{aligned}\Gamma(z) - \Gamma(w) &= (z - w)\check{G}(w) \cdot G(z), \\ \Gamma(\bar{z}) &= \Gamma(z)^*\end{aligned}$$

and such that $\Theta + \Gamma(z)$ is boundedly invertible would suffice; moreover hypothesis (2.1) is not necessary (see [16], Theorem 2.1); note that, once Θ is

given, $\Gamma(z)$ univocally defines $(-A_\Theta + z)^{-1}$ and hence A_Θ itself. For alternative choices of $\Gamma(z)$ we refer to [16]; also see [17] where it is shown how, under the hypotheses $\text{Kernel } A = \{0\}$ and $\|\tau\phi\|_0 \leq c\|A\phi\|$, it is always possible to take $z_0 = 0$ in Theorem 2.1 (at the expense of having then ϕ_\star in the completion of $D(A)$ with respect to the norm $\phi \mapsto \|A\phi\|$). However we remark that any different choice (either of z_0 or of the operator $\Gamma(z)$ itself) does not change the family of extensions as a whole.

REMARK 2.5. In the case A has a non-empty real resolvent set, by [16], Remark 2.7, if in Theorem 2.1 one consider only the sub-family of extensions in which the Θ 's have bounded inverses, then one can take $z_0 \in \mathbb{R} \cap \rho(A)$. More generally one can take $z_0 \in W_\Theta$ independently of the invertibility of Θ ; however this could give rise to implicit conditions (related to the location of the spectrum of A_Θ) on the choice of z_0 .

3. – Extensions by Additive Perturbations

We define the pre-Hilbert space $\tilde{\mathcal{H}}_-$ as the set \mathcal{H} equipped with the scalar product

$$\langle \phi_1, \phi_2 \rangle_- := \langle (A^2 + 1)^{-1/2} \phi_1, (A^2 + 1)^{-1/2} \phi_2 \rangle.$$

We denote then by \mathcal{H}_- the Hilbert space given by the completion of $\tilde{\mathcal{H}}_-$. We will avoid to identify \mathcal{H}_+ and \mathcal{H}_- with their duals; indeed, see Lemma 3.1 below, we will identify \mathcal{H}'_+ with \mathcal{H}_- .

As usual \mathcal{H} will be treated as a (dense) subspace of \mathcal{H}_- by means of the canonical embedding

$$I_- : \mathcal{H} \rightarrow \mathcal{H}_-$$

which associates to ϕ the set of all the Cauchy sequences converging to ϕ . Considering also the canonical embedding (with dense range)

$$I_+ : \mathcal{H}' \rightarrow \mathcal{H}'_+, \quad I_+ \lambda(\phi) := \langle C^{-1} \lambda, \phi \rangle,$$

we can then define the conjugate linear operator

$$C_- : \mathcal{H}'_+ \rightarrow \mathcal{H}_-$$

as the unique bounded extension of

$$I_- \cdot C^{-1} \cdot I_+^{-1} : I_+(\mathcal{H}') \subseteq \mathcal{H}'_+ \rightarrow \mathcal{H}_-.$$

Analogously we define the conjugate linear operator

$$C_+ : \mathcal{H}_- \rightarrow \mathcal{H}'_+$$

as the unique bounded extension of

$$I_+ \cdot C \cdot I_-^{-1} : I_-(\mathcal{H}) \subseteq \mathcal{H}_- \rightarrow \mathcal{H}'_+.$$

These definitions immediately lead to the following

LEMMA 3.1. *One has*

$$C_+ = C_-^{-1}, \quad C_- = C_+^{-1},$$

so that

$$\mathcal{H}'_+ \simeq \mathcal{H}_-.$$

We will denote by

$$(\cdot, \cdot) : \mathcal{H}_- \times \mathcal{H}_+ \rightarrow \mathbb{C}, \quad (\varphi, \phi) := C_+ \varphi(\phi)$$

the pairing between \mathcal{H}_- and \mathcal{H}_+ . It is nothing else than the extension of the scalar product of \mathcal{H} , being

$$(I_- \phi_1, \phi_2) = \langle \phi_1, \phi_2 \rangle.$$

We consider now the linear operator

$$I_- \cdot A : \mathcal{H}_+ \subseteq \mathcal{H} \rightarrow \mathcal{H}_-.$$

Since

$$\|(A^2 + 1)^{-1/2} A \phi\| \leq \|\phi\|,$$

the operator $I_- \cdot A$ has a unique extension

$$\bar{A} : \mathcal{H} \rightarrow \mathcal{H}_-, \quad \bar{A} \in \mathcal{B}(\mathcal{H}, \mathcal{H}_-).$$

LEMMA 3.2. *Let $A' : \mathcal{H}' \rightarrow \mathcal{H}'_+$ be the adjoint of the linear operator A when viewed as an element of $\mathcal{B}(\mathcal{H}_+, \mathcal{H})$. Then one has*

$$\bar{A} = C_- \cdot A' \cdot C.$$

PROOF. Being I_- injective, by continuity and density the thesis follows from the identity

$$A = A^* \equiv C^{-1} \cdot A' \cdot C. \quad \square$$

REMARK 3.3. If we use the symbol A_+ to denote the linear operator A when we consider it as an element of $\mathcal{B}(\mathcal{H}_+, \mathcal{H})$, and if we use C_- as a substitute of $C_{\mathcal{H}_+}^{-1}$, then by Lemma 3.2 and a slight abuse of notations we can write

$$\bar{A} = A_+^*.$$

By the same abuse of notations we define $\tau^* \in \mathcal{B}(\mathfrak{h}, \mathcal{H}_-)$ by

$$\tau^* := C_- \cdot \tau' \cdot C_{\mathfrak{h}}.$$

Now we can reformulate Theorem 2.1 in terms of additive perturbations:

THEOREM 3.4. *Define*

$$D(\tilde{A}) := \{\phi \in \mathcal{H} : \phi = \phi_\star + G_\star Q_\phi, \phi_\star \in D(A), Q_\phi \in \mathfrak{h}\},$$

$$\tilde{A} : D(\tilde{A}) \rightarrow \mathcal{H}_-, \quad \tilde{A} := \bar{A} + T,$$

where

$$T : D(\tilde{A}) \rightarrow \mathcal{H}_-, \quad T\phi := \tau^\star Q_\phi.$$

Then the linear operator \tilde{A} is \mathcal{H} -valued and coincides with A_Θ when restricted to $D(A_\Theta)$, i.e. when a boundary condition of the kind $\tau\phi_\star = \Theta Q_\phi$ holds for some self-adjoint operator Θ . Therefore, posing $T_\Theta := T|_{D(A_\Theta)}$, one has

$$A_\Theta : D(A_\Theta) \rightarrow \mathcal{H}, \quad A_\Theta := \bar{A} + T_\Theta,$$

and, in the case Θ has a bounded inverse,

$$A_\Theta : D(A_\Theta) \rightarrow \mathcal{H}, \quad A_\Theta\phi = \bar{A}\phi + V_\Theta\phi_\star,$$

where

$$V_\Theta : \mathcal{H}_+ \rightarrow \mathcal{H}_-, \quad (V_\Theta\phi_1, \phi_2) := \langle \Theta^{-1}\tau\phi_1, \tau\phi_2 \rangle_{\mathfrak{h}}.$$

PROOF. By the definition of \bar{A} , τ^\star and G_\star one has, for any $\phi \in D(\tilde{A})$,

$$\begin{aligned} \bar{A}\phi &= I_- \cdot A\phi_\star + C_- \cdot A' \cdot C \cdot G_\star Q_\phi \\ &= I_- \cdot A\phi_\star + \frac{1}{2} C_- \cdot A' \cdot R(\bar{z}_0)' \cdot \tau' \cdot C_{\mathfrak{h}} Q_\phi \\ &\quad + \frac{1}{2} C_- \cdot A' \cdot R(z_0)' \cdot \tau' \cdot C_{\mathfrak{h}} Q_\phi \\ &= I_- \cdot (A\phi_\star + \operatorname{Re}(z_0) G_\star Q_\phi + i \operatorname{Im}(z_0) G_\diamond Q_\phi) - T Q_\phi. \end{aligned}$$

The proof is then concluded by Theorem 2.1. \square

REMARK 3.5. In the case $0 \in \rho(A)$ and Θ is boundedly invertible, by Theorem 2.1 and Remark 2.4 (taking $z_0 = 0$) one can define A_Θ either by $A_\Theta\phi := A\phi_\star$ or, equivalently, by

$$A_\Theta^{-1} = A^{-1} + G \cdot \Theta^{-1} \cdot \check{G},$$

where $G := G(0)$, $\check{G} := \check{G}(0)$. Since, for any $\phi_1, \phi_2 \in \mathcal{H}$, one has

$$\begin{aligned} \langle \bar{A}^{-1} \cdot V_\Theta \cdot A^{-1}\phi_1, \phi_2 \rangle &= \langle V_\Theta A^{-1}\phi_1, A^{-1}\phi_2 \rangle \\ &= \langle \Theta^{-1}\tau \cdot A^{-1}\phi_1, \tau \cdot A^{-1}\phi_2 \rangle_{\mathfrak{h}} = \langle \Theta^{-1}\check{G}\phi_1, \check{G}\phi_2 \rangle_{\mathfrak{h}} \\ &= \langle G \cdot \Theta^{-1} \cdot \check{G}\phi_1, \phi_2 \rangle, \end{aligned}$$

the self-adjoint extension A_Θ could be defined directly in terms of V_Θ by

$$A_\Theta^{-1} = A^{-1} + \bar{A}^{-1} \cdot V_\Theta \cdot A^{-1}.$$

This reproduces the formulae appearing in [2], Lemma 2.3, where however no additive representation of the extension A_Θ is given, and in [10] where an additive representation is obtained only when \mathcal{N} is closed in $D(A^{1/2})$.

4. – The connection with von Neumann's Theory

In this section we explore the connection between the results given in the previous sections and von Neumann's theory of self-adjoint extensions [15]. Such a theory (see e.g. [5], Section 13, for a very compact exposition) tells us that

$$D(A_{\mathcal{N}}^*) = \mathcal{N} \oplus \mathcal{K}_+ \oplus \mathcal{K}_-, \quad A_{\mathcal{N}}^*(\phi_0 + \phi_+ + \phi_-) = A\phi_0 + i\phi_+ - i\phi_-,$$

the direct sum decomposition being orthogonal with respect to the graph inner product of $A_{\mathcal{N}}^*$; any self-adjoint extension A_U of $A_{\mathcal{N}}$ is then obtained by restricting $A_{\mathcal{N}}^*$ to a subspace of the kind $\mathcal{N} \oplus \text{Graph } U$, where $U : \mathcal{K}_+ \rightarrow \mathcal{K}_-$ is unitary.

For simplicity in the next theorem we will consider only the case $z_0 = i$ and we put $G_{\pm} := G(\pm i)$ and $\Gamma := \Gamma(i)$.

THEOREM 4.1. *Let $\tilde{A} = \bar{A} + T$ as defined in Theorem 3.4. Then*

$$\tilde{A} = A_{\mathcal{N}}^*.$$

The linear operator

$$G_{\pm} : \mathfrak{h} \rightarrow \mathcal{K}_{\pm}$$

is a continuous bijection which becomes unitary when one puts on \mathfrak{h} the scalar product

$$\langle Q_1, Q_2 \rangle_{\Gamma} := \langle \sqrt{-i\Gamma} Q_1, \sqrt{-i\Gamma} Q_2 \rangle_{\mathfrak{h}}.$$

The linear operator

$$U : \mathcal{K}_+ \rightarrow \mathcal{K}_-, \quad U := -G_- \cdot (1 + 2(\Theta - \Gamma)^{-1} \cdot \Gamma) \cdot G_+^{-1}$$

is unitary and the corresponding von Neumann's extension A_U coincides with the self-adjoint operator A_{Θ} defined in Theorems 2.1 and 3.4.

PROOF. By the definition of $\check{G}_{\pm} \equiv \check{G}(\pm i)$ one has

$$\text{Range}(-A_{\mathcal{N}} \pm i) = \text{Kernel } \check{G}_{\pm}$$

and so, since

$$\mathcal{K}_{\pm} = \text{Range}(-A_{\mathcal{N}} \mp i)^{\perp}$$

and

$$\text{Range } G_{\pm}^{\perp} = \text{Kernel } \check{G}_{\mp},$$

in conclusion there follows

$$\text{Range } G_{\pm} = \mathcal{K}_{\pm}$$

if and only if $\text{Range } G_{\pm}$ is closed. By the closed range theorem $\text{Range } G_{\pm}$ is closed if and only if $\text{Range } \check{G}_{\pm}$ is closed, and this is equivalent to the range of τ being closed. Being τ surjective, G_{\pm} is injective with a closed range and so

$$G_{\pm} : \mathfrak{h} \rightarrow \mathcal{K}_{\pm}$$

is a bijection.

By von Neumann's theory we know that any $\phi \in D(A_{\mathcal{N}}^*)$ can be univocally decomposed as

$$\phi = \phi_0 + \phi_+ + \phi_-, \quad \phi_0 \in \mathcal{N}, \quad \phi_{\pm} \in \mathcal{K}_{\pm},$$

i.e.

$$\phi = \phi_0 + G_+ Q_+ + G_- Q_-, \quad \phi_0 \in \mathcal{N}, \quad Q_{\pm} \in \mathfrak{h}.$$

The above decomposition can be then rearranged as

$$\begin{aligned} \phi &= \phi_0 + \frac{1}{2} (G_+ - G_-) Q_+ + \frac{1}{2} (G_+ + G_-) Q_+ \\ &\quad + \frac{1}{2} (G_- - G_+) Q_- + \frac{1}{2} (G_- + G_+) Q_- \\ &= \phi_0 + \frac{1}{2} (G_- - G_+) (Q_- - Q_+) + G_{\star} (Q_- + Q_+). \end{aligned}$$

By (2.4) one has

$$(4.1) \quad G_{\mp} - G_{\pm} = \pm 2i R(\mp i) \cdot G_{\pm}.$$

Since the scalar product of \mathcal{H}_+ can be equivalently written as

$$\langle \phi_1, \phi_2 \rangle_+ = \langle (-A + i)\phi_1, (-A + i)\phi_2 \rangle,$$

one has

$$G_- - G_+ = 2i R(-i) \cdot R(-i)^* \cdot \tau^* = 2i \tau^*.$$

This implies, since $\text{Range } G_+$ is closed,

$$\text{Range } (G_- - G_+) = \text{Range } \tau^* = \text{Kernel } \tau^{\perp}.$$

Thus, being $\mathcal{H}_+ = \mathcal{N} \oplus \mathcal{N}^{\perp}$, the vector

$$\phi_0 + \frac{1}{2} (G_- - G_+) (Q_- - Q_+)$$

is a generic element of $D(A)$ and we have shown that $D(\tilde{A}) = D(A_{\mathcal{N}}^*)$. It is then straightforward to check that $\tilde{A} = A_{\mathcal{N}}^*$.

By (4.1) one has

$$\Gamma = \pm \frac{1}{2} \tau \cdot (G_{\mp} - G_{\pm}) = i \tau \cdot R(\mp i) \cdot G_{\pm} = i \check{G}_{\mp} \cdot G_{\pm} = i G_{\pm}^* \cdot G_{\pm}.$$

This implies

$$\|G_{\pm} Q\| = \|\sqrt{-i\Gamma} Q\|_{\mathfrak{h}},$$

thus $U = -G_- \cdot (1 + 2(\Theta - \Gamma)^{-1} \cdot \Gamma) \cdot G_+^{-1}$ is isometric if and only if

$$\forall Q \in \mathfrak{h}, \quad \|\sqrt{-i\Gamma} \cdot \tilde{U} Q\|_{\mathfrak{h}} = \|\sqrt{-i\Gamma} Q\|_{\mathfrak{h}},$$

where $\tilde{U} := G_-^{-1} \cdot U \cdot G_+$. By using the identities $\Gamma^* = -\Gamma$ and

$$(4.2) \quad (\Theta - \Gamma)^{-1} - (\Theta + \Gamma)^{-1} = 2(\Theta + \Gamma)^{-1} \cdot \Gamma \cdot (\Theta - \Gamma)^{-1},$$

one has

$$\begin{aligned} & i \|\sqrt{-i\Gamma} \cdot (1 + 2(\Theta - \Gamma)^{-1} \cdot \Gamma) Q\|_{\mathfrak{h}} \\ &= \langle \Gamma Q + 2\Gamma \cdot (\Theta - \Gamma)^{-1} \cdot \Gamma Q, Q + 2(\Theta - \Gamma)^{-1} \cdot \Gamma Q \rangle \\ &= \langle \Gamma Q, Q \rangle + 2\langle \Gamma Q, (\Theta - \Gamma)^{-1} \cdot \Gamma Q \rangle + 2\langle \Gamma \cdot (\Theta - \Gamma)^{-1} \cdot \Gamma Q, Q \rangle \\ & \quad + 4\langle \Gamma \cdot (\Theta - \Gamma)^{-1} \cdot \Gamma Q, (\Theta - \Gamma)^{-1} \cdot \Gamma Q \rangle \\ &= \langle \Gamma Q, Q \rangle + 2\langle \Gamma Q, ((\Theta - \Gamma)^{-1} - (\Theta + \Gamma)^{-1}) \cdot \Gamma Q \rangle \\ & \quad - 4\langle \Gamma Q, (\Theta + \Gamma)^{-1} \cdot \Gamma \cdot (\Theta - \Gamma)^{-1} \cdot \Gamma Q \rangle \\ &= \langle \Gamma Q, Q \rangle = i \|\sqrt{-i\Gamma} Q\|_{\mathfrak{h}}, \end{aligned}$$

and so U is an isometry. By again using identity (4.2) one can check that U has an inverse defined by

$$U^{-1} := -G_+ \cdot (1 - 2(\Theta + \Gamma)^{-1} \cdot \Gamma) \cdot G_-^{-1}.$$

Thus U is unitary. Let us now take $G_- Q_- = U G_+ Q_+$. Then

$$-2(\Theta - \Gamma)^{-1} \cdot \Gamma Q_+ = Q_- + Q_+$$

and so $Q_- + Q_+ \in D(\Theta)$ and

$$\tau \left(\phi_0 + \frac{1}{2} (G_- - G_+) (Q_- - Q_+) \right) \equiv \Gamma(Q_- - Q_+) = \Theta(Q_- + Q_+). \quad \square$$

REMARK 4.2. Note that when Θ is bounded, in the previous theorem one can re-write the unitary U as

$$U = -G_- \cdot (\Theta - \Gamma)^{-1} \cdot (\Theta + \Gamma) \cdot G_+^{-1}.$$

Being Θ always bounded when $\dim \mathcal{K}_{\pm} = n$, the previous theorem gives an analogue of Theorem 3.1.2 in [3] avoiding however the use of an admissible matrix R (see [3], Definition 3.1.2).

The previous theorem has the following converse:

THEOREM 4.3. *Let A_U be a self-adjoint extension of A_N as given by von Neumann's theory. Suppose that $D(A_U) \cap D(A) = \mathcal{N}$ and let $U_A := (-A + i) \cdot (-A - i)^{-1}$ be the Cayley transform of A . Then the set*

$$D(\Theta) := \text{Range } G_-^{-1} \cdot (U + U_A)$$

is dense,

$$\Theta : D(\Theta) \subseteq \mathfrak{h} \rightarrow \mathfrak{h}, \quad \Theta := i \check{G}_+ \cdot (U - U_A) \cdot (U + U_A)^{-1} \cdot G_- ,$$

is self-adjoint and the corresponding self-adjoint operator A_Θ , defined in Theorems 2.1 and 3.4, coincides with A_U .

PROOF. By (4.1) one has

$$G_- \cdot G_+^{-1} = 1 + 2i R(-i) = U_A .$$

Thus, by inverting the relation $U = -G_- \cdot (1 + 2(\Theta - \Gamma)^{-1} \cdot \Gamma) \cdot G_+^{-1}$ given in the previous theorem, one obtains

$$\begin{aligned} \Theta &= \Gamma \cdot (G_-^{-1} \cdot U \cdot G_+ - 1) \cdot (G_-^{-1} \cdot U \cdot G_+ + 1)^{-1} \\ &= \Gamma \cdot G_-^{-1} \cdot (U - G_- \cdot G_+^{-1}) \cdot (U + G_- \cdot G_+^{-1})^{-1} \cdot G_- \\ &= \Gamma \cdot G_-^{-1} \cdot (U - U_A) \cdot (U + U_A)^{-1} \cdot G_- . \end{aligned}$$

Since $U = -U_{A_U}$ and $1 \notin \sigma_p(U_A \cdot U_{A_U}^{-1})$ if and only if $D(A_U) \cap D(A) = \mathcal{N}$ (see e.g. [6], Lemma 1), the range of $U + U_A$ is dense and thus Θ is densely defined as G_- is a continuous bijection. By (4.1) one has

$$\Gamma \cdot G_-^{-1} = i\tau \cdot R(i) \equiv i\check{G}_+$$

and so, since $\check{G}_+^* = G_-$ and $G_-^* = \check{G}_+$, Θ is self-adjoint if and only if

$$(U^* + U_A^*) \cdot (U - U_A) = -(U^* - U_A^*) \cdot (U + U_A) .$$

Such an equality is then an immediate consequence of the unitarity of both U and U_A . \square

COROLLARY 4.4. $\tilde{A} = \bar{A} + T$ as defined in Theorem 3.4 coincides with a self-adjoint extension \hat{A} of A_N such that $D(\hat{A}) \cap D(A) = \mathcal{N}$ if and only if the boundary condition $\tau\phi_\star = \Theta Q_\phi$ holds for some self-adjoint operator $\Theta : D(\Theta) \subseteq \mathfrak{h} \rightarrow \mathfrak{h}$.

5. – Examples

EXAMPLE 5.1. Finite rank perturbations. Suppose $\dim \mathcal{K}_\pm = n$, so that $\mathfrak{h} \simeq \mathbb{C}^n$ and $\tau \in \mathcal{B}(\mathcal{H}_+, \mathbb{C}^n)$. Then necessarily

$$\tau : \mathcal{H}_+ \rightarrow \mathbb{C}^n, \quad \tau \phi = \{(\varphi_j, \phi)\}_1^n,$$

with $\varphi_1, \dots, \varphi_n \in \mathcal{H}_-$. Hypotheses (2.1) and (2.2) correspond to

$$\exists \phi_1, \dots, \phi_n \in \mathcal{H}_+ \quad \text{s.t.} \quad (\varphi_i, \phi_j) = \delta_{ij},$$

and

$$\sum_{j=1}^n c_j \varphi_j \in \mathcal{H} \quad \text{iff} \quad c_1 = \dots = c_n = 0.$$

Considering then an Hermitean invertible matrix $\Theta = (\theta_{ij})$ with inverse $\Theta^{-1} = (t_{ij})$, by Theorem 3.4 one can define the self-adjoint operator

$$A_\Theta \phi := \bar{A}\phi + \sum_{i,j=1}^n t_{ij}(\varphi_i, \phi_\star)\varphi_j$$

with

$$D(A_\Theta) := \left\{ \phi \in \mathcal{H} : \phi = \phi_\star + \sum_{j=1}^n Q_j R_\star \varphi_j, \phi_\star \in D(A), Q \in \mathbb{C}^n, (\varphi_i, \phi_\star) = \sum_{j=1}^n \theta_{ij} Q_j \right\},$$

where

$$R_\star := \frac{1}{2} (\hat{R}(z_0) + \hat{R}(\bar{z}_0)), \quad \hat{R}(z) : \mathcal{H}_- \rightarrow \mathcal{H}, \quad \langle \hat{R}(z)\varphi, \phi \rangle := (\varphi, R(\bar{z})\phi).$$

According to Theorem 2.1 its resolvent is given by

$$(-A_\Theta + z)^{-1} = (-A + z)^{-1} + \sum_{i,j=1}^n (\Theta + \Gamma(z))_{ij}^{-1} \hat{R}(z)\varphi_i \hat{R}(\bar{z})\varphi_j,$$

where

$$\Gamma(z)_{ij} = \frac{1}{2} (\varphi_i, (\hat{R}(z_0) + \hat{R}(z_0) - 2\hat{R}(z))\varphi_j).$$

The operator A_Θ above coincides with a generic finite rank perturbation of the self-adjoint operator A as defined in [3], Section 3.1. In order to realize that the resolvent written above (in the case $z_0 = i$) is the same given there, the identity

$$\frac{1}{2} (R(i) + R(-i) - 2R(z)) = (1 + zA) \cdot (A - z)^{-1} \cdot (A^2 + 1)^{-1}$$

has to be used.

The previous construction can be applied to the case of so-called point interactions in three dimensions (see [1] and references therein). Since in Example 5.2 below we will consider the case of infinitely many point interactions, here we just treat the simplest situation in which only one point interaction (placed at the origin) is present. In this case we take $A = \Delta$, $\mathcal{H} = L^2(\mathbb{R}^3)$, $\mathcal{H}_+ = H^2(\mathbb{R}^3)$, $\mathcal{H}_- = H^{-2}(\mathbb{R}^3)$, and $\varphi = \delta_0$. Therefore τ is simply the evaluation map at the origin

$$\tau : H^2(\mathbb{R}^3) \rightarrow \mathbb{C}, \quad \tau\phi = \phi(0),$$

and we have the family of self-adjoint operators Δ_θ , $\theta \in \mathbb{R} \setminus \{0\}$, defined as (we take $z_0 = i$)

$$\Delta_\theta\phi := \Delta\phi + \theta^{-1}\phi_\star(0)\delta_0$$

on the domain

$$D(\Delta_\theta) := \{\phi \in L^2(\mathbb{R}^3) : \phi = \phi_\star + Q\mathcal{G}_\star, \phi_\star \in H^2(\mathbb{R}^3), Q \in \mathbb{C}, \phi_\star(0) = \theta Q\},$$

where

$$\mathcal{G}_\star(x) = \cos \frac{|x|}{\sqrt{2}} \frac{e^{-|x|/\sqrt{2}}}{4\pi|x|}.$$

This reproduces the family given in [3], Section 1.5.1, and coincides with the family Δ_α given in [1], Section I.1.1, when one takes $\alpha = \theta - (4\pi\sqrt{2})^{-1}$. The case $\alpha = -(4\pi\sqrt{2})^{-1}$ can be then recovered by directly using Theorem 3.4 in the case $\theta = 0$.

EXAMPLE 5.2. Infinite rank perturbations. Suppose $\dim \mathcal{K}_\pm = +\infty$. Then (we suppose \mathcal{H} is separable) $\mathfrak{h} \simeq \ell^2(\mathbb{N})$, $\tau \in \mathcal{B}(\mathcal{H}_+, \ell^2(\mathbb{N}))$ and necessarily

$$\tau : \mathcal{H}_+ \rightarrow \ell^2(\mathbb{N}), \quad \tau\phi = \{(\varphi_j, \phi)\}_1^\infty,$$

with $\{\varphi_j\}_1^\infty \subset \mathcal{H}_-$. The generalization of the finite rank case to this situation is then evident. As concrete example one can consider infinitely many point interactions in three dimensions by taking $A = \Delta$, $\mathcal{H} = L^2(\mathbb{R}^3)$, $\mathcal{H}_+ = H^2(\mathbb{R}^3)$, $\mathcal{H}_- = H^{-2}(\mathbb{R}^3)$ as before and an infinite and countable set $Y \subset \mathbb{R}^3$ such that

$$\inf_{y \neq \tilde{y}} |y - \tilde{y}| = d > 0.$$

Defining then $\varphi_y := \delta_y$, by [1] (see page 172) one has

$$\tau \in \mathcal{B}(H^2(\mathbb{R}^3), \ell^2(Y)),$$

where

$$\tau : H^2(\mathbb{R}^3) \rightarrow \ell^2(Y), \quad \tau\phi = \{\phi(y)\}_{y \in Y},$$

and hypotheses (2.1) and (2.2) are an immediate consequence of the discreteness of Y (see [16], Example 3.4). By Theorem 3.4, given any invertible infinite

Hermitean matrix $\Theta = (\theta_{y\tilde{y}})$ with a bounded inverse $\Theta^{-1} = (t_{y\tilde{y}})$, one can then define the family of self-adjoint operators

$$\Delta_{\Theta}\phi := \Delta\phi + \sum_{y, \tilde{y} \in Y} t_{y\tilde{y}} \phi_{\star}(y) \delta_{\tilde{y}}$$

on the domain

$$D(\Delta_{\Theta}) := \left\{ \phi \in L^2(\mathbb{R}^3) : \phi = \phi_{\star} + \sum_{y \in Y} Q_y \mathcal{G}_{\star}^y, \phi_{\star} \in H^2(\mathbb{R}^3), Q \in D(\Theta), \phi_{\star}(y) = \sum_{\tilde{y} \in Y} \theta_{y\tilde{y}} Q_{\tilde{y}} \right\},$$

where $\mathcal{G}_{\star}^y(x) := \mathcal{G}_{\star}(x - y)$. When

$$\theta_{yy} = \alpha + \frac{1}{4\pi\sqrt{2}}, \quad \theta_{y\tilde{y}} = -\mathcal{G}_{\star}(y - \tilde{y}), \quad y \neq \tilde{y},$$

the self-adjoint extension Δ_{Θ} coincides with the operator $\Delta_{\alpha, Y}$ given in [1], Section III.1.1 (also see [16], Example 3.4).

In more general situations where the set Y is not discrete the use of the unitary isomorphism $\mathfrak{h} \simeq \ell^2(\mathbb{N})$ given no advantages and, how the following example shows, it is better to work with \mathfrak{h} itself.

Let $A = \Psi$, $\mathcal{H} = L^2(\mathbb{R}^n)$, $\mathcal{H}_+ = H^s(\mathbb{R}^n)$, $\mathcal{H}_- = H^{-s}(\mathbb{R}^n)$, where the self-adjoint pseudo-differential operator Ψ is defined by

$$\Psi : H^s(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n), \quad \Psi\phi := F^{-1}(\psi F\phi),$$

with ψ is a real-valued function such that

$$\frac{1}{c} (1 + |x|^2)^{s/2} \leq 1 + |\psi(x)| \leq c (1 + |x|^2)^{s/2}, \quad c > 0.$$

We want now to define the self-adjoint extensions of the restriction of Ψ to functions vanishing on a d -set, with $0 < n - d < 2s$. A Borel set $M \subset \mathbb{R}^n$ is called a d -set, $d \in (0, n]$, if

$$\exists c_1, c_2 > 0 : \forall x \in M, \forall r \in (0, 1), \quad c_1 r^d \leq \mu_d(B_r(x) \cap M) \leq c_2 r^d,$$

where μ_d is the d -dimensional Hausdorff measure and $B_r(x)$ is the closed n -dimensional ball of radius r centered at the point x (see [7], Section 1.1, Chapter VIII). Examples of d -sets are d -dimensional Lipschitz submanifolds and (when d is not an integer) self-similar fractals of Hausdorff dimension d (see [7], Chapter II, Example 2). We take as the linear operator τ the unique continuous surjective (thus (2.1) holds true) map

$$\tau_M : H^s(\mathbb{R}^n) \rightarrow B_{\alpha}^{2,2}(M), \quad \alpha = s - \frac{n-d}{2}$$

such that, for μ_d -a.e. $x \in M$,

$$\tau_M \phi(x) \equiv \{\phi_M^{(j)}(x)\}_{|j| < \alpha} = \left\{ \lim_{r \downarrow 0} \frac{1}{\lambda_n(r)} \int_{B_r(x)} dy D^j \phi(y) \right\}_{|j| < \alpha},$$

where $j \in \mathbb{Z}_+^n$, $|j| := j_1 + \dots + j_n$, $D^j := \partial_{j_1} \dots \partial_{j_n}$ and $\lambda_n(r)$ denotes the n -dimensional Lebesgue measure of $B_r(x)$. We refer to [7], Theorems 1 and 3, Chapter VII, for the existence of the map τ_M ; obviously it coincides with the usual evaluation along M when restricted to smooth functions. The definition of the Besov-like space $B_\alpha^{2,2}(M)$ is quite involved and we will not reproduce it here (see [7], Section 2.1, Chapter V). However, in the case $0 < \alpha < 1$ (i.e. $2(s-1) < n-d < 2s$), $B_\alpha^{2,2}(M)$ can be alternatively defined (see [7], Section 1.1, Chapter V) as the Hilbert space of $f \in L^2(F; \mu_M)$ having finite norm

$$\|f\|_{B_\alpha^{2,2}(M)}^2 := \|f\|_{L^2(M)}^2 + \int_{|x-y| < 1} d\mu_M(x) d\mu_M(y) \frac{|f(x) - f(y)|^2}{|x-y|^{d+2\alpha}},$$

where μ_M denotes the restriction of the d -dimensional Hausdorff measure μ_d to the set M .

The adjoint map τ_M^* gives rise, for any $Q \in B_\alpha^{2,2}(M)$, to the signed measure $\nu_M(Q) \in H^{-s}(\mathbb{R}^n)$ defined by

$$(\nu_M(Q), \phi) = \langle Q, \tau_M \phi \rangle_{B_\alpha^{2,2}(M)}.$$

Since $\nu_M(Q)$ has support given by the closure of M , hypothesis (2.2) is always verified when the closure of M has zero Lebesgue measure. Defining then

$$\mathcal{G}_\star^\psi := \operatorname{Re} F^{-1} \frac{1}{-\psi + z_0},$$

one has

$$G_\star : B_\alpha^{2,2}(M) \rightarrow L^2(\mathbb{R}^n), \quad G_\star Q := \mathcal{G}_\star^\psi * \nu_M(Q).$$

Therefore, given any self-adjoint $\Theta : D(\Theta) \subseteq B_\alpha^{2,2}(M) \rightarrow B_\alpha^{2,2}(M)$, one has the family of self-adjoint extensions

$$D(\Psi_\Theta) := \{\phi \in L^2(\mathbb{R}^n) : \phi = \phi_\star + \mathcal{G}_\star^\psi * \nu_M(Q_\phi), \phi_\star \in H^s(\mathbb{R}^n), Q_\phi \in D(\Theta), \tau_M \phi_\star = \Theta Q_\phi\},$$

$$\Psi_\Theta \phi := F^{-1}(\psi F \phi) + \nu_M(Q_\phi)$$

(see [16], Example 3.6, [17], Section 4, for alternative definitions).

When M is a compact Riemannian manifold, Δ_{LB} the Laplace-Beltrami operator, one has

$$B_\alpha^{2,2}(M) \simeq H^\alpha(M) = \{Q \in L^2(M) : (-\Delta_{LB})^{\alpha/2} Q \in L^2(M)\}$$

and

$$v_M(Q) = ((-\Delta_{LB})^\alpha Q) \delta_M,$$

where, for any $\tilde{Q} \in H^{-\alpha}(M) \equiv H^\alpha(M)'$,

$$\tilde{Q}\delta_M(\phi) := \int_M dv(-\Delta_{LB})^{-\alpha/2} \tilde{Q}(-\Delta_{LB})^{\alpha/2} \tau_M \phi,$$

dv denoting the volume element of M . Therefore in this case, when $\alpha \geq 1$ (i.e. $0 < n - d \leq 2$), taking $\psi(k) = |k|^2$, $\Theta = (-\Delta_{LB})^{\alpha-1}$, one can define the self-adjoint extension

$$-\Delta_M \phi := -\Delta \phi - \Delta_{LB} \cdot \tau_M \phi \star \delta_M,$$

and so the construction given here generalizes the examples given in [8] and [9]. Also see [17], Example 14, for an alternative definition.

REFERENCES

- [1] S. ALBEVERIO – F. GESZTESY – R. HØEGH-KROHN – H. HOLDEN, “Solvable Models in Quantum Mechanics”, Springer-Verlag, Berlin, Heidelberg, New York, 1988.
- [2] S. ALBEVERIO – W. KARWOWSKI – V. KOSHMANENKO, *Square Powers of Singularly Perturbed Operators*, Math. Nachr. **173** (1995), 5-24.
- [3] S. ALBEVERIO, P. KURASOV, “Singular Perturbations of Differential Operators”, Cambridge Univ. Press, Cambridge, 2000.
- [4] V. A. DERKACH – M. M. MALAMUD, *Generalized Resolvents and the Boundary Value Problem for Hermitian Operators with Gaps*, J. Funct. Anal. **95** (1991), 1-95.
- [5] W. G. FARIS, “Self-Adjoint Operators”, Lecture Notes in Mathematics **433**, Springer-Verlag, Berlin, Heidelberg, New York, 1975.
- [6] F. GESZTESY – K. A. MAKAROV – E. TSEKANOVSKII, *An Addendum to Kreĭn’s Formula*, J. Math. Anal. Appl. **222** (1998), 594-606.
- [7] A. JONSSON – H. WALLIN, *Function Spaces on Subsets of \mathbb{R}^n* , Math. Reports **2** (1984), 1-221.
- [8] W. KARWOWSKI – V. KOSHMANENKO – S. ÔTA, *Schrödinger Operators Perturbed by Operators Related to Null Sets*, Positivity **2** (1998), 77-99.
- [9] S. KONDEJ, *Singular Perturbations of Laplace Operator in Terms of Boundary Conditions*, To appear in *Positivity*.
- [10] V. KOSHMANENKO, *Singular Operators as a Parameter of Self-Adjoint Extensions*, Oper. Theory Adv. Appl. **118** (2000), 205-223.
- [11] M. G. KRĚŇ, *On Hermitian Operators with Deficiency Indices One*, Dokl. Akad. Nauk SSSR **43** (1944), 339-342 (in Russian).
- [12] M. G. KRĚŇ, *Resolvents of Hermitian Operators with Defect Index (m, m)* , Dokl. Akad. Nauk SSSR **52** (1946), 657-660 (in Russian).
- [13] M. G. KRĚŇ – V. A. YAVRYAN, *Spectral Shift Functions that arise in Perturbations of a Positive Operator*, J. Operator Theory **81** (1981), 155-181.

- [14] P. KURASOV – S. T. KURODA, *Kreĭn's Formula and Perturbation Theory*, Preprint, Stockholm University, 2000.
- [15] J. VON NEUMANN, *Allgemeine Eigenwerttheorie Hermitescher Funktionaloperatoren*, Math. Ann. **102** (1929-30), 49-131.
- [16] A. POSILICANO, *A Kreĭn-like Formula for Singular Perturbations of Self-Adjoint Operators and Applications*, J. Funct. Anal. **183** (2001), 109-147.
- [17] A. POSILICANO, *Boundary Conditions for Singular Perturbations of Self-Adjoint Operators*, Oper. Theory Adv. Appl. **132** (2002), 333-346.
- [18] SH. N. SAAKJAN, *On the Theory of Resolvents of a Symmetric Operator with Infinite Deficiency Indices*, Dokl. Akad. Nauk Arm. SSSR **44** (1965), 193-198 (in Russian).

Dipartimento di Scienze
Università dell'Insubria
22100 Como (Italy)
andreap@uninsubria.it