

# Self-Cleavable Chemiluminescent Probes Suitable for Protease Sensing

**Jean-Alexandre Richard,<sup>a,d</sup> Ludovic Jean,<sup>a,b</sup> Caroline Schenkels,<sup>a,d</sup> Marc Massonneau,<sup>d</sup> Anthony Romieu<sup>\*a,b</sup> and Pierre-Yves Renard<sup>\*a,b,c</sup>**


<sup>a</sup> *Equipe de Chimie Bio-Organique, COBRA - CNRS UMR 6014 & FR 3038, rue Lucien Tesnière, 76131 Mont-Saint-Aignan, France. E-mail: [pierre-yves.renard@univ-rouen.fr](mailto:pierre-yves.renard@univ-rouen.fr) or [anthony.romieu@univ-rouen.fr](mailto:anthony.romieu@univ-rouen.fr); Fax: +33 2-35-52-29-59*

<sup>b</sup> *Université de Rouen, Place Emile Blondel, 76821 Mont-Saint-Aignan, France*

<sup>c</sup> *Institut Universitaire de France*

<sup>d</sup> *QUIDD, Technopôle du Madrillet, 50, rue Ettore Bugatti, 76800 Saint-Etienne du Rouvray, France*

## Supporting Information



**Fig. 6** Luminescence intensity (area under the emission curve in the range 490-570 nm) recorded every 45 s with probe **29** after incubation with recombinant PGA ( $1.05 \cdot 10^{-2}$  U, incubation time: 10 min).

**Table 1** Influence of experimental conditions on aspartate epimerisation during the coupling between tetrapeptide **19** and PABA linker.

Entry	T (°C)	Conditions	Coupling reagent	<b>30a</b> : <b>30b</b> ratio <sup>a</sup>
1	20 °C	2 equiv. DIEA, DMF	EDC	55 : 45
2	20 °C	3 equiv. DIEA, DMF	BOP	66 : 34
3	20°C	1 equiv. HOBt, DMF	EDC	75 : 25
4	20 °C	DMF	EDC	75 : 25
5	0 °C to 20 °C	DMF	EEDQ	80 : 20
6	0 °C to 20 °C	DMF	EDC	90 : 10

<sup>a</sup> determined by RP-HPLC.