

SELF-INJECTIVE SEMIGROUP RINGS FOR FINITE INVERSE SEMIGROUPS¹

R. WENGER

The purpose of this article is the proof of the following theorem. R will always denote a ring with identity, and $R(S)$ the semigroup ring (contracted if S has a zero) of a semigroup S over R .

THEOREM. *Let S be a finite inverse semigroup. Then $R(S)$ is self-injective (s.i.) if, and only if, R is s.i.*

This is an extension of Theorem 8.3 in [3], of part 1 of Theorem 4.1 in [2], and of the corollary to Theorem 1 in [4]. The results in [2] and [3] are used in its proof. As R has an identity, S is assumed embedded in $R(S)$. $|X|$ will denote the cardinality of the set X and $X \setminus Y$ will denote the complement of a set Y in a set X . Terminology and definitions are given in [1].

1. An identity for $R(S)$ and the main theorem. Let S be an inverse semigroup, i.e. a regular semigroup in which idempotents commute. Let E be the set of idempotents in S . Then E is a commutative idempotent subsemigroup of S and each principal left (right) ideal of S has a unique idempotent generator [1, Theorem 1.17, p. 28]. Then $R(S)$ has an identity if $R(E)$ has one. If Z denotes the ring of integers and $Z(E)$ has an identity, then $R(E)$ has an identity. Note that E has a zero if $|E|$ is finite.

THEOREM 1. *If E is a finite commutative idempotent semigroup, then $Z(E)$ has an identity.*

PROOF. The proof is by induction on $|E|$. If $|E|$ is 1 or 2, the result is clear. Since $|E|$ is finite there exists an element $u (\neq 0)$ in E such that $uE = \{u, 0\}$. Then $I = \{u, 0\}$ is an ideal of E . $|E/I| < |E|$ and E/I , the Rees factor semigroup, is a commutative idempotent semigroup. Let $\phi': E \rightarrow E/I$ be the natural homomorphism and extend ϕ' linearly to the ring epimorphism $\phi: Z(E) \rightarrow Z(E/I)$ with kernel $Z(I)$. Let $\phi(a)$ be denoted by \bar{a} . E and E/I are assumed embedded in $Z(E)$ and $Z(E/I)$ respectively. Let $e^* = \sum_{a \in E \setminus I} \alpha(\bar{a})\bar{a}$, $\alpha(a) \in Z$, be the identity of $Z(E/I)$ insured by the induction hypothesis. Let $e' = \sum_{a \in E \setminus I} \alpha(a)a$

Received by the editors January 22, 1967 and, in revised form, September 8, 1967.

¹ This work was supported by a University of Delaware Summer Faculty Fellowship. The author wishes to thank the referee for his important comments.

$\in Z(E)$. Then $\phi(e') = e^*$ and $e^* \bar{x} = \bar{x}$ for $x \in E$ so $e'x \in x + Z(I)$. For $x \in E$, let $f(x) \in Z$ such that $e'x = x + f(x)u$. If $xu = u$ and $yu = u$, then $f(x) = f(y)$ since

$$(e'x)y = (x + f(x)u)y = xy + f(x)u$$

and

$$(e'y)x = (y + f(y)u)x = yx + f(y)u$$

and E is commutative. Thus there is an $\alpha \in Z$ such that $f(x) = \alpha$ for each $x \in E$ such that $xu = u$. If $f(x) \neq 0$ there is a $w \in E$ such that $xw = u$ so $xu = u$ as E is a semigroup of idempotents. Let $e = e' - \alpha u$. If $x \in E$, then $ex = e'x - \alpha ux = x + f(x)u - \alpha ux$. If $xu = u$, then $f(x) = \alpha$ and $ex = x$. If $xu = 0$, then $f(x) = 0$ and again $ex = x$. Hence, $Z(E)$ has an identity and the induction argument is complete.

This result can be extended somewhat. In what follows, E and E' will denote commutative idempotent semigroups. Let $M(E) = \{a : a \in E \text{ and } x \in E, ax = a \text{ imply } x = a\}$. Suppose $M(E)$ nonempty and $M(E)E = E$. Then for each $x \in E$ there is an $a \in M(E)$ for which $ax = x$. This condition is satisfied, for example, if in the set $P(E)$ of principal ideals of E , each chain has an upper bound (with respect to the partial ordering of $P(E)$ by set inclusion). If E is finite $M(E)E = E$. $M(E)E = E$ will be assumed for each idempotent semigroup in this article. Clearly, $M(E)$ is contained in any set of generators of E . E is said to be *unrefined* if $M(E)$ is a set of generators for E . A semigroup E' is called a *refinement* of E if E is a subsemigroup of E' and $M(E) = M(E')$. In general, E' is a refinement of the subsemigroup E generated by $M(E')$ and E is an unrefined semigroup with $M(E) = M(E')$. If $e \in Z(E)$ is an identity for $Z(E) \subseteq Z(E')$ and $x \in E'$, there is an $a \in M(E') = M(E) \subseteq E$ such that $ax = x$. Then $ex = e(ax) = (ea)x = ax = x$, so e is an identity for $Z(E')$ also. Thus, only unrefined semigroups need be considered in order to seek the existence of an identity. For example, let $E' = \{k : k \text{ a positive integer}\}$ and define $k * m = \max\{k, m\}$. Then $M(E') = \{1\}$ and E' is a refinement of $E = \{1\}$.

LEMMA 1. *If E is unrefined, then E is finite if, and only if, $M(E)$ is finite.*

PROOF. If $|M(E)| = m$, $|E| \leq 2^m - 1$.

LEMMA 2. *If $M(E')$ is finite, then E' is a refinement of a finite unrefined semigroup E .*

COROLLARY (TO THEOREM 1). *$Z(E)$ has an identity if, and only if, $M(E)$ is finite.*

PROOF. Assume $M(E)$ is finite and let E' be the subsemigroup generated by $M(E)$. Then E' is unrefined and $Z(E')$ has an identity by Lemma 1. Then $Z(E)$ has an identity by the remarks above as $M(E)E = E$.

Conversely, suppose $e = \sum_{i=1}^p n_i a_i$, $n_i \in \mathbb{Z}$, $a_i \in E$, is an identity for $Z(E)$. Then it is an identity on $M(E)$. If there exists an $a \in M(E)$ with $a \neq a_i$ for each i , then $a_i a \neq a$ for each i so $ea \neq a$; a contradiction. Thus, $M(E)$ is finite. (Note that if $M(E)$ is finite and e is the identity of $Z(E)$, then each $a \in M(E)$ occurs with a nonzero coefficient in e .)

THEOREM 2. *Let S be an inverse semigroup with idempotent semigroup E . Then $R(S)$ has an identity if, and only if, $M(E)$ is finite.*

PROOF. If $M(E)$ is finite, $Z(E)$ has an identity, say $e' = \sum_{i=1}^p n_i a_i$, $n_i \in \mathbb{Z}$, $a_i \in E$. Let 1 be the identity of R and let $e = \sum_{i=1}^p (n_i 1) a_i$. Then e is an identity for $R(E)$. If $s \in S$, there are elements a and b in E such that $as = s$ and $sb = s$ so e is an identity for $R(S)$. Conversely, if $e = \sum_{i=1}^p r_i x_i$, $r_i \in R$, $x_i \in S$, is an identity for $R(S)$, then, in particular, $ea = a$ for each $a \in M(E)$. Thus $M(E)$ must be finite by the argument used in the proof of the corollary.

THEOREM 3. *Let S be a finite inverse semigroup. Then $R(S)$ is s.i. if, and only if, R is s.i.*

PROOF. Let $S = S_0 \supset S_1 \supset \dots \supset S_{n+1}$ be a principal series for S with $S_{n+1} = \{0\}$ if S has a zero and S_{n+1} empty otherwise. S_i/S_{i+1} is a Brandt semigroup by [1, Exercise 3, p. 103], for each $i = 0, 1, \dots, n$. The proof is by induction on n . If $n = 0$, $S \cong S_0/S_1$ is a Brandt semigroup so $S \cong M^0(G; m; m; \Delta)$, an $m \times m$ Rees matrix semigroup over a group with zero G^0 and with the $m \times m$ identity matrix Δ as a sandwich matrix [1, Theorem 3.9, p. 102]. Then $R(S) \cong M_m(R(G))$, the ring of $m \times m$ matrices over $R(G)$. $M_m(R(G))$ is s.i. if, and only if, $R(G)$ is s.i. by [3, Theorem 8.3]. As G is finite, $R(G)$ is s.i. if, and only if, R is s.i. by [2, Theorem 4.1].

As suppose $n > 0$. Then $R(S/S_n)$ is s.i. if, and only if, R is s.i. and S/S_n is a finite inverse semigroup and has a principal series of length less than n . S_n is a Brandt semigroup (so an inverse semigroup) so $R(S_n) (\subseteq R(S))$ has an identity, say f . If $x \in R(S)$, xf and fx are in $R(S_n)$ so $xf = f(xf) = (fx)f = fx$ and f is central in $R(S)$. Let e be the identity of $R(S)$ insured by Theorem 2. Then $R(S) = R(S)(e-f) \oplus R(S)f$, a ring direct sum. $R(S)$ is s.i. if, and only if, both $R(S)(e-f)$ and $R(S)f$ are s.i. [4, Lemma 1]. $R(S)f = R(S_n)$ so $R(S)(e-f) \cong R(S)/R(S_n) \cong R(S/S_n)$. If R is s.i., then $R(S/S_n)$ and $R(S_n)$ are s.i. so $R(S)$ is. If $R(S)$ is s.i., then $R(S_n)$ is s.i. so R is s.i. as S_n is a Brandt

semigroup and the argument used in the case $n=0$ applies. This completes the induction.

Note that $M(E)$ finite is not sufficient to yield this result [2, Theorem 4.1].

A group G is an inverse semigroup; so if G is finite, $R(G)$ is s.i. if, and only if, R is s.i. This is part of Connell's result. Utumi's result is that $M_m(R)$, the ring of $m \times m$ matrices over R , is s.i. if, and only if, R is s.i. But $M_m(R) = R(S)$, where $S = \{e_{ij}: 1 \leq i, j \leq m\} \cup \{0\}$ with $e_{ij}e_{pq} = \delta_{jp}e_{iq}$, δ_{jp} the Kronecker delta, is an inverse semigroup.

BIBLIOGRAPHY

1. A. H. Clifford and G. B. Preston, *Algebraic theory of semigroups*, Vol. I, Math. Surveys, No. 7, Amer. Math. Soc., Providence, R. I., 1961.
2. Ian Connell, *On the group ring*, Canad. J. Math. 15 (1963), 650–685.
3. Yuzo Utumi, *On continuous rings and self-injective rings*, Trans. Amer. Math. Soc. 118 (1965), 158–173.
4. R. Wenger, *Semigroups having quasi-Fröbenius algebras*. I, Proc. London Math. Soc. 18 (1968), 484–494.

UNIVERSITY OF DELAWARE