

Self-Recognition of CD1 by γ/δ T Cells: Implications for Innate Immunity

By Franca M. Spada,* Ethan P. Grant,* Peter J. Peters,[‡] Masahiko Sugita,* Agustín Melián,* David S. Leslie,* Hoi K. Lee,[§] Elly van Donselaar,[‡] Dennis A. Hanson,^{||} Alan M. Krensky,[¶] Otto Majdic,** Steven A. Porcelli,* Craig T. Morita,[§] and Michael B. Brenner*

From the *Division of Rheumatology, Immunology, and Allergy, Department of Medicine, Brigham and Women's Hospital at Harvard Medical School, Boston, Massachusetts 02115; [‡]The Netherlands Cancer Institute, 1066 CX Amsterdam, The Netherlands; the [§]Division of Rheumatology, Department of Internal Medicine and Interdisciplinary Graduate Program in Immunology, University of Iowa, Iowa City, Iowa 52242; the ^{||}Virginia Mason Research Center, Seattle, Washington 98101; the [¶]Division of Immunology and Transplantation Biology, Department of Pediatrics, Stanford University, Stanford, California 94305; and the **Institute of Immunology, University of Vienna, A-1090 Vienna, Austria

Abstract

The specificity of immunoglobulins and α/β T cell receptors (TCRs) provides a framework for the molecular basis of antigen recognition. Yet, evolution has preserved a separate lineage of γ/δ antigen receptors that share characteristics of both immunoglobulins and α/β TCRs but whose antigens remain poorly understood. We now show that T cells of the major tissue γ/δ T cell subset recognize nonpolymorphic CD1c molecules. These T cells proliferated in response to CD1⁺ presenter cells, lysed CD1c⁺ targets, and released T helper type 1 (Th1) cytokines. The CD1c-reactive γ/δ T cells were cytotoxic and used both perforin- and Fas-mediated cytotoxicity. Moreover, they produced granulysin, an important antimicrobial protein. Recognition of CD1c was TCR mediated, as recognition was transferred by transfection of the γ/δ TCR. Importantly, all CD1c-reactive γ/δ T cells express V δ 1 TCRs, the TCR expressed by most tissue γ/δ T cells. Recognition by this tissue pool of γ/δ T cells provides the human immune system with the capacity to respond rapidly to nonpolymorphic molecules on professional antigen presenting cells (APCs) in the absence of foreign antigens that may activate or eliminate the APCs. The presence of bactericidal granulysin suggests these cells may directly mediate host defense even before foreign antigen-specific T cells have differentiated.

Key words: T lymphocytes • T cell antigen receptors γ/δ • CD1 • cytolysis • granulysin

Introduction

Studies in mice and humans have implicated γ/δ T cells in host defense. For example, γ/δ T cells modulate the severity of disease or mediate a component of protective immunity in murine models of listeriosis (1), tuberculosis (2–4), malaria (5), and HSV-1 encephalitis (6). γ/δ T cells are critically important in preventing death from airway infection with *Nocardia asteroides* (7). In humans, large expan-

sions of γ/δ T cells during infections suggest their importance. γ/δ T cells increase from normal levels of 4% of all circulating T cells to a mean of 12, 14, 29, and 57% of all circulating T cells during infection with *Listeria monocytogenes*, *Mycobacterium tuberculosis*, *Brucella melitensis*, and *Ehrlichia chaffeensis*, respectively (8–11). Despite the growing list of experimental models and human diseases with γ/δ T cell expansions, only a few examples of well-characterized antigen-specific γ/δ T cells have been noted, and these do not support MHC-restricted peptide-specific recognition (12, 13). Instead, in mice, γ/δ T cell recognition of an intact (unprocessed) cell surface glycoprotein of HSV was

Address correspondence to Michael B. Brenner, Division of Rheumatology, Immunology, and Allergy, Department of Medicine, Brigham and Women's Hospital, 1 Jimmy Fund Way, Boston, MA 02115. Phone: 617-525-1000; Fax: 617-525-1010; E-mail: mbrenner@rics.bwh.harvard.edu

noted in one case, whereas in other cases, alloreactive γ/δ T cell recognition of TL and I-E was not dependent on bound peptides (14–16). These studies, together with analysis of CDR3 length distribution of the TCR δ chain (17) and the structure of a V δ domain revealing shared features with both immunoglobulins and α/β TCRs (18), led to the suggestion that, in some cases, γ/δ recognition may be more like that of immunoglobulins than of α/β TCRs.

Major advances in the study of antigen recognition by human γ/δ T cells were made with the discovery that most circulating human γ/δ T cells that bear V γ 2/V δ 2 (V γ 2 is also termed V γ 9) TCRs could be stimulated by nonpeptide antigens (for a review, see reference 19) derived from *Mycobacteria* (20–24). These nonpeptide antigens were identified as isopentenyl pyrophosphate (IPP)¹ and related prenyl pyrophosphate molecules (24, 25) and the more recently characterized alkylamine antigens (26). Both the phosphate and the amine antigens are small molecules consisting of short (typically one to five carbons) straight or branched aliphatic chains and either a phosphate or an amine moiety. They are important products of microbes as well as self-antigens. The mechanism by which these antigens are presented is not known, but it does not involve the known MHC class I or MHC class II peptide antigen-presenting molecules (23, 27, 28), and we have suggested they may be recognized much as haptens are recognized by either immunoglobulins or TCRs (23, 29). Namely, their requirement for an antigen-presenting element is unclear, but their recognition is critically dependent on the CDR3 sequence of the γ/δ TCR (29, 30). The recognition of these small aliphatic phosphate and amine organic molecules was exclusively found among γ/δ T cells of the V δ 2 subset.

In contrast to V γ 2/V δ 2⁺ T cells that are the major circulating pool of γ/δ T cells, human γ/δ T cells bearing V δ 1-encoded TCRs account for the vast majority of γ/δ T cells in tissues such as intestine and spleen (31). Yet, little is known about the antigen reactivity of these tissue γ/δ T cells. Recently, examples of γ/δ T cells in this subset were found to recognize the MHC-encoded proteins, MICA and MICB (32). Recognition was through the activating NKG2D C-type lectin (33) with an unclear contribution from the γ/δ TCR. MICA and MICB class I molecules identify stressed cells and have a very restricted pattern of expression primarily limited to the intestine. MICA and MICB probably do not present peptides, as they have a limited peptide binding groove (34). Instead, these molecules may function in innate immunity as important targets for V δ 1⁺ γ/δ T cell killing of stressed cells (32).

Here, we provide evidence that an important TCR-mediated reactivity of tissue γ/δ T cells is against CD1 molecules. Remarkably, all of the V δ 1 cells studied were focused specifically on CD1c, one member of a family of nonpolymorphic CD1 molecules, expressed exclusively on APCs, that present lipid and glycolipid foreign antigens to

T cells (35). However, the γ/δ T cell lines and clones showed highly specific, direct reactivity to CD1c proteins that was not dependent on the presence of an exogenous foreign antigen. These CD1c-specific γ/δ T cells produced inflammatory cytokines, killed CD1c-bearing targets, and contained bactericidal granulysin.

Materials and Methods

mAbs. The following mAbs were used for flow cytometry and blocking experiments: P3 (IgG control) (22), SPV-T3b (anti-CD3) (36), anti-TCR- δ 1 (pan anti-C δ) (37), δ TCS1 (anti-V δ 1/J δ 1) (38), Ti γ A (anti-V γ 2) (39, 40), 9.3 (anti-CD28) (41), OKT4 (anti-CD4; American Type Culture Collection), OKT8 (anti-CD8 α ; American Type Culture Collection), DX1 (anti-NKR-P1A; provided by Dr. L. Lanier, DNAX, Palo Alto, CA), BMAO31 (pan anti-TCR- α/β ; provided by Dr. R.G. Kurrele, Boehringerwerke, Marburg, Germany), 7C6 (anti-CD1c) (42), F10/21A3 (anti-CD1c) (42a), BCD1b3.2 (anti-CD1b) (43), 10H3.9.3 (anti-CD1a) (44), W6/32 (anti-MHC class I; American Type Culture Collection), L243 (anti-HLA-DR; American Type Culture Collection), NS4.1 (IgM control; American Type Culture Collection), 4A11 (anti-V γ 1.4) (45), CD95 Fas ZB4 clone (anti-Fas; Immunotech), δ G9 (antiperforin; Ancell), DH2 (antigranulysin) (46), and MPC11 (IgG2b control; American Type Culture Collection).

Immunofluorescence Analysis. Cells were incubated with mouse mAbs on ice for 30 min, washed, and stained with FITC-conjugated F(ab')₂ goat anti-mouse Ig (Tago) for an additional 30 min on ice. After washing, the cells were resuspended in propidium iodide and analyzed by flow cytometry (FACSsort™; Becton Dickinson). Results were expressed as percentage of positive cells compared with negative cells stained with isotype-matched control mAbs.

T Cell Lines and Clones. Lymphocytes were isolated from the blood of random healthy donors by Ficoll-Hypaque centrifugation. γ/δ T cells were enriched by staining with the anti-TCR- α/β mAb, BMAO31, followed by depletion of α/β T cells with magnetic beads coated with goat anti-mouse IgG (Dynal). T cell lines JR.2 and XV.1 were established from two separate donors by culturing 1×10^6 freshly isolated γ/δ T cells in 1-ml culture wells with 1×10^6 irradiated (5,000 rads) autologous CD1⁺ monocyte-derived dendritic cells as APCs and the organic phase of a chloroform/methanol/water (2:1:1) extract of *M. tuberculosis* prepared as described (47) at 1:5,000 final dilution. After 2 wk of culture, viable cells were recovered and the residual α/β T cells were depleted with BMAO31 mAb and magnetic beads (Dynal). The resulting population was restimulated with heterologous CD1⁺ dendritic cells, an organic phase extract of *M. tuberculosis*, and rIL-2 (1 nM; Ajinomoto Co.). T cells were maintained by restimulation every 2 wk with irradiated heterologous CD1⁺ dendritic cells and rIL-2. T cell clones were derived from lines JR.2 and XV.1 by limiting dilution culture using PHA stimulation. T cells were seeded at 1 and 5 cells per well in 96-well round-bottomed plates in a volume of 0.2 ml, with 5×10^4 irradiated (4,000 rads) heterologous PBMCs and 5×10^4 irradiated (5,000 rads) EBV-transformed B lymphoblastoid cells as feeders in RPMI medium supplemented with PHA-P (1:4,000 final dilution; Difco) and IL-2 (2 nM).

APC Lines. Monocyte-derived dendritic cells were generated from human blood monocytes that were isolated from the byproducts of platelet pheresis and induced to differentiate and express CD1a, CD1b, and CD1c by incubation with GM-CSF

¹Abbreviations used in this paper: IPP, isopentenyl pyrophosphate; MEP, monoethyl phosphate; UTR, untranslated region.

and IL-4 as described (48). The lymphoblastoid cell lines C1R were transfected with the expression vector pSR α -Neo into which cDNAs encoding either CD1a, CD1b, or CD1c were inserted as described previously (48). HeLa cells were either mock transfected (HeLa Mock) or CD1c transfected (HeLa CD1c) (49).

Cytolytic Assays. Cytolytic assays were performed in a 4-h chromium-release assay (43). The targets used were CD1-expressing GM-CSF/IL-4-treated blood monocytes; C1R lymphoblastoid cells either mock transfected (C1R Mock), CD1b transfected (C1R CD1b), or CD1c transfected (C1R CD1c); and HeLa Mock or HeLa CD1c cells (49). In the mAb blocking experiments, targets were incubated with mAbs (20 μ g/ml) for 30 min at room temperature before adding effector cells. In experiments where both anti-Fas mAb and strontium ions were used, the targets were incubated for 1 h with anti-Fas mAb (1 μ g/ml) before adding the effectors. The effector T cells were treated by incubating with 25 mM strontium chloride hexahydrate (Aldrich Chemical Co.) at 37°C overnight, washing, and then counting after trypan blue incubation to determine their viability. For the *M. tuberculosis*-specific CD1b-restricted α/β T cell line DN1 (47), the target C1R/CD1b was incubated overnight at 37°C with *M. tuberculosis* total sonicate at 1 μ g/ml before T cell addition. For V γ 2/V δ 2 T cell clones specific for IPP and related compounds, the antigen monoethyl phosphate (MEP) was added together with the effectors T cells and targets (1:300) (24). Assays were performed in triplicate, and results were expressed as percent specific ^{51}Cr release \pm SEM.

Proliferation Assays. 5×10^4 T cells were plated in triplicate in 96-well flat-bottomed plates with either 5×10^4 CD1 $^+$ dendritic cells or 5×10^4 mitomycin C-treated HeLa Mock and HeLa CD1c cells as APCs. In the mAb blocking experiments, mAbs were added as ascites (final dilution of 1:200) or purified mAbs (20 μ g/ml). Cultures were incubated at 37°C, pulsed with 1 μ Ci of [^3H]thymidine (2 Ci/mmol; Amersham Pharmacia Biotech) on day 2, and harvested 16 h later using a Tomtec harvester. The filter papers were counted on a Betaplate scintillation counter (Wallac). Results were expressed as cpm \pm SEM.

Cytokine Assay. 5×10^5 T cells from the JR.2 and XV.1 lines were cultured with 5×10^5 CD1 $^+$ monocyte-derived dendritic cells in the absence or presence of an anti-CD1c mAb (7C6) or an IgM mAb control (NS4.1). PHA-P (1:4,000) was added as a positive control. Supernatants were harvested after 24 and 48 h of culture. Cytokine release was determined for IL-2, IL-4, IL-10, and IFN- γ by sandwich ELISA assay (50) using antibody pairs purchased from PharMingen (IL-2, IL-4, IL-10) or from Endogen (IFN- γ). Results were expressed as ng/ml \pm SEM.

Confocal Microscopy Analysis. For confocal microscopy analysis, cells were fixed and permeabilized as described (49). For the JR.2 line, cells were incubated with either antiperforin directly conjugated with FITC (δ G9 mAb, 1–4 ng/ml) or with antigranulysin (DH2 mAb [51], 2.5–8 μ g/ml) followed by incubation with goat anti-mouse IgG-Cy3 (4 μ g/ml; Jackson ImmunoResearch Labs), in single and double staining. Negative controls were MPC11-FITC mAb (IgG $_{2b}$ isotype control for δ G9 mAb) together with DH2-Cy3, and P3-Cy3 mAb (IgG $_1$ isotype control for DH2 mAb) together with δ G9-FITC. For the 12G12 clone, cells were incubated either with δ G9 mAb (3 μ g/ml) or with DH2 mAb followed by incubation with anti-mouse Ig-Texas red (2.5–5 μ g/ml; Molecular Probes) and goat anti-mouse IgG1-FITC (2.5–5 μ g/ml; Southern Biotechnology Associates), respectively, in single and double staining. Controls were δ G9-Texas red with P3-FITC and DH2-FITC with MPC11-Texas red, double staining. Each incubation was performed at 4°C for 1 h;

in the double staining, cells were incubated for 10 min with 10% mouse serum between the first and second mAb staining.

Immunoelectron Microscopy. For electron microscopy, T cells were harvested from culture and dead cells were removed by Ficoll-Hypaque density gradient centrifugation. $0.5\text{--}1 \times 10^7$ live T cells were fixed at room temperature with 2% glutaraldehyde (Polysciences) in 0.1 M phosphate buffer for 2 h. The fixed cells were then collected into 0.1 M phosphate buffer containing 0.2% paraformaldehyde (Electron Microscopy Sciences) and processed for ultrathin cryosectioning as described (52). Cryosections were incubated with antigranulysin mAb, DH2, for 45 min, washed, and then incubated with protein A-gold (Electron Microscopy Laboratory, Utrecht University, Utrecht, The Netherlands) for 30 min. Labeled sections were viewed with a JOEL 1010 electron microscope at 80 kV.

TCR Analysis by PCR. Total RNA was isolated as described (53) from six JR.2 clones obtained from the JR.2 line. The γ and δ chains were amplified by PCR as described (29) using the following primers: V δ 1 chain, 5'-GGGCTCGAGCTTCAGGCAGCA-CAACT-3' (5' untranslated region [UTR]) and 5'-GGGAGAT-CTTGGCAGCTCTTTGATGGTGGTTGCTTTGGTTT-3' (C δ region); V γ 2 chain, 5'-GGGGTCGACCTGGTGAAGTCATACAGT-3' (V γ 2 internal region) and 5'-GGGXCTAGAGTGAG-GTTCTCTGTGT-3' (C γ 3' UTR).

The PCR products were cloned into pBluescript II (Stratagene), and the sequences of the V-J junctional region were determined using an automated sequencer (PE Applied Biosystems).

TCR Transfection. Full-length cDNAs encoding the JR.2 TCR were amplified by PCR using the following primers: 5'-GGGCTCGAGCTTCAGGCAGCACAACT-3' (V δ 1 5' UTR), 5'-GGGGATCCGGAGTGTAGCTTCTCAT-3' (V δ 1 3' UTR), 5'-GGGGGTACCTGCCCTGGCAGAAAGCA-3' (V γ 2 5' UTR), and 5'-GGGCTCGAGATGGCCTCCTTGTC-3' (V γ 2 3' UTR).

TCR reconstitution was carried out essentially as described (54). The γ and δ cDNAs were cloned into pREP7 and pREP9 (Invitrogen), respectively. J.RT3-T3 cells, Jurkat derivatives that lack cell surface TCR expression (55), were obtained from American Type Culture Collection. J.RT3-T3.5 cells were transfected with 20 μ g each of pREP7-JR.2 γ and pREP-JR.2 δ by electroporation at 250 V, 960 μ F with a Gene Pulser (Bio-Rad). Control transfectants were electroporated with 20 μ g each of the pREP7 and pREP9 vectors alone. After 48 h, transfectants were placed into medium containing 1 mg/ml geneticin (Life Technologies) and 0.5 mg/ml hygromycin B (Life Technologies). After 2 wk, transfectants were analyzed for cell surface TCR expression by flow cytometry using anti-CD3 (SPV-T3b) and anti-C δ TCR (anti-TCR- δ 1) mAbs and used as responders in T cell stimulation assays. The JR.2/J.RT3 derived in this fashion had 35% TCR- γ/δ^+ cells that stained with anti-TCR- δ 1 to a mean fluorescence intensity of \sim 100 on flow cytometry.

T Cell Transfectant Stimulation Assay. J.RT3-T3.5 transfectants (1×10^5 /well) were cultured in 96-well flat-bottomed microtiter plates either in the presence of 10 ng/ml PMA alone, or in the presence of PMA plus CD1 $^+$ monocyte-derived dendritic cells with or without the following mAbs (at a concentration of 20 μ g/ml): P3 (isotype control), 10H3.9.3 (anti-CD1a), BCD1b3.2 (anti-CD1b), or F10/21A3.1 (anti-CD1c) in a total volume of 200 μ l/well. 50 μ g aliquots of the culture supernatants were collected after 20–24 h culture at 37°C and diluted 1:2 with culture medium. To measure the amount of IL-2 released into the supernatants, HT-2 cells (5,000/well) were added and cultured 25–30 h at 37°C. [^3H]Thymidine (1 μ Ci/well, 6.7 Ci/mmol; New En-




Figure 1. Flow cytometric analysis of JR.2 and XV.1 lines. The mAbs used were P3 (IgG control), SPV-T3b (anti-CD3), OKT4 (anti-CD4), OKT8 (anti-CD8 α), 9.3 (anti-CD28), DX1 (anti-NKR-P1A), anti-TCR- δ 1 (pan anti-C δ TCR), δ TCS1 (anti-V δ 1/J δ 1), Ti γ A (anti-V γ 2), and 4A11 (anti-V γ 1.4).

gland Nuclear) was added during the final 5–6 h of culture. Plates were harvested on a Tomtec plate harvester, and [3 H]thymidine incorporation was measured in a Betaplate liquid scintillation counter (Wallac).

Results

Derivation of CD1-reactive T Cell Lines. The limited germ-line diversity of the γ/δ TCR had led to the suggestion that these cells might recognize nonpolymorphic antigen-presenting molecules (56). Since CD1 molecules resemble MHC but lack polymorphism, we pursued efforts to isolate CD1-restricted γ/δ T cells. To derive γ/δ T cell lines, normal human PBMCs were depleted of α/β T cells by antibody staining and magnetic bead separation. The γ/δ T cell-enriched PBMCs were cultured in vitro with an organic phase extract of *M. tuberculosis* in the presence of CD1-expressing monocyte-derived dendritic cells (see Materials and Methods). After several restimulations and expansions, two independently derived T cell lines, JR.2 and XV.1, were obtained. Flow cytometric analyses revealed that both lines were of the V δ 1 γ/δ T cell subset. The JR.2 line expressed V γ 2 (mAb Ti γ A $^+$) and V δ 1/J δ 1 (mAb δ TCS1 $^+$), whereas the XV.1 line expressed V γ 1.4 (mAb 4A11 $^+$) and V δ 1/J δ 1 (mAb δ TCS1 $^+$). Both cell lines lacked expression of NKR-P1A, CD28, and CD4. As is common for γ/δ T cells (57), 27% (JR.2) and 32% (XV.1) of the cells expressed the CD8 α chain but not the CD8 β chain (Fig. 1, and data not shown).

γ/δ T Cells Specifically Recognize CD1 $^+$ APCs in Proliferation and Cytolytic Assays. Although the cell lines were expanded in the presence of *M. tuberculosis* antigens, the T cells did not specifically recognize *M. tuberculosis* antigens. Instead, JR.2 cells proliferated in the presence of CD1 $^+$ dendritic cells (64,675 cpm), even without *M. tuberculosis* antigens. This response was not blocked by mAbs against CD1a or CD1b, but was blocked to 307 cpm by mAb against CD1c. Nearly identical results were obtained for XV.1 cells (Fig. 2 A). Addition of a standard *M. tuberculosis* lipid/lipoglycan extract (47) did not augment proliferation by either line (data not shown). To confirm the specificity suggested by mAb blocking, CD1c transfectant cells were studied. JR.2 γ/δ T cells proliferated markedly to HeLa CD1c (49) but not to HeLa Mock cells (Fig. 2 B). Note that JR.2 T cells did not appear to recognize MICA, as these T cells did not recognize the HeLa Mock cell line that expresses MICA (58).

As it is known that many γ/δ cell lines are capable of cell-mediated cytotoxicity (13), JR.2 and XV.1 T cells were examined for their ability to lyse transfected C1R lymphoblastoid cell lines C1R CD1b, C1R CD1c, or C1R Mock (35). JR.2 and XV.1 T cells failed to lyse C1R CD1b or C1R Mock targets, but efficiently lysed C1R CD1c targets (70 and 62% specific lysis, respectively; Fig. 2 C). In each case, lysis was specifically blocked (99 and 63% inhibition) by anti-CD1c mAb. Similar results were obtained in cytolytic assays using CD1-expressing monocyte-derived




Figure 2. CD1c-specific recognition by JR.2 and XV.1 γ/δ T cell lines. (A) The proliferative response of JR.2 and XV.1 lines to CD1c $^+$ dendritic cells was inhibited by anti-CD1c mAb (F10/2A3) but not by anti-MHC class I, anti-MHC class II, anti-CD1a, and anti-CD1b mAbs. (B) The JR.2 T cell line proliferated in response to HeLa CD1c cells but not to HeLa Mock cells. Background proliferation of HeLa Mock and HeLa CD1c cells alone was 872 and 1,197 cpm, respectively. (C) C1R Mock, C1R CD1b, and C1R CD1c targets were 51 Cr labeled and tested in cytolytic assays with JR.2 and XV.1 T cell lines at different E/T ratios (30:1 is shown) in the presence of anti-CD1b (BCD1b3.2) or anti-CD1c (F10/213) mAbs. Note that JR.2 and XV.1 lysed only C1R CD1c transfectant cells and that lysis was blocked by anti-CD1c mAb. White bars, no mAb; black bars, anti-CD1b; gray bars, anti-CD1c.

dendritic cells (data not shown). Thus, JR.2 and XV.1 γ/δ T cells were CD1c specific as revealed in the proliferation and cytolytic analyses of CD1 transfectant cell lines and proliferation assays with CD1⁺ monocyte-derived dendritic cells. Note that these experiments were performed without addition of *M. tuberculosis* antigens.

CD1c-reactive γ/δ T Cells Are Th1-like. To assess the ability to produce cytokines upon exposure to CD1c, cells from the JR.2 line and the XV.1.14 clone were cocultured with CD1-expressing human dendritic cells, and cytokine levels were measured. Substantial levels of IFN- γ (10 and 0.28 ng/ml, respectively) and IL-2 (19 and 17 ng/ml, respectively) were produced after the T cell lines were exposed to CD1⁺ dendritic cells. Cytokines also were secreted at somewhat higher levels after stimulation with the mitogen PHA (36 and 2.4 ng/ml IFN- γ , and 45 and 9.2 ng/ml IL-2, respectively). The production of these cytokines to CD1c-expressing dendritic cells was blocked completely by the presence of an anti-CD1c mAb (Fig. 3). In contrast to these cytokines, IL-4 and IL-10 were undetectable under these conditions (detection limits were <16 pg/ml IL-4 and <24 pg/ml IL-10 for cytokine ELISA). Thus, the profile of cytokine production by CD1c-reactive γ/δ T cells was Th1-like.

γ/δ T Cells Use Perforin-dependent and Fas-dependent Cytotoxicity Pathways. CD1-restricted, microbial-specific CD8⁺ α/β T cells lyse targets using perforin-mediated pathways and impart a bactericidal effect mediated by granulysin, whereas CD1-restricted CD4⁻CD8⁻ α/β T cells lyse targets through Fas-FasL interaction (51, 59). As γ/δ T cells typically display cytolytic potential, we examined the cellular cytotoxicity mechanisms that CD1c-reactive γ/δ T cells used. Fas-dependent lysis can be inhibited by certain anti-Fas mAbs, whereas exhaustive granule secretion induced by strontium ions renders T cells dependent on perforin and granule-related enzymes temporarily incapable of killing target cells (59). To determine the cellular cytotoxicity mechanism of γ/δ T cells, a panel of γ/δ T cells was examined




Figure 3. JR.2 and XV.1 γ/δ T cell lines show a Th1-like cytokines profile. JR.2.1 and XV.1.14 clones were incubated with CD1c⁺ dendritic cells for 24 and 48 h. The culture supernatants were collected and assayed for IL-2, IFN- γ , IL-4, and IL-10. Note that the JR.2.1 and XV.1.14 clones produced IFN- γ and IL-2 in response to exposure to CD1⁺ dendritic cells. IL-4 and IL-10 were not detected (data not shown).

including the JR.2 line and IDP2 clone that are CD1c reactive (60) and previously studied V γ 2/V δ 2-bearing clones, 12G12, DG.SF68, HD.108, and CP.1.15 (25, 57), that are specific for small phosphate antigens such as MEP (24). 12G12 and CP.1.15 are CD8⁺, whereas HD.108 and DG.SF68 are CD4⁻CD8⁻. The antigen-specific lysis of targets by all of these γ/δ T cell clones was consistently and efficiently blocked by treatment with strontium ions (60–84% average inhibition; Fig. 4). Inhibition was observed when cells were treated with anti-Fas mAb as well, but this inhibition was variable. In contrast to γ/δ T cells, the TCR- α/β -expressing CD1b-restricted CD4⁻CD8⁻ T cell line, DN1 (47), lysed antigen-pulsed targets in a manner that was unaffected by pretreatment with strontium ions but was efficiently blocked by mAb against Fas (Fig. 4). These data indicate that cytotoxicity mediated by both V δ 1⁺ CD1c-reactive γ/δ T cells and V δ 2⁺ prenyl pyrophosphate-specific γ/δ T cells can be both perforin mediated and Fas mediated, like that described for CD8⁺ MHC class I-restricted α/β T cells (61).

γ/δ T Cells Express the Antimicrobial Protein, Granulysin. Granulysin is a protein present in the cytotoxic granules of cytolytic α/β T cells and NK cells (62) that has potent antimicrobial activity against a variety of bacteria including *M. tuberculosis*, *Staphylococcus aureus*, and *Listeria monocytogenes* (51). However, granulysin can kill intracellular bacteria only after the infected cells are lysed by perforin (51). To determine if CD1c-reactive γ/δ T cells express granulysin and therefore had the potential to kill intracellular bacteria,




Figure 4. Analysis of cytotoxicity mechanisms used by CD1c-reactive and prenyl pyrophosphate-specific γ/δ T cells. CD1c-reactive V γ 2/V δ 1 T cells, JR.2 and IDP2, and prenyl pyrophosphate antigen-specific V γ 2/V δ 2 T cell clones, 12G12, HD.108, CP.1.15, and DG.SF68, were used in cytolytic assays performed in the presence or absence of anti-Fas mAb to block Fas-mediated cytotoxicity or after treatment with strontium ions to block perforin-mediated cytotoxicity. The mycolic acid-specific, CD1b-restricted, CD4⁻CD8⁻ α/β T cell line, DN1, is shown as a control. MEP is an alkyl phosphate analogue of the IPP antigen recognized by V γ 2/V δ 2 T cells and was used to stimulate the prenyl pyrophosphate-specific clones. The E/T ratio was 10:1. Targets for CD1c-reactive and prenyl pyrophosphate-specific γ/δ T cells were C1R CD1c cells. Targets for the CD1b-restricted clone were C1R CD1b cells that had been incubated for 16 h at 37°C with *M. tuberculosis* sonicate at 1 μ g/ml. Note that the cytotoxic activity mediated by γ/δ T cells was blocked by both treatments and thus depends on both granule secretion and Fas-FasL interactions.





Figure 5. Expression of granulysin by γ/δ T cells. (A) Colocalization of granulysin and perforin in the cytolitic granules. Differential interference contrast images of γ/δ T cells are shown on the left. Confocal microscopic analysis of fluorescent immunostaining of perforin and granulysin of the JR.2 CD1c-reactive γ/δ T cell line and 12G12 phosphoantigen-specific γ/δ T cell clone are shown as labeled. Superimposed figures are shown in the right panels. Note that the counterstaining were exchanged such that perforin is green in the top panel but red in the bottom panel, and granulysin is red in the top panel but green in the bottom panel. (B) Localization of granulysin to the cytolitic granules of γ/δ T cells by immunoelectron microscopy. The presence of granulysin in the cytolitic granules of JR.2 (bottom) and 12G12 (top) is demonstrated by the gold labeling (small particles) seen with antigranulysin mAb staining. c, centriole; e, endosome; P, plasma membrane; m, mitochondria; n, nucleus; G, Golgi apparatus. Bar, 200 nm.

we examined the JR.2 cell line for expression of granulysin and perforin. JR.2 (CD1c-reactive) as well as 12G12 (phosphoantigen-specific) γ/δ T cell lines expressed both granulysin and perforin. These proteins were found to colocalize in cytolitic granules by confocal microscopy (Fig. 5 A). IgG control mAbs showed no significant staining regardless of isotype (data not shown). Further studies by immunoelectron microscopy confirmed the presence of granulysin in the cytolitic granules of both the JR.2 line and the 12G12 clone (Fig. 5 B) as demonstrated by the presence of immunogold particles in cytolitic granules after staining with an antigranulysin mAb.

TCR- γ/δ -mediated Recognition of CD1c. Although studies using mAb blocking and transfection of CD1c-encoding expression vectors into recipient targets showed directly that CD1c is the critical molecule recognized by $V\delta 1^+$ γ/δ T cells, we sought to determine the role of the TCR in this recognition. To demonstrate that the TCR- γ/δ mediates recognition of CD1c, a series of mAbs blocking experiments and TCR transfection studies was carried out. mAbs against TCR- α/β and TCR- γ/δ were tested for their ability to block JR.2 T cell lysis of C1R CD1c transfectant cells. Only the mAb against the TCR- γ/δ blocked killing (83% inhibition), suggesting that CD1c recognition was TCR mediated (Fig. 6 A).




Figure 6. TCR- γ/δ mediates CD1c recognition. (A) JR.2.1 T cell clone lysed C1R CD1c cells. Lysis was blocked by anti-CD1c mAb (F10/2A3; not shown) as well by anti-C δ TCR mAb (anti-TCR- δ 1). E/T 30:1. (B) JR.2 V γ 2/V δ 1 transfectant recognizes CD1c. The TCR- J.RT3-T3 cell line was transfected with cDNAs encoding the V γ 2/V δ 1 TCR from the JR.2 cell line or was mock transfected. The resulting JR.2/J.RT3 or Mock/J.RT3 cell lines were cultured with CD1⁺ dendritic cells in the presence or absence of mAbs to CD1a, CD1b, and CD1c, and the supernatants were harvested after 24 h. IL-2 release was assessed by the proliferation of the IL-2-dependent HT-2 T cell line. Note that the JR.2/J.RT3 but not the Mock/J.RT3 cell line released IL-2 when cultured with CD1⁺ monocyte-derived dendritic cells. This IL-2 release was completely blocked by the addition of an anti-CD1c mAb.

To formally demonstrate that the TCR plays a direct role in the recognition of CD1c, we cloned the TCR γ and δ chains from the CD1c-reactive JR.2 T cell clone by reverse transcription PCR, transfected the JR.2 V γ 2/V δ 1 TCR into the TCR-deficient recipient Jurkat T cell tumor cell line, and measured the ability of the transfectants to produce IL-2 in response to CD1⁺ dendritic cells. When cultured with CD1⁺ dendritic cells, the V γ 2/V δ 1 TCR transfectant (JR.2/J.RT3) secreted IL-2 as judged in bioassays by the proliferation of HT-2 cells (34,000 cpm; Fig. 6 B). The production of IL-2 was inhibited completely by adding anti-CD1c mAb, but not anti-CD1a or anti-CD1b mAbs. Mock transfectant J.RT3 re-

ipient cells (Mock/J.RT3) did not produce IL-2 in response to CD1⁺ presenting cells. Thus, γ/δ T cell specificity for CD1c is determined by the V δ 1-containing TCR, as transfection of this TCR confers reactivity upon the recipient.

CD1c-reactive γ/δ T Cells Use Diverse TCRs. The recognition of CD1d antigen-presenting molecules by α/β T cells in the absence of exogenous foreign antigens has been observed for a unique subset of CD4⁺ mouse NKT cells expressing a canonical V α 14-J α 281 rearrangement (63) as well as for human T cells expressing the homologous V α 24-J α 18 rearrangement (50). The CD1c-reactive γ/δ T cell lines derived here express γ/δ TCRs encoded by V δ 1 gene segments (Fig. 1). To determine the nature of γ/δ TCR diversity involved in recognition of CD1c, we amplified and sequenced the junctional regions of the γ and δ chains of a JR.2 subclone. These sequences were compared with those of γ/δ T cell line IDP2 (64, 65), previously characterized as specific for CD1c (60). Both IDP2 and JR.2 rearranged V δ 1 to D δ 2, D δ 3, and J δ 1 gene segments to encode the junctional region of V δ 1 chain. However the amino acid sequences of the V δ 1 chains in the CDR3 region of the two T cells have almost no homology and are of substantially different lengths. The V γ 2 chains of the clones exhibited limited N region diversity but used different J γ segments (J γ 1.2 for JR.2 and J γ 2.3 for IDP2; Fig. 7). V γ usage differed in the XV.1 cell line that used the V γ 1.4 gene segment and another CD1c-reactive V δ 1⁺ T cell clone (66) that used either V γ 1.3 or V γ 1.4 with J γ 2.3. Thus, CD1c-reactive γ/δ T cells have divergent V δ 1 junctional regions rather than a canonical TCR as found on CD1d-reactive NKT cells and can use different V γ gene segments. However, since only a limited number of CD1c-reactive TCRs has been analyzed, more TCRs will need to be sequenced to determine the diversity of TCR junctions used by CD1c-reactive γ/δ T cells.

Discussion

We have described and functionally analyzed γ/δ T cell lines and clones derived from normal donors that directly




Figure 7. Diverse CDR3 junction region of two CD1c-reactive V γ 2/V δ 1 T cell clones, showing nucleotide sequence analysis of the γ and the δ gene rearrangements of JR.2 CD1c-reactive clone. Deduced amino acid sequences are in the single letter code. IDP2 sequence data are taken from Hata et al. (reference 64) and Krangel et al. (reference 65) and are included for comparison.

recognize human CD1c molecules. Compared with other CD1-reactive T cells described in humans, these are remarkable because their recognition was not dependent on the presence of foreign lipid or glycolipid antigens. These CD1c-reactive T cells constitute a subset of directly reactive T cells that may be part of the innate immune system. A certain similarity exists with murine CD4⁺ α/β NKT cells reactive to murine CD1.1, a homologue of human CD1d (63, 67). These murine α/β T cells express markers of the NK locus (NK1.1) and express nondiverse, canonical V α 14-J α 281 TCR α chains frequently paired with diverse V β 8 chains. The production of IL-4 and IFN- γ by these NKT cells has been implicated as an early immunoregulatory response in mice (63, 68–70), and activation of these cells in vivo by exogenous antigen directs conventional T cells to the Th2 phenotype (71, 72). The secretion of large amounts of IFN- γ in response to exogenous antigen can also inhibit ongoing Th2 responses (69). In contrast, the γ/δ T cells described here are reactive to CD1c, another CD1 family member. Recognition of CD1c is through the γ/δ TCR as directly demonstrated by transfection of the TCR (Fig. 6). TCR γ and δ gene sequences from these T cells revealed the incorporation of template-independent N nucleotides resulting in diverse noncanonical V-(D)-J junctions. Moreover, CD1c-reactive γ/δ T cells lacked expression of NKR-P1A (Fig. 1), a human member of the family of NK receptors analogous to murine NK1.1 (73).

It is not clear if the CD1c-reactive γ/δ T cells are responding to CD1c molecules alone or to CD1c molecules containing a self-lipid molecule. The recent findings that insect-derived murine CD1d molecules contain a bound lipid in their antigen binding pocket (74), that mammalian CD1 can bind cellular glycosylphosphatidylinositol (75), and that murine and human V α 14 invariant chain, NKR⁺ α/β T cells respond to autologous CD1d molecules loaded with glycosylceramides (76, 77) suggest that the recognition of murine CD1d involves CD1d-bound lipid. Thus, it seems likely that the γ/δ T cells described here are recognizing a self-lipid presented by CD1c. However, such a lipid must be broadly distributed, being present in B cells, HeLa cells, and dendritic cells, as CD1c expressed on these various cell sources was recognized in each case (Fig. 2, A and B).

The recognition of CD1c by human γ/δ T cells may be an important form of antigen recognition by these T cells. Recognition of CD1c by γ/δ T cells in the absence of exogenous foreign antigens could be compared with the alloreactive and autoreactive recognition of MHC class I and class II proteins by α/β T cells that are easily detected in MLRs (mixed lymphocyte reactions) and AMLRs (autologous mixed lymphocyte reactions) (78). Thus, autoreactivity to restricting elements may be a common theme between TCRs that recognize foreign antigens in the context of MHC or CD1.

The best-studied human γ/δ T cell reactivity is to small aliphatic phosphate molecules typically composed of an isoprenoid chain linked to a pyrophosphate moiety (24). Recently, we have characterized another class of aliphatic molecules typically consisting of four or five carbons chains

linked to a primary amine (26). Recognition of these small aliphatic compounds may represent a form of pattern recognition by the major circulating γ/δ T cell subset defined by expression of V δ 2 TCR. The V γ 2/V δ 2 subset of γ/δ T cells accounts for the vast majority of human γ/δ T cells in the circulation.

Here, we characterized recognition by the V δ 1 subset of γ/δ T cells, the major population in tissues such as the intestine and the spleen. We suggest that this subset of γ/δ T cells focuses on recognition of nonpolymorphic cell surface molecules related in structure to classical MHC molecules. Groh and Spies have recently demonstrated that some members of this γ/δ T cell subset killed target cells expressing stress-induced MICA and MICB MHC-encoded structures (32) through reactivity with the NKG2D NK receptor (33). Although mAbs specific for the γ/δ TCR blocked γ/δ T cell killing of MICA⁺ targets (32), the blocking was partial and TCR gene transfer was not done. It remains to be definitely determined if V δ 1⁺ TCRs can recognize MICA. Here, we show that some members of this same receptor subset recognize nonpolymorphic CD1c. Two other described γ/δ T cell clones reactive with human CD1c also expressed TCRs encoded by the V δ 1 gene segment (60, 66). Moreover, we have derived and partially characterized two additional V δ 1⁺ T cell lines that recognize CD1c (data not shown). Thus, these findings suggest that CD1c reactivity may be common among V δ 1⁺ T cells. However, it is clear that many V δ 1⁺ T cells are not CD1c reactive (data not shown), and further studies will be needed to determine the frequency of CD1c-reactive γ/δ T cells. Additionally, since the above γ/δ T cell lines were derived by in vitro culture, primary resident CD1c-reactive γ/δ T cells in situ may not be able to respond to CD1c due to anergy.

Direct recognition of CD1c by V δ 1⁺ T cells may represent a bridge between innate and adaptive immunity in a similar fashion to recognition of CD1d by murine and human NK⁺ α/β T cells (50, 63). Such CD1d-reactive α/β T cells are critically required to mediate IL-12-induced tumor immunity in mice (68), can polarize other T cells to a Th2 phenotype when activated by exogenous antigen (71, 72), and play regulatory roles in listerial infection (79) and Th2 responses (69). We speculate that CD1c-reactive γ/δ T cells also have unique immunological functions. Evidence from in vivo and in vitro experiments has demonstrated that γ/δ T cells play roles complementary to those of conventional α/β T cells in the host defense against infectious agents, in autoimmune diseases, and in tumor surveillance. We demonstrate here that both the CD1c- and the phosphate antigen-reactive γ/δ T cells are cytotoxic and can use the perforin-dependent cytotoxicity pathway. The perforin pathway plays an important role in antiviral immunity, resistance to intracellular bacteria, tumor surveillance, and other immune functions because it is the primary cytotoxic effector mechanism in host defense (61). Further underscoring the potential functional importance of γ/δ T cells in vivo, we show for the first time that γ/δ T cells express granulysin, a potent antimicrobial protein that in conjunction with perforin kills intracellular *M. tuberculosis*

organisms (51). Granulysin also kills a wide variety of microbes in vitro (51, 62). Moreover, the CD1c-reactive γ/δ T cells possess a Th1-like cytokine profile, producing significant amount of IFN- γ and IL-2.

CD1 molecules can be directly induced on monocytes as they differentiate into dendritic cells in response to GM-CSF or indirectly in response to agents such as bacteria or inflammatory products that induce the secretion of GM-CSF. Such de novo expression of CD1c can be seen on dendritic cells in granulomas induced by *Mycobacterium leprae* infection in leprosy (80). We speculate that this new expression of CD1c, or stress-induced endogenous lipid antigens presented by CD1c, could then activate tissue V δ 1⁺ T cells that are specific for CD1c. After their activation, the CD1c-reactive γ/δ T cells could play a role in directing conventional α/β T cells to the acquisition of a Th1 phenotype through their secretion of IFN- γ . The early secretion of cytokines by murine γ/δ T cells during responses to infectious agents provides evidence for such a role (81). These cells, then, could serve an analogous functional role to that proposed for V α 14⁺NK1.1⁺ T cells, except CD1c recognition by γ/δ T cells would lead to IFN- γ production rather than to IFN- γ and IL-4 production that is seen with most NK1.1⁺ T cells. As CD1c-specific γ/δ T cells are cytotoxic and express granulysin, they could also lyse infected dendritic cells via the perforin pathway and kill released intracellular bacteria by secretion of granulysin. In addition, as CD1c is expressed on some circulating B cells and on mantle zone B cells in lymph nodes (although at low levels) (82–84), V δ 1⁺ γ/δ T cells may play a role in T cell–B cell interactions.

Since various professional APCs, including B cells, tissue dendritic cells, activated monocytes, and activated macrophages in humans all express CD1c and since γ/δ T cells recognizing CD1c appear to do so in the absence of foreign antigens before foreign antigen-specific T cells differentiate and expand, upregulation of CD1c could represent an innate danger signal through the activation of CD1c-specific γ/δ T cells. Thus, such γ/δ T cells may play an early role by activating APCs via Th1 cytokines or may play a role in host defense by directly killing microbes via the combination of perforin and granulysin.

We thank Jack F. Bukowski for helpful suggestions.

This work was supported by the National Institutes of Health (M.B. Brenner, S.A. Porcelli, A.M. Krensky, and C.T. Morita), the Arthritis Foundation (S.A. Porcelli and C.T. Morita), and the American College of Rheumatology (C.T. Morita). A.M. Krensky is a Burroughs Wellcome Scholar in Experimental Therapeutics and the Shelagh Galligan Professor of Pediatrics.

Submitted: 15 September 1999

Revised: 10 November 1999

Accepted: 17 November 1999

Released online: 13 March 2000

References

1. Hiromatsu, K., Y. Yoshikai, G. Matsuzaki, S. Ohga, K. Muramori, K. Matsumoto, J.A. Bluestone, and K. Nomoto. 1992. A protective role of γ/δ T cells in primary infection with *Listeria monocytogenes* in mice. *J. Exp. Med.* 175:49–56.
2. Ladel, C.H., C. Blum, A. Dreher, K. Reifenberg, and S.H.E. Kaufmann. 1995. Protective role of γ/δ T cells and α/β T cells in tuberculosis. *Eur. J. Immunol.* 25:2877–2881.
3. Ladel, C.H., J. Hess, S. Daugelet, P. Mombaerts, S. Tonegawa, and S.H.E. Kaufmann. 1995. Contribution of α/β and γ/δ T lymphocytes to immunity against *Mycobacterium bovis* bacillus Calmette Guérin: studies with T cell receptor-deficient mutant mice. *Eur. J. Immunol.* 25:838–846.
4. D'Souza, C.D., A.M. Cooper, A.A. Frank, R.J. Mazzaccaro, B.R. Bloom, and I.M. Orme. 1997. An anti-inflammatory role for γ/δ T lymphocytes in acquired immunity to *Mycobacterium tuberculosis*. *J. Immunol.* 158:1217–1221.
5. Tsuji, M., P. Mombaerts, L. Lefrancois, R.S. Nussenzweig, F. Zavala, and S. Tonegawa. 1994. γ/δ T cells contribute to immunity against the liver stages of malaria in α/β T-cell-deficient mice. *Proc. Natl. Acad. Sci. USA.* 91:345–349.
6. Sciammas, R., P. Kodukula, Q. Tang, R.L. Hendricks, and J.A. Bluestone. 1997. T cell receptor- γ/δ cells protect mice from herpes simplex virus type 1-induced lethal encephalitis. *J. Exp. Med.* 185:1969–1975.
7. King, D.P., D.M. Hyde, K.A. Jackson, D.M. Novosad, T.N. Ellis, L. Putney, M.Y. Stovall, L.S. Van Winkle, B.L. Beaman, and D.A. Ferrick. 1999. Protective response to pulmonary injury requires γ/δ T lymphocytes. *J. Immunol.* 162: 5033–5036.
8. Jouen-Beades, F., E. Paris, C. Dieulois, J.-F. Lemeland, V. Barre-Dezelus, S. Marret, G. Humbert, J. Leroy, and F. Tron. 1997. *In vivo* and *in vitro* activation and expansion of γ/δ T cells during *Listeria monocytogenes* infection in humans. *Infect. Immun.* 65:4267–4272.
9. Balbi, B., M.T. Valle, S. Oddera, D. Giunti, F. Manca, G.A. Rossi, and L. Allegra. 1993. T-lymphocytes with γ/δ ⁺ V δ 2⁺ antigen receptors are present in increased proportions in a fraction of patients with tuberculosis or with sarcoidosis. *Am. Rev. Respir. Dis.* 148:1685–1690.
10. Bertotto, A., R. Gerli, F. Spinozzi, C. Muscat, F. Scalise, G. Castellucci, M. Sposito, F. Candio, and R. Vaccaro. 1993. Lymphocytes bearing the γ/δ T cell receptor in acute *Brucella melitensis* infection. *Eur. J. Immunol.* 23:1177–1180.
11. Caldwell, C.W., E.D. Everett, G. McDonald, Y.W. Yesus, and W.E. Roland. 1995. Lymphocytosis of γ/δ T cells in human ehrlichiosis. *Am. J. Clin. Pathol.* 103:761–766.
12. Haas, W., P. Pereira, and S. Tonegawa. 1993. Gamma/delta cells. *Annu. Rev. Immunol.* 11:637–685.
13. Porcelli, S., M.B. Brenner, and H. Band. 1991. Biology of the human γ/δ T-cell receptor. *Immunol. Rev.* 120:137–183.
14. Johnson, R.M., D.W. Lancki, A.I. Sperling, R.F. Dick, P.G. Spear, F.W. Fitch, and J.A. Bluestone. 1992. A murine CD4⁻, CD8⁻ T cell receptor- γ/δ T lymphocyte clone specific for herpes simplex virus glycoprotein I. *J. Immunol.* 148:983–988.
15. Schild, H., N. Mavaddat, C. Litzenger, E.W. Ehrlich, M.M. Davis, J.A. Bluestone, L. Matis, R.K. Draper, and Y.-h. Chien. 1994. The nature of major histocompatibility complex recognition by γ/δ T cells. *Cell.* 76:29–37.
16. Crowley, M.P., Z. Reich, N. Mavaddat, J.D. Altman, and Y.-h. Chien. 1997. The recognition of the nonclassical major histocompatibility complex (MHC) class I molecule, T10, by the γ/δ T cell, G8. *J. Exp. Med.* 185:1223–1230.
17. Rock, E.P., P.R. Sibbald, M.M. Davis, and Y.-h. Chien. 1994. CDR3 length in antigen-specific immune receptors. *J. Exp. Med.* 179:323–328.

18. Li, H., M.I. Lebedeva, A.S. Llera, B.A. Fields, M.B. Brenner, and R.A. Mariuzza. 1998. Structure of the V δ domain of a human $\gamma\delta$ T-cell antigen receptor. *Nature*. 391:502–506.
19. Morita, C.T., H.K. Lee, D.S. Leslie, Y.T. Tanaka, J.F. Bukowski, and E. Märker-Hermann. 1999. Recognition of nonpeptide prenyl pyrophosphate antigens by human $\gamma\delta$ T cells. *Microbes Infect.* 1:175–186.
20. Kabelitz, D., A. Bender, S. Schondelmaier, B. Schoel, and S.H.E. Kaufmann. 1990. A large fraction of human peripheral blood γ/δ^+ T cells is activated by *Mycobacterium tuberculosis* but not by its 65-kD heat shock protein. *J. Exp. Med.* 171:667–679.
21. Pfeffer, K., B. Schoel, H. Gulle, S.H.E. Kaufmann, and H. Wagner. 1990. Primary responses of human T cells to mycobacteria: a frequent set of γ/δ T cells are stimulated by protease-resistant ligands. *Eur. J. Immunol.* 20:1175–1179.
22. Panchamoorthy, G., J. McLean, R.L. Modlin, C.T. Morita, S. Ishikawa, M.B. Brenner, and H. Band. 1991. A predominance of the T cell receptor V γ 2/V δ 2 subset in human mycobacteria-responsive T cells suggests germline gene encoded recognition. *J. Immunol.* 147:3360–3369.
23. Morita, C.T., E.M. Beckman, J.F. Bukowski, Y. Tanaka, H. Band, B.R. Bloom, D.E. Golan, and M.B. Brenner. 1995. Direct presentation of nonpeptide prenyl pyrophosphate antigens to human $\gamma\delta$ T cells. *Immunity*. 3:495–507.
24. Tanaka, Y., C.T. Morita, Y. Tanaka, E. Nieves, M.B. Brenner, and B.R. Bloom. 1995. Natural and synthetic non-peptide antigens recognized by human $\gamma\delta$ T cells. *Nature*. 375:155–158.
25. Tanaka, Y., S. Sano, E. Nieves, G. De Libero, D. Roca, R.L. Modlin, M.B. Brenner, B.R. Bloom, and C.T. Morita. 1994. Nonpeptide ligands for human $\gamma\delta$ T cells. *Proc. Natl. Acad. Sci. USA*. 91:8175–8179.
26. Bukowski, J.F., C.T. Morita, and M.B. Brenner. 1999. Human $\gamma\delta$ T cells recognize alkylamines derived from microbes, edible plants, and tea: implications for innate immunity. *Immunity*. 11:57–65.
27. Lang, F., M.A. Peyrat, P. Constant, F. Davodeau, J. David-Ameline, Y. Poquet, H. Vié, J.J. Fournié, and M. Bonneville. 1995. Early activation of human V γ 9V δ 2 T cell broad cytotoxicity and TNF production by nonpeptidic mycobacterial ligands. *J. Immunol.* 154:5986–5994.
28. Holoshitz, J., N.C. Romzek, Y. Jia, L. Wagner, L.M. Vila, S.-J. Chen, J.M. Wilson, and D.R. Karp. 1993. MHC-independent presentation of mycobacteria to human $\gamma\delta$ T cells. *Int. Immunol.* 5:1437–1443.
29. Bukowski, J.F., C.T. Morita, Y. Tanaka, B.R. Bloom, M.B. Brenner, and H. Band. 1995. V γ 2V δ 2 TCR-dependent recognition of non-peptide antigens and Daudi cells analyzed by TCR gene transfer. *J. Immunol.* 154:998–1006.
30. Bukowski, J.F., C.T. Morita, H. Band, and M.B. Brenner. 1998. Crucial role of TCR γ chain junctional region in prenyl pyrophosphate antigen recognition by $\gamma\delta$ T cells. *J. Immunol.* 161:286–293.
31. Falini, B., L. Flenghi, S. Pileri, P. Pelicci, M. Fagioli, M.F. Martelli, L. Moretta, and E. Ciccone. 1989. Distribution of T cells bearing different forms of the T cell receptor γ/δ in normal and pathological human tissues. *J. Immunol.* 143:2480–2488.
32. Groh, V., A. Steinle, S. Bauer, and T. Spies. 1998. Recognition of stress-induced MHC molecules by intestinal epithelial $\gamma\delta$ T cells. *Science*. 279:1737–1740.
33. Bauer, S., V. Groh, J. Wu, A. Steinle, J.H. Phillips, L.L. Lanier, and T. Spies. 1999. Activation of NK cells and T cells by NKG2D, a receptor for stress-inducible MICA. *Science*. 285:727–729.
34. Li, P., S.T. Willie, S. Bauer, D.L. Morris, T. Spies, and R.K. Strong. 1999. Crystal structure of the MHC class I homolog MIC-A, a $\gamma\delta$ T cell ligand. *Immunity*. 10:577–584.
35. Beckman, E.M., A. Melian, S.M. Behar, P.A. Sieling, D. Chatterjee, S.T. Furlong, R. Matsumoto, J.P. Rosat, R.L. Modlin, and S.A. Porcelli. 1996. CD1c restricts responses of mycobacteria-specific T cells. Evidence for antigen presentation by a second member of the human CD1 family. *J. Immunol.* 157:2795–2803.
36. Spits, H., G. Keizer, J. Borst, C. Terhorst, A. Hekman, and J.E. de Vries. 1983. Characterization of monoclonal antibodies against cell surface molecules associated with cytotoxic activity of natural and activated killer cells and cloned CTL lines. *Hybridoma*. 2:423–437.
37. Band, H., F. Hochstenbach, J. McLean, S. Hata, M.S. Krangel, and M.B. Brenner. 1987. Immunochemical proof that a novel rearranging gene encodes the T cell receptor δ subunit. *Science*. 238:682–684.
38. Wu, Y.J., W.T. Tian, R.M. Snider, C. Rittershaus, P. Rogers, L. LaManna, and S.H. Ip. 1988. Signal transduction of γ/δ T cell antigen receptor with a novel mitogenic anti- δ antibody. *J. Immunol.* 141:1476–1479.
39. Jitsukawa, S., F. Faure, M. Lipinski, F. Triebel, and T. Hercend. 1987. A novel subset of human lymphocytes with a T cell receptor- γ complex. *J. Exp. Med.* 166:1192–1197.
40. Band, H., F. Hochstenbach, C.M. Parker, J. McLean, M.S. Krangel, and M.B. Brenner. 1989. Expression of human T cell receptor- $\gamma\delta$ structural forms. *J. Immunol.* 142:3627–3633.
41. Linsley, P.S., J.L. Greene, P. Tan, J. Bradshaw, J.R. Ledbetter, C. Anasetti, and N.K. Damle. 1992. Coexpression and functional cooperation of CTLA-4 and CD28 on activated T lymphocytes. *J. Exp. Med.* 176:1595–1604.
42. Martin, L.H., F. Calabi, F.A. Lefebvre, C.A. Bilsland, and C. Milstein. 1987. Structure and expression of the human thymocyte antigens CD1a, CD1b, and CD1c. *Proc. Natl. Acad. Sci. USA*. 84:9189–9193.
- 42a. Grant, E.P., M. Degano, J.P. Rosat, S. Stenger, R.L. Modlin, I.A. Wilson, S.A. Porcelli, and M.B. Brenner. 1999. Molecular recognition of lipid antigens by T cell receptors. *J. Exp. Med.* 189:195–205.
43. Behar, S.M., S.A. Porcelli, E.M. Beckman, and M.B. Brenner. 1995. A pathway of costimulation that prevents anergy in CD28⁻ T cells: B7-independent costimulation of CD1-restricted T cells. *J. Exp. Med.* 182:2007–2018.
44. Olive, D., P. Dubreuil, and C. Mawas. 1984. Two distinct TL-like molecular subsets defined by monoclonal antibodies on the surface of human thymocytes with different expression on leukemia lines. *Immunogenetics*. 20:253–264.
45. De Libero, G., G. Casorati, C. Giachino, C. Carbonara, N. Migone, P. Matzinger, and A. Lanzavecchia. 1991. Selection by two powerful antigens may account for the presence of the major population of human peripheral γ/δ T cells. *J. Exp. Med.* 173:1311–1322.
46. Hanson, D.A., A. Kasper, F.R. Poulain, and A.M. Krensky. 1999. Biosynthesis of granulysin, a novel cytolytic molecule. *Mol. Immunol.* 36:413–422.
47. Beckman, E.M., S.A. Porcelli, C.T. Morita, S.M. Behar, S.T. Furlong, and M.B. Brenner. 1994. Recognition of a lipid antigen by CD1-restricted $\alpha\beta^+$ T cells. *Nature*. 372:691–694.
48. Porcelli, S., C.T. Morita, and M.B. Brenner. 1992. CD1b restricts the response of human CD4⁻8⁻ T lymphocytes to a microbial antigen. *Nature*. 360:593–597.

49. Sugita, M., R.M. Jackman, E. van Donselaar, S.M. Behar, R.A. Rogers, P.J. Peters, M.B. Brenner, and S.A. Porcelli. 1996. Cytoplasmic tail-dependent localization of CD1b antigen-presenting molecules to MHCs. *Science*. 273:349–352.
50. Exley, M., J. Garcia, S.P. Balk, and S. Porcelli. 1997. Requirements for CD1d recognition by human invariant V α 24⁺ CD4⁻CD8⁻ T cells. *J. Exp. Med.* 186:109–120.
51. Stenger, S., D.A. Hanson, R. Teitelbaum, P. Dewan, K.R. Niazi, C.J. Froelich, T. Ganz, S. Thoma-Uszynski, A. Melian, C. Bogdan, et al. 1998. An antimicrobial activity of cytolytic T cells mediated by granulysin. *Science*. 282:121–125.
52. Peters, P.J., J.J. Neefjes, V. Oorschot, H.L. Ploegh, and H.J. Geuze. 1991. Segregation of MHC class II molecules from MHC class I molecules in the Golgi complex for transport to lysosomal compartments. *Nature*. 349:669–676.
53. Chomczynski, P., and N. Sacchi. 1987. Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal. Biochem.* 162:156–159.
54. Brawley, J.V., and P. Concannon. 1996. Modulation of promiscuous T cell receptor recognition by mutagenesis of CDR2 residues. *J. Exp. Med.* 183:2043–2051.
55. Weiss, A., and J.D. Stobo. 1984. Requirement for the coexpression of T3 and the T cell antigen receptor on a malignant human T cell line. *J. Exp. Med.* 160:1284–1299.
56. Brenner, M.B., J.L. Strominger, and M.S. Krangel. 1988. The $\gamma\delta$ T cell receptor. *Adv. Immunol.* 43:133–192.
57. Morita, C.T., S. Verma, P. Aparicio, C. Martinez-A, H. Spits, and M.B. Brenner. 1991. Functionally distinct subsets of human γ/δ T cells. *Eur. J. Immunol.* 21:2999–3007.
58. Zwirner, N.W., M.A. Fernández-Viña, and P. Stastny. 1998. MICA, a new polymorphic HLA-related antigen, is expressed mainly by keratinocytes, endothelial cells, and monocytes. *Immunogenetics*. 47:139–148.
59. Stenger, S., R.J. Mazzaccaro, K. Uyemura, S. Cho, P.F. Barnes, J.P. Rosat, A. Sette, M.B. Brenner, S.A. Porcelli, B.R. Bloom, and R.L. Modlin. 1997. Differential effects of cytolytic T cell subsets on intracellular infection. *Science*. 276:1684–1687.
60. Porcelli, S., M.B. Brenner, J.L. Greenstein, S.P. Balk, C. Terhorst, and P.A. Bleicher. 1989. Recognition of cluster of differentiation 1 antigens by human CD4⁻CD8⁻ cytolytic T lymphocytes. *Nature*. 341:447–450.
61. Kagi, D., B. Ledermann, K. Burki, R.M. Zinkernagel, and H. Hengartner. 1996. Molecular mechanisms of lymphocyte-mediated cytotoxicity and their role in immunological protection and pathogenesis *in vivo*. *Annu. Rev. Immunol.* 14:207–232.
62. Pena, S.V., D.A. Hanson, B.A. Carr, T.J. Goralski, and A.M. Krensky. 1997. Processing, subcellular localization, and function of 519 (granulysin), a human late T cell activation molecule with homology to small, lytic, granule proteins. *J. Immunol.* 158:2680–2688.
63. Bendelac, A., M.N. Rivera, S.H. Park, and J.H. Roark. 1997. Mouse CD1-specific NK1 T cells: development, specificity, and function. *Annu. Rev. Immunol.* 15:535–562.
64. Hata, S., M. Brenner, and M. Krangel. 1987. Identification of putative human T cell receptor δ complementary DNA clones. *Science*. 238:678–682.
65. Krangel, M., H. Band, S. Hata, J. McLean, and M. Brenner. 1987. Structurally divergent human T cell receptor proteins encoded by distinct C γ genes. *Science*. 237:64–67.
66. Faure, F., S. Jitsukawa, C. Miossec, and T. Hercend. 1990. CD1c as a target recognition structure for human T lymphocytes: analysis with peripheral blood γ/δ cells. *Eur. J. Immunol.* 20:703–706.
67. Porcelli, S.A. 1995. The CD1 family: a third lineage of antigen-presenting molecules. *Adv. Immunol.* 59:1–98.
68. Cui, J., T. Shin, T. Kawano, H. Sato, E. Kondo, I. Toura, Y. Kaneko, H. Koseki, M. Kanno, and M. Taniguchi. 1997. Requirement for V α 14 NKT cells in IL-12-mediated rejection of tumors. *Science*. 278:1623–1626.
69. Cui, J., N. Watanabe, T. Kawano, M. Yamashita, T. Kamata, C. Shimizu, M. Kimura, E. Shimizu, J. Koike, H. Koseki, et al. 1999. Inhibition of T helper cell type 2 cell differentiation and immunoglobulin E response by ligand-activated V α 14 natural killer T cells. *J. Exp. Med.* 190:783–792.
70. Carnaud, C., D. Lee, O. Donnars, S.H. Park, A. Beavis, Y. Koezuka, and A. Bendelac. 1999. Cross-talk between cells of the innate immune system: NKT cells rapidly activate NK cells. *J. Immunol.* 163:4647–4650.
71. Burdin, N., L. Brossay, and M. Kronenberg. 1999. Immunization with α -galactosylceramide polarizes CD1-reactive NK T cells towards Th2 cytokine synthesis. *Eur. J. Immunol.* 29:2014–2025.
72. Singh, N., S. Hong, D.C. Scherer, I. Serizawa, N. Burdin, M. Kronenberg, Y. Koezuka, and L. Van Kaer. 1999. Activation of NK T cells by CD1d and α -galactosylceramide directs conventional T cells to the acquisition of a Th2 phenotype. *J. Immunol.* 163:2373–2377.
73. Lanier, L.L., C. Chang, and J.H. Phillips. 1994. Human NKR-P1A. A disulfide-linked homodimer of the C-type lectin superfamily expressed by a subset of NK and T lymphocytes. *J. Immunol.* 153:2417–2428.
74. Zeng, Z.-H., A.R. Castaño, B.W. Segelke, E.A. Stura, P.A. Peterson, and I.A. Wilson. 1997. Crystal structure of mouse CD1: an MHC-like fold with a large hydrophobic binding groove. *Science*. 277:339–345.
75. Joyce, S., A.S. Woods, J.W. Yewdell, J.R. Bennink, A.D. De Silva, A. Boesteanu, S.P. Balk, R.J. Cotter, and R.R. Brutkiewicz. 1998. Natural ligand of mouse CD1d1: cellular glycosylphosphatidylinositol. *Science*. 279:1541–1544.
76. Kawano, T., J. Cui, Y. Koezuka, I. Toura, Y. Kaneko, K. Motoki, H. Ueno, R. Nakagawa, H. Sato, E. Kondo, et al. 1997. CD1d-restricted and TCR-mediated activation of V α 14 NKT cells by glycosylceramides. *Science*. 278:1626–1629.
77. Spada, F.M., Y. Koezuka, and S.A. Porcelli. 1998. CD1d-restricted recognition of synthetic glycolipid antigens by human natural killer T cells. *J. Exp. Med.* 188:1529–1534.
78. Sherman, L.A., and S. Chattopadhyay. 1993. The molecular basis of allorecognition. *Annu. Rev. Immunol.* 11:385–402.
79. Szalay, G., C.H. Ladel, C. Blum, L. Brossay, M. Kronenberg, and S.H. Kaufmann. 1999. Anti-CD1 monoclonal antibody treatment reverses the production patterns of TGF- β 2 and Th1 cytokines and ameliorates listeriosis in mice. *J. Immunol.* 162:6955–6958.
80. Sieling, P.A., D. Jullien, M. Dahlem, T.F. Tedder, T.H. Rea, R.L. Modlin, and S.A. Porcelli. 1999. CD1 expression by dendritic cells in human leprosy lesions: correlation with effective host immunity. *J. Immunol.* 162:1851–1858.
81. Ferrick, D.A., M.D. Schrenzel, T. Mulvania, B. Hsieh, W.G. Ferlin, and H. Lepper. 1995. Differential production of interferon- γ and interleukin-4 in response to Th1- and Th2-stimulating pathogens by $\gamma\delta$ T cells *in vivo*. *Nature*. 373:255–257.
82. Small, T.N., R.W. Knowles, C. Keever, N.A. Kernan, N. Collins, R.J. O'Reilly, B. Dupont, and N. Flomenberg.

1987. M241 (CD1) expression on B lymphocytes. *J. Immunol.* 138:2864–2868.
83. Small, T.N., C. Keever, N. Collins, B. Dupont, R.J. O'Reilly, and N. Flomenberg. 1989. Characterization of B cells in severe combined immunodeficiency disease. *Hum. Immunol.* 25:181–193.
84. Cattoretti, G., E. Berti, C. Parravicini, M. Buscaglia, F. Cap-
pio, R. Caputo, A. Cerri, L. Crosti, D. Delia, G. Gaiera, and N. Polli. 1989. Expression of CD1 molecules on dendritic cells: ontogeny, epitope analysis on normal and malignant cells, and tissue distribution. *In Leukocyte Typing IV.* W. Knapp, B. Dorken, W.R. Gilks, E.P. Rieber, R.E. Schmidt, H. Stein, and A.E.G.K. von dem Borne, editors. Oxford University Press, Oxford. 263–264.