

Aplikace matematiky

Abraham Charnes; William Wager Cooper; Kenneth O. Kortanek
Semi-infinite programming, differentiability and geometric programming: Part II

Aplikace matematiky, Vol. 14 (1969), No. 1, 15–22

Persistent URL: <http://dml.cz/dmlcz/103204>

Terms of use:

© Institute of Mathematics AS CR, 1969

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

SEMI-INFINITE PROGRAMMING, DIFFERENTIABILITY AND GEOMETRIC PROGRAMMING: PART II

A. CHARNES*, W. W. COOPER**, and K. O. KORTANEK***

(Received March 20, 1967)

We propose to specialize the CCK duality theory¹), which associates as dual problems minimization of an arbitrary convex function over an arbitrary convex set in n -space with maximization of a linear function in non-negative variables of a generalized finite sequence space subject to a finite system of linear equations, to derive Kuhn-Tucker Theorem²) extensions in situations involving (partial) differentiability of objective and constraint functions. There are several ways to procure such generalizations as, for example, by means of non-differentiable analogs of quasi-saddle point conditions or in terms of a saddle point criterion itself. Since we are interested here in exploring extensions which involve some differentiability conditions, we shall proceed via the former course especially since these conditions themselves are analogs of first order conditions of the saddle point criterion.³)

For our purposes then, let $f(u)$, and $G(u) = (g_1(u), g_2(u), \dots, g_m(u))$ be defined over an open convex set K in R_n . We shall say that $f(u)$ is simple piecewise differentiably convex if $f(u) = \max_{j=1,2,\dots,N} \{f^{(j)}(u)\}$, where $f^{(j)}(u)$ is continuously differentiable and convex over K . We shall assume that $G(u)$ is continuously differentiable and concave, but the extension to simple piecewise concave functions will become apparent during the course of proof for functions of this class.

*) Northwestern University.

**) Carnegie-Mellon University.

***) Cornell University. Part of the research underlying this report was undertaken for the Pilot Program in Environmental Systems Analysis, NIH, 1 P10 ES 00098-01, at Cornell, and for the Office of Naval Research projects (Contract Nonr-1228(10), Project NR 047-021 at Northwestern University; Contract Nonr-760(24), Project NR 047-48 at Carnegie-Mellon) and for the U.S. Army Research Office-Durham, Contract No. DA-31-124-ARO-D-322 at Northwestern University.

¹) See Charnes-Cooper-Kortanek [4] and [5].

²) See Kuhn-Tucker [7] and Arrow-Hurwicz-Uzawa [1].

³) See Arrow-Hurwicz-Uzawa, ibid., where the authors show that in the case of differentiability the quasi-saddle point condition implies the saddle point condition.

Theorem. (Generalized Quasi-Saddle Point Theorem for Simple Piecewise Differentially Convex Functions). Let $f(u)$ and $G(u)$ have the properties defined above and consider the minimization problem

$$\min f(u)$$

subject to

$$G(u) \geqq 0 .$$

Assume the constraint set $C = \{u \mid G(u) \geqq 0\}$ has an interior point.⁴⁾ Then u^* in C is an optimal solution to the minimization problem if and only if there exists positive vectors

$$\eta^* = (\eta_*^{(1)}, \eta_*^{(2)}, \dots, \eta_*^{(N)}) \quad \text{and} \quad \lambda^* = (\lambda_*^{(1)}, \dots, \lambda_*^{(m)})$$

such that the following properties hold:

$$(1) \quad - \sum_{j=1}^N (\partial f^{(j)}|_{u^*}) \eta_*^{(j)} + \sum_{i=1}^m (\partial g^{(i)}|_{u^*}) \lambda_*^{(i)} = 0$$

$$(2) \quad \sum_{j \in J} \eta_*^{(j)} = 1$$

$$(3) \quad G(u^*)^T \lambda^* = 0 , \quad \text{and} \quad G(u^*) \geqq 0^5), \quad \text{where} \quad J = \{j \mid f^{(j)}(u^*) = f(u^*)\}$$

Preliminary Lemmas on Canonical Closure for Differential Systems. By introducing support systems for both objective and constraint functions, we obtain the following equivalent semi-infinite problem (I) with semi-infinite dual (II), which, for the moment, we write in general form.

$$\begin{array}{ll} (I) & (II) \\ \min z & \max \sum_{\alpha} d_{\alpha} \eta_{\alpha} + \sum_i c_i \lambda_i \\ z - u^T Q_{\alpha} \geqq d_{\alpha}, \alpha \in A & \sum_{\alpha} \eta_{\alpha} = 1 \\ u^T P_i \geqq c_i, i \in I & - \sum_{\alpha} Q_{\alpha} \eta_{\alpha} + \sum_i P_i \lambda_i = 0 \\ & \eta_{\alpha}, \lambda_i \geqq 0 . \end{array}$$

⁴⁾ This type of constraint qualification has strong intuitive appeal especially in the case of non-differentiability. However, it is known that non-differentiable analogs to the most general constraint qualification for which differentiable Lagrangian techniques are valid (see [6]) involve support systems which are themselves Farkas-Minkowski systems. (See [4] and [5]).

⁵⁾ Notationally speaking, $\partial f|_{u^*}$ is the gradient of f evaluated at u^* . We use superscripts to correspond to functions and subscripts to correspond to elements in the index set. Thus, $\partial f_{\alpha}^{(j)}$ denotes the gradient of $f(j)$ evaluated at the point $\alpha \in A$. For convenience, " $\alpha \in A$ " may be identified with " $u_{\alpha} \in A'$ ", when $A \subseteq R_n$.

Recall that a system of linear inequalities is *canonically closed* if it has interior points and the coefficient set is compact.⁶⁾ We need the following lemma.

Lemma 1. Suppose that the system is canonically closed and that u_* solves (I), i.e., the minimum $z_* = f(u)_*$ is attained. Then in the dual expression, (II), for z_* , the only supports which arise are those passing through the point (z_*, u_*) , i.e., the only support planes with $\eta_\alpha^* \neq 0$ and $\lambda_i^* \neq 0$ are those for which $z_* = u_*^T Q_\alpha + d_\alpha$ and $u_*^T P_i = c_i$.

Proof. By the extended dual theorem, there exist η, λ such that

$$z_* = \sum_\alpha d_\alpha \eta_\alpha^* + \sum_i c_i \lambda_i^*.$$

We must show that if $\eta_\alpha^* > 0$, then $z_* = u_*^T Q_\alpha + d_\alpha$ and if $\lambda_i^* > 0$, then $u_*^T P_i = c_i$. First, $z_* - u_*^T Q_\alpha \geq d_\alpha$, for all α . Hence

$$\sum_\alpha d_\alpha \eta_\alpha^* \leq \sum_\alpha z_* \eta_\alpha^* - \sum_\alpha (u_*^T Q_\alpha) \eta_\alpha^* = z_* - \sum_\alpha (u_*^T Q_\alpha) \eta_\alpha^*.$$

Therefore,

$$z_* = \sum_\alpha d_\alpha \eta_\alpha^* + \sum_i c_i \lambda_i^* \leq z_* - \sum_\alpha (u_*^T Q_\alpha) \eta_\alpha^* + \sum_i c_i \lambda_i^*.$$

i.e.,

$$(A) \quad -\sum_\alpha (u_*^T Q_\alpha) \eta_\alpha^* + \sum_i c_i \lambda_i^* \geq 0.$$

On the other hand, by dual feasibility,

$$u_*^T \left[-\sum_\alpha Q_\alpha \eta_\alpha^* + \sum_i P_i \lambda_i^* \right] = u_*^T(0) = 0.$$

However, since $u_*^T P_i \geq c_i$ for all i , we can rewrite this as follows:

$$(B) \quad 0 = \sum_\alpha -u_*^T Q_\alpha \eta_\alpha^* + \sum_i u_*^T P_i c_i \geq -\sum_\alpha u_*^T Q_\alpha \eta_\alpha^* + \sum_i c_i \lambda_i^*.$$

Therefore combining (A) and (B) we have,

$$\sum_\alpha u_*^T Q_\alpha \eta_\alpha^* = \sum_i c_i \lambda_i^*.$$

Two conclusions follow:

$$(C_1) \quad z_* = \sum_\alpha [u_*^T Q_\alpha + d_\alpha] \eta_\alpha^*, \text{ where } \sum_\alpha \eta_\alpha^* = 1, \eta \geq 0, \text{ and}$$

$$z_* \geq u_*^T Q_\alpha + d_\alpha. \text{ Hence } z_* = u_*^T Q_\alpha + d_\alpha \text{ for every } \alpha \text{ with } \eta_\alpha^* > 0.$$

$$(C_2) \quad \sum_i u_*^T P_i \lambda_i^* = \sum_i c_i \lambda_i^* \Rightarrow \sum_i (u_*^T P_i - c_i) \lambda_i^* = 0$$

Hence $\lambda_i^* > 0$ implies $u_*^T P_i = c_i$.

⁶⁾ See [4] and [5]. Note that canonical closure is a sufficient condition but not necessary for the validity of the extended dual theorem as pointed out in [4].

Proof of Theorem. With respect to the minimization problem of the Theorem, consider the particular semi-infinite equivalent

$$(I) \quad \begin{aligned} & \min z \\ & \text{subject to } z - u^T \partial f_\alpha^{(j)} \geq f^{(j)}(u_\alpha) - u_\alpha^T \partial f_\alpha^{(j)}, \quad j = 1, 2, \dots, N \\ & \quad u^T \partial g_\alpha^{(i)} \geq -g^{(i)}(u_\alpha) + u_\alpha^T \partial g^{(i)}(u_\alpha), \quad i = 1, 2, \dots, m \end{aligned}$$

for all $\alpha \in A$, where A is some index set in R_n (e.g. the convex constraint set C). Since C has interior points, it follows that this linear inequality system also does. Form a *canonical normalization*⁷⁾, (i.e., divide each inequality by a positive constant to make the sum of the absolute values of the coefficients sum to 1), to obtain an equivalent system with bounded coefficients and interiority.

$$(\hat{I}) \quad \begin{aligned} & \min z \\ & \text{subject to } \mu_\alpha^{(j)} z - u^T \partial f_\alpha^{(j)} \mu_\alpha^{(j)} \geq f^{(j)}(u_\alpha) \mu_\alpha^{(j)} - u_\alpha^T \partial f_\alpha^{(j)} \mu_\alpha^{(j)}, \quad \mu_\alpha^{(j)} > 0 \\ & \quad u^T \partial g_\alpha^{(i)} v_\alpha^{(i)} \geq -g^{(i)}(u_\alpha) v_\alpha^{(i)} + u_\alpha^T \partial g^{(i)}(u_\alpha) v_\alpha^{(i)}, \quad v_\alpha^{(i)} > 0 \end{aligned}$$

where $j = 1, 2, \dots, m$, and $\alpha \in A$.

Now form a canonical closure by possibly enlarging the index set to $\bar{A} \supseteq A$ and adjoining the corresponding limiting inequalities which are of the form:

$$\begin{aligned} \mu_\alpha^{(j)} z - u^T Q_\alpha^{(j)} &\geq d_\alpha^{(j)} \\ u^T P_\alpha^{(i)} &\geq c_\alpha^i \quad \text{for } \alpha \in \bar{A} - A. \end{aligned}$$

Let (\bar{I}) denote this new canonically closed equivalent (which differs from (\hat{I}) by only these possibly adjoined inequalities and also has interior points).

Now if u^* is optimal for (I) it is also optimal for the canonically closed equivalent (\bar{I}) and lemma 1 applies. However, any of the possibly *newly* adjoined inequalities which are actively involved in the dual are positive multiples of differential hyperplanes already in the system, for suppose one of them has a $\lambda^{(j)} > 0$, say, $\mu_\alpha^{(j)} z - u^T Q_\alpha^{(j)} \geq d_\alpha^{(j)}$ with $\alpha \in \bar{A} - A$. Then by lemma 1, the support plane $\mu_\alpha^{(j)} z - u^T Q_\alpha^{(j)} = d_\alpha^{(j)}$ contains the point $(u^*, f(u^*))$ i.e., $\mu_\alpha^{(j)} f(u^*) = \mu_\alpha^{(j)} z_* = u^{*T} Q_\alpha^{(j)} + d_\alpha^{(j)}$ or equivalently, the plane $\mu_\alpha^{(j)} z = u^T Q_\alpha^{(j)} + d_\alpha^{(j)}$ is tangent to the surface $z = f^{(j)}(u)$ at the point u^* . Since $f^{(j)}(u)$ is continuously differentiable, and since $\mu_\alpha^{(j)} z \geq u^T Q_\alpha^{(j)} + d_\alpha^{(j)}$ over C , this tangent plane is unique up to a constant *positive* multiple, and therefore we do not need to adjoin these additional inequalities. A similar argument obviously holds for the constraint functions.

⁷⁾ See [5], p. 114.

We now present the semi-infinite dual ($\bar{\Pi}$) and derive the conditions of the theorem.

$$\max \sum_j \sum_{\alpha} [f^{(j)}(u_{\alpha}) \mu_{\alpha}^{(j)} - u_{\alpha}^T \partial f_{\alpha}^{(j)} \mu_{\alpha}^{(j)}] \bar{\eta}_{\alpha}^{(j)} + \sum_j \sum_{\alpha} [-g^{(i)}(u_{\alpha}) v_{\alpha}^{(i)} + u_{\alpha}^T \partial g_{\alpha}^{(i)}(u_{\alpha}) v_{\alpha}^{(i)}] \bar{\lambda}_{\alpha}^{(i)}$$

subject to

$$\begin{aligned} \sum_j \mu_{\alpha}^{(j)} \bar{\eta}_{\alpha}^{(j)} &= 1 \\ -\sum_j \sum_{\alpha} (\partial f_{\alpha}^{(j)} \mu_{\alpha}^{(j)}) \bar{\eta}_{\alpha}^{(j)} + \sum_i \sum_{\alpha} \partial g_{\alpha}^{(i)} v_{\alpha}^{(i)} \bar{\lambda}_{\alpha}^{(i)} &= 0 \end{aligned}$$

and

$$\bar{\eta}, \bar{\lambda} \geq 0.$$

By the dual theorem there exists a dual optimal solution $(\bar{\eta}^*, \bar{\lambda}^*)$. By lemma 1 $\bar{\eta}^*$ has non-zero coordinates corresponding only to support planes passing through the optimum (u^*, z_*) , i.e., those gradient tangent planes at this point, one for each function $f^{(j)}$. This also applies to $\bar{\lambda}^*$ and constraint functions $g^{(i)}$, and therefore we may write $\bar{\eta}^* = (\bar{\eta}_*^{(1)}, \dots, \bar{\eta}_*^{(N)})$ and $\bar{\lambda}^* = (\bar{\lambda}_*^{(1)}, \dots, \bar{\lambda}_*^{(m)})$. Thus, upon setting $\eta_*^{(j)} = \mu_*^{(j)} \bar{\eta}_*^{(j)}$ for $j = 1, \dots, N$ and $\lambda_*^{(i)} = v_*^{(i)} \bar{\lambda}_*^{(i)}$ for $i = 1, \dots, m$, we obtain the following dual optimal conditions:

$$(1) \quad -\sum_j \partial f_*^{(j)} \eta_*^{(j)} + \sum_i \partial g_*^{(i)} \lambda_*^{(i)} = 0$$

and

$$(2) \quad \sum_j \eta_*^{(j)} = 1$$

where all $\eta_*^{(j)}$ and $\lambda_*^{(i)} \geq 0$.

The equality of dual functionals yields,

$$\begin{aligned} f(u^*) = z_* &= \sum_j f^{(j)}(u^*) \eta_*^{(j)} - \sum_j u^{*T} \partial f_*^{(j)} \eta_*^{(j)} + \sum_i u^{*T} \partial g_*^{(i)} \lambda_*^{(i)} + \sum_i (-g^{(i)}(u^*) \lambda_*^{(i)}) \\ &= \sum_j f^{(j)}(u^*) \eta_*^{(j)} - \sum_i g^{(i)}(u^*) \lambda_*^{(i)}. \end{aligned}$$

Since $f^{(j)}(u^*) \leq f(u^*)$ for all j and $g^{(i)}(u^*) \geq 0$ for all i , it therefore follows that, (3) $\sum_i g^{(i)}(u^*) \lambda_*^{(i)} = 0$. Furthermore, since $f(u^*) = \max \{f^{(j)}(u^*)\}$, it follows that $\eta_*^{(j)} = 0$ whenever $f^{(j)}(u^*) < f(u^*)$, giving condition (2). Thus, the three conditions of the theorem are proved.

On the other hand, given positive vectors η^* and λ^* satisfying conditions (1), (2), and (3) with respect to u^* , then since $\mu_*^{(j)} \neq 0$ and $\lambda_*^{(i)} \neq 0$ in the canonically closed system ($\bar{\Pi}$), we obtain dual feasible solutions upon setting $\bar{\eta}_*^{(j)} = \eta_*^{(j)} / \mu_*^{(j)}$ and $\bar{\lambda}_*^{(i)} = \lambda_*^{(i)} / v_i^*$. Furthermore, the dual objective value is $\sum_j f^{(j)}(u^*) \eta_*^{(j)}$, and condition (2) implies that $f(u^*) = \sum_j f^{(j)}(u^*) \eta_*^{(j)}$ giving dual equality of objective functions, thereby proving that $(f(u^*), u^*)$ is optimal.

Our generalization of the quasi-saddle point version of the Kuhn-Tucker Theorem is not as general as we may possibly get, but it does indicate a unified approach to study these equivalences under more general circumstances. In fact, we are already obtaining results for generalized saddle-point equivalence theorems for arbitrary convex functions over R_n . This is the subject of another paper and will be reported on elsewhere.

Already these methods have shown that the crucial property of the constraint functions is the Farkas-Minkowski property, which is a property of the functions themselves expressed in terms of finite positive linear combinations of their "gradients". Geometric qualifications are sufficient restrictions on the constraining functions to permit such Farkas-Minkowski expressions. In general however, it may be necessary to go beyond the natural gradient inequalities provided by the constraint functions to obtain strong duality results.

In conclusion, we illustrate this now by constructing a canonically closed equivalent for the one-variable Slater example by adjoining a new variable to the gradient inequality system following the methods of our regularization procedures for semi-infinite programs.⁸⁾ Restating the Slater example, we have:

$$(I) \quad \begin{aligned} & \min x \\ & \text{subject to } -(1-x)^2 \geq 0 \end{aligned}$$

with unique optimum $x_* = 1$. Introducing a differential system of supports to contain the optimum, we obtain the equivalent problem:

$$\begin{aligned} (I) \quad & \min x \\ & \text{subject to } 2(1-\alpha)x \geq 1 - \alpha^2 \end{aligned}$$

for $0 \leq \alpha \leq 2$.

Let M and V be large positive numbers, either real or non-Archimedean, i.e. larger than any real number⁹⁾, and construct the following semi-infinite dual regularizations.

$$\begin{aligned} (I_R) \quad & \min Mt + x \\ & \text{subject to } t + 2(1-\alpha)x \geq 1 - \alpha^2, \quad 0 \leq \alpha \leq 2 \\ & \quad x \geq -V \\ & \quad -x \geq -V \end{aligned}$$

⁸⁾ See Slater [7], [4] p. 216, and [5], p. 119.

⁹⁾ See [3] pp. 756-7.

$$\begin{aligned}
& (\Pi_R) \\
& \max \sum_{\alpha} (1 - \alpha^2) \lambda_{\alpha} - V\lambda^+ - V\lambda^- \\
& \text{subject to } \sum_{\alpha} \lambda_{\alpha} = M \\
& \quad \sum_{\alpha} 2(1 - \alpha) \lambda_{\alpha} + \lambda^+ - \lambda^- = 1 \\
& \quad \lambda's \geq 0 .
\end{aligned}$$

Observe that problem (I_R) is canonically closed and that $t \geq 0$ is included in the inequality system and corresponds to the index point $\alpha = 1$. As stated above, M may be viewed as real or non-Archimedean, and therefore we shall derive dual optimal solutions for (I_R) and (Π_R) in a manner which is valid for either case.

We know that $(t, x) = (0, 1)$ is (I_R) -feasible with functional value 1. Thus, we search for a solution (t_*, x_*) with objective value < 1 , if it exists, and therefore we assume $x_* < 1$. By lemma 1, this optimum involves only support planes which are tangent to it and therefore involves only its own gradient inequality with index point $\alpha_* = x_*$. But this implies $t_* = (1 - \alpha_*)^2$ yielding (I_R) -objective value $M(1 - \alpha_*)^2 + \alpha_*$. Applying the usual differential methods for finding a minimum to this function yields the Taylor expansion,

$$M(1 - \alpha_*)^2 + \alpha_* = \frac{4M - 1}{4M} + M \left(\alpha_* - \frac{2M - 1}{2M} \right)^2 \quad \text{for } 0 \leq \alpha_* \leq 2 ,$$

an equation which is obviously valid for arbitrary M . This tells us to take $\alpha_* = (2M - 1)/2M$ to obtain minimum objective value $(4M - 1)/4M < 1$. Furthermore, the point $(t_*, x_*) = (1/4M^2, (2M - 1)/2M)$ is (I_R) -feasible because

$$t \geq \frac{1}{4M^2} - \left(\alpha - \frac{2M - 1}{2M} \right)^2 = 1 - \alpha^2 - 2(1 - \alpha)x_* \quad \text{for } 0 \leq \alpha \leq 2 ,$$

which is a restatement of (I_R) -feasibility. But taking $\lambda_{\alpha_*} = M$, the dual variable associated with the binding constraint, and $\lambda_{\alpha} = 0$ for $\alpha \neq \alpha_*$ and $\lambda^+ = \lambda^- = 0$, yields a dual (Π_R) -solution with equality of dual objective functions, and therefore shows that in fact the two solutions form dual optimal solutions for problems (I_R) , (Π_R) whether M is viewed as real or non-Archimedean.

Observe that the dual solution, λ^* , is an extreme point of the associated generalized finite sequence space¹⁰) and as such the non-zero coordinate is linear and homogeneous in M ¹¹), in particular, $\lambda_{\alpha_*} = M$. Two courses of action with respect to M

¹⁰) See [4] p. 211.

¹¹) See [2], where this statement was first proved for finite linear programming over non-Archimedean ordered fields.

are now open to us. *First*, if M is real, we may let $M \rightarrow \infty$ so that $(t_*, x_*) \rightarrow (0, 1)$, the solution to the Slater problem, with corresponding dual variable characterized by $\lambda_{x_*} \rightarrow \infty$. *Second*, viewing M as non-Archimedean, we obtain dual optimal solutions in Hilbert's field with common objective value $1 - 1/4M$ which in the extended ordering is *larger* than any real number less than 1, but itself is less than 1.

References

- [1] Arrow, K. J., L. Hurwicz and Uzawa, H.: "Constraint Qualifications in Maximization Problems," Naval Research Logistics Quarterly, Vol. 8, No. 2, June 1961.
- [2] Charnes, A., and W. W. Cooper: "The Strong Minkowski Farkas-Weyl Theorem for Vector spaces over Ordered Fields," Proceedings of Nat. Acad. Sciences, Vol. 44, No. 9, pp. 914–916, Sept. 1958.
- [3] Charnes, A., and W. W. Cooper: Management Models and Industrial Applications of Linear Programming, Vols. I and II, New York, J. Wiley and Sons, 1961.
- [4] Charnes, A., Cooper, W. W. and Kortanek, K.: "Duality in Semi-Infinite Programs and Some works of Haar and Caratheodory," Management Science, Vol. 9, No. 2, January, 1963, 209–228.
- [5] Charnes, A., Cooper, W. W., and Kortanek, K.: "On Representations of Semi-Infinite Programs Which Have No Duality Gaps," Management Science Vol. 12, No. 1, September, 1965.
- [6] Kortanek, K.: "Duality, Semi-Infinite Programming, and Some Aspects of Control in Business and Economic Systems," Ph. D. Thesis, Northwestern University, Evanston, Ill., 1964.
- [7] Kuhn, H. W., and Tucker, A. W.: "Non-Linear Programming," Proc. 2nd Berkeley Symp. Math. Stat. and Prob., J. Neyman (ed.), U. Calif. Press, Berkeley, Calif., 1951, pp. 481–492.
- [8] Slater, M.: "Lagrange Multipliers Revisited: A Contribution to Non-Linear Programming," Cowles Commission Paper, Math. No. 403, New Haven, Nov. 1950.

Souhrn

SEMIINFINITNÍ PROGRAMOVÁNÍ, DIFERENCOVATELNOST A GEOMETRICKÉ PROGRAMOVÁNÍ: ČÁST II

A. CHARNES, W. W. COOPER, K. O. KORTANEK

Autoři se v článku zabývají jistou specializací své teorie duality na případ, kdy cílová funkce je spojitě diferencovatelná a konvexní na množině K přípustných řešení a funkce omezení definující K jsou spojitě diferencovatelné a konkávní. V článku je dále ukázána cesta jak výklad zobecnit na případ, kdy funkce v omezeních problému jsou po částečně diferencovatelné a konkávní. Získané podmínky lze chápat jako rozšíření Kuhn-Tucherovy věty.