

SEMI-MAGIC SQUARES AND ELLIPTIC CURVES

EDRAY HERBER GOINS

ABSTRACT. We show that, for all odd natural numbers N , the N -torsion points on an elliptic curve may be placed in an $N \times N$ grid such that the sum of each column and each row is the point at infinity.

1. INTRODUCTION

Let N be a positive integer, and consider the integers $1, 2, \dots, N^2$. An $N \times N$ grid containing these consecutive integers such that the sum of each column and each row is the same is called a magic square. (This is usually called a semi-magic square in the literature because we do not assume that the sum of both diagonals is also equal to the sum of the columns and rows [6].) For example, when $N = 3$, we have the grids

3	5	7
8	1	6
4	9	2

8	1	6
3	5	7
4	9	2

where the sum of each column and each row is 15. (The one on the right is a magic square in the classical sense; the one on the left does not have diagonals which sum to 15.)

We need not limit ourselves to a grid with integer entries. The author of [1], inspired by the discussion in [2, Section 1.4], considered the problem of arranging the 9 points of inflection on an elliptic curve in a 3×3 magic square. That is, it is possible to arrange the points of order 3 in a 3×3 grid so that the sum of each row and each column is the same, namely the point at infinity. We generalize this result.

Theorem 1. *Let $N \geq 1$ be an odd integer, let E be an elliptic curve defined over an algebraically closed field with characteristic not dividing N . Then the N^2 points of order N on E can be placed in an $N \times N$ magic square such that the sum of each column and each row is the point at infinity \mathcal{O} .*

We construct such a grid using Lehmer's Uniform Step Method, as motivated by the discussion in [4]. In particular, the theorem holds for any group G such that the N -torsion $G[N] \simeq (\mathbb{Z}/N\mathbb{Z}) \times (\mathbb{Z}/N\mathbb{Z})$.

2. SEMI-MAGIC SQUARES OVER ABELIAN GROUPS

As stated above, we define a magic square to be an $N \times N$ grid containing the consecutive integers 1 through N^2 such that the sum of each column and each row is the same. Strictly speaking, this is a semi-magic square, but we abuse notation slightly for the sake of brevity. We do not limit ourselves to constructing magic squares with integer entries. Indeed, we will construct an $N \times N$ magic square for a certain class of abelian groups.

Let G be an abelian group under \oplus . Given $P \in G$, denote $[-1]P$ as its inverse and $[0]P = \mathcal{O}$ as the identity. For each nonzero integer m , denote $[m]P$ as $[\pm 1]P$ added to itself $|m|$ times, where “ \pm ” is chosen as the sign of m . Denote $G[m] \subseteq G$ as that subgroup consisting of points $P \in G$ such that $[m]P = \mathcal{O}$. We will always assume that G is chosen such that for some positive integer N there is a group isomorphism

$$\psi : (\mathbb{Z}/N\mathbb{Z}) \times (\mathbb{Z}/N\mathbb{Z}) \xrightarrow{\sim} G[N]. \tag{2.1}$$

We have a bijection $\{1, 2, \dots, N^2\} \rightarrow (\mathbb{Z}/N\mathbb{Z}) \times (\mathbb{Z}/N\mathbb{Z})$ given by

$$\phi : k \mapsto \left(k - 1 \pmod{N}, \left\lfloor \frac{k - 1}{N} \right\rfloor \pmod{N} \right) \tag{2.2}$$

where $\lfloor \cdot \rfloor$ is the greatest integer function. That is, if $1 \leq k \leq N^2$ then we can write $k - 1 = m + Nn$ for some unique $0 \leq m, n < N$, and so we map $k \mapsto (m, n)$. This means we have a bijection of sets

$$\psi \circ \phi : \{1, 2, \dots, N^2\} \xrightarrow{\sim} G[N].$$

We will use this identification to place the elements in $G[N]$ in an $N \times N$ magic square.

There are two examples in particular which will be of interest to us. Upon fixing N , the group $G = (\mathbb{Z}/N\mathbb{Z}) \times (\mathbb{Z}/N\mathbb{Z})$ satisfies the criterion above. As another example, fix an algebraically closed field F and let E be an elliptic curve defined over F . We may choose $G = E(F)$ as the F -rational points on E , where we have a non-canonical isomorphism $G[N] \simeq (\mathbb{Z}/N\mathbb{Z}) \times (\mathbb{Z}/N\mathbb{Z})$ only when the characteristic of F does not divide N . (For more properties of elliptic curves, see [3].)

3. UNIFORM STEP METHOD

Fix a positive integer N . Let G be an abelian group under \oplus , and assume

$$G[N] = \{R_1, R_2, \dots, R_k, \dots, R_{N^2}\} \simeq (\mathbb{Z}/N\mathbb{Z}) \times (\mathbb{Z}/N\mathbb{Z}).$$

We wish to place these N^2 elements in an $N \times N$ grid such that the sum of each row and the sum of each column, as an element in G , is the same. We use an idea of D. H. Lehmer from 1929, known as the Uniform Step Method. To this end, we are motivated by the discussion in [4, Chapter 4].

Given an $N \times N$ grid, we consider its entries in Cartesian coordinates. For the moment, fix integers $a, b, c,$ and $d,$ and consider placing the element $R_k \in G[N]$ in the (x_k, y_k) position. After arbitrarily placing R_1 in the (x_1, y_1) -position, we will define x_k and y_k by the recursive sequence

$$\begin{aligned} x_k &\equiv x_1 + a(k-1) + b \left\lfloor \frac{k-1}{N} \right\rfloor \pmod{N} \\ y_k &\equiv y_1 + c(k-1) + d \left\lfloor \frac{k-1}{N} \right\rfloor \pmod{N} \end{aligned} \quad \text{for } 1 \leq k \leq N^2.$$

We will exhibit conditions on these integers $a, b, c,$ and d such that the sequences above indeed generate a magic square.

Proposition 2. *If N is odd and relatively prime to $ad - bc,$ then the sequence (x_k, y_k) places exactly one R_k in each of the N^2 cells of the $N \times N$ grid.*

Proof. It suffices to show that $(x_{k_1}, y_{k_1}) = (x_{k_2}, y_{k_2})$ only when $k_1 = k_2;$ for then we would have N^2 different points so they must fill in the entire grid. Using the bijection ϕ as in (2.2) note that we may write

$$\begin{aligned} x_k &\equiv x_1 + a m + b n \pmod{N} \\ y_k &\equiv y_1 + c m + d n \pmod{N} \end{aligned} \quad \text{where } (m, n) = \phi(k).$$

Write $(m_1, n_1) = \phi(k_1)$ and $(m_2, n_2) = \phi(k_2),$ so that

$$(x_{k_1}, y_{k_1}) = (x_{k_2}, y_{k_2}) \iff \begin{aligned} a(m_1 - m_2) + b(n_1 - n_2) &\equiv 0 \pmod{N} \\ c(m_1 - m_2) + d(n_1 - n_2) &\equiv 0 \pmod{N}. \end{aligned}$$

Since $ad - bc \pmod{N}$ is invertible, we see that this happens if and only if

$$\phi(k_1) = (m_1, n_1) = (m_2, n_2) = \phi(k_2)$$

and so $k_1 = k_2.$ □

Proposition 3. *If N is relatively prime to a and $b,$ then the sum of the entries in the i th column is $\mathcal{O}.$ If N is relatively prime to c and $d,$ then the sum of the entries in the j th row is $\mathcal{O}.$*

Proof. The entries in the i th column consist of those R_k corresponding to k such that $x_k = i.$ Similarly, the entries in the j th row consist of those R_k corresponding to k such that $y_k = j.$ Hence, the sum of the entries in the i th column and j th row are

$$\sum_{x_k=i} R_k \quad \text{and} \quad \sum_{y_k=j} R_k, \quad \text{respectively.}$$

First we determine the values of k which occur in the i th column. Since N is relatively prime to a and $b,$ there are exactly N pairs $(m, n) \in (\mathbb{Z}/N\mathbb{Z}) \times (\mathbb{Z}/N\mathbb{Z})$ satisfying $am + bn \equiv i - x_1 \pmod{N};$ indeed, given any m we

can solve for n , and vice-versa. Hence, there are exactly N integers $k \equiv 1 + m + Nn \pmod{N^2}$ such that $x_k = i$, which we denote by k_α . If we denote $(m_\alpha, n_\alpha) = \phi(k_\alpha)$ using the bijection in (2.2), then it is clear we have $\{\dots, m_\alpha, \dots\} = \{\dots, n_\alpha, \dots\} = \mathbb{Z}/N\mathbb{Z}$.

Now we compute the sum of the values in the i th column. Using the group isomorphism in (2.1), denote $P = \psi((1, 0))$ and $Q = \psi((0, 1))$ so that we have $R_k = [m]P \oplus [n]Q$ when $(m, n) = \phi(k)$. This gives the sum

$$\sum_{x_k=i} R_k = \sum_{\alpha} R_{k_\alpha} = \sum_{\alpha} ([m_\alpha]P \oplus [n_\alpha]Q) = [m']P \oplus [n']Q,$$

where we have set

$$m' \equiv n' \equiv \sum_{\alpha} m_\alpha \equiv \sum_{\alpha} n_\alpha \equiv \sum_{m \in \mathbb{Z}/N\mathbb{Z}} m \equiv \frac{N(N-1)}{2} \pmod{N}.$$

Since N is assumed odd, this sum is a multiple of N so that $[m']P = [n']Q = \mathcal{O}$. Hence, the sum of the entries in the i th column is indeed \mathcal{O} .

A similar argument works for the j th row. □

We summarize this as follows.

Theorem 4. *Let G be an abelian group under \oplus , and assume that there is a positive odd integer N such that*

$$G[N] = \{R_1, R_2, \dots, R_k, \dots, R_{N^2}\} \simeq (\mathbb{Z}/N\mathbb{Z}) \times (\mathbb{Z}/N\mathbb{Z}).$$

Fix integers a, b, c , and d relatively prime to N such that $ad - bc$ is also relatively prime to N , and consider the sequence (x_k, y_k) defined by

$$\begin{aligned} x_k &\equiv x_1 + a(k-1) + b \left\lfloor \frac{k-1}{N} \right\rfloor \pmod{N} \\ y_k &\equiv y_1 + c(k-1) + d \left\lfloor \frac{k-1}{N} \right\rfloor \pmod{N} \end{aligned} \quad \text{for } 1 \leq k \leq N^2.$$

The $N \times N$ grid formed by placing R_k in the (x_k, y_k) position is a magic square, where the sum of each column and each row is the identity \mathcal{O} .

We remark that this method does not exhaust all ways in which a magic square can be generated. For example, this method does not seem to work for N even. Indeed, the sum of each column and each row involves the expression $N(N-1)/2$, which in general is not a multiple of N . Also, when $N = 4$, we have the magic square

16	3	2	13
5	10	11	8
9	6	7	12
4	15	14	1

It is easy to check that such a square cannot be generated by a sequence (x_k, y_k) for any a, b, c , or d . This first appeared in 1514 in an engraving by Albrecht Dürer entitled “Melencolia.”

4. APPLICATIONS

We can specialize a, b, c , and d to generate examples of magic squares.

Corollary 5. *Let G be an abelian group under \oplus , and assume that there is an odd positive integer N such that*

$$G[N] = \{R_1, R_2, \dots, R_k, \dots, R_{N^2}\} \simeq (\mathbb{Z}/N\mathbb{Z}) \times (\mathbb{Z}/N\mathbb{Z}).$$

Then these elements can be placed in an $N \times N$ magic square such that the sum of each column and each row is the identity \mathcal{O} .

Proof. We follow the construction using a method first outlined by De la Loubère in 1693. (An example of how this method works follows at the end of the paper.) Using Theorem 4, set $a = 1, b = c = -1$, and $d = 2$. As N is odd, it is relatively prime to these integers as well as the determinant $ad - bc = 1$. □

Remark. Theorem 1 follows from this corollary, since the group $E[N]$ of N -torsion points on an elliptic curve E is isomorphic to $(\mathbb{Z}/N\mathbb{Z}) \times (\mathbb{Z}/N\mathbb{Z})$.

The following was pointed out to the author by J.-K. Yu. Upon choosing the basis $\{P, Q\}$ for $G[N]$ given by $P = \psi((1, 0))$ and $Q = \psi((0, 1))$, we may write $R_k = [m]P \oplus [n]Q$ when $(m, n) = \phi(k)$. (Here, we use the maps defined in (2.1) and (2.2).) In this way, we may identify R_k with (m, n) . If we choose $a = d = 1$ and $b = c = 0$, then we have a magic square upon placing $(m, n) = \phi(k)$ in the (x_k, y_k) -position. In general, if for odd N we have an $N \times N$ Latin Square with the (m, n) -position having entry a_{mn} then we may place (m, a_{mn}) in the (x_k, y_k) -position. (For more on Latin squares, see [5].)

We discuss a specific example by considering the 3-torsion on elliptic curves; to this end, set $N = 3$. We explain how this construction generalizes that in [1]. Consider an elliptic curve defined over the complex numbers \mathbb{C} , and let $G = E(\mathbb{C})$ be the group of complex points on the curve. Then it is well-known that we can express the 3-torsion as

$$E[3] = \{A, B, C, D, [-1]A, [-1]B, [-1]C, [-1]D, \mathcal{O}\} \simeq (\mathbb{Z}/3\mathbb{Z}) \times (\mathbb{Z}/3\mathbb{Z})$$

where $B = A \oplus D$ and $[-1]B = C \oplus D$. If we label these points as

$$\begin{array}{lll} R_1 = \mathcal{O}, & R_4 = D, & R_7 = [-1]D, \\ R_2 = [-1]B, & R_5 = [-1]A, & R_8 = C, \\ R_3 = B, & R_6 = [-1]C, & R_9 = A; \end{array}$$

then we can use the magic square from the introduction to place the 3-torsion in a magic square:

$$\begin{array}{|c|c|c|} \hline 3 & 5 & 7 \\ \hline 8 & 1 & 6 \\ \hline 4 & 9 & 2 \\ \hline \end{array} \implies \begin{array}{|c|c|c|} \hline B & [-1]A & [-1]D \\ \hline C & \mathcal{O} & [-1]C \\ \hline D & A & [-1]B \\ \hline \end{array}$$

We can also compute this magic square using the method in the proof of the corollary. Choosing the basis $P = [-1]B$ and $Q = D$; it can be easily checked that $R_k = [m]P \oplus [n]Q$ when $(m, n) = \phi(k)$. If we also choose $(x_1, y_1) = (2, 2)$ as the center of the 3×3 grid, then R_k may be placed in the (x_k, y_k) -position, where

$$\begin{aligned}
 x_k &\equiv x_1 + (k - 1) - \left\lfloor \frac{k - 1}{N} \right\rfloor \pmod{N} \\
 y_k &\equiv y_1 - (k - 1) + 2 \left\lfloor \frac{k - 1}{N} \right\rfloor \pmod{N}
 \end{aligned}
 \quad \text{for } 1 \leq k \leq N^2.$$

As mentioned before, this is known as De la Loubère’s method or the Siamese method. Following a comment of the referee, if we choose $(x_1, y_1) = (1, 2)$ as the center of the 3×3 grid, then we find the (classically) magic square

8	1	6
3	5	7
4	9	2

5. ACKNOWLEDGEMENT

The author would like to thank Jiu-Kang Yu for helpful comments and the referee for useful suggestions.

REFERENCES

[1] E. Brown, Magic squares, finite planes, and points of inflection on elliptic curves, *College Mathematics Journal*, **32** (2001), 260–267.
 [2] V. Prasolov and Y. Solovyev, *Elliptic Functions and Elliptic Integrals*, Volume 170 of Translations of Mathematical Monographs, translated from the Russian manuscript by D. Leites, American Mathematical Society, Providence, RI, 1997.
 [3] J. H. Silverman, *The Arithmetic of Elliptic Curves*, Springer-Verlag, New York-Berlin, 1986.
 [4] H. M. Stark, *An Introduction to Number Theory*, MIT Press, Cambridge, MA, 1978.
 [5] E. W. Weisstein, Latin square, <http://mathworld.wolfram.com/LatinSquare.html>.
 [6] E. W. Weisstein, Magic square, <http://mathworld.wolfram.com/MagicSquare.html>.

MSC2000: 14H52, 11Axx, 05Bxx.

PURDUE UNIVERSITY, DEPARTMENT OF MATHEMATICS, MATHEMATICAL SCIENCES BUILDING, 150 NORTH UNIVERSITY STREET, WEST LAFAYETTE, IN 47907-2067
E-mail address: egoins@math.purdue.edu