

SEMI-RIEMANNIAN SUBMERSIONS WITH TOTALLY GEODESIC FIBRES

GABRIEL BĂDIȚOIU

(Received August 19, 2002, revised August 4, 2003)

Abstract. We classify semi-Riemannian submersions with connected totally geodesic fibres from a real pseudo-hyperbolic space onto a semi-Riemannian manifold under the assumption that the dimension of the fibres is less than or equal to three. Also, we obtain the classification of semi-Riemannian submersions with connected complex totally geodesic fibres from a complex pseudo-hyperbolic space onto a semi-Riemannian manifold under the assumption that the dimension of the fibres is less than or equal to two. We prove that there are no semi-Riemannian submersions with connected quaternionic fibres from a quaternionic pseudo-hyperbolic space onto a Riemannian manifold.

1. Introduction and main results. Riemannian submersions, introduced by O’Neill [One1] and Gray [Gra], have been used by many authors to construct specific Riemannian metrics. A systematic exposition can be found in Besse’s book [Bes]. In this paper, we obtain classification results for semi-Riemannian submersions with totally geodesic fibres.

We first recall briefly some related work on the classification problem of semi-Riemannian submersions. Escobales [Esc1, Esc2] and Ranjan [Ran1] classified Riemannian submersions with connected totally geodesic fibres from an n -sphere S^n , and with connected complex totally geodesic fibres from a complex projective n -space CP^n , respectively. Ucci [Ucc] showed that there are no Riemannian submersions with fibres CP^3 from the complex projective space CP^7 onto $S^8(4)$, and with fibres HP^1 from the quaternionic projective space HP^3 onto $S^8(4)$. In [Ran2], Ranjan obtained a classification theorem for Riemannian submersions with connected totally geodesic fibres from a compact simple Lie group. Gromoll and Grove obtained in [G-G1] that, up to equivalence, the only Riemannian submersions of spheres (with connected fibres) are the Hopf fibrations, except possibly for fibrations of the 15-sphere by homotopy 7-spheres. This classification was invoked in the proof of the Diameter Rigidity Theorem (see [G-G2]) and of the Radius Rigidity Theorem (see [Wil]). Using an approach different from Gromoll and Grove [G-G1], Wilking [Wilk] proved that a Riemannian submersion $\pi : S^m \rightarrow B^b$ is metrically equivalent to the Hopf fibration for $(m, b) = (15, 8)$ and obtained an improved version of the Diameter Rigidity Theorem as a consequence of his classification theorem.

2000 *Mathematics Subject Classification.* Primary 53C50.

Key words and phrases. Semi-Riemannian submersions, isotropic semi-Riemannian manifolds, totally geodesic submanifolds, Ehresmann connections.

In comparison, there are few classification results for semi-Riemannian submersions, and the consequences seem to be at least as important as those for Riemannian submersions. In [Mag], Magid proved that the only semi-Riemannian submersions with totally geodesic fibres from an anti-de Sitter space onto a Riemannian manifold are the canonical semi-Riemannian submersions $H_1^{2m+1} \rightarrow CH^m$. In [Ba-Ia], the present author and Stere Ianuș classified semi-Riemannian submersions with connected totally geodesic fibres from a pseudo-hyperbolic space onto a Riemannian manifold, and with connected complex totally geodesic fibres from a complex pseudo-hyperbolic space onto a Riemannian manifold.

The aim of this work is to prove new classification results in the theory of semi-Riemannian submersions analogous to those in Riemannian geometry. It is my pleasure to thank Professor Stere Ianuș for useful discussions on this subject.

Now, we list the main results proved in this paper.

THEOREM 1.1. *Let $\pi : H_{s+r'}^{n+r} \rightarrow B_s^n$ be a semi-Riemannian submersion with connected totally geodesic fibres from a pseudo-hyperbolic space onto a semi-Riemannian manifold. If the dimension of the fibres is less than or equal to 3, then π is equivalent to one of the following canonical semi-Riemannian submersions:*

- (a) $H_{2t+1}^{2m+1} \rightarrow CH_t^m$, $0 \leq t \leq m$.
- (b) $H_{4t+3}^{4m+3} \rightarrow HH_t^m$, $0 \leq t \leq m$.

THEOREM 1.2. *Let $\pi : H_{s+r'}^{n+r} \rightarrow B_s^n$ be a semi-Riemannian submersion with connected totally geodesic fibres from a pseudo-hyperbolic space onto a semi-Riemannian manifold. Assume that one of the following conditions is satisfied:*

- (A) B is an isotropic semi-Riemannian manifold, which means that for any $x \in B_s^n$ and any real number t , the group of isometries $\mathbf{I}(B_s^n, g')$ preserving x acts transitively on the set of all nonzero tangent vectors X at x for which $g'(X, X) = t$, or
- (B) $\text{index}(B) \in \{0, \dim B\}$.

Then π is equivalent to one of the following canonical semi-Riemannian submersions:

- (a) $H_{2t+1}^{2m+1} \rightarrow CH_t^m$, $0 \leq t \leq m$.
- (b) $H_{4t+3}^{4m+3} \rightarrow HH_t^m$, $0 \leq t \leq m$.
- (c) $H_{7+8t}^{15} \rightarrow H_{8t}^8(-4)$, $t \in \{0, 1\}$.

THEOREM 1.3. *Let $\pi : CH_s^n \rightarrow B$ be a semi-Riemannian submersion from a complex pseudo-hyperbolic space onto a semi-Riemannian manifold. Assume that the fibres are connected complex totally geodesic submanifolds, and one of the following conditions is satisfied:*

- (A) *The real dimension of the fibres is $r \leq 2$, or*
- (B) B is an isotropic semi-Riemannian manifold, or
- (C) $\text{index}(B) \in \{0, \dim B\}$.

Then π is equivalent to the canonical semi-Riemannian submersion

$$CH_{2t+1}^{2m+1} \rightarrow HH_t^m, \quad 0 \leq t \leq m.$$

THEOREM 1.4. *There exist no semi-Riemannian submersions $\pi : \mathbf{H}\mathbf{H}_s^n \rightarrow B$ with connected quaternionic fibres from a quaternionic pseudo-hyperbolic space onto an isotropic semi-Riemannian manifold or onto a semi-Riemannian manifold of $\text{index}(B) \in \{0, \dim(B)\}$.*

2. Preliminaries and examples. In this section we recall several notions and results which will be needed throughout the paper. We also exhibit the construction of canonical semi-Riemannian submersions.

DEFINITION 2.1. Let (M, g) be an $(n+r)$ -dimensional connected semi-Riemannian manifold of index $s+r'$, and (B, g') an n -dimensional connected semi-Riemannian manifold of index s , where $0 \leq s \leq n$, $0 \leq r' \leq r$. A *semi-Riemannian submersion* (see [One2]) is a smooth map $\pi : M \rightarrow B$ which is surjective and satisfies the following axioms:

- (a) $\pi_*|_p$ is surjective for all $p \in M$;
- (b) the fibres $\pi^{-1}(b)$, $b \in B$, are semi-Riemannian submanifolds of M ;
- (c) π_* preserves scalar products of vectors normal to fibres.

We shall always assume that the fibres are connected, the dimension of the fibres $\dim M - \dim B > 0$ and $\dim B > 0$. The vectors tangent to fibres are called vertical and those normal to fibres are called horizontal. We denote by \mathcal{V} the vertical distribution and by \mathcal{H} the horizontal distribution.

The geometry of semi-Riemannian submersions is characterized by O'Neill's tensors T , A (see [One1], [One2]) defined for vector fields E, F on M by

$$\begin{aligned} A_E F &= h\nabla_{hE} vF + v\nabla_{hE} hF, \\ T_E F &= h\nabla_{vE} vF + v\nabla_{vE} hF, \end{aligned}$$

where ∇ is the Levi-Civita connection of g , and v and h denote the orthogonal projections on \mathcal{V} and \mathcal{H} , respectively. For basic properties of O'Neill's tensors see [One1], [One2], [Bes] or [Ian].

DEFINITION 2.2. (i) A vector field X on M is said to be *basic* if X is horizontal and π -related to a vector field X' on B .

(ii) A vector field X along the fibre $\pi^{-1}(x)$, $x \in B$, is said to be *basic along $\pi^{-1}(x)$* if X is horizontal and $\pi_{*p}X(p) = \pi_{*q}X(q)$ for every $p, q \in \pi^{-1}(x)$.

We notice that each vector field X' on B has a unique horizontal lift X to M which is basic. For a vertical vector field V and a basic vector field X we have $h\nabla_V X = A_X V$ (see [One1]). We denote by R, R' and \hat{R} the Riemann curvature tensors of M, B and of the fibre $\pi^{-1}(x)$, $x \in B$, respectively. We choose the convention for the curvature tensor $R(E, F) = \nabla_E \nabla F - \nabla_F \nabla E - \nabla_{[E, F]}$. The Riemann curvature tensor is defined by

$$R(E, F, G, H) = g(R(G, H)F, E).$$

For O'Neill's equations of a semi-Riemannian submersion we refer to [One1] or [Bes].

DEFINITION 2.3. Two semi-Riemannian submersions $\pi, \pi' : (M, g) \rightarrow (B, g')$ are said to be *equivalent* if there exists an isometry f of M which induces an isometry \tilde{f} of B so that $\pi' \circ f = \tilde{f} \circ \pi$. The pair (f, \tilde{f}) is called a *bundle isometry*.

We shall need the following theorem, which is the semi-Riemannian version of Theorem 2.2 in [Esc1].

THEOREM 2.4. Let $\pi_1, \pi_2 : M \rightarrow B$ be semi-Riemannian submersions from a complete connected semi-Riemannian manifold M onto a semi-Riemannian manifold B . Assume that the fibres of these submersions are connected and totally geodesic. Let f be an isometry of M satisfying the following properties at a given point $p \in M$:

- (1) $f_{*p} : T_p M \rightarrow T_{f(p)} M$ maps \mathcal{H}_{1p} onto $\mathcal{H}_{2f(p)}$, where \mathcal{H}_i denote the horizontal distributions of π_i for $i \in \{1, 2\}$.
- (2) $f_* A_{1E} F = A_{2f_* E} f_* F$ for every $E, F \in T_p M$, where A_i are the integrability tensors associated with π_i .

Then f induces an isometry \tilde{f} of B so that the pair (f, \tilde{f}) is a bundle isometry between π_1 and π_2 . In particular, π_1 and π_2 are equivalent.

Escobales's proof of Theorem 2.2 in [Esc1], also works in this semi-Riemannian case. He proves that for any $b \in B$ which can be joined with $\pi_1(p)$ by a geodesic we have:

- (i) for every $x \in \pi_1^{-1}(b)$, $f_{*x} : T_x M \rightarrow T_{f(x)} M$ maps \mathcal{H}_{1x} onto $\mathcal{H}_{2f(x)}$, and
- (ii) f maps the fibre $\pi_1^{-1}(b)$ into the fibre $\pi_2^{-1}(\pi_2(f(x)))$ with $x \in \pi_1^{-1}(b)$.

We notice that for any $x \in \pi_1^{-1}(b)$ with $b \in B$, which can be joined with $\pi_1(p)$ by a geodesic, the conditions (1) and (2) are also satisfied for the point x . Since M is connected, B is also connected. Therefore, any point $b \in B$ can be joined with $\pi_1(p)$ by a broken geodesic. Repeating the argument above, for any corner point of this broken geodesic, we see that for any $b \in B$, f maps the fibre $\pi_1^{-1}(b)$ into a fibre.

DEFINITION 2.5. Let $\langle \cdot, \cdot \rangle$ be the symmetric bilinear form on \mathbf{R}^{m+1} given by

$$\langle x, y \rangle = - \sum_{i=0}^s x_i y_i + \sum_{i=s+1}^m x_i y_i$$

for $x = (x_0, \dots, x_m), y = (y_0, \dots, y_m) \in \mathbf{R}^{m+1}$. For any $c < 0$ and any positive integer s , let $H_s^m(c) = \{x \in \mathbf{R}^{m+1} \mid \langle x, x \rangle = 1/c\}$ be the semi-Riemannian submanifold of

$$\mathbf{R}_{s+1}^{m+1} = (\mathbf{R}^{m+1}, ds^2 = -dx^0 \otimes dx^0 - \dots - dx^s \otimes dx^s + dx^{s+1} \otimes dx^{s+1} + \dots + dx^m \otimes dx^m).$$

$H_s^m(c)$ is called the m -dimensional (*real*) *pseudo-hyperbolic space* of index s .

We notice that $H_s^m(c)$ has constant sectional curvature c , whose curvature tensor is given by $R(X, Y, X, Y) = c(g(X, X)g(Y, Y) - g(X, Y)^2)$. We shall denote simply $H_s^m = H_s^m(-1)$. It should be remarked that H_s^m can be written as a homogeneous space, namely $H_s^m = SO(s+1, m-s)/SO(s, m-s)$, $H_{2s+1}^{2m+1} = SU(s+1, m-s)/SU(s, m-s)$, and $H_{4s+3}^{4m+3} = Sp(s+1, m-s)/Sp(s, m-s)$ (see [Wol]).

DEFINITION 2.6. Let (\cdot, \cdot) be the Hermitian form on \mathbf{C}^{m+1} given by

$$(z, w) = -\sum_{i=0}^s z_i \bar{w}_i + \sum_{i=s+1}^m z_i \bar{w}_i$$

for $z = (z_0, \dots, z_m), w = (w_0, \dots, w_m) \in \mathbf{C}^{m+1}$. For $c < 0$, let $M(c)$ be the real hypersurface of \mathbf{C}^{m+1} given by $M(c) = \{z \in \mathbf{C}^{m+1} \mid (z, z) = 4/c\}$, which is endowed with the induced metric of

$$(\mathbf{C}^{m+1}, ds^2 = -dz^0 \otimes d\bar{z}^0 - \dots - dz^s \otimes d\bar{z}^s + dz^{s+1} \otimes d\bar{z}^{s+1} + \dots + dz^m \otimes d\bar{z}^m).$$

The natural action of $S^1 = \{e^{i\theta} \mid \theta \in \mathbf{R}\}$ on \mathbf{C}^{m+1} induces an action on $M(c)$. Let $CH_s^m(c) = M(c)/S^1$ endowed with the unique indefinite Kähler metric of index $2s$ such that the projection $M(c) \rightarrow M(c)/S^1$ becomes a semi-Riemannian submersion (see [Ba-Ro]). $CH_s^m(c)$ is called the *complex pseudo-hyperbolic space*.

Notice that $CH_s^m(c)$ has constant holomorphic sectional curvature c , whose curvature tensor is given by $R(X, Y, X, Y) = (c/4)(g(X, X)g(Y, Y) - g(X, Y)^2 + 3g(I_0 X, Y)^2)$, where I_0 is the natural complex structure on $CH_s^m(c)$. We shall denote simply $CH_s^m = CH_s^m(-4)$. It is well-known that CH_s^m is a homogeneous space, namely $CH_s^m = SU(s+1, m-s)/S(U(1)U(s, m-s))$ and $CH_{2s+1}^{2m+1} = Sp(s+1, m-s)/U(1)Sp(s, m-s)$ (see [Wol]).

We shall denote by HH_s^n the quaternionic pseudo-hyperbolic space of real dimension $4n$, and of quaternionic index s with quaternionic sectional curvature -4 , and by S^n and $S^n(4)$ the spheres with sectional curvature 1 and 4, respectively.

By a standard construction (see Theorem 9.80 in [Bes]), one can obtain many examples of semi-Riemannian submersions with totally geodesic fibres of type $\pi : G/K \rightarrow G/H$, where G is a Lie group and K, H are closed Lie subgroups of G with $K \subset H$. In this way the following *canonical semi-Riemannian submersions*, also called *generalized Hopf fibrations*, are obtained:

EXAMPLE 1. Let $G = SU(t+1, m-t), H = S(U(1)U(t, m-t)), K = SU(t, m-t)$. For every $0 \leq t \leq m$, we have the semi-Riemannian submersion

$$H_{2t+1}^{2m+1} = SU(t+1, m-t)/SU(t, m-t) \rightarrow CH_t^m = SU(t+1, m-t)/S(U(1)U(t, m-t)).$$

EXAMPLE 2. Let $G = Sp(t+1, m-t), H = Sp(1)Sp(t, m-t), K = Sp(t, m-t)$. For every $0 \leq t \leq m$, we get the semi-Riemannian submersion

$$H_{4t+3}^{4m+3} = Sp(t+1, m-t)/Sp(t, m-t) \rightarrow HH_t^m = Sp(t+1, m-t)/Sp(1)Sp(t, m-t).$$

EXAMPLE 3. a) Let $G = Spin(1, 8), H = Spin(8), K = Spin(7)$. Then we have the semi-Riemannian submersion (see [Ba-Ia])

$$H_7^{15} = Spin(1, 8)/Spin(7) \rightarrow H^8(-4) = Spin(1, 8)/Spin(8).$$

b) Let $G = Spin(9), H = Spin(8), K = Spin(7)$. Then we have the semi-Riemannian submersion (see [Bes])

$$S^{15} = Spin(9)/Spin(7) \rightarrow S^8(4) = Spin(9)/Spin(8).$$

EXAMPLE 4. Let $G = Sp(t+1, m-t)$, $H = Sp(1)Sp(t, m-t)$, $K = U(1)Sp(t, m-t)$. For every $0 \leq t \leq m$, we obtain the semi-Riemannian submersion

$$CH_{2t+1}^{2m+1} = Sp(t+1, m-t)/U(1)Sp(t, m-t) \rightarrow \mathbf{H}H_t^m = Sp(t+1, m-t)/Sp(1)Sp(t, m-t).$$

In order to prove Theorem 1.2, we need the following nonexistence proposition, which is the semi-Riemannian version of Proposition 5.1 in [Ran1].

PROPOSITION 2.7. *There exist no semi-Riemannian submersions $\pi : H_{7+8t}^{23} \rightarrow CaH_t^2$, $t \in \{0, 1, 2\}$, with totally geodesic fibres from the 23-dimensional pseudo-hyperbolic space of index $7 + 8t$ onto the Cayley pseudo-hyperbolic plane of Cayley index t .*

We notice that the case $t = 2$ is Proposition 5.1 in [Ran1]. For the case $t = 0$, see [Ba-Ia]. Here we only recall some details of Ranjan's proof and suggest its modification to the semi-Riemannian case. Ranjan's argument in [Ran1], which leads to a contradiction to the assumption of the existence of such a submersion, is based on finding for every $X \in \mathcal{H}_p$, $g(X, X) \neq 0$, an irreducible $Cl(\mathcal{V}_p)$ -submodule S of \mathcal{H}_p passing through X . Here $Cl(\mathcal{V}_p)$ denotes the Clifford algebra of $(\mathcal{V}_p, \tilde{g}_p)$, where $\tilde{g}(U, V) = -g(U, V)$ for every $U, V \in \mathcal{V}_p$. \mathcal{H}_p becomes a $Cl(\mathcal{V}_p)$ -module by considering the extension of the map $\mathcal{U} : \mathcal{V}_p \rightarrow \text{End}(\mathcal{H}_p)$ defined by $\mathcal{U}(V)(X) = A_X V$ to the Clifford algebra $Cl(\mathcal{V}_p)$. Since \tilde{g}_p is positive definite, we have $Cl(\mathcal{V}_p) \simeq \mathbf{R}(8) \oplus \mathbf{R}(8)$. Hence, \mathcal{H}_p splits into two 8-dimensional irreducible $Cl(\mathcal{V}_p)$ -modules. Since the induced metrics on fibres are negative definite, we obtain in a manner similar to Ranjan's proof that

(i) for $g(X, X) > 0$, $\pi^{-1}(CaH^1)$ is totally geodesic in H_{7+8t}^{23} and is isometric to H_7^{15} , where CaH^1 denotes the Cayley hyperbolic line through $\pi_* X$, and

(ii) for $g(X, X) < 0$, $\pi^{-1}(CaH_1^1)$ is totally geodesic in H_{7+8t}^{23} and is isometric to H_{15}^{15} , where CaH_1^1 denotes the negative definite Cayley hyperbolic line through $\pi_* X$.

We choose S to be the horizontal space of the restricted submersion $\tilde{\pi} : H_7^{15} \rightarrow CaH^1$ if $g(X, X) > 0$ or $\tilde{\pi} : H_{15}^{15} \rightarrow CaH_1^1$ if $g(X, X) < 0$.

3. Proof of the main results. The next lemma gives useful properties of O'Neill's integrability tensor.

LEMMA 3.1. *Let $\pi : M \rightarrow B$ be a semi-Riemannian submersion with connected totally geodesic fibres from a semi-Riemannian manifold M with constant curvature $c \neq 0$. Then the following assertions are true:*

(a) *If X is a horizontal vector such that $g(X, X) \neq 0$, then the map $A_X : \mathcal{V} \rightarrow \mathcal{H}$ given by $A_X(V) = A_X V$ is injective and the map $A_X^* : \mathcal{H} \rightarrow \mathcal{V}$ given by $A_X^*(Y) = A_X Y$ is surjective.*

(b) *If X, Y are the horizontal liftings along the fibre $\pi^{-1}(\pi(p))$, $p \in M$, of two vectors $X', Y' \in T_{\pi(p)}B$ respectively, $g'(X', X') \neq 0$ and $(A_X Y)(p) = 0$, then $A_X Y = 0$ along the fibre $\pi^{-1}(\pi(p))$.*

PROOF. (a) By O'Neill's equations, we get

$$g(A_X V, A_X W) = c g(X, X) g(V, W)$$

for a horizontal vector field X and for vertical vector fields V and W . Thus $A_X^* A_X V = -cg(X, X)V$ for every vertical vector field V . Therefore $A_X : \mathcal{V} \rightarrow \mathcal{H}$ is injective and $A_X^* : \mathcal{H} \rightarrow \mathcal{V}$ is surjective.

(b) By O'Neill's equations, we have

$$-3g(A_X Y, A_X Z) = c[g(X, X)g(Y, Z) - g(X, Y)g(X, Z)] - R'(\pi_* X, \pi_* Y, \pi_* X, \pi_* Z)$$

for horizontal vector fields X, Y and Z .

If X, Y, Z are basic vector fields, then $g(A_X Y, A_X Z)$ is constant along the fibre $\pi^{-1}(\pi(p))$. Therefore, $g(A_X A_X Y, Z) = 0$ along the fibre $\pi^{-1}(\pi(p))$ for every basic vector field Z . Hence $A_X A_X Y = 0$ along $\pi^{-1}(\pi(p))$. Since $A_X : \mathcal{V} \rightarrow \mathcal{H}$ is injective, it follows that $A_X Y = 0$ along the fibre $\pi^{-1}(\pi(p))$. \square

LEMMA 3.2. *If $\pi : M \rightarrow B$ is a semi-Riemannian submersion with connected totally geodesic fibres from a semi-Riemannian manifold M with constant curvature $c \neq 0$ onto a semi-Riemannian manifold B , then the tangent bundle of any fibre is trivial.*

PROOF. Let $x \in B$ and $p \in \pi^{-1}(x)$. Let $\{v_{1p}, \dots, v_{rp}\}$ be an orthonormal basis in \mathcal{V}_p . Let Y_1, Y_2, \dots, Y_r be the horizontal liftings along the fibre $\pi^{-1}(\pi(p))$ of $(1/c)\pi_* A_X v_{1p}, (1/c)\pi_* A_X v_{2p}, \dots, (1/c)\pi_* A_X v_{rp}$, respectively. Let $v_i = A_X Y_i$ for each $i \in \{1, \dots, r\}$. Since

$$\begin{aligned} g(v_j, v_l) &= g(A_X Y_j, A_X Y_l) \\ &= (1/3)(R'(\pi_* X, \pi_* Y_j, \pi_* X, \pi_* Y_l) - cg(X, X)g(Y_j, Y_l) + cg(X, Y_j)g(X, Y_l)) \end{aligned}$$

is constant along the fibre $\pi^{-1}(\pi(p))$ and

$$g(A_X Y_j, A_X Y_l)(p) = \frac{1}{c^2}g(A_X A_X v_{jp}, A_X A_X v_{lp}) = g(X, X)^2 g(v_{jp}, v_{lp}) = \varepsilon_j \delta_{jl},$$

we see that $\{v_1, v_2, \dots, v_r\}$ is a global orthonormal basis of the tangent bundle of the fibre $\pi^{-1}(x)$, which makes the tangent bundle trivial. \square

We suppose that the curvature of the total space is negative. The case of positive curvature can be reduced to the negative one by changing simultaneously the signs of the metrics on the base and on the total space. We establish relations between the dimensions and the indices of fibres and of base spaces, and see how the geometry of base spaces looks like.

THEOREM 3.3. *Let $\pi : M \rightarrow B$ be a semi-Riemannian submersion with connected totally geodesic fibres from an $(n + r)$ -dimensional semi-Riemannian manifold M of index $s + r'$ with constant negative curvature c onto an n -dimensional semi-Riemannian manifold B of index s . Then the following hold:*

- (1) $n = k(r + 1)$ for some positive integer k and $s = q_1(r' + 1) + q_2(r - r')$ for some nonnegative integers q_1, q_2 with $q_1 + q_2 = k$.
- (2) *If, moreover, M is a simply connected complete semi-Riemannian manifold and the dimension of fibres is less than or equal to 3, then B is an isotropic semi-Riemannian manifold and $r \in \{1, 3\}$.*

PROOF. Normalizing the metric on M , we can suppose $c = -1$. Let $p \in M$. Since the tangent bundle of the fibre $\pi^{-1}(\pi(p))$ is trivial, we can choose a global orthonormal frame $\{v_1, v_2, \dots, v_r\}$ for the tangent bundle of $\pi^{-1}(\pi(p))$. We have $g(v_i, v_j) = \varepsilon_i \delta_{ij}$, $\varepsilon_i \in \{-1, 1\}$, and $\text{card}\{i | \varepsilon_i < 0\} = r'$.

(1) Let X be the horizontal lifting along the fibre $\pi^{-1}(\pi(p))$ of a vector $X' \in T_{\pi(p)}B$, so that $g(X', X') \in \{-1, 1\}$. By O'Neill's equations, we have

$$g(A_Y V, A_Y V) = -g(Y, Y)g(V, V)$$

for a horizontal vector field Y and for a vertical vector field V . Along the fibre $\pi^{-1}(\pi(p))$ we obtain for every $i, j \in \{1, \dots, r\}$

$$\begin{aligned} g(A_X v_i, A_X v_j) &= -g(X, X)g(v_i, v_j) = -g(X, X)\varepsilon_i \delta_{ij}, \\ g(X, A_X v_i) &= -g(A_X X, v_i) = 0. \end{aligned}$$

Thus $\{X, A_X v_1, \dots, A_X v_r\}$ is an orthonormal system. Hence $n \geq r + 1$.

Let $L_0 = X$. For every integer α such that $1 \leq \alpha < n/(r + 1)$, let L_α be a horizontal vector field along the fibre $\pi^{-1}(\pi(p))$ such that L_α is the horizontal lifting of some unit vector (i.e., $g(L_\alpha, L_\alpha) \in \{-1, 1\}$), that L_α is orthogonal to $L_0, L_1, \dots, L_{\alpha-1}$ and that $L_\alpha(p) \in \ker A_{L_0(p)}^* \cap \ker A_{L_1(p)}^* \cap \dots \cap \ker A_{L_{\alpha-1}(p)}^*$. Then, by Lemma 3.1, $L_\alpha(q)$ belongs to $\ker A_{L_0(q)}^* \cap \ker A_{L_1(q)}^* \cap \dots \cap \ker A_{L_{\alpha-1}(q)}^*$ for every $q \in \pi^{-1}(\pi(p))$. Therefore, for $j \in \{1, \dots, r\}$ and $\alpha, \beta \geq 0$, we get

$$g(A_{L_\alpha} v_j, L_\beta) = -g(v_j, A_{L_\alpha} L_\beta) = 0$$

along the fibre $\pi^{-1}(\pi(p))$.

By O'Neill's equations, we obtain

$$\begin{aligned} R(X, U, Y, V) &= g((\nabla_U A)_X Y, V) + g(A_X U, A_Y V) \\ &= g(\nabla_U A_X Y, V) - g(A_{\nabla_U X} Y, V) - g(A_X \nabla_U Y, V) + g(A_X U, A_Y V) \\ (3.1) \quad &= g(\nabla_U A_X Y, V) + g(A_Y A_X U, V) - g(A_X A_Y U, V) - g(A_Y A_X U, V) \\ &= g(\nabla_U A_X Y, V) + g(A_Y U, A_X V) \end{aligned}$$

for basic vector fields X, Y and for vertical vector fields U, V . Thus, along the fibre $\pi^{-1}(\pi(p))$ we get for every $\alpha, \beta \geq 0$ and $j, l \in \{1, \dots, r\}$

$$\begin{aligned} g(A_{L_\alpha} v_j, A_{L_\beta} v_l) &= R(L_\alpha, v_l, L_\beta, v_j) - g(\nabla_{v_l} A_{L_\alpha} L_\beta, v_j) \\ &= -g(L_\alpha, L_\beta)g(v_l, v_j) - v_l(g(A_{L_\alpha} L_\beta, v_j)) + g(A_{L_\alpha} L_\beta, \nabla_{v_l} v_j). \end{aligned}$$

Since $A_{L_\alpha} L_\beta = 0$ along the fibre $\pi^{-1}(\pi(p))$, it follows that

$$g(A_{L_\alpha} v_j, A_{L_\beta} v_l) = -g(L_\alpha, L_\beta)g(v_l, v_j) = -g(L_\alpha, L_\beta)\varepsilon_l \delta_{lj}.$$

We proved that for some positive integer k ,

$$\mathcal{L} = \{L_0, A_{L_0} v_1, \dots, A_{L_0} v_r, \dots, L_{k-1}, A_{L_{k-1}} v_1, \dots, A_{L_{k-1}} v_r\}$$

is an orthonormal basis of \mathcal{H} along the fibre $\pi^{-1}(\pi(p))$. Thus $\dim B = (1 + \dim \text{fibre})k$ for some positive integer k . Counting the timelike vectors in \mathcal{L} , we get $\text{index}(B) = q_1(r' + 1) + q_2(r - r')$ for some nonnegative integers q_1, q_2 with $q_1 + q_2 = k$.

(2) Let $x \in B$ and $X', Y' \in T_x B$ such that $g'(X', X') = g'(Y', Y') \neq 0$. We shall construct an isometry $\tilde{f} : B \rightarrow B$ such that $\tilde{f}(x) = x$ and $\tilde{f}_* X' = Y'$. Note that we may assume that $g'(X', X') = g'(Y', Y') = \pm 1$. Let X, Y be the horizontal liftings along the fibre $\pi^{-1}(x)$ of X' and Y' , respectively. Take $p \in \pi^{-1}(x)$. Let

$$\begin{aligned}\mathcal{L} &= \{L_0, A_{L_0}v_1, \dots, A_{L_0}v_r, \dots, L_{k-1}, A_{L_{k-1}}v_1, \dots, A_{L_{k-1}}v_r\}, \\ \mathcal{L}' &= \{L'_0, A_{L'_0}v'_1, \dots, A_{L'_0}v'_r, \dots, L'_{k-1}, A_{L'_{k-1}}v'_1, \dots, A_{L'_{k-1}}v'_r\}\end{aligned}$$

be two orthonormal bases constructed as above such that $L_0 = X, L'_0 = Y, g(L_\alpha, L_\alpha) = g(L'_\alpha, L'_\alpha)$ for $\alpha \in \{1, \dots, k-1\}$, and that $\{v_1 = A_X Y_1, \dots, v_r = A_X Y_r\}$ and $\{v'_1 = A_Y Y'_1, \dots, v'_r = A_Y Y'_r\}$ are orthonormal bases of the tangent bundle of the fibre $\pi^{-1}(\pi(p))$, where Y_1, \dots, Y_r and Y'_1, \dots, Y'_r are the horizontal liftings along $\pi^{-1}(\pi(p))$ of the vectors $\pi_* A_X v_{1p}, \dots, \pi_* A_X v_{rp}$ and $\pi_* A_Y v'_{1p}, \dots, \pi_* A_Y v'_{rp}$, respectively (as in Lemma 3.1), for which $g(v_i, v_j) = g(v'_i, v'_j)$ for $i, j \in \{1, \dots, r\}$. Let $\phi : T_p M \rightarrow T_p M$ be the linear map given by $\phi(L_\alpha) = L'_\alpha, \phi(v_j) = v'_j, \phi(A_{L_\alpha} v_j) = A_{L'_\alpha} v'_j$ for every $\alpha \in \{0, \dots, k-1\}$ and $j \in \{1, \dots, r\}$. Since both $\mathcal{L}, \mathcal{L}'$ are orthonormal bases, we see that ϕ is a linear isometry.

We shall apply Theorem 2.4. Thus we need to prove that $\phi(A_E F) = A_{\phi(E)}\phi(F)$ for every $E, F \in T_p M$. Indeed, we obtain for $\alpha, \beta \in \{0, \dots, k-1\}$ and $j, l \in \{1, \dots, r\}$,

$$\begin{aligned}\phi(A_{L_\alpha} L_\beta) &= \phi(0) = 0 = A_{L'_\alpha} L'_\beta = A_{\phi(L_\alpha)}\phi(L_\beta), \\ g(v_j, A_{L_\alpha} A_{L_\beta} v_l) &= -g(A_{L_\alpha} v_j, A_{L_\beta} v_l) = -g(L_\alpha, L_\beta)g(v_j, v_l) \\ &= -g(L'_\alpha, L'_\beta)g(v'_j, v'_l) = g(v'_j, A_{L'_\alpha} A_{L'_\beta} v'_l).\end{aligned}$$

Hence $\phi(A_{L_\alpha} A_{L_\beta} v_l) = A_{\phi(L_\alpha)}\phi(A_{L_\beta} v_l)$. \square

LEMMA 3.4. $A_{L_\alpha} v_j$ is a basic vector field along the fibre $\pi^{-1}(\pi(p))$ for every $1 \leq j \leq r$ and $\alpha \geq 0$.

PROOF OF LEMMA 3.4. We have $g(A_X v_j, Z) = g(A_X A_X Y_j, Z) = -g(A_X Y_j, A_X Z)$. For every basic vector field Z along the fibre $\pi^{-1}(\pi(p))$ we know that $g(A_X Y_j, A_X Z)$ is constant along the fibre $\pi^{-1}(\pi(p))$. Hence $A_X v_j$ is a basic vector field along the fibre $\pi^{-1}(\pi(p))$.

Now we assume $\alpha \geq 1$. Since $\dim(\ker A_X^* \cup \ker A_{L_\alpha}^*) = \dim \ker A_X^* + \dim \ker A_{L_\alpha}^* - \dim(\ker A_X^* \cap \ker A_{L_\alpha}^*) = (n-r) + (n-r) - (n-2r) = n$, it follows that $\ker A_X^* + \ker A_{L_\alpha}^* = \mathcal{H}$. Hence $A_{L_\alpha} v_j$ is a basic vector field along the fibre $\pi^{-1}(\pi(p))$ if and only if the following conditions are satisfied: $g(A_{L_\alpha} v_j, Z_1)$ is constant along $\pi^{-1}(\pi(p))$ for every $Z_1 \in \ker A_X^*$, which is a basic vector field along $\pi^{-1}(\pi(p))$, and $g(A_{L_\alpha} v_j, Z_2)$ is constant along the fibre $\pi^{-1}(\pi(p))$ for every $Z_2 \in \ker A_{L_\alpha}^*$, which is a basic vector field along $\pi^{-1}(\pi(p))$. If $Z_2 \in \ker A_{L_\alpha}^*$, then $A_{L_\alpha}^* Z_2 = 0$ along $\pi^{-1}(\pi(p))$. So $g(A_{L_\alpha} v_j, Z_2) = -g(v_j, A_{L_\alpha} Z_2) = 0$ along $\pi^{-1}(\pi(p))$. If $Z_1 \in \ker A_X^*$, then $A_X^* Z_1 = 0$ along $\pi^{-1}(\pi(p))$. By O'Neill's equations, we

get along the fibre $\pi^{-1}(\pi(p))$

$$\begin{aligned} R'(\pi_* X, \pi_* Y_j, \pi_* L_\alpha, \pi_* Z_1) &= R(X, Y_j, L_\alpha, Z_1) + 2g(A_X Y_j, A_{L_\alpha} Z_1) \\ &\quad - g(A_{Y_j} L_\alpha, A_X Z_1) - g(A_{L_\alpha} X, A_{Y_j} Z_1) \\ &= -g(X, L_\alpha)g(Y_j, Z_1) + g(X, Z_1)g(Y_j, L_\alpha) \\ &\quad + 2g(v_j, A_{L_\alpha} Z_1), \end{aligned}$$

since $A_{L_\alpha} X = -A_X L_\alpha = 0$ and $A_X Z_1 = 0$. Hence $g(v_j, A_{L_\alpha} Z_1) = -g(A_{L_\alpha} v_j, Z_1)$ is constant along $\pi^{-1}(\pi(p))$ for every $Z_1 \in \ker A_X^*$, which is a basic vector field along $\pi^{-1}(\pi(p))$.

We proved that $A_{L_\alpha} v_j$ is a basic vector field along $\pi^{-1}(\pi(p))$ for every $\alpha \geq 0$ and $j \in \{1, \dots, r\}$. \square

We denote by $\hat{\nabla}$ the induced Levi-Civita connection on the fibre $\pi^{-1}(\pi(p))$.

$$\text{LEMMA 3.5. } A_{A_{L_\alpha} v_i} A_{L_\beta} v_j = g(L_\alpha, L_\beta) \hat{\nabla}_{v_i} v_j.$$

PROOF OF LEMMA 3.5. By the relation (3.1) together with Lemma 3.4, we obtain for $i, j, l \in \{1, \dots, r\}$ and $\alpha, \beta \geq 0$ that

$$\begin{aligned} g(A_{A_{L_\alpha} v_i} A_{L_\beta} v_j, v_l) &= -g(A_{A_{L_\alpha} v_i} v_l, A_{L_\beta} v_j) \\ &= -R(L_\beta, v_l, A_{L_\alpha} v_i, v_j) + g(\nabla_{v_l} A_{L_\beta} A_{L_\alpha} v_i, v_j) \\ &= g(L_\beta, A_{L_\alpha} v_i)g(v_l, v_j) + v_l g(A_{L_\beta} A_{L_\alpha} v_i, v_j) \\ &\quad - g(A_{L_\beta} A_{L_\alpha} v_i, \nabla_{v_l} v_j) \\ &= -v_l g(A_{L_\alpha} v_i, A_{L_\beta} v_j) + g(A_{L_\alpha} v_i, A_{L_\beta} v_l)g(\nabla_{v_l} v_j, v_i) \varepsilon_l \\ &= -g(L_\alpha, L_\beta)g(\hat{\nabla}_{v_l} v_j, v_i) \\ &= g(L_\alpha, L_\beta)g(\hat{\nabla}_{v_i} v_j, v_l). \end{aligned}$$

In the last equality we used the fact that $v_j = A_X Y_j$ is a Killing vector field along the fibre $\pi^{-1}(\pi(p))$ (see [Bis] or [Bes]). Thus

$$A_{A_{L_\alpha} v_i} A_{L_\beta} v_j = g(L_\alpha, L_\beta) \hat{\nabla}_{v_i} v_j.$$

\square

LEMMA 3.6. *The following assertions are true:*

- (a) $r \neq 2$.
- (b) *If* $r = 1$, *then* $A_{A_{L_\alpha} v_1} A_{L_\beta} v_1 = 0$.
- (c) *If* $r = 3$ *and if we set* $v_{3p} = g(X, X)^{-1}(\hat{\nabla}_{v_1} v_2)(p)$, *then* $v_3 = \hat{\nabla}_{v_1} v_2$ *and*

$$g(\hat{\nabla}_{v_i} v_j, v_k) = \begin{cases} 0 & \text{if two of } i, j, k \text{ are equal,} \\ \varepsilon \binom{1 \ 2 \ 3}{i \ j \ k} g(v_3, v_3) & \text{if } \{i, j, k\} = \{1, 2, 3\}, \end{cases}$$

where $\varepsilon \binom{1 \ 2 \ 3}{i \ j \ k}$ is the signature of the permutation $\binom{1 \ 2 \ 3}{i \ j \ k}$.

PROOF OF LEMMA 3.6. Since v_1, \dots, v_r are Killing vector fields along $\pi^{-1}(\pi(p))$ and $g(v_i, v_i) \in \{-1, 1\}$ for every i , we get

$$g(\hat{\nabla}_{v_i} v_j, v_i) = g(\hat{\nabla}_{v_i} v_i, v_j) = g(\hat{\nabla}_{v_j} v_i, v_i) = 0$$

for every $i, j \in \{1, \dots, r\}$.

(a) The case $r = 2$ is not possible. Indeed, if $r = 2$, then the relation $g(\nabla_{v_1} v_2, v_1) = g(\nabla_{v_1} v_2, v_2) = 0$ implies $\nabla_{v_1} v_2 = 0$. On the other hand,

$g(\nabla_{v_1} v_2, \nabla_{v_1} v_2) = -g(\hat{\nabla}_{v_1} \hat{\nabla}_{v_2} v_2, v_1) + \hat{R}(v_1, v_2, v_1, v_2) = -g(v_1, v_1)g(v_2, v_2) \in \{-1, 1\}$, since $\hat{\nabla}_{v_2} v_2 = g(X, X)^{-1} A_{A_X v_2} A_X v_2 = 0$ and each fibre has constant curvature -1 . So we get a contradiction.

(b) If $r = 1$, then $A_{A_{L_\alpha} v_1} A_{L_\beta} v_1 = 0$ for every α and β , because $0 = A_{A_X v_1} A_X v_1 = g(X, X) \nabla_{v_1} v_1$ implies $\nabla_{v_1} v_1 = 0$.

(c) In the case $r = 3$ we shall prove $g(\hat{\nabla}_{v_1} v_2, v_3)$ is constant along the fibre $\pi^{-1}(\pi(p))$. Since O'Neill's integrability tensor A is skew-symmetric, it follows that $\hat{\nabla}_{v_i} v_j = -\hat{\nabla}_{v_j} v_i$. Then $\hat{\nabla}_{v_i} v_j = (1/2)[v_i, v_j]$ is a Killing vector field along $\pi^{-1}(\pi(p))$. We then obtain

$$\begin{aligned} v_1 g(\hat{\nabla}_{v_1} v_2, v_3) &= g(\hat{\nabla}_{v_1} \hat{\nabla}_{v_1} v_2, v_3) + g(\hat{\nabla}_{v_1} v_2, \hat{\nabla}_{v_1} v_3) \\ &= -g(\hat{\nabla}_{v_3} \hat{\nabla}_{v_1} v_2, v_1) + g(\hat{\nabla}_{v_1} v_2, \hat{\nabla}_{v_1} v_3) \\ &= -v_3 g(\hat{\nabla}_{v_1} v_2, v_1) + g(\hat{\nabla}_{v_1} v_2, \hat{\nabla}_{v_1} v_3 + \hat{\nabla}_{v_3} v_1) = 0. \end{aligned}$$

Analogously, we get $v_2 g(\hat{\nabla}_{v_1} v_2, v_3) = -v_2 g(\hat{\nabla}_{v_2} v_1, v_3) = 0$. We also obtain

$$v_3 g(\hat{\nabla}_{v_1} v_2, v_3) = g(\hat{\nabla}_{v_3} \hat{\nabla}_{v_1} v_2, v_3) + g(\hat{\nabla}_{v_1} v_2, \hat{\nabla}_{v_3} v_3) = 0,$$

since $\hat{\nabla}_{v_3} v_3 = 0$ and $\hat{\nabla}_{v_1} v_2$ is a Killing vector field along $\pi^{-1}(\pi(p))$. It is easy to see that

$$\begin{aligned} g(\hat{\nabla}_{v_1} v_2, v_3) &= -g(\hat{\nabla}_{v_2} v_1, v_3) = g(\hat{\nabla}_{v_2} v_3, v_1) \\ &= -g(\hat{\nabla}_{v_3} v_2, v_1) = g(\hat{\nabla}_{v_3} v_1, v_2) = -g(\hat{\nabla}_{v_1} v_3, v_2). \end{aligned}$$

Thus $g(\hat{\nabla}_{v_i} v_j, v_l)$ is constant along the fibre $\pi^{-1}(\pi(p))$ for each $i, j, l \in \{1, 2, 3\}$. Therefore

$$g(A_X A_{A_X v_i} A_X v_j, A_X v_l) = -g(X, X) g(A_{A_X v_i} A_X v_j, v_l) = -g(X, X)^2 g(\hat{\nabla}_{v_i} v_j, v_l)$$

is constant along $\pi^{-1}(\pi(p))$. Also, we compute for $\alpha \geq 1$

$$g(A_X A_{A_X v_i} A_X v_j, A_{L_\alpha} v_l) = -g(A_{A_X v_i} A_X v_j, A_X A_{L_\alpha} v_l) = 0,$$

$$g(A_X A_{A_X v_i} A_X v_j, L_\alpha) = -g(A_{A_X v_i} A_X v_j, A_X L_\alpha) = 0.$$

Hence $A_X A_{A_X v_i} A_X v_j = g(X, X) A_X \hat{\nabla}_{v_i} v_j$ is a basic vector field for each $i, j \in \{1, \dots, r\}$.

We choose $v_{3p} = (g(X, X)^{-1} \hat{\nabla}_{v_1} v_2)(p)$. Since $A_X \hat{\nabla}_{v_1} v_2$ is a basic vector field along $\pi^{-1}(\pi(p))$, we get the horizontal lifting along $\pi^{-1}(\pi(p))$ of $\pi_*(g(X, X)^{-1} A_X \hat{\nabla}_{v_1} v_2)(p) = \pi_* A_X v_{3p}$ is $g(X, X)^{-1} A_X \hat{\nabla}_{v_1} v_2$. On the other hand, Y_3 is, by definition, the horizontal lifting along $\pi^{-1}(\pi(p))$ of $\pi_* A_X v_{3p}$. It follows that $Y_3 = g(X, X)^{-1} A_X \hat{\nabla}_{v_1} v_2$ along $\pi^{-1}(\pi(p))$. Thus $v_3 = A_X Y_3 = g(X, X)^{-1} A_X A_X \hat{\nabla}_{v_1} v_2 = \hat{\nabla}_{v_1} v_2$ along the fibre $\pi^{-1}(\pi(p))$. \square

For $r = 3$, we choose $v'_{3p} = g(Y, Y)^{-1}(\hat{\nabla}_{v'_1} v'_2)(p)$. If we repeat the argument above for the basis $\{v'_1, v'_2, v'_3\}$, by Lemma 3.6, we get $v'_3 = \hat{\nabla}_{v'_1} v'_2$ along the fibre $\pi^{-1}(\pi(p))$. It follows that $g(\hat{\nabla}_{v_i} v_j, v_l) = g(\hat{\nabla}_{v'_i} v'_j, v'_l)$ for each $i, j, l \in \{1, 2, 3\}$.

Returning to the computation of $g(A_{A_{L_\alpha} v_i} A_{L_\beta} v_j, v_l)$, in both cases $r = 1$ and $r = 3$, we get for every $\alpha, \beta \geq 0$ and $i, j, k \in \{1, \dots, r\}$

$$\begin{aligned} g(A_{A_{L_\alpha} v_i} A_{L_\beta} v_j, v_l) &= g(L_\alpha, L_\beta) g(\hat{\nabla}_{v_i} v_j, v_l) \\ &= g(L'_\alpha, L'_\beta) g(\hat{\nabla}_{v'_i} v'_j, v'_l) = g(A_{A_{L'_\alpha} v'_i} A_{L'_\beta} v'_j, v'_l). \end{aligned}$$

Hence $\phi(A_{A_{L_\alpha} v_i} A_{L_\beta} v_j) = A_{\phi(A_{L_\alpha} v_i)} \phi(A_{L_\beta} v_j)$ and $\phi(A_{A_{L_\alpha} v_i} v_j) = A_{\phi(A_{L_\alpha} v_i)} \phi(v_j)$.

By Corollary 2.3.14 in [Wol] we see that $\phi : T_p M \rightarrow T_p M$ extends to an isometry on M , denoted by $f : M \rightarrow M$, such that $f(p) = p$ and $f_{*p} = \phi$. Hence $f_{*p} X = Y$ and $f_*(\mathcal{H}_p) = \mathcal{H}_p$. Since $f_* A_E F = A_{f_* E} f_* F$ for every $E, F \in T_p M$, we see, by Theorem 2.4, that there is an isometry $\tilde{f} : B \rightarrow B$ such that $\tilde{f} \circ \pi = \pi \circ f$. Thus $\tilde{f}_* X' = \tilde{f}_* \pi_* X = \pi_* f_* X = \pi_* Y = Y'$ and $\tilde{f}(x) = \tilde{f}(\pi(p)) = \pi(f(p)) = \pi(p) = x$.

Therefore B is an isotropic semi-Riemannian manifold. This completes the proof of Theorem 3.3. \square

If the metric on the base space is negative definite, the following lemma follows from Theorem 3.3.

LEMMA 3.7. *If $\pi : M \rightarrow B$ is a semi-Riemannian submersion with connected totally geodesic fibres from an $(n+r)$ -dimensional semi-Riemannian manifold M of index $r'+n$ and of constant negative curvature onto an n -dimensional semi-Riemannian manifold B of index n , then $r' = r$.*

PROOF. By Theorem 3.3, we have $n = q_1(r'+1) + q_2(r-r') = (q_1+q_2)(r+1)$ for some nonnegative integers q_1 and q_2 . Hence $0 = q_1(r-r') + q_2(r'+1)$. Since the right hand side is the sum of two non-negative numbers, it follows that $q_1(r-r') = 0$ and $q_2(r'+1) = 0$. Therefore $q_2 = 0$. This implies $r' = r$. \square

REMARK. Changing simultaneously the signs of metrics on the total space and on the base space, any semi-Riemannian submersion, under the assumptions of Lemma 3.7, becomes a Riemannian submersion with totally geodesic fibres from a sphere onto a Riemannian manifold. This case was completely classified by Escobales (see [Esc1]) and Ranjan (see [Ran1]).

PROPOSITION 3.8. *Let $\pi : M \rightarrow B$ be a semi-Riemannian submersion with connected totally geodesic fibres from a complete simply connected semi-Riemannian manifold M onto a semi-Riemannian manifold B . Then B is simply connected and complete.*

PROOF. If M is geodesically complete, then so is B (see [Bes] or [Ba-Ia]). Since M is a complete semi-Riemannian manifold and the fibres are totally geodesic, any fibre is also geodesically complete. By a theorem in [Rec], it follows that the horizontal distribution \mathcal{H} is an Ehresmann connection. Therefore, by [Ehr], we see that π is a fibre bundle. So we obtain

an exact homotopy sequence:

$$\cdots \rightarrow \pi_2(M) \rightarrow \pi_2(B) \rightarrow \pi_1(\text{fibre}) \rightarrow \pi_1(M) \rightarrow \pi_1(B) \rightarrow 0.$$

Thus $\pi_1(B) = 0$. □

By Theorem 12.3.2 in [Wol], we know that any connected, simply connected isotropic semi-Riemannian manifold is isometric to one of the following semi-Riemannian manifolds:

(i) \mathbf{R}_t^m or the universal semi-Riemannian covering of the pseudo-hyperbolic space $H_t^m(c)$ with constant sectional curvature $c < 0$, or of the pseudo-sphere $S_t^m(c)$ with constant sectional curvature $c > 0$.

(ii) The complex pseudo-hyperbolic space $\mathbf{C}H_t^m(c)$ with constant holomorphic sectional curvature $c < 0$, or the complex pseudo-projective space $\mathbf{C}P_t^m(c)$ with constant holomorphic sectional curvature $c > 0$.

(iii) The quaternionic pseudo-hyperbolic space $\mathbf{H}H_t^m(c)$ with constant quaternionic sectional curvature $c < 0$, or the quaternionic pseudo-projective space $\mathbf{H}P_t^m(c)$ with constant quaternionic sectional curvature $c > 0$.

(iv) The Cayley pseudo-hyperbolic plane $\mathbf{C}aH_t^2(c)$ with Cayley sectional curvature $c < 0$, or the Cayley pseudo-projective plane $\mathbf{C}aP_t^2(c)$ with Cayley sectional curvature $c > 0$.

LEMMA 3.9. (a) *If B is a semi-Riemannian manifold isometric to one of the semi-Riemannian manifolds $\mathbf{C}P_t^m(c)$, $\mathbf{H}P_t^m(c)$, $\mathbf{C}aP_t^2(c)$ ($c > 0$), then the curvature tensor satisfies the inequality*

$$(3.2) \quad R'(X', Y', X', Y') \geq \frac{c}{4}(g'(X', X')g'(Y', Y') - g'(X', Y')^2)$$

for each tangent vectors X', Y' of B .

(b) *If B is a semi-Riemannian manifold isometric to one of the semi-Riemannian manifolds $\mathbf{C}H_t^m(c)$, $\mathbf{H}H_t^m(c)$, $\mathbf{C}aH_t^2(c)$ ($c < 0$), then the curvature tensor satisfies the inequality*

$$(3.3) \quad R'(X', Y', X', Y') \leq \frac{c}{4}(g'(X', X')g'(Y', Y') - g'(X', Y')^2)$$

for each tangent vectors X', Y' of B .

PROOF. For each tangent vectors X', Y' of B , we have the following formulas for the curvature tensors:

(i) If $B \in \{\mathbf{C}P_t^m(c), \mathbf{C}H_t^m(c)\}$ and I_0 is the natural complex structure on B , then

$$(3.4) \quad R'(X', Y', X', Y') = \frac{c}{4}(g'(X', X')g'(Y', Y') - g'(X', Y')^2 + 3g'(X', I_0Y')^2).$$

(ii) If $B \in \{\mathbf{H}P_t^m(c), \mathbf{H}H_t^m(c)\}$ and I_0, J_0, K_0 are local almost complex structures which give rise to the quaternionic structure on B , then

$$(3.5) \quad R'(X', Y', X', Y') = (c/4)(g'(X', X')g'(Y', Y') - g'(X', Y')^2 + 3g'(X', I_0Y')^2 + 3g'(X', J_0Y')^2 + 3g'(X', K_0Y')^2).$$

(iii) If $B \in \{\mathcal{C}aP_t^2(c), \mathcal{C}aH_t^2(c)\}$ and $I_0, J_0, K_0, M_0, M_0I_0, M_0J_0, M_0K_0$ are local almost complex structures which give rise to the Cayley structure on B , then

$$(3.6) \quad \begin{aligned} R'(X', Y', X', Y') &= (c/4)(g'(X', X')g'(Y', Y') - g'(X', Y')^2 \\ &\quad + 3g'(X', I_0Y')^2 + 3g'(X', J_0Y')^2 + 3g'(X', K_0Y')^2 \\ &\quad + 3g'(X', M_0Y')^2 + 3g'(X', M_0I_0Y')^2 + 3g'(X', M_0J_0Y')^2 \\ &\quad + 3g'(X', M_0K_0Y')^2). \end{aligned}$$

By these explicit formulas for curvature tensors, in all cases we obtain the inequalities (3.2) and (3.3). \square

First, we shall discuss the case of a base space with nonconstant curvature.

LEMMA 3.10. *If $\pi : H_{s+r}^{n+r} \rightarrow B_s^n$ is a semi-Riemannian submersion with connected totally geodesic fibres from an $(n+r)$ -dimensional pseudo-hyperbolic space H_{s+r}^{n+r} of index $s+r' > 1$ onto an n -dimensional isotropic semi-Riemannian manifold B_s^n of index s with nonconstant curvature, then the induced metrics on the fibres are negative definite and B is isometric to one of the following semi-Riemannian manifolds:*

- (i) $\mathcal{C}H_t^m, m > 1,$
- (ii) $\mathcal{H}H_t^m, m > 1,$
- (iii) $\mathcal{C}aH_t^2.$

PROOF. Since $\dim \mathcal{H} = k(\dim \mathcal{V} + 1)$ for some positive integer k , we get $\dim \mathcal{H} \geq \dim \mathcal{V} + 1$. Let X be a horizontal vector field along a fibre $\pi^{-1}(\pi(p))$ such that $g(X, X) \neq 0$ and X is the horizontal lifting of some tangent vector of B .

First, we shall prove that

$$\dim \mathcal{H} > \dim \mathcal{V} + 1.$$

Suppose that $\dim \mathcal{H} = \dim \mathcal{V} + 1$. Then $A_X : \mathcal{V} \rightarrow X^\perp = \{Y \in \mathcal{H} \mid g(X, Y) = 0\}$ is bijective. For every $Y \in X^\perp$ we get $Y = A_X V$ for some vertical vector V . It follows that

$$\begin{aligned} g(A_X Y, A_X Y) &= g(A_X A_X V, A_X A_X V) = g(X, X)^2 g(V, V), \\ g(Y, Y) &= g(A_X V, A_X V) = -g(X, X)g(V, V). \end{aligned}$$

Thus $g(A_X Y, A_X Y) = -g(X, X)g(Y, Y)$ for every $Y \in X^\perp$. By O'Neill's equations, we have

$$\begin{aligned} R'(\pi_* X, \pi_* Y, \pi_* X, \pi_* Y) &= -g(X, X)g(Y, Y) + g(X, Y)^2 + 3g(A_X Y, A_X Y) \\ &= -4(g(X, X)g(Y, Y) - g(X, Y)^2) \end{aligned}$$

for every horizontal vector field Y along $\pi^{-1}(\pi(p))$. Hence B has constant curvature, a contradiction.

We established that $\dim \mathcal{H} > \dim \mathcal{V} + 1$. So we can find a horizontal vector field Z along the fibre $\pi^{-1}(\pi(p))$ such that $Z \in \ker A_X^*$, $g(X, Z) = 0$, $g(Z, Z) \neq 0$ and Z is the horizontal

lifting of some $Z' \in T_{\pi(p)}B$. We then have

$$\begin{aligned} R'(\pi_*X, \pi_*Z, \pi_*X, \pi_*Z) &= -g(X, X)g(Z, Z) + g(X, Z)^2 + 3g(A_XZ, A_XZ) \\ &= -g(X, X)g(Z, Z). \end{aligned}$$

Since B is a simply connected isotropic semi-Riemannian manifold with nonconstant curvature, we see that B is isometric to one of the following semi-Riemannian manifolds:

- (a) $CP_t^m(c)$, $HP_t^m(c)$, $CaP_t^2(c)$, or
- (b) $CH_t^m(c)$, $HH_t^m(c)$, $CaH_t^2(c)$.

We shall prove that only the case (b) is possible.

First, we suppose that B is isometric to one of the following semi-Riemannian manifolds:

$$CP_t^m(c), \quad HP_t^m(c), \quad CaP_t^2(c) \quad (c > 0).$$

By the inequality (3.2), we get

$$\begin{aligned} R'(\pi_*X, \pi_*A_XV, \pi_*X, \pi_*A_XV) &= -4g(X, X)g(A_XV, A_XV) \\ &= 4g(X, X)^2g(V, V) \geq -(c/4)g(X, X)^2g(V, V). \end{aligned}$$

Therefore

$$(3.7) \quad g(V, V) \geq 0$$

for every vertical vector V . Since X and Z are basic vector fields along $\pi^{-1}(\pi(p))$ with $g(X, Z) = 0$ and $A_XZ = 0$ along $\pi^{-1}(\pi(p))$, it follows from the relation (3.1) that $A_ZV \in \ker A_X^*$. On the other hand, by the inequality (3.2), we get

$$\begin{aligned} R'(\pi_*X, \pi_*Z, \pi_*X, \pi_*Z) &= -g(X, X)g(Z, Z) \geq (c/4)g(X, X)g(Z, Z), \\ R'(\pi_*X, \pi_*A_ZV, \pi_*X, \pi_*A_ZV) &= -g(X, X)g(A_ZV, A_ZV) \\ &\geq (c/4)g(X, X)g(A_ZV, A_ZV). \end{aligned}$$

Hence $g(X, X)g(Z, Z) \leq 0$ and $g(X, X)g(A_ZV, A_ZV) \leq 0$. Thus

$$0 \leq g(Z, Z)g(A_ZV, A_ZV) = -g(Z, Z)^2g(V, V).$$

So for any vertical vector V we get

$$(3.8) \quad g(V, V) \leq 0.$$

Since the induced metrics on fibres are nondegenerate, it is not possible to have both (3.7) and (3.8). So we obtain the required contradiction. It follows that B is isometric to one of the following semi-Riemannian manifolds:

$$CH_t^m(c), \quad HH_t^m(c), \quad CaH_t^2(c) \quad (c < 0).$$

We shall now prove that $c = -4$. Suppose $(c/4) + 1 \neq 0$. By the inequality (3.3), we get

$$(3.9) \quad \begin{aligned} R'(\pi_*X, \pi_*Z, \pi_*X, \pi_*Z) &= -g(X, X)g(Z, Z) \leq (c/4)g(X, X)g(Z, Z), \\ R'(\pi_*X, \pi_*A_ZV, \pi_*X, \pi_*A_ZV) &= -g(X, X)g(A_ZV, A_ZV) \\ &\leq (c/4)g(X, X)g(A_ZV, A_ZV). \end{aligned}$$

Hence

$$(3.10) \quad ((c/4) + 1)^2 g(X, X)^2 g(Z, Z) g(A_ZV, A_ZV) \geq 0,$$

from which follows that $0 \leq g(Z, Z)g(A_ZV, A_ZV) = -g(Z, Z)^2 g(V, V)$. Therefore $g(V, V) \leq 0$ for every vertical vector field V . In particular, we have $g(A_XY, A_XY) \leq 0$, which implies

$$(3.11) \quad R'(\pi_*X, \pi_*Y, \pi_*X, \pi_*Y) \leq g(X, X)g(Y, Y) - g(X, Y)^2$$

for every horizontal vectors X and Y . We have the following cases:

Case (a) $0 < \text{index } B < \dim B$. We can choose vector fields X', Y' on B such that $g'(X', X')g'(Y', Y') < 0$ and that one of the following conditions is satisfied:

(i) $Y' \in \{X', I_0X'\}^\perp$ if $B = \mathbf{CH}_s^m(c)$, where I_0 is the natural complex structure on $\mathbf{CH}_s^m(c)$,

(ii) $Y' \in \{X', I_0X', J_0X', K_0X'\}^\perp$ if $B = \mathbf{HH}_s^m(c)$, where $\{I_0, J_0, K_0\}$ are local almost complex structures which give rise to the quaternionic structure on $\mathbf{HH}_s^m(c)$, or

(iii) $Y' \in \{X', I_0X', J_0X', K_0X', M_0X', M_0I_0X', M_0J_0X', M_0K_0X'\}^\perp$ if $B = \mathbf{CaH}_t^2(c)$, where $\{I_0, J_0, K_0, M_0, M_0I_0, M_0J_0, M_0K_0\}$ are local almost complex structures which give rise to the Cayley structure on $\mathbf{CaH}_t^2(c)$.

Let X, Y be the horizontal liftings of X', Y' . The inequality (3.11) then implies

$$\frac{c}{4}g(X, X)g(Y, Y) \leq -g(X, X)g(Y, Y).$$

Hence $((c/4) + 1)g(X, X)g(Y, Y) \leq 0$. Therefore $(c/4) + 1 > 0$. On the other hand, we can choose horizontal vector fields X, Z such that $g(X, Z) = 0$, $Z \in \ker A_X^*$ and $g(X, X)g(Z, Z) < 0$, because $0 < \text{index } B < \dim B$. Then the inequality (3.9) becomes $(c/4) + 1 < 0$. So we get a contradiction.

Case (b) $\text{index } B \in \{0, \dim B\}$. Similarly, we can choose vector fields X', Y' on B such that $g'(X', Y') = 0$ and $R'(X', Y', X', Y') = (c/4)g'(X', X')g'(Y', Y')$. The inequality (3.11) then implies $((c/4) + 1)g'(X', X')g'(Y', Y') \leq 0$. By the hypothesis of Case (b), we get $(c/4) + 1 \leq 0$. On the other hand, the inequality (3.9) becomes $(c/4) + 1 > 0$. So we get a contradiction.

We have proved $c = -4$. The inequality (3.3) then becomes

$$(3.12) \quad R'(X', Y', X', Y') \leq -g'(X', X')g'(Y', Y') + g'(X', Y')^2$$

for tangent vector fields X', Y' on B . Then we have

$$R'(\pi_*X, \pi_*A_XV, \pi_*X, \pi_*A_XV) = -4g(X, X)g(A_XV, A_XV) \leq -g(X, X)g(A_XV, A_XV)$$

for a vertical vector field V and for a horizontal vector field X . Hence

$$0 \leq g(X, X)g(A_X V, A_X V) = -g(X, X)^2 g(V, V).$$

Therefore the induced metrics on fibres are negative definite. \square

By Lemma 3.10, we deduce the following proposition.

PROPOSITION 3.11. *If $\pi : H_{s+r'}^{n+r} \rightarrow B_s^n$ is a semi-Riemannian submersion with connected totally geodesic fibres from an $(n+r)$ -dimensional pseudo-hyperbolic space $H_{s+r'}^{n+r}$ of index $s+r'$ onto an n -dimensional isotropic semi-Riemannian manifold B_s^n of index s with nonconstant curvature, then one of the following holds:*

(1) $n = 2m > 2$, $s = 2t$, $r = r' = 1$ for some non-negative integers m, t , and B_s^n is isometric to $\mathbf{C}H_t^m$.

(2) $n = 4m > 4$, $s = 4t$, $r = r' = 3$ for some non-negative integers m, t , and B_s^n is isometric to $\mathbf{H}H_t^m$.

(3) $n = 16$, $s \in \{0, 8, 16\}$, $r = r' = 7$, and B_s^n is isometric to $\mathbf{C}aH_{s/8}^2$.

PROOF. First, we shall discuss the case $s+r' > 1$. By Lemma 3.10, B is isometric to one of the semi-Riemannian manifolds $\mathbf{C}H_t^m$, $\mathbf{H}H_t^m$, $\mathbf{C}aH_t^2$ for some $m > 1$.

Let $x \in B$ and let $X' \in T_x B$ such that $g'(X', X') \neq 0$, and let $\mathcal{F}_{X'}$ be the subspace in $T_x B$ given by

$$\mathcal{F}_{X'} = \{Y' \in T_x B \mid R'(X', Y')X' = -g'(X', Y')X' + g'(X', X')Y'\}.$$

Let $p \in \pi^{-1}(x)$ and let X be the horizontal lifting vector at p of X' . By O'Neill's equations, we have $R'(\pi_* X, \pi_* Y, \pi_* X, \pi_* Z) = R(X, Y, X, Z) + 3g(A_X^* Y, A_X^* Z)$ for horizontal vectors Y, Z . Since $A_X^* : \mathcal{H}_p \rightarrow \mathcal{V}_p$ is surjective and since the induced metrics on fibres are nondegenerate, we get $Y \in \ker A_X^*$ if and only if $\pi_* Y \in \mathcal{F}_{X'}$. Thus

$$\dim \ker A_X^* = \dim \mathcal{H} - \dim \mathcal{V} = \dim \mathcal{F}_{X'}.$$

We have the following possibilities:

(1) B_s^n is isometric to $\mathbf{C}H_t^m$. So $n = 2m$, $s = 2t$. From the geometry of the complex pseudo-hyperbolic space (see relation (3.4)), we get $\dim \mathcal{F}_{X'} = \dim \mathcal{H} - 1$. It follows that $r = r' = \dim \mathcal{V} = 1$.

(2) B_s^n is isometric to $\mathbf{H}H_t^m$. So $n = 4m$, $s = 4t$. From the geometry of the quaternionic pseudo-hyperbolic space (see relation (3.5)), we get $\dim \mathcal{F}_{X'} = \dim \mathcal{H} - 3$. It follows that $r = r' = \dim \mathcal{V} = 3$.

(3) B_s^n is isometric to the Cayley pseudo-hyperbolic plane $\mathbf{C}aH_t^2$. So $n = 16$, $s \in \{0, 8, 16\}$. From the geometry of the Cayley pseudo-hyperbolic plane (see relation (3.6)), we obtain $\dim \mathcal{F}_{X'} = \dim \mathcal{H} - 7$. Hence $r = r' = \dim \mathcal{V} = 7$.

Now, we discuss the remaining case $s+r' = 1$. From $s+r' = 1$, we have either

- (i) $s = 0$, $r' = 1$, or
- (ii) $s = 1$, $r' = 0$.

If $s = 0$, $r' = 1$, then $\pi : H_1^{n+r} \rightarrow B^n$ is a semi-Riemannian submersion with totally geodesic fibres from an anti-de Sitter space onto a Riemannian manifold. In this case, investigated by Magid in [Mag], it follows that B is isometric to the complex hyperbolic space CH^m and $r = r' = 1$.

For $s = 1$, $r' = 0$, we get, by Theorem 3.3, $1 = q_1 + q_2r \geq q_1 + q_2$ with $q_1 + q_2 = k = n/(r + 1)$. Thus $q_1 + q_2 = 1$. It follows that $n = r + 1$. Hence $A_X : \mathcal{V} \rightarrow X^\perp$ is bijective. Since $R'(\pi_*X, \pi_*A_XV, \pi_*X, \pi_*A_XV) = -4g(X, X)g(A_XV, A_XV)$, we see that B has constant curvature -4 , which contradicts our assumption of nonconstant curvature of the base space. \square

We shall now discuss the case where the base space is of constant curvature. We give the following obstruction to the existence of semi-Riemannian submersions in terms of the index of base space.

LEMMA 3.12. *There are no semi-Riemannian submersions $\pi : H_{s+r}^{n+r} \rightarrow B_s^n$ with connected totally geodesic fibres from an $(n + r)$ -dimensional pseudo-hyperbolic space of index $s + r'$ onto an n -dimensional semi-Riemannian manifold of index s with constant curvature, where $0 < s < n$.*

PROOF. Let X be a horizontal vector and V a vertical vector such that $g(X, X) \neq 0$ and $g(V, V) \neq 0$. By O'Neill's equations, we have

$$\begin{aligned} R'(\pi_*X, \pi_*A_XV, \pi_*X, \pi_*A_XV) &= -g(X, X)g(A_XV, A_XV) + g(X, A_XV)^2 \\ &\quad + 3g(A_XA_XV, A_XA_XV) \\ &= -g(X, X)g(A_XV, A_XV) + 3g(X, X)^2g(V, V) \\ &= -4g(X, X)g(A_XV, A_XV). \end{aligned}$$

If B has constant curvature, then the curvature of B should be -4 . Therefore, by O'Neill's equations (see (9.29c) in [Bes]), for a horizontal vector Y we get

$$(3.13) \quad g(A_XY, A_XY) = -g(X, X)g(Y, Y) + g(X, Y)^2.$$

By polarization of (3.13), we get

$$g(A_XY, A_XZ) = -g(X, X)g(Y, Z) + g(X, Y)g(X, Z)$$

for horizontal vectors Y and Z . Therefore $A_XA_X^*Y = g(X, X)Y - g(X, Y)X$.

Let $A_X^{*,\perp} : X^\perp \rightarrow \mathcal{V}$ be the restriction of A_X^* to the orthogonal complement of X in \mathcal{H} . Then $A_XA_X^{*,\perp}Y = g(X, X)Y$ for every $Y \in X^\perp$, from which follows that $A_X^{*,\perp}$ is injective. Hence $\dim \mathcal{H} - 1 \leq \dim \mathcal{V}$. By Lemma 3.1, we get $A_X : \mathcal{V} \rightarrow X^\perp$ is injective. So $\dim \mathcal{H} - 1 \geq \dim \mathcal{V}$.

We proved that $\dim \mathcal{H} - 1 = \dim \mathcal{V}$, which means that $n = r + 1$. If $0 < s < n$, then we can choose horizontal vectors X, Y such that $g(X, X) = 1$ and $g(Y, Y) = -1$. Let $\{v_1, v_2, \dots, v_r\}$ be an orthonormal basis of the tangent bundle of the fibre $\pi^{-1}(\pi(p))$, $p \in M$. Since $\{X, A_Xv_1, \dots, A_Xv_r\}$ and $\{Y, A_Yv_1, \dots, A_Yv_r\}$ are orthonormal bases, by the proof of

Theorem 3.3, for every $i \in \{1, \dots, r\}$ we have

$$\begin{aligned} g(A_X v_i, A_X v_i) &= -g(X, X)g(v_i, v_i) = -\varepsilon_i, \\ g(A_Y v_i, A_Y v_i) &= -g(Y, Y)g(v_i, v_i) = \varepsilon_i. \end{aligned}$$

It follows that $s = r - r'$ and $s = r' + 1$. Therefore $r = 2r' + 1$ and $n = 2r' + 2$.

We shall prove that there are no semi-Riemannian submersions $\pi : H_{2r'+1}^{4r'+3} \rightarrow B_{r'+1}^{2r'+2}$ with totally geodesic fibres. Since the fibre $F = \pi^{-1}(\pi(p))$ is a $(2r' + 1)$ -dimensional semi-Riemannian manifold of index r' and since B is a $(2r' + 2)$ -dimensional semi-Riemannian manifold of index $r' + 1$, we can choose orthonormal spacelike vectors $Y_{2r'+2}, \dots, Y_{3r'+2}$ in $T_p F$ (i.e., $g(Y_i, Y_j) = \delta_{ij}$ for $i, j \in \{2r' + 2, \dots, 3r' + 2\}$) and orthonormal spacelike vectors $Y'_{3r'+3}, \dots, Y'_{4r'+3}$ in $T_{\pi(p)} B$ (i.e., $g'(Y'_\alpha, Y'_\beta) = \delta_{\alpha\beta}$ for $\alpha, \beta \in \{3r' + 3, \dots, 4r' + 3\}$). Let $Y_{3r'+3}, \dots, Y_{4r'+3}$ be the horizontal liftings of $Y'_{3r'+3}, \dots, Y'_{4r'+3}$, respectively. Let $Y_1, \dots, Y_{2r'+1}$ be orthonormal timelike vectors in $T_p H_{2r'+1}^{4r'+3}$ (i.e., $g(Y_l, Y_t) = -\delta_{lt}$ for $l, t \in \{1, \dots, 2r' + 1\}$) such that they are orthogonal to $Y_{2r'+2}, \dots, Y_{4r'+3}$. Then $\{Y_1, \dots, Y_{4r'+3}\}$ is an orthonormal basis in $T_p H_{2r'+1}^{4r'+3}$. By definition,

$$H_{2r'+1}^{4r'+3} = \{(x_0, x_1, \dots, x_{4r'+3}) \in \mathbf{R}^{4r'+4} \mid -x_0^2 - x_1^2 - \dots - x_{2r'+1}^2 + x_{2r'+2}^2 + \dots + x_{4r'+3}^2 = -1\}.$$

Let \tilde{H} be the semi-Riemannian submanifold in $H_{2r'+1}^{4r'+3}$ defined by

$$\tilde{H} = \{(x_0, 0, 0, \dots, 0, x_{2r'+2}, \dots, x_{4r'+3}) \in \mathbf{R}^{4r'+4} \mid -x_0^2 + x_{2r'+2}^2 + \dots + x_{4r'+3}^2 = -1, x_0 > 0\}.$$

It is easy to see that \tilde{H} is a complete totally geodesic submanifold in $H_{2r'+1}^{4r'+3}$ and that \tilde{H} is isometric to a hyperbolic space. Let $\{X_{2r'+2}, \dots, X_{4r'+3}\}$ be an orthonormal basis in $T_{\tilde{p}} \tilde{H}$, $\tilde{p} \in \tilde{H}$, and let $\{X_1, \dots, X_{2r'+1}\} \subset T_{\tilde{p}} H_{2r'+1}^{4r'+3}$ be an orthonormal basis of the normal bundle of the submanifold \tilde{H} . Since $H_{2r'+1}^{4r'+3}$ is a frame-homogeneous space (see [One2], or a strong isotropic manifold, cf. [Wol]), we have an isometry $\phi : H_{2r'+1}^{4r'+3} \rightarrow H_{2r'+1}^{4r'+3}$ such that $\phi(\tilde{p}) = p$ and $\phi(X_i) = Y_i$ for each $i \in \{1, \dots, 4r' + 3\}$. Hence $H = \phi(\tilde{H})$ is a complete, totally geodesic submanifold in $H_{2r'+1}^{4r'+3}$ and $T_p H = \text{span}\{Y_{2r'+2}, \dots, Y_{4r'+3}\}$. Therefore H has constant curvature -1 . By Lemma 14 on page 105 in [One2], one sees that H is a unique complete, totally geodesic semi-Riemannian submanifold such that $T_p H = \text{span}\{Y_{2r'+2}, \dots, Y_{4r'+3}\}$.

If $r' \geq 1$, then the base space B is simply connected by Proposition 3.8. Hence B is isometric to a pseudo-hyperbolic space. Let B' be a unique semi-Riemannian submanifold of the base space $B_{r'+1}^{2r'+2}$ such that B' is complete, totally geodesic in $B_{r'+1}^{2r'+2}$, and that $\pi(p) \in B'$ and $T_{\pi(p)} B' = \text{span}\{Y'_{3r'+3}, \dots, Y'_{4r'+3}\}$, constructed as above if $r' > 0$, or chosen to be the image of the spacelike geodesic in B passing through $\pi(p)$, with velocity in $\pi(p)$ equal to Y'_3 , if $r' = 0$. Then B' is isometric to $H^{r'+1}(-4)$.

First, we shall prove that $\pi(H) = B'$. Let $q' \in B'$. There is a unique geodesic τ' in B' joining $\pi(p)$ with q' , and satisfying $\tau'(0) = \pi(p)$, $\tau'(1) = q'$. Since B' is totally geodesic in B , τ' is a geodesic in B . Let τ be the horizontal lifting of τ' , with $p = \tau(0)$. Since $\dot{\tau}(0)$ is the horizontal lifting in p of $\dot{\tau}'(0) \in T_{\pi(p)} B' = \text{span}\{Y'_{3r'+3}, \dots, Y'_{4r'+3}\}$, we

obtain $\dot{\tau}(0) \in T_p H$. Since H was chosen to be totally geodesic in $H_{2r'+1}^{4r'+3}$, it follows that τ is contained in H . Thus $q' = \pi(\tau(1)) \in \pi(H)$. Therefore $B' \subset \pi(H)$.

Let $q \in H$. Since H is isometric to the real hyperbolic space $H^{2r'+2}$, there is a unique geodesic γ in H joining p with q . We denote by $X = h(\dot{\gamma}(0))$ and $V = v(\dot{\gamma}(0))$ the horizontal and vertical components of $\dot{\gamma}(0)$, respectively. If $X = 0$, then the geodesic γ is contained in the fibre $\pi^{-1}(\pi(p))$, since the fibres are totally geodesic. Hence $\pi(q) = \pi(p) \in B'$. If $V = 0$, then γ is a horizontal geodesic and hence $\pi \circ \gamma$ is a geodesic in B . Since B' is totally geodesic in B , the geodesic $\pi \circ \gamma$ is contained in B' . Therefore $\pi(q) \in B'$.

For $X \neq 0, V \neq 0$, we denote by γ_0 the geodesic given by the initial conditions $\gamma_0(0) = p$ and $\dot{\gamma}_0(0) = h(\dot{\gamma}(0)) \in T_p H$. Let H_2 be the unique complete totally geodesic submanifold in H passing through p with $T_p H_2 = \text{span}\{X, V\}$. Since H_2 is isometric to a real hyperbolic plane, we can choose a point $q_0 \in H_2$ on the geodesic γ_0 such that there exists a unique geodesic in H_2 , denoted by γ_1 , joining q_0 with q , and having the velocity vector at q_0 equal to the parallel translation along γ_0 of the vector wV for some constant $w \in \mathbf{R}$. Since H_2 is totally geodesic in H and H is totally geodesic in $H_{2r'+1}^{4r'+3}$, it follows that γ_1 is a geodesic in $H_{2r'+1}^{4r'+3}$. Since the fibres are totally geodesic, we obtain that γ_1 is contained in $\pi^{-1}(\pi(q))$. Thus $\pi(q_0) = \pi(q)$. Since γ_0 is a horizontal geodesic, $\pi \circ \gamma_0$ is a geodesic joining $\pi(p)$ with $\pi(q_0) = \pi(q)$, which has the initial velocity $\pi_* \dot{\gamma}_0(0) \in T_{\pi(p)} B'$. The geodesic $\pi \circ \gamma_0$ is contained in B' , because B' is totally geodesic. Hence $\pi(q) \in B'$. We proved that $\pi(H) = B'$.

Let $\tilde{\pi} : H \rightarrow B'$ be the restriction of π to H . It is easy to see that $\tilde{\pi}$ is a Riemannian submersion. We need to prove that the fibre $\tilde{\pi}^{-1}(\pi(p))$ is totally geodesic in p . Let γ be a geodesic in H such that $\gamma(0) = p$ and $\dot{\gamma}(0) \in T_p \tilde{\pi}^{-1}(\pi(p)) = \ker \tilde{\pi}_*|_p$. Then $0 = \tilde{\pi}_*(\dot{\gamma}(0)) = \pi_*(\dot{\gamma}(0))$. Hence $\dot{\gamma}(0)$ is vertical. Since the fibres of π are totally geodesic, γ is contained in $\pi^{-1}(\pi(p))$. Therefore $\gamma(t) \in \tilde{\pi}^{-1}(\pi(p))$ for every $t \in \mathbf{R}$. Since $\tilde{\pi} : H \rightarrow B'$ is a Riemannian submersion with the fibre $\tilde{\pi}^{-1}(\pi(p))$ totally geodesic in p , we get, by O'Neill's equations (see (9.29b) in [Bes]) applied at p , that

$$0 \leq g(A_X V, A_X V) = -g(X, X)g(V, V) \leq 0$$

for every $X \in \tilde{\mathcal{H}}_p, V \in \tilde{\mathcal{V}}_p$. So we get a contradiction. □

From Lemma 3.12, we obtain the following proposition.

PROPOSITION 3.13. *If $\pi : H_{s+r'}^{n+r'} \rightarrow B_s^n$ is a semi-Riemannian submersion with connected totally geodesic fibres from an $(n+r)$ -dimensional pseudo-hyperbolic space of index $s+r'$ onto an n -dimensional semi-Riemannian manifold of index s with constant curvature, then one of the following holds:*

- (1) $n = s = 2^t, r = r' = n - 1, B$ is isometric to $H_{2^t}^{2^t}(-4)$ and $t \in \{1, 2, 3\}$.
- (2) $n = 2^t, s = 0, r = r' = n - 1, B$ is isometric to $H^{2^t}(-4)$ and $t \in \{1, 2, 3\}$.

PROOF. If B has constant curvature, then the curvature of B is -4 and $n = r + 1$. By Lemma 3.12, we then get $\text{index}(B) \in \{0, \dim B\}$.

If $\text{index}(B) = \dim B$, then, by Lemma 3.7, we obtain $r = r'$. Hence, by [Ran1], we have (1).

If $\text{index}(B) = 0$, then, by [Ba-Ia], we have (2).

The idea of the proof in [Ran1] and [Ba-Ia] is to see that the tangent bundle of any fibre is trivial and that fibres are diffeomorphic to spheres, and then to apply a well-known result of Adams which claims that the spheres of dimensions 1, 3 and 7 are the only spheres with trivial tangent bundle. \square

The next theorems solve the equivalence problem of semi-Riemannian submersions from real and complex pseudo-hyperbolic spaces.

THEOREM 3.14. *If $\pi_1, \pi_2 : H_{s+r'}^{n+r} \rightarrow B_s^n$ are two semi-Riemannian submersions with connected totally geodesic fibres from a pseudo-hyperbolic space of index $s + r' > 1$ and the dimension of the fibres is $r \in \{1, 3\}$, then π_1 and π_2 are equivalent.*

PROOF. Let $p, q \in H_{s+r'}^{n+r}$. Let

$$\mathcal{L} = \{L_0, A_{1L_0}v_1, \dots, A_{1L_0}v_r, \dots, L_{k-1}, A_{1L_{k-1}}v_1, \dots, A_{1L_{k-1}}v_r\},$$

$$\mathcal{L}' = \{L'_0, A_{2L'_0}v'_1, \dots, A_{2L'_0}v'_r, \dots, L'_{k-1}, A_{2L'_{k-1}}v'_1, \dots, A_{2L'_{k-1}}v'_r\}$$

be two orthonormal bases of \mathcal{H}_1 along $\pi_1^{-1}(\pi_1(p))$ and of \mathcal{H}_2 along $\pi_2^{-1}(\pi_2(q))$ constructed as in the proof of Theorem 3.3 such that $g_p(L_\alpha, L_\beta) = g_q(L'_\alpha, L'_\beta) = \varepsilon_\alpha \delta_{\alpha\beta}$ for $\alpha, \beta \in \{0, \dots, k-1\}$, $g_p(v_i, v_j) = g_q(v'_i, v'_j) = \varepsilon_i \delta_{ij}$ for $i, j \in \{1, \dots, r\}$ and for $r = 3$, $v_{3p} = (\hat{\nabla}_{v_1} v_2)(p)$ and $v'_{3q} = (\hat{\nabla}_{v'_1} v'_2)(q)$.

Let $\phi : T_p H_{s+r'}^{n+r} \rightarrow T_q H_{s+r'}^{n+r}$ be the linear map given by $\phi(L_\alpha) = L'_\alpha$, $\phi(A_{1L_\alpha} v_i) = A_{2L'_\alpha} v'_i$, $\phi(v_i) = v'_i$ for every α and i . In a manner similar to the proof of Theorem 3.3, we obtain $\phi(A_{1E} F) = A_{2\phi(E)} \phi(F)$ for every $E, F \in T_p H_{s+r'}^{n+r}$. By Corollary 2.3.14 in [Wol], ϕ extends to an isometry on $H_{s+r'}^{n+r}$, denoted by $f : H_{s+r'}^{n+r} \rightarrow H_{s+r'}^{n+r}$, satisfying $f(p) = q$ and $f_{*p} = \phi$. From Theorem 2.4 it follows that f induces an isometry \tilde{f} on B , such that $\tilde{f} \circ \pi = \pi \circ f$. Hence π_1 and π_2 are equivalent. \square

THEOREM 3.15. *If $\pi_1, \pi_2 : CH_{2s+1}^{2n+1} \rightarrow HH_s^n$ are two semi-Riemannian submersions with connected complex totally geodesic fibres from a complex pseudo-hyperbolic space, then π_1 and π_2 are equivalent.*

PROOF. Let $\theta : H_{4s+3}^{4n+3} \rightarrow CH_{2s+1}^{2n+1}$ be the canonical semi-Riemannian submersion. By Theorem 2.5 in [Esc2], we see that $\tilde{\pi}_1 = \pi_1 \circ \theta : H_{4s+3}^{4n+3} \rightarrow HH_s^n$ and $\tilde{\pi}_2 = \pi_2 \circ \theta : H_{4s+3}^{4n+3} \rightarrow HH_s^n$ are semi-Riemannian submersions with totally geodesic fibres. We denote by $\tilde{A}_1, \tilde{A}_2, A_1, A_2, A$ O'Neill's integrability tensors of $\tilde{\pi}_1, \tilde{\pi}_2, \pi_1, \pi_2, \theta$, respectively. In order to reduce the proof of the equivalence theorem of semi-Riemannian submersions from a complex pseudo-hyperbolic space to that from a pseudo-hyperbolic space, we need to establish relations among the integrability tensors \tilde{A}_1, A_1, A .

First, we prove that $\theta_* \tilde{A}_1 X Y = A_{1\theta_* X} \theta_* Y$ for $\tilde{\pi}_1$ -basic vector fields X and Y . Let $p \in H_{4s+3}^{4n+3}$. Let w'_1, w'_2 be two orthonormal π_1 -vertical vectors in $T_{\theta(p)} CH_{2s+1}^{2n+1}$ and let $w_1,$

w_2 be the θ -horizontal liftings at p of w'_1, w'_2 , respectively. Let w_3 be a unit θ -vertical vector in $T_p H_{4s+3}^{4n+3}$. Then $\{w_1, w_2, w_3\}$ gives an orthonormal basis of $\tilde{\mathcal{V}}_{1p}$. Since the induced metrics on the fibres of $\tilde{\pi}_1$ are negative definite, we have

$$\tilde{A}_{1X}Y = -g(\nabla_X Y, w_1)w_1 - g(\nabla_X Y, w_2)w_2 - g(\nabla_X Y, w_3)w_3.$$

Thus

$$\theta_* \tilde{A}_{1X}Y = -g'(\nabla'_{\theta_* X} \theta_* Y, \theta_* w_1) \theta_* w_1 - g'(\nabla'_{\theta_* X} \theta_* Y, \theta_* w_2) \theta_* w_2 = A_{1\theta_* X} \theta_* Y$$

for $\tilde{\pi}_1$ -basic vector fields X and Y , where g' denotes the metric on CH_{2s+1}^{2n+1} and ∇' is the Levi-Civita connection of g' .

Let X be the $\tilde{\pi}_1$ -horizontal lifting along the fibre $\tilde{\pi}_1^{-1}(\tilde{\pi}_1(p))$ of some unit vector in $T_{\tilde{\pi}_1(p)} \mathbf{H}H_s^n$. Let Y_1, Y_2, Y_3 be the $\tilde{\pi}_1$ -horizontal liftings along the fibre $\tilde{\pi}_1^{-1}(\tilde{\pi}_1(p))$ of $\tilde{\pi}_{1*} \tilde{A}_{1X} w_1, \tilde{\pi}_{1*} \tilde{A}_{1X} w_2, \tilde{\pi}_{1*} \tilde{A}_{1X} w_3$, respectively. Let $v_i = \tilde{A}_{1X} Y_i$ for $i \in \{1, 2, 3\}$. As in Theorem 3.3, we choose $w_3 = g(X, X)^{-1} (\nabla_{v_1} v_2)(p)$, which implies that $v_3 = \nabla_{v_1} v_2$ (see Lemma 3.6).

We remark that $v_3 = \tilde{A}_{1X} Y_3$ is a θ -vertical vector field along the fibre $\theta^{-1}(\theta(p))$. Indeed, we have

$$\begin{aligned} \theta_*(\tilde{A}_{1X} Y_3(p')) &= (A_{1\theta_* X} \theta_* Y_3)(\theta(p')) = (A_{1\theta_* X} \theta_* Y_3)(\theta(p)) = \theta_*(\tilde{A}_{1X} Y_3(p)) \\ &= \theta_*(A_{1X} A_{1X} w_3) = g(X, X) \theta_* w_3 = 0 \end{aligned}$$

for any $p' \in \theta^{-1}(\theta(p))$.

Since v_1, v_2 are orthogonal to the vertical vector field v_3 along $\theta^{-1}(\theta(p))$, we see that v_1, v_2 are θ -horizontal. Since $\theta_*(\tilde{A}_{1X} Y_i(p')) = (A_{1\theta_* X} \theta_* Y_i)(\theta(p'))$ for $p' \in \theta^{-1}(\theta(p))$ and for $i \in \{1, 2\}$, we obtain that v_1, v_2 are θ -basic vector fields along $\theta^{-1}(\theta(p))$. Thus $h\nabla_{v_3} v_1 = A_{v_1} v_3$ along $\theta^{-1}(\theta(p))$. Here h and v denote the θ -horizontal and θ -vertical projections, respectively. We also obtain that $v\nabla_{v_3} v_1 = -g(\nabla_{v_3} v_1, v_3)v_3 = 0$. Therefore, $A_{v_1} v_3 = \nabla_{v_3} v_1 = v_2$ along $\theta^{-1}(\theta(p))$.

We shall prove that $\tilde{A}_{1X} v_3 = A_X v_3$ along $\theta^{-1}(\theta(p))$ for every $\tilde{\pi}_1$ -basic vector field X along $\tilde{\pi}_1^{-1}(\tilde{\pi}_1(p))$. We first obtain along $\theta^{-1}(\theta(p))$ that

$$\begin{aligned} \tilde{A}_{1X} v_3 &= \nabla_X v_3 + g(\nabla_X v_3, v_1)v_1 + g(\nabla_X v_3, v_2)v_2 + g(\nabla_X v_3, v_3)v_3, \\ g(\nabla_X v_3, v_1) &= g(A_X v_3, v_1) = -g(v_3, A_X v_1) = g(v_3, A_{v_1} X) = -g(A_{v_1} v_3, X) \\ &= -g(v_2, X) = 0 \end{aligned}$$

for a $\tilde{\pi}_1$ -basic vector field X along $\tilde{\pi}_1^{-1}(\tilde{\pi}_1(p))$. Analogously, we get $g(\nabla_X v_3, v_2) = 0$. Thus

$$\tilde{A}_{1X} v_3 = \nabla_X v_3 + g(\nabla_X v_3, v_3)v_3 = A_X v_3$$

along $\theta^{-1}(\theta(p))$ for every $\tilde{\pi}_1$ -basic vector field X along $\tilde{\pi}_1^{-1}(\tilde{\pi}_1(p))$.

Let $\tilde{\mathcal{L}} = \{L_0 = X, \tilde{A}_{1L_0} v_1, \tilde{A}_{1L_0} v_2, \tilde{A}_{1L_0} v_3, \dots, L_{n-1}, \tilde{A}_{1L_{n-1}} v_1, \tilde{A}_{1L_{n-1}} v_2, \tilde{A}_{1L_{n-1}} v_3\}$ be an orthonormal basis of $\tilde{\mathcal{H}}_1$ along the fibre $\tilde{\pi}_1^{-1}(\tilde{\pi}_1(p))$ constructed as in Theorem 3.3, for the semi-Riemannian submersion $\tilde{\pi}_1$. From the proof of Theorem 3.3, we have

$$g(\tilde{A}_{1\tilde{A}_{1L_j} v_1} v_3, \tilde{A}_{1L_l} v_2) = 0$$

for $j \neq l$, and

$$g(\tilde{A}_{1\tilde{A}_{1L_j}v_1}v_3, L_t) = 0$$

for $0 \leq j, t \leq n-1$. We then obtain along $\tilde{\pi}_1^{-1}(\tilde{\pi}_1(p))$ that

$$\begin{aligned} g(\tilde{A}_{1\tilde{A}_{1L_j}v_1}v_3, \tilde{A}_{1L_j}v_2) &= -g(v_3, \tilde{A}_{1\tilde{A}_{1L_j}v_1}\tilde{A}_{1L_j}v_2) \\ &= -g(v_3, \nabla_{v_1}v_2)g(L_j, L_j) \\ &= -g(v_3, v_3)g(L_j, L_j) = -g(v_2, v_2)g(L_j, L_j) \\ &= g(\tilde{A}_{1L_j}v_2, \tilde{A}_{1L_j}v_2), \end{aligned}$$

from which follows $\tilde{A}_{1L_j}v_2 = \tilde{A}_{1\tilde{A}_{1L_j}v_1}v_3$. Hence $\tilde{A}_{1L_j}v_2 = A_{\tilde{A}_{1L_j}v_1}v_3$, because $\tilde{A}_{1L_j}v_1$ is $\tilde{\pi}_1$ -basic. We also have $\tilde{A}_{1L_j}v_3 = A_{L_j}v_3$.

Let $\mathcal{L} = \tilde{\mathcal{L}} \cup \{v_1, v_2\}$. Summarizing all the above, we obtain that

$$\mathcal{L} = \{L_0, A_{L_0}v_3, \tilde{A}_{1L_0}v_1, A_{\tilde{A}_{1L_0}v_1}v_3, \dots, L_{n-1}, A_{L_{n-1}}v_3, \tilde{A}_{1L_{n-1}}v_1, A_{\tilde{A}_{1L_{n-1}}v_1}v_3, v_1, A_{v_1}v_3\}$$

is an orthonormal basis of the θ -horizontal space \mathcal{H} along the fibre $\theta^{-1}(\theta(p))$ and \mathcal{L} satisfies all conditions imposed in the construction of the basis \mathcal{L} in the proof of Theorem 3.3. We notice that $v_3 = A_X Y_3$ along $\theta^{-1}(\theta(p))$, and that along $\theta^{-1}(\theta(p))$, Y_3 is equal to the θ -horizontal lifting of $\theta_* A_X w_3$.

Let $q \in H_{4s+3}^{4n+3}$. Let

$$\tilde{\mathcal{L}}' = \{L'_0, \tilde{A}_{2L'_0}v'_1, \tilde{A}_{2L'_0}v'_2, \tilde{A}_{2L'_0}v'_3, \dots, L'_{n-1}, \tilde{A}_{2L'_{n-1}}v'_1, \tilde{A}_{2L'_{n-1}}v'_2, \tilde{A}_{2L'_{n-1}}v'_3\}$$

be an orthonormal basis of $\tilde{\mathcal{H}}_2$ along $\tilde{\pi}_2^{-1}(\tilde{\pi}_2(q))$ constructed in the same way as $\tilde{\mathcal{L}}$, but for the semi-Riemannian submersion $\tilde{\pi}_2$ (see the proof of Theorem 3.3), in such a way that $g_p(L_\alpha, L_\beta) = g_q(L'_\alpha, L'_\beta)$ for $0 \leq \alpha, \beta \leq n-1$, $g_p(v_i, v_j) = g_q(v'_i, v'_j)$ for $1 \leq i, j \leq 3$, and $v'_3(q) = (\nabla_{v'_1}v'_2)(q)$. Let $\phi : T_p H_{4s+3}^{4n+3} \rightarrow T_q H_{4s+3}^{4n+3}$ be the linear map given by $\phi(v_i) = v'_i$, $\phi(\tilde{A}_{1L_\alpha}v_i) = \tilde{A}_{2L'_\alpha}v'_i$ for $0 \leq \alpha \leq n-1$ and for $1 \leq i \leq 3$.

By Corollary 2.3.14 in [Wol], ϕ extends to an isometry $f : H_{4s+3}^{4n+3} \rightarrow H_{4s+3}^{4n+3}$ such that $f(p) = q$ and $f_*p = \phi$. By the proof of Theorem 3.3, we have $f_*\tilde{A}_{1E}F = \tilde{A}_{2f_*E}f_*F$ for every $E, F \in T_p H_{4s+3}^{4n+3}$. By the proof of Theorem 3.14 and by Theorem 2.4, f induces an isometry on $\mathcal{C}H_{2s+1}^{2n+1}$, denoted by $\tilde{f} : \mathcal{C}H_{2s+1}^{2n+1} \rightarrow \mathcal{C}H_{2s+1}^{2n+1}$, such that $\theta \circ f = \tilde{f} \circ \theta$. Since the π_1 -vertical space at $\theta(p)$ is spanned by $\{\theta_*v_1, \theta_*v_2\}$, since the π_2 -vertical space at $\theta(q)$ is spanned by $\{\theta_*v'_1, \theta_*v'_2\}$, and since $\tilde{f}_*(\theta_*v_i) = \theta_*v'_i$, for $i \in \{1, 2\}$, we see that \tilde{f}_* maps the π_1 -vertical space at $\theta(p)$ into the π_2 -vertical space at $\theta(q)$. For $\tilde{\pi}_1$ -horizontal vectors X and Y we obtain

$$\begin{aligned} \tilde{f}_*A_{1\theta_*X}\theta_*Y &= \tilde{f}_*\theta_*\tilde{A}_{1X}Y = \theta_*f_*\tilde{A}_{1X}Y \\ &= \theta_*\tilde{A}_{2f_*X}f_*Y = A_{2\theta_*f_*X}\theta_*f_*Y \\ &= A_{2\tilde{f}_*(\theta_*X)}\tilde{f}_*(\theta_*Y). \end{aligned}$$

Therefore, by Theorem 2.4, we see that π_1 and π_2 are equivalent. \square

REMARK. We notice that our equivalence theorems can be applied, in particular, to Riemannian submersions from a sphere with totally geodesic fibres of dimension less than or equal to 3, and for Riemannian submersions with complex totally geodesic fibres from a complex projective space. Unlike those in [Esc1], [Esc2], [Ran1], our proofs of the equivalence theorems are intrinsic, we do not need to assume the existence of any specific structure on the base space, such as complex or quaternionic one. In Theorem 3.15, we need to assume only that the fibres are 2-dimensional and that the induced metrics on fibres are negative definite.

Summarizing all results above, we now prove the main theorems.

PROOF OF THEOREM 1.1. If $s + r' > 1$, then $H_{s+r'}^{m+r}$ is simply connected and hence, by Theorem 3.3, B is an isotropic semi-Riemannian manifold and $r \in \{1, 3\}$. By Propositions 3.11 and 3.13, we see that the base space of the semi-Riemannian submersion is isometric to a complex pseudo-hyperbolic space if the dimension of fibres is one, or to a quaternionic pseudo-hyperbolic space if the dimension of fibres is 3. In Theorem 3.14 we solved the equivalence problem. The existence problem is solved by the explicit construction given in the preliminaries (see Examples 1 and 2).

If $s + r' = 1$, then either (i) $s = 1, r' = 0$, or (ii) $s = 0, r' = 1$.

(i) If $s = 1, r' = 0$, then, by the proof of Proposition 3.11, B has constant curvature. By Theorem 3.3, (1) we get $n = k(r + 1) \geq 2$, since $r \geq 1$. Hence, by Lemma 3.12, there are no such semi-Riemannian submersions.

(ii) If $s = 0, r' = 1$, then π is a semi-Riemannian submersion from an anti-de Sitter space onto a Riemannian manifold. By [Mag], π is equivalent to the canonical submersion $\pi : H_1^{2m+1} \rightarrow CH^m$. This falls in the case (a). \square

PROOF OF THEOREM 1.2. If the dimension of the fibres is less than or equal to 3, then, by Theorem 1.1, π is equivalent to the canonical semi-Riemannian submersions:

- (a) $H_{2t+1}^{2m+1} \rightarrow CH_t^m, 0 \leq t \leq m$, or
- (b) $H_{4t+3}^{4m+3} \rightarrow HH_t^m, 0 \leq t \leq m$.

Now we assume that the dimension of the fibres is greater than or equal to 4.

(A) If we assume that the dimension of the fibres is greater than or equal to 4 and B is an isotropic semi-Riemannian manifold with non-constant curvature, then, by Proposition 3.11, B is isometric to $CaH_t^2, t \in \{0, 1, 2\}$, and the dimension of the fibres is $r = r' = 7$. By Proposition 2.7, there are no such semi-Riemannian submersions with base space CaH_t^2 . Therefore, the assumptions (A) and $r \geq 4$ imply that B has constant curvature, and hence, by Lemma 3.12, we obtain $s = \text{index}(B) \in \{0, \dim(B)\}$.

(B) If $\text{index}(B) = 0$ and $r \geq 4$, then, by [Ba-Ia], the semi-Riemannian submersion π is equivalent to the canonical semi-Riemannian submersion $H_7^{15} \rightarrow H^8(-4)$. If $\text{index}(B) = \dim(B)$, then, by Lemma 3.7, we get $r' = r$. By changing the signs of the metrics on the base and on the total space, π becomes a Riemannian submersion with connected totally geodesic fibres from a sphere onto a Riemannian manifold. So, by [Esc1] and [Ran1], one obtains the conclusion. \square

PROOF OF THEOREM 1.3. Let $\theta : H_{2s+1}^{2n+1} \rightarrow CH_s^n$ be the canonical semi-Riemannian submersion. By Theorem 2.5 in [Esc2], one obtains that $\pi \circ \theta : H_{2s+1}^{2n+1} \rightarrow B$ is a semi-Riemannian submersion with connected totally geodesic fibres.

(A) If the dimension of the fibres of π is r and $1 \leq r \leq 2$, then the dimension of the fibres of the semi-Riemannian submersion $\pi \circ \theta$ is less than or equal to 3 and greater than or equal to 2. By Theorem 1.1, B is isometric to HH_t^m and $2n+1 = 4m+3$, $2s+1 = 4t+3$. Then $n = 2m+1$, $s = 2t+1$. By Theorem 3.15, we see that $\pi : CH_{2t+1}^{2m+1} \rightarrow HH_t^m$ is equivalent to the canonical semi-Riemannian submersion.

(B) and (C) If B is an isotropic semi-Riemannian manifold or if $\text{index}(B) \in \{0, \dim B\}$, then, by Theorem 1.2, $\pi \circ \theta$ is equivalent to one of the following canonical semi-Riemannian submersions:

$$\begin{aligned} H_{2t+1}^{2m+1} &\rightarrow CH_t^m, & 0 \leq t \leq m; \\ H_{4t+3}^{4m+3} &\rightarrow HH_t^m, & 0 \leq t \leq m; \\ H_{7+8t}^{15} &\rightarrow H_{8t}^8(-4), & t \in \{0, 1\}. \end{aligned}$$

If the dimension of the fibres of π is greater than or equal to 3, then the dimension of the fibres of $\pi \circ \theta$ is greater than or equal to 4. Hence, in this case, $\pi \circ \theta$ is equivalent to $H_{7+8t}^{15} \rightarrow H_{8t}^8(-4)$, $t \in \{0, 1\}$. For $t = 1$, the semi-Riemannian submersion π is, after a change of signs of the metrics on the total space and on the base space, of type $\pi : CP^7 \rightarrow S^8(4)$. For $t = 0$, π is of type $\pi : CH_3^7 \rightarrow H^8(-4)$. In [Ran1] (for case $t = 1$) and [Ba-Ia] (for case $t = 0$), it is proved that there are no such semi-Riemannian submersions with totally geodesic fibres. We proved that the dimension of fibres of π is less than or equal to 2. \square

PROOF OF THEOREM 1.4. We suppose that there are such semi-Riemannian submersions. It is well-known that any quaternionic submanifold in HH_s^n is totally geodesic. Let $\eta : H_{4s+3}^{4n+3} \rightarrow HH_s^n$, $\xi : CH_{2s+1}^{2n+1} \rightarrow HH_s^n$, be the canonical semi-Riemannian submersions. By Theorem 2.5 in [Esc2], we see that $\pi \circ \eta : H_{4s+3}^{4n+3} \rightarrow B$ is a semi-Riemannian submersion with connected totally geodesic fibres. We remark that the dimension of the fibres of $\pi \circ \eta$ is greater than or equal to 4. Thus, by Theorem 1.2, we see that $\pi \circ \eta$ is equivalent to the canonical semi-Riemannian submersion

$$H_7^{15} \rightarrow H^8(-4), \quad \text{or} \quad H_{15}^{15} \rightarrow H_8^8(-4).$$

It follows that π is one of the following types:

- (i) $\pi : HH_1^3 \rightarrow H^8(-4)$, or
- (ii) $\pi : HH_3^3 \rightarrow H_8^8(-4)$.

In [Ucc], Ucci proved that there are no Riemannian submersions with fibres HP^1 from HP^3 onto $S^8(4)$. Therefore, Case (ii) is not possible.

The fibres of semi-Riemannian submersion $\pi \circ \xi : CH_3^7 \rightarrow H^8(-4)$ are totally geodesic by Theorem 2.5 in [Esc2], and complex submanifolds, since the horizontal lifting of

the tangent space of the quaternionic line $\pi^{-1}(\pi(p))$ is invariant under the canonical complex structure on CH_3^7 . By [Ba-Ia], there are no semi-Riemannian submersions with complex totally geodesic fibres from CH_3^7 onto $H^8(-4)$. Thus Case (i) is impossible. \square

REFERENCES

- [Ba-Ia] G. BĂDIȚOIU AND S. IANUȘ, Semi-Riemannian submersions from real and complex pseudo-hyperbolic spaces, *Differential Geom. Appl.* 16 (2002), 79–94.
- [Ba-Ro] M. BARROS AND A. ROMERO, Indefinite Kähler manifolds, *Math. Ann.* 261 (1982), 55–62.
- [Bes] A. L. BESSE, *Einstein manifolds*, Springer-Verlag, Berlin, 1987.
- [Bis] R. L. BISHOP, Clairaut submersions, *Differential geometry (in honor of K. Yano)*, 21–31, Kinokuniya, Tokyo, 1972.
- [Ehr] C. EHRESMANN, Les connexions infinitésimales dans un espace fibré différentiable, *Colloque de Topologie, Bruxelles 1950*, p. 29–55, Georges Thone, Liège; Masson et Cie., Paris, 1951.
- [Esc1] R. ESCOBALES, Riemannian submersions with totally geodesic fibers, *J. Differential Geom.* 10 (1975), 253–276.
- [Esc2] R. ESCOBALES, Riemannian submersions from complex projective spaces, *J. Differential Geom.* 13 (1978), 93–107.
- [Gra] A. GRAY, Pseudo-Riemannian almost product manifolds and submersions, *J. Math. Mech.* 16 (1967), 715–737.
- [G-G1] D. GROMOLL AND K. GROVE, The low-dimensional metric foliations of Euclidean spheres, *J. Differential Geom.* 28 (1988), 143–156.
- [G-G2] D. GROMOLL AND K. GROVE, A generalization of Berger's rigidity theorem for positively curved manifolds, *Ann. Sci. École Norm. Sup. (4)* 20 (1987), 227–239.
- [Ian] S. IANUȘ, *Differential geometry with applications to the theory of relativity*, (in romanian) Ed. Academiei Române, București, 1983.
- [Mag] M. A. MAGID, Submersions from anti-de Sitter space with totally geodesic fibres, *J. Differential Geom.* 16 (1981), 323–331.
- [One1] B. O'NEILL, The fundamental equations of a submersion, *Michigan Math. J.* 13 (1966), 459–469.
- [One2] B. O'NEILL, *Semi-Riemannian geometry with applications to relativity*, Academic Press, New York, London, 1983.
- [Ran1] A. RANJAN, Riemannian submersions of spheres with totally geodesic fibres, *Osaka J. Math.* 22 (1985), 243–260.
- [Ran2] A. RANJAN, Riemannian submersions of compact simple Lie groups with connected totally geodesic fibres, *Math. Z.* 191 (1986), 239–246.
- [Rec] H. RECKZIEGEL, A fibre bundle theorem, *Manuscripta Math.* 76 (1992), 105–110.
- [Ucc] J. UCCI, On the nonexistence of Riemannian submersions from $CP(7)$ and $HP(3)$, *Proc. Amer. Math. Soc.* 88 (1983), 698–700.
- [Wil] F. WILHELM, The radius rigidity theorem for manifolds of positive curvature, *J. Differential Geom.* 44 (1996), 634–665.
- [Wilk] B. WILKING, Index parity of closed geodesics and rigidity of Hopf fibrations, *Invent. Math.* 144 (2001), 281–295.
- [Wol] J. WOLF, *Spaces of constant curvature*, McGraw-Hill Inc., New York, 1967.

INSTITUTE OF MATHEMATICS OF THE ROMANIAN ACADEMY
P.O. BOX 1–764
BUCHAREST 70700
ROMANIA