

Pacific Journal of Mathematics

SEMIGROUPS SATISFYING IDENTITY $xy = f(x, y)$

TAKAYUKI TAMURA

SEMIGROUPS SATISFYING IDENTITY $xy = f(x, y)$

TAKAYUKI TAMURA

Dedicated to Professor Keizo Asano on his Sixtieth Birthday

Let $f(x, y)$ be a word of length greater than 2 starting in y and ending in x . The purpose of this paper is to prove that a semigroup satisfies an identity $xy = f(x, y)$ if and only if it is an inflation of a semilattice of groups satisfying the same identity. As its consequence we find a necessary and sufficient condition for $xy = f(x, y)$ to imply commutativity.

Recently E. J. Tully has proved [7] that if a semigroup S satisfies an identity of the form $xy = y^m x^n$ then S is an inflation of a semilattice of abelian groups G_α 's satisfying $x^k = 1$ for all $x \in G_\alpha$ where k is the greatest common divisor of $m - 1$ and $n - 1$; hence $xy = y^m x^n$ implies commutativity. This paper is to consider the general case of the right side of $xy = y^m x^n$ with the left side unchanged.

Let $f(x, y)$ denote a word involving both x and y , and $|f(x, y)|$ be the length of the word $f(x, y)$; $|x|_f$ be the number of x 's which appear in $f(x, y)$; $|y|_f$ be also defined for y . For example if $f(x, y) = x^3 y^2 x y$, $|f(x, y)| = 7$, $|x|_f = 4$, $|y|_f = 3$. Throughout this paper we assume $|f(x, y)| > 2$, equivalently $|x|_f > 1$ or $|y|_f > 1$ or both.

Consider an identity of the form

$$(1) \quad xy = f(x, y)$$

in semigroups. A question is raised: Under what condition on $f(x, y)$ does (1) imply commutativity $xy = yx$? What we can say immediately is that $f(x, y)$ has to start in y . Because if $f(x, y)$ starts in x , then left zero semigroups of order > 1 satisfy (1) but are not commutative. For the similar reason $f(x, y)$ must end in x . From now on we assume $f(x, y)$ in (1) has the form:

$$(2) \quad \begin{cases} f(x, y) = y^{m_1} x^{n_1} \cdots y^{m_s} x^{n_s}, m_i > 0, n_i > 0, i = 1, \dots, s, \\ \text{and } |f(x, y)| > 2. \end{cases}$$

A semigroup D is called an inflation of a semigroup T if T is a subsemigroup of D and there is a mapping φ of D into T such that

$$\varphi(x) = x \quad \text{for } x \in T$$

and

$$xy = \varphi(x)\varphi(y) \quad \text{for } x, y \in D.$$

Let L be a semilattice. A semigroup S is called a semilattice L of semigroups S_α , $\alpha \in L$, if S is a disjoint union of $\{S_\alpha; \alpha \in L\}$ and

$$S_\alpha S_\beta \subseteq S_{\alpha\beta} \quad \text{for all } \alpha, \beta \in L.$$

Needless to say an identity is preserved by homomorphic images and by subsemigroups; in particular the identity $xy = f(x, y)$ with (2) is satisfied by any semilattice.

THEOREM. *A semigroup S satisfies an identity $xy = f(x, y)$ with (2) if and only if S is an inflation of a semilattice of groups satisfying the same identity.*

COROLLARY. *Let $f(x, y)$ be a word involving both x and y , and let $|f(x, y)| > 2$. $xy = f(x, y)$ implies $xy = yx$ in semigroups if and only if*

- (3.1) $f(x, y)$ starts in y and ends in x , and
- (3.2) $xy = f(x, y)$ implies $xy = yx$ in groups.

The statement in the theorem can be replaced by another statement. In the following proposition we do not assume $xy = f(x, y)$.

PROPOSITION. *The following three statements are equivalent:*

- (4.1) S^2 is a semilattice L of groups $G_\alpha, \alpha \in L$.
- (4.2) S is a semilattice L of semigroups $S_\alpha, \alpha \in L$, each of which is an inflation of a group G_α and

$$x_\alpha y_\beta \in G_{\alpha\beta} \quad \text{for } x_\alpha \in S_\alpha, y_\beta \in S_\beta.$$

- (4.3) S is an inflation of a semilattice L of groups $G_\alpha, \alpha \in L$.

Let a_1, \dots, a_k be a finite number of elements of a semigroup S . All the elements x of S each of which is the product of all of a_1, \dots, a_k (admitting repeated use) form a subsemigroup of S . It is called the content (of a_1, \dots, a_k) in S and denoted by $C(a_1, \dots, a_k)$ or C . The elements a_1, \dots, a_k are not required to be distinct. For example $C(a, a) = \{a^i; i > 1\}$ but $C(a) = \{a^i; i \geq 1\}$. The number k is called the rank of $C(a_1, \dots, a_k)$.

Any semigroup S has a smallest semilattice-congruence (\mathcal{S} -congruence) ρ_0 , that is, S/ρ_0 is a semilattice and if S/ρ is a semilattice, then $\rho_0 \subseteq \rho$. The decomposition of S induced by ρ_0 is called the greatest \mathcal{S} -decomposition of S . If $\rho_0 = S \times S$, S is called \mathcal{S} -indecomposable. An \mathcal{S} -decomposition $S = \bigcup_{\alpha \in L} S_\alpha$ of S is greatest if and only if each S_α is \mathcal{S} -indecomposable [4], [5], [6].

LEMMA 1. *A content is \mathcal{S} -indecomposable. (See [6].)*

LEMMA 2. *$a\rho_0 b$ if and only if there is a finite sequence of contents*

C_1, \dots, C_t such that

$$(5) \quad a \in C_1, \quad C_i \cap C_{i+1} \neq \emptyset \quad (i = 1, \dots, t - 1), \quad b \in C_t.$$

Proof. Define ρ_1 as follows: $a\rho_1b$ if and only if there is a finite sequence of contents C_1, \dots, C_t satisfying (5). We will prove $\rho_0 = \rho_1$. It is easily shown that ρ_1 is an equivalence relation. To prove compatibility, suppose $a\rho_1b$. There exists a sequence of contents satisfying (5), more specifically

$$(5') \quad \begin{cases} C_i = C(a_{i1}, \dots, a_{ik_i}) & i = 1, \dots, t \\ a \in C_1, \quad d_i \in C_i \cap C_{i+1} & (i = 1, \dots, t - 1), \quad b \in C_t. \end{cases}$$

Let $c \in S$. Consider a sequence of contents

$$C'_i = C(c, a_{i1}, \dots, a_{ik_i}) \quad (i = 1, \dots, t - 1).$$

Then $ca \in C'_1, cd_i \in C'_i \cap C'_{i+1} (i = 1, \dots, t - 1), cb \in C'_t$. Hence $ca\rho_1cb$. Likewise $ac\rho_1bc$. Thus ρ_1 is a congruence. Since $a, a^2 \in C(a)$ and $ab, ba \in C(a, b)$, we see $a\rho_1a^2, ab\rho_1ba$ for all $a, b \in S$, that is, ρ_1 is an \mathcal{S} -congruence on S . Let ρ be an \mathcal{S} -congruence on S . We will prove $\rho_1 \subseteq \rho$. Let $a\rho_1b$. There is a sequence C_1, \dots, C_t described in (5'). Since C_i is \mathcal{S} -indecomposable ($i = 1, \dots, t$) by Lemma 1, we have $a\rho d_1, d_1\rho d_2, \dots, d_{t-1}\rho b$, hence $a\rho b$. Since ρ_1 is the smallest \mathcal{S} -congruence, we have $\rho_0 = \rho_1$.

Let

$$S = \bigcup_{\alpha \in L} S_\alpha, \quad S_\alpha S_\beta \subseteq S_{\alpha\beta}, \quad S_\alpha \cap S_\beta = \emptyset, \quad \alpha \neq \beta$$

be the greatest \mathcal{S} -decomposition of S . We notice that if a and b are in S_α then the contents $C_i (i = 1, \dots, t)$ described in (5) are contained in S_α .

LEMMA 3. *An ideal of an \mathcal{S} -indecomposable semigroup is \mathcal{S} -indecomposable.*

Proof. An equivalent statement is proved in [4]. However, we will prove this by using Lemma 2. Let I be an ideal of an \mathcal{S} -indecomposable semigroup S . Let $a, b \in I, a \neq b$. By Lemma 2, there is a sequence of contents $C_i = C(a_{i1}, \dots, a_{ik_i}) \subseteq S (i = 1, \dots, t)$ such that $a \in C_1, d_i \in C_i \cap C_{i+1} (i = 1, \dots, t - 1), b \in C_t$. Consider a sequence of contents:

$$\begin{aligned} C'_0 &= C(a), \quad C'_i = C(a, a_{i1}, \dots, a_{ik_i}) & (i = 1, \dots, t) \\ C'_{2t+1} &= C(b), \quad C'_{t+i} = C(b, a_{i1}, \dots, a_{ik_i}) & (i = 1, \dots, t). \end{aligned}$$

Then

$$\begin{aligned} a &\in C'_0, \quad a^2 \in C'_0 \cap C'_1, \quad ad_i \in C'_i \cap C'_{i+1} \quad (i = 1, \dots, t-1) \\ ab &\in C'_i \cap C'_{i+1}, \quad d_i b \in C'_{i+1} \cap C'_{i+2} \quad (i = 1, \dots, t-1) \\ b^2 &\in C'_{2t} \cap C'_{2t+1}, \quad b \in C'_{2t+1} \end{aligned}$$

and all $C'_j (j = 0, \dots, 2t+1)$ are in I . By Lemma 2, we have proved that I is \mathcal{S} -indecomposable.

Proof of proposition. The proof of $(4.2) \Rightarrow (4.1)$ and $(4.3) \Rightarrow (4.1)$ is immediate. We will prove only $(4.1) \Rightarrow (4.2)$ and (4.3) . Assume $S^2 = \bigcup_{\alpha \in L} G_\alpha$ where G_α is a group for each α . This is the greatest \mathcal{S} -decomposition of S^2 because the groups G_α are \mathcal{S} -indecomposable. Let

$$(6) \quad S = \bigcup_{\xi \in L'} S_\xi$$

be the greatest \mathcal{S} -decomposition of S . Since S_ξ^2 is an ideal of the \mathcal{S} -indecomposable semigroup S_ξ , S_ξ^2 is also \mathcal{S} -indecomposable by Lemma 3. For each $\alpha \in L$ there is a unique α' in L' such that $G_\alpha \subseteq S_{\alpha'}$; for each $\eta \in L'$ there is a unique η'' in L such that $S_\eta^2 \subseteq G_{\eta''}$. We define two mappings f and $g, f: L \rightarrow L', g: L' \rightarrow L$ by $\alpha' = f(\alpha)$ and $\eta'' = g(\eta)$, respectively; in other words

$$G_\alpha \subseteq S_{f(\alpha)}, \quad S_\eta^2 \subseteq G_{g(\eta)}.$$

This implies $G_\alpha \subseteq S_{f(\alpha)}^2 \subseteq G_{gf(\alpha)}$. It follows that $gf(\alpha) = \alpha$; thus gf is the identity mapping on L . Likewise fg is the identity mapping on L' . Hence f and g are one-to-one and onto, $g = f^{-1}$.

Identifying α with $f(\alpha)$, and L with L' , we have

$$(7) \quad S = \bigcup_{\alpha \in L} S_\alpha$$

$$(8) \quad S^2 = \bigcup_{\alpha \in L} S_\alpha^2, \quad S_\alpha^2 = G_\alpha.$$

We notice that $G_\alpha \subseteq S_\alpha$, hence $S^2 \cap S_\alpha = S_\alpha^2 = G_\alpha$. By the assumption on (6), (7) is the greatest \mathcal{S} -decomposition of S .

Let e_α be an identity element of G_α . Since $S_\alpha^2 = G_\alpha$, e_α is a unique idempotent of S_α . We will prove (9) through (12) below:

$$(9) \quad e_{\alpha\beta}e_\alpha = e_{\alpha\beta} = e_\alpha e_{\alpha\beta} \quad \text{for all } \alpha, \beta \in L.$$

Noticing that $e_{\alpha\beta}e_\alpha \in G_{\alpha\beta}$,

$$(e_{\alpha\beta}e_\alpha)(e_{\alpha\beta}e_\alpha) = (e_{\alpha\beta}e_\alpha e_{\alpha\beta})e_\alpha = (e_{\alpha\beta}e_{\alpha\beta}e_\alpha)e_\alpha = e_{\alpha\beta}e_\alpha.$$

Thus $e_{\alpha\beta}e_\alpha$ is an idempotent, hence $e_{\alpha\beta}e_\alpha = e_{\alpha\beta}$. The proof of the remaining part is done in the same way.

$$(10) \quad x_\alpha e_\beta = e_\beta x_\alpha \quad \text{for all } x_\alpha \in S_\alpha, \text{ all } \alpha, \beta \in L .$$

By using (9),

$$\begin{aligned} x_\alpha e_\beta &= (x_\alpha e_\beta) e_{\alpha\beta} = x_\alpha (e_\beta e_{\alpha\beta}) = x_\alpha e_{\alpha\beta} = e_{\alpha\beta} x_\alpha e_{\alpha\beta} \\ &= e_{\alpha\beta} (e_{\alpha\beta} x_\alpha) = e_{\alpha\beta} x_\alpha = (e_{\alpha\beta} e_\beta) x_\alpha = e_{\alpha\beta} (e_\beta x_\alpha) = e_\beta x_\alpha . \end{aligned}$$

$$(11) \quad e_\alpha e_\beta = e_{\alpha\beta} = e_\beta e_\alpha \quad \text{for all } \alpha, \beta \in L .$$

We have $e_\alpha e_\beta = e_\beta e_\alpha$ by (10). It can be easily proved that $e_\alpha e_\beta$ is an idempotent. Therefore $e_\alpha e_\beta = e_{\alpha\beta}$.

$$(12) \quad x_\alpha y_\beta = (x_\alpha e_\alpha)(y_\beta e_\beta) \quad \text{for all } x_\alpha \in S_\alpha, y_\beta \in S_\beta, \text{ all } \alpha, \beta \in L .$$

By (10) and (11) we have

$$(x_\alpha e_\alpha)(y_\beta e_\beta) = x_\alpha (e_\alpha y_\beta) e_\beta = x_\alpha (y_\beta e_\alpha) e_\beta = (x_\alpha y_\beta)(e_\alpha e_\beta) = (x_\alpha y_\beta) e_{\alpha\beta} = x_\alpha y_\beta .$$

Let mappings $\varphi_\alpha: S_\alpha \rightarrow G_\alpha (\alpha \in L)$ be defined by

$$\varphi_\alpha(x_\alpha) = x_\alpha e_\alpha = e_\alpha x_\alpha .$$

Then each φ_α is a homomorphism of S_α onto G_α such that $\varphi_\alpha(x_\alpha) = x_\alpha$ for all $x_\alpha \in G_\alpha$ and

$$(13) \quad x_\alpha y_\beta = \varphi_\alpha(x_\alpha) \varphi_\beta(y_\beta), \quad x_\alpha \in S_\alpha, \quad y_\beta \in S_\beta .$$

Consequently S is an inflation of $G = \bigcup_{\alpha \in L} G_\alpha$ and S_α is an inflation of G_α . Thus we have proved that (4.1) \Rightarrow (4.2) and (4.3).

REMARK. In the proof of the proposition, if we use the fact that a semilattice of groups is an inverse semigroup, we immediately have (11), hence (9). However, we proved these directly without using the property of inverse semigroups.

Proof of theorem. We will need two lemmas to prove the theorem.

LEMMA 4. *If S satisfies $xy = f(x, y)$ with (2), then every content of rank greater than 1 is a group.*

Proof. Let $C = C(a_1, \dots, a_k), k > 1$. Each element of C is expressed as a word involving all the letters a_1, \dots, a_k . Let $w \in C$ be decomposed into the product of two words w_1, w_2 of a_i 's, namely elements w_1, w_2 of S : $w = w_1 w_2$. Replacing w by $f(w_1, w_2)$ repeatedly we can arrange w such that $|a_i|_w > 1$ for all i . First we will prove that each element w has a form $a_i v', v' \in C$. If a_i is the initial letter of w , then we have already the form since $|a_i|_w > 1$ and hence $v' \in C$. Suppose

$$w = w_1 a_i w_2, \quad w_1, w_2 \in S.$$

Replacing $w_1(a_i w_2)$ by $f(w_1, a_i w_2)$ we have a form $w = a_i v'$. Since $|a_j|_w > 1$ for all j , $|a_i|_{v'} \geq 1$ and $|a_j|_{v'} > 1$ for $j \neq i$. Hence $v' \in C$. Again we can arrange v' such that $|a_j|_{v'} > 1$ for all j . Now let w' be any element of C :

$$w' = a_{i_1} a_{i_2} \cdots a_{i_l}.$$

By repeating the same procedure

$$w = a_{i_1} w_{i_1} = a_{i_1} a_{i_2} w_{i_2} = \cdots = a_{i_1} a_{i_2} \cdots a_{i_l} w_{i_l} = w' w_{i_l}$$

where $w_{i_j} \in C (j = 1, \dots, l)$. In the same way we have $w = w'_{i_l} w'$ for some $w'_{i_l} \in C$. Thus we have proved the right and left divisibility; therefore $C(a_1, \dots, a_k)$ is a group.

LEMMA 5. Let $S = \bigcup_{\alpha \in L} S_\alpha$ be greatest \mathcal{S} -decomposition of a semigroup S . If S satisfies $xy = f(x, y)$ with (2), then S_α^2 is a group and $S^2 = \bigcup_{\alpha \in L} S_\alpha^2$.

Proof. Let $a \in S_\alpha^2, a = xy$ for some $x, y \in S_\alpha$. Then a is in the content $C(x, y)$ which is a subgroup of S_α^2 by Lemma 4. Thus S_α^2 is a union of subgroups, hence a disjoint union of maximal subgroups of S_α . We will prove that for two distinct arbitrary elements a and b of S_α^2 there are subgroups G_a and G_b which contain a and b , respectively, such that $G_a \cap G_b \neq \emptyset$. Then S_α^2 will be a group. Since $a, b \in S_\alpha$ there is a finite sequence of contents C_1, C_2, \dots, C_t in S such that

$$a \in C_1, \quad C_i \cap C_{i+1} \neq \emptyset \quad (i = 1, \dots, t-1), \quad b \in C_t.$$

As remarked in §1, $C_i \subseteq S_\alpha (i = 1, \dots, t)$. Since $a, b \in S_\alpha^2$, we may assume C_1 and C_t are of rank > 1 . Also we may assume C_2, \dots, C_{t-1} are of rank > 1 for the following reason. Suppose $C_i, 1 < i < t$, has rank 1, $C_i = [x]$, the cyclic subsemigroup generated by x . If x is in either $C_i \cap C_{i-1}$ or $C_i \cap C_{i+1}$, then $C_i \subseteq C_{i-1}$ or $C_i \subseteq C_{i+1}$, respectively; so C_i can be excluded from the sequence. If x is in C_i but not in $(C_i \cap C_{i-1}) \cup (C_i \cap C_{i+1})$, then we can replace C_i by $\{x^i; i > 1\}$ of rank > 1 . By Lemma 4 all the $C_i (i = 1, \dots, t)$ are subgroups of S_α . It is easy to prove that $C_i \cap C_{i+1}$ and $C_{i+1} \cap C_{i+2}$ are also subgroups; hence $C_i \cap C_{i+2} \neq \emptyset$ since the identity element of the group C_{i+1} has to lie in C_i and C_{i+2} . Continuing this procedure we have $C_1 \cap C_t \neq \emptyset$ as desired. Thus it has been proved that S_α^2 is a group. Let $G_\alpha = S_\alpha^2$ for each $\alpha \in L$. G_α is the greatest subgroup in S_α , a maximal subgroup in S . Now let $z \in S^2, z = xy$ for some $x \in S_\alpha, y \in S_\beta$, so $z \in S_{\alpha\beta}$. The element z is in a subgroup $C(x, y) \subset S_{\alpha\beta}^2$, hence $z \in G_{\alpha\beta}$. Thus we have

$$S^2 = \bigcup_{\alpha \in L} G_\alpha, \quad G_\alpha = S_\alpha^2 .$$

Proof of theorem. By Lemma 5 and the proposition, it has been proved that if S satisfies $xy = f(x, y)$ with (2) then S is an inflation of $\bigcup_{\alpha \in L} G_\alpha$, a semilattice L of groups G_α , in which G_α satisfies the identity $xy = f(x, y)$. We will prove the converse of the theorem. Suppose that S is an inflation of $\bigcup_{\alpha \in L} G_\alpha$, where each G_α is a group satisfying $xy = f(x, y)$ with (2). Let $x_\alpha \in S_\alpha, y_\beta \in S_\beta$. By proposition $x_\alpha y_\beta \in G_{\alpha\beta}$. By using (10) and by recalling the form of $f(x, y)$ we have

$$x_\alpha y_\beta = x_\alpha y_\beta e_{\alpha\beta} = (x_\alpha e_{\alpha\beta})(y_\beta e_{\alpha\beta}) = f(x_\alpha e_{\alpha\beta}, y_\beta e_{\alpha\beta}) = f(x_\alpha, y_\beta) e_{\alpha\beta} = f(x_\alpha, y_\beta) .$$

This proves that S satisfies the same identity. Thus the proof of the theorem has been completed.

Proof of corollary. Suppose $xy = f(x, y)$ implies $xy = yx$ in semigroups. Then (3.2) is obvious and the necessity of (3.1) is already proved in § 1. It remains to prove the sufficiency of (3.1) and (3.2). The theorem describes the structure of S satisfying the identity $xy = f(x, y)$ in which $|f(x, y)| > 2$ and $f(x, y)$ satisfies (3.1), that is, $f(x, y)$ has the form (2). Now additionally assume (3.2). By using (10)

$$x_\alpha y_\beta = (x_\alpha e_{\alpha\beta})(y_\beta e_{\alpha\beta}) = (y_\beta e_{\alpha\beta})(x_\alpha e_{\alpha\beta}) = y_\beta x_\alpha .$$

Examples and problems. As the application of corollary we give a few examples below:

EXAMPLE 1. (Tully). $xy = y^m x^n, m \geq 1, n \geq 1,$

EXAMPLE 2. $xy = (yx)^m, m \geq 1,$

EXAMPLE 3. $xy = y^{m_1} x^{n_1} \dots y^{m_h} x^{n_h}, m = \sum_{i=1}^h m_i, n = \sum_{i=1}^h n_i,$ the greatest common divisor of $m - 1$ and $n - 1$ is 2 or 1.

Each identity of Examples 1, 2 and 3 implies $xy = yx$ in semigroups because it can be easily proved that commutativity follows in groups.

EXAMPLE 4. $xy = (y^2 x^2)^3.$

In groups this identity is equivalent to the identity

$$x^3 = e, e \text{ the identity element .}$$

We know that the free group G generated by a and b subject to $x^3 = e$ is a finite noncommutative group [2] [3]. Therefore commutativity

does not follow.

Finally we give examples which satisfy (3.2) but not (3.1).

EXAMPLE 5. $xy = x^2y^3x$,

EXAMPLE 6. $xy = yx^3y^2$,

EXAMPLE 7. $xy = xy^3$.

Each of Examples 5, 6, and 7 implies commutativity in groups. Accordingly we can say:

The conditions (3.1) and (3.2) of the corollary are independent.

ADDENDA. 1. Let $f(x, y)$ and $g(x, y)$ be words involving both x and y and let $|f(x, y)| \geq 2$ and $|g(x, y)| \geq 2$.

If $g(x, y) = f(x, y)$ implies $xy = yx$ in semigroups¹, then one of $g(x, y)$ and $f(x, y)$ is xy and the other satisfies the condition (3.1); hence this case is reduced to that of the corollary.

Suppose $|g(x, y)| > 2$ and $|f(x, y)| > 2$. Let F be the free semigroup generated by the two letters a and b , and I be the ideal of F consisting of all words with length more than 2. Let $S = F/I$. We see $S = \{0, a, b, a^2, b^2, ab, ba\}$. S satisfies $g(x, y) = f(x, y)$ but it is not commutative as $ab \neq ba$. We may assume the identity is $xy = f(x, y)$ which is the condition in the corollary.

2. Let $f(x, y)$ be a word involving both x and y .

$x^2 = f(x, y)$ does not imply $xy = yx$ in general.

For the reason used for (3.1), we may assume $f(x, y)$ starts in y and ends in y . Let F be the free semigroup generated by a and b , and I be the set of all words which involve either at least two a 's or at least two b 's. I is an ideal. $S = F/I = \{0, a, b, ab, ba\}$. S satisfies $x^2 = f(x, y)$, but is not commutative.

PROBLEMS. 1. Let $f(x, y) = y^{m_1}x^{n_1} \dots y^{m_h}x^{n_h}$ with (2). How can we describe explicitly (3.2) in terms of $m_1, n_1, \dots, m_h, n_h$?

2. Determine the structure of semigroups satisfying an identity of the form $xy = y^{m_1}x^{n_1} \dots y^{m_{h-1}}x^{n_{h-1}}y^{m_h}$, $m_1 \neq 0$, $m_h \neq 0$, $h \geq 2$.

3. Determine the structure of semigroups satisfying an identity of the form $xy = x^{m_1}y^{n_1} \dots x^{m_{h-1}}y^{n_{h-1}}$, $m_1 \neq 0$, $n_{h-1} \neq 0$.

¹ The author owes this result to Dr. D. G. Mead's helpful suggestion.

4. Let $f(x, y) = y^{m_1}x^{n_1} \dots y^{m_h}x^{n_h}y^{m_{h+1}}$, $m_i, m_{h+1} \neq 0$, $h \neq 0$. Under what condition on $f(x, y)$ does the identity $x = f(x, y)$ imply $xy = yx$?

A semigroup S satisfying the identity $x = f(x, y)$ is a group.

In fact S is a union of groups. By [1] S is a semilattice L of completely simple semigroups $S_\alpha (\alpha \in L)$

$$S = \bigcup_{\alpha \in L} S_\alpha .$$

We can easily prove that $|L| = 1$; S is a completely simple semigroup, that is, a rectangular band B of groups. However we can prove that $|B| = 1$.

A partial answer follows:

Let
$$\text{g.c.d.} \left(\sum_{i=1}^{h+1} m_i, \sum_{i=1}^h n_i - 1 \right) = g .$$

If $g = 2$, the answer is affirmative.

References

1. A. H. Clifford and G. B. Preston, *The algebraic theory of semigroups*, Vol. 1, Math. Surveys No. 7, Amer. Math. Soc., Providence, R.I., 1961.
2. J. A. Green and D. Rees, *On semigroups in which $x^r = x$* , Cambridge Philos. Soc. **48** (1952), 35-40.
3. M. Hall, *The theory of groups*, Macmillan Company, New York, 1959.
4. Mario Petrich, *The maximal semilattice decomposition of a semigroup*, Math. Zeit. **85** (1964), 68-82.
5. T. Tamura, *The theory of construction of finite semigroups I*, Osaka Math. J. **8** (1956), 243-261.
6. ———, *Another proof of a theorem concerning the greatest semilattice decomposition of a semigroup*, Proc. Jap. Acad. **40** (1964), 777-780.
7. E. J. Tully, *Semigroups satisfying an identity of the form $xy = y^m x^n$* (to be published.)

Received May 25, 1968.

UNIVERSITY OF CALIFORNIA
DAVIS, CALIFORNIA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. ROYDEN
Stanford University
Stanford, California

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

RICHARD PIERCE
University of Washington
Seattle, Washington 98105

BASIL GORDON
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. 36, 1539-1546. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

Pacific Journal of Mathematics

Vol. 31, No. 2

December, 1969

Efraim Pacillas Armendariz, <i>Quasi-injective modules and stable torsion classes</i>	277
J. Adrian (John) Bondy, <i>On Ulam's conjecture for separable graphs</i>	281
Vasily Cateforis and Francis Louis Sandomierski, <i>On commutative rings over which the singular submodule is a direct summand for every module</i>	289
Rafael Van Severen Chacon, <i>Approximation of transformations with continuous spectrum</i>	293
Raymond Frank Dickman and Alan Zame, <i>Functionally compact spaces</i>	303
Ronald George Douglas and Walter Rudin, <i>Approximation by inner functions</i>	313
John Walter Duke, <i>A note on the similarity of matrix and its conjugate transpose</i>	321
Micheal Neal Dyer and Allan John Sieradski, <i>Coverings of mapping spaces</i>	325
Donald Campbell Dykes, <i>Weakly hypercentral subgroups of finite groups</i>	337
Nancy Dykes, <i>Mappings and realcompact spaces</i>	347
Edmund H. Feller and Richard Laham Gantos, <i>Completely injective semigroups</i>	359
Irving Leonard Glicksberg, <i>Semi-square-summable Fourier-Stieltjes transforms</i>	367
Samuel Irving Goldberg and Kentaro Yano, <i>Integrability of almost cosymplectic structures</i>	373
Seymour Haber and Charles Freeman Osgood, <i>On the sum $\sum \langle n\alpha \rangle^{-1}$ and numerical integration</i>	383
Sav Roman Harasymiv, <i>Dilations of rapidly decreasing functions</i>	395
William Leonard Harkness and R. Shantaram, <i>Convergence of a sequence of transformations of distribution functions</i>	403
Herbert Frederick Kreimer, Jr., <i>A note on the outer Galois theory of rings</i>	417
James Donald Kuelbs, <i>Abstract Wiener spaces and applications to analysis</i>	433
Roland Edwin Larson, <i>Minimal T_0-spaces and minimal T_D-spaces</i>	451
A. Meir and Ambikeshwar Sharma, <i>On Ilyeff's conjecture</i>	459
Isaac Namioka and Robert Ralph Phelps, <i>Tensor products of compact convex sets</i>	469
James L. Rovnyak, <i>On the theory of unbounded Toeplitz operators</i>	481
Benjamin L. Schwartz, <i>Infinite self-interchange graphs</i>	497
George Szeto, <i>On the Brauer splitting theorem</i>	505
Takayuki Tamura, <i>Semigroups satisfying identity $xy = f(x, y)$</i>	513
Kenneth Tolo, <i>Factorizable semigroups</i>	523
Mineko Watanabe, <i>On a boundary property of principal functions</i>	537
James Juei-Chin Yeh, <i>Singularity of Gaussian measures in function spaces with factorable covariance functions</i>	547