

Semiring identities of the Brandt monoid*

Mikhail Volkov
 Institute of Natural Sciences and Mathematics
 Ural Federal University
 m.v.volkov@urfu.ru

Abstract

The 6-element Brandt monoid B_2^1 admits a unique addition under which it becomes an additively idempotent semiring. We show that this addition is a term operation of B_2^1 as an inverse semigroup. As a consequence, we exhibit an easy proof that the semiring identities of B_2^1 are not finitely based.

We assume the reader's acquaintance with basic concepts of universal algebra such as an identity and a variety; see, e.g., [1, Chapter II].

The 6-element Brandt monoid B_2^1 can be represented as a semigroup of the following zero-one 2×2 -matrices

$$\begin{array}{cccccc} \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} & \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} & \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} & \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} & \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} & \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \\ 0 & E & E_{12} & E_{21} & E_{11} & E_{22} \end{array} \quad (1)$$

under the usual matrix multiplication \cdot or as a monoid with presentation

$$\langle E_{12}, E_{21} \mid E_{12}E_{21}E_{12} = E_{12}, E_{21}E_{12}E_{21} = E_{21}, E_{12}^2 = E_{21}^2 = 0 \rangle.$$

Quoting from a recent paper [3], 'This Brandt monoid is perhaps the most ubiquitous harbinger of complex behaviour in all finite semigroups'. In particular, (B_2^1, \cdot) has no finite basis for its identities (Perkins [13, 14]) and is one of the four smallest semigroups with this property (Lee and Zhang [10]).

The monoid (B_2^1, \cdot) has a natural involution that swaps E_{12} and E_{21} and fixes all other elements. In terms of the matrix representation (1) this involution is nothing but the usual matrix transposition; we will, however, use the notation $x \mapsto x^{-1}$ for

*Supported by the Ministry of Science and Higher Education of the Russian Federation (Ural Mathematical Center project No. 075-02-2020-1537/1)

the involution, emphasizing that x^{-1} is the unique inverse of x . Recall that elements x, y of a semigroup (S, \cdot) are said to be *inverses* of each other if $xyx = x$ and $yxy = y$. A semigroup is called *inverse* if every its element has a unique inverse; inverse semigroups can therefore be thought of as algebras of type $(2, 1)$. Being considered as an inverse semigroup, the monoid $(B_2^1, \cdot, {}^{-1})$ retains its complex equational behaviour: B_2^1 has no finite basis for its inverse semigroup identities (Kleiman [6]) and is the smallest inverse semigroup with this property (Kleiman [5, 6]).

In the present note we consider equational properties of yet another enhancement of the monoid (B_2^1, \cdot) with an additional operation, this time binary. Recall that an *additively idempotent semiring* an algebra $(S, +, \cdot)$ of type $(2, 2)$ such that the additive reduct $(S, +)$ is a *semilattice* (that is, a commutative idempotent semigroup), the multiplicative reduct (S, \cdot) is a semigroup, and multiplication distributes over addition on the left and on the right, that is, $(S, +, \cdot)$ satisfies the identities $x(y + z) \approx xy + xz$ and $(y + z)x \approx yx + zx$. In papers which motivation comes from semigroup theory, objects of this sort sometimes appear under the name *semilattice-ordered semigroups*, see, e.g., [8] or [12]. We will stay with the term ‘additively idempotent semiring’, abbreviated to ‘ai-semiring’ in the sequel.

Our key observation is the following:

Lemma 1. *Let $(S, \cdot, {}^{-1})$ be an inverse semigroup satisfying the identity*

$$x^n \approx x^{n+1} \tag{2}$$

for some n . Define

$$x \oplus y := (xy^{-1})^n x.$$

Then (S, \cdot, \oplus) is an ai-semiring.

Proof. Let $E(S)$ stand for the set of all idempotents of S . The relation

$$\leq := \{(a, b) \in S \times S \mid a = eb \text{ for some } e \in E(S)\}$$

is a partial order on S referred to as the *natural partial order*; see [15, Section II.1] or [9, pp. 21–23]. We need two basic properties of the natural partial order:

- 1) \leq is compatible with both multiplication and inversion;
- 2) $a \leq b$ if and only if $a = bf$ for some $f \in E(S)$.

Take any $a, b \in S$ and suppose that $c \leq a$ and $c \leq b$. Then $c^{-1} \leq b^{-1}$ whence by the compatibility with multiplication

$$c = (cc^{-1})^n c \leq (ab^{-1})^n a = a \oplus b.$$

In presence of the identity (2), $(ab^{-1})^n = (ab^{-1})^{n+1} = \dots = (ab^{-1})^{2n}$. Hence

$$a \oplus b = (ab^{-1})^n \cdot a \leq a.$$

Further,

$$\begin{aligned} a \oplus b &= (ab^{-1})^n a = (ab^{-1})^{n+1} a = \dots = (ab^{-1})^{2n-1} a = \\ &= (ab^{-1})^n \cdot (ab^{-1})^{n-1} a = (ab^{-1})^n \cdot a(b^{-1}a)^{n-1} = \text{(using } b^{-1} = b^{-1}bb^{-1}\text{)} \\ &= (ab^{-1})^n \cdot b \cdot (b^{-1}a)^n \leq b \cdot (b^{-1}a)^n \leq b \end{aligned}$$

since $(b^{-1}a)^n \in E(S)$. We see that $a \oplus b$ is nothing but the infimum of $\{a, b\}$ with respect to the natural partial order. Thus, (S, \oplus) is a semilattice. It is known [16, Proposition 1.22], see also [9, Proposition 19] that if a subset $H \subseteq S$ possesses an infimum under the natural partial order, then so do the subsets sH and Hs for any $s \in S$, and $\inf(sH) = s(\inf H)$, $\inf(Hs) = (\inf H)s$. This implies that multiplication distributes over \oplus on the left and on the right. \square

Remark 1. The essence of Lemma 1 is known. Leech, in the course of his comprehensive study of inverse monoids $(S, \cdot, {}^{-1}, 1)$ that are inf-semilattices under the natural partial order, has verified that (S, \leq) is a inf-semilattice whenever S is a periodic combinatorial¹ inverse monoid; see [11, Example 1.21(d), item (iv)]. Of course, the requirement of S being a monoid is not essential: if a semigroup S periodic and combinatorial then so is the monoid S^1 obtained by adjoining a formal identity to S . Clearly, if a semigroup satisfies (2), then it is both periodic and combinatorial whence Leech's observation applies. We have preferred the above direct proof of Lemma 1 because we need a $(\cdot, {}^{-1})$ -term for the semilattice operation, and such a term is not explicitly present in [11].

Obviously, the 6-element Brandt monoid satisfies the identity $x^2 \approx x^3$. Thus, Lemma 1 applies, and (B_2^1, \oplus, \cdot) is an ai-semiring. It is known (and easy to verify) that \oplus is the only addition on B_2^1 under which B_2^1 becomes an ai-semiring.

Our main result states that, similarly to the plain semigroup (B_2^1, \cdot) and the inverse semigroup $(B_2^1, \cdot, {}^{-1})$, the ai-semiring (B_2^1, \oplus, \cdot) admits no finite identity basis. Its proof employs a series of inverse semigroups C_n , $n = 2, 3, \dots$, constructed in [6] as semigroups of partial one-to-one transformations. Here, to align with the matrix representation chosen for the B_2^1 , we describe them as semigroups of zero-one matrices.

The set R_m of all zero-one $m \times m$ -matrices which have at most one entry equal to 1 in each row and column forms an inverse monoid under usual matrix multipli-

¹A semigroup S is *periodic* if all monogenic subsemigroups of S are finite and *combinatorial* if all subgroups of S are trivial.

cation \cdot and transposition. The inverse monoid R_m is called the *rook monoid*² as its matrices encode placements of nonattacking rooks on an $m \times m$ chessboard.

Let $m = 2n + 1$ and define $m \times m$ -matrices c_1, \dots, c_n by

$$c_k := E_{k+1k} + E_{n+k \ n+k+1}, \quad k = 1, \dots, n,$$

where, as usual, E_{ij} denotes the $m \times m$ -matrix unit with an entry 1 in the (i, j) position and 0's elsewhere. For instance, if $n = 2$, then c_1 and c_2 are the following 5×5 -matrices:

$$c_1 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad c_2 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

Let C_n be the inverse subsemigroup of the rook monoid R_m generated by the matrices c_1, \dots, c_n . As a plain subsemigroup, C_n is generated by c_1, \dots, c_n and their inverses (i.e., transposes) $c_1^{-1}, \dots, c_n^{-1}$.

The next lemma collects properties of the semigroups C_n that we need.

Lemma 2. (i) *The semigroup (C_n, \cdot) does not belong to the semigroup variety generated by the monoid (B_2^1, \cdot) .*

(ii) *The semigroup (C_n, \cdot) satisfies the identity $x^2 \approx x^3$.*

(iii) *For each $k = 1, \dots, n$, $M_k(n) := C_n \setminus \{c_k, c_k^{-1}\}$ forms an inverse subsemigroup of the inverse semigroup $(C_n, \cdot, {}^{-1})$.*

(iv) *For each $k = 1, \dots, n$, the inverse semigroup $(M_k(n), \cdot, {}^{-1})$ belongs to the inverse semigroup variety generated by the inverse monoid $(B_2^1, \cdot, {}^{-1})$.*

Proof. (i) This property was established in [7, Lemma 3] by exhibiting, for each $n \geq 2$, a semigroup identity that holds in (B_2^1, \cdot) and fails in (C_n, \cdot) .

(ii) This is easy to verify (and also follows from the proof of Lemma 1 in [6]).

(iii) This is clear (and is a part of Lemma 1 in [6]).

(iv) This is Property (C) in [6]. □

Remark 2. Items (i)–(iii) of Lemma 2 are easy. In contrast, the proof of (iv) in [6] is long and complicated. We mention in passing that now the proof can be radically simplified by using a deep result by Kaĉourek [4] who provided an effective membership test for the inverse semigroup variety generated by $(B_2^1, \cdot, {}^{-1})$.

²The rook monoid is nothing but the matrix representation of the *symmetric inverse monoid*; see [15, Section IV.1] or [9, p. 6]. The name ‘rook monoid’ was suggested by Solomon [17].

Theorem 3. *The semiring identities of the additively idempotent semiring (B_2^1, \oplus, \cdot) admit no basis involving only finitely many variables, and hence, no finite basis.*

Proof. Arguing by contradiction, assume that (B_2^1, \oplus, \cdot) has an identity basis Σ such that each identity $u \approx v$ in Σ involves less than n variables. Consider the inverse semigroup $(C_n, \cdot, ^{-1})$. By Lemmas 1 and 2(ii), the addition defined by $x \oplus y := (xy^{-1})^2x$ makes (C_n, \oplus, \cdot) an ai-semiring. Consider an arbitrary evaluation ε of variables x_1, \dots, x_ℓ involved in the identity $u \approx v$ in this ai-semiring. By the pigeonhole principle, there exists an index $k \in \{1, \dots, n\}$ such that neither c_k nor c_k^{-1} belongs to the set $\{\varepsilon(x_1), \dots, \varepsilon(x_\ell)\}$ as this set contains at most $\ell < n$ elements. Thus, $\{\varepsilon(x_1), \dots, \varepsilon(x_\ell)\} \subset M_k(n)$.

Since $x \oplus y$ expresses as $(\cdot, ^{-1})$ -term, one can rewrite the identity $u \approx v$ into an identity $u' \approx v'$ in which u' and v' are $(\cdot, ^{-1})$ -terms. Since $u \approx v$ holds in (B_2^1, \oplus, \cdot) , the rewritten identity $u' \approx v'$ holds in the inverse semigroup $(B_2^1, \cdot, ^{-1})$. By Lemma 2(iv) the latter identity holds also in the inverse semigroup $(M_k(n), \cdot, ^{-1})$, and so u' and v' take the same value under every evaluation of the variables x_1, \dots, x_ℓ in $M_k(n)$. Hence $\varepsilon(u) = \varepsilon(u') = \varepsilon(v') = \varepsilon(v)$. We conclude that the identity $u \approx v$ holds in the ai-semiring (C_n, \oplus, \cdot) . Since an arbitrary identity from Σ holds in (C_n, \oplus, \cdot) , this ai-semiring belongs to the ai-semiring variety generated by (B_2^1, \oplus, \cdot) . This, however, contradicts Lemma 2(i), according to which even the semigroup reduct (C_n, \cdot) , does not belong to semigroup variety generated by (B_2^1, \cdot) . \square

Remark 3. To the best of my knowledge, the result of Theorem 3 has not been published up to now. However, after preparing the present article I have learnt that the result has also been obtained by colleagues in Xi'an and Melbourne but with an entirely unrelated proof.

I mention also a related paper by Dolinka [2] where he introduces a 7-element ai-semiring denoted Σ_7 and proves that its identities are not finitely based. The semigroup reduct of Σ_7 is just the monoid B_2^1 with an extra zero adjoined so that $(\Sigma_7, \cdot, ^{-1})$ and $(B_2^1, \cdot, ^{-1})$ satisfy the same inverse semigroup identities. However, the addition in Σ_7 is not derived from its inverse semigroup structure, and one can easily see that the semiring identities of $(\Sigma_7, +, \cdot)$ and (B_2^1, \oplus, \cdot) are essentially different. It should be also mentioned that in [2] Dolinka actually considers ai-semirings with 0 as algebras of type $(2,2,0)$.

Remark 4. Leech [11] defined an *inverse algebra* as an algebra $(A, \cdot, \wedge, ^{-1}, 1)$ of type $(2,2,1,0)$ such that the reduct $(A, \cdot, ^{-1}, 1)$ is an inverse monoid, the reduct (A, \wedge) is a meet semilattice, and the natural partial order of the inverse monoid coincides with that of the semilattice. Clearly, $(B_2^1, \cdot, \oplus, ^{-1}, E)$ constitutes an inverse algebra in Leech's sense, and the above proof of Theorem 3 can be easily adapted to show that $(B_2^1, \cdot, \oplus, ^{-1}, E)$ has no finite identity basis also as such algebra.

References

- [1] S. Burris and H.P. Sankappanavar, *A Course in Universal Algebra*. Springer-Verlag, Berlin-Heidelberg-New York (1981)
- [2] I. Dolinka, A nonfinitely based finite semiring. *Int. J. Algebra Comput.* **17**(8), 1537–1551 (2007)
- [3] M. Jackson and W.T. Zhang, From A to B to Z . *Semigroup Forum* (in print)
- [4] J. Kađourek, On varieties of combinatorial inverse semigroups. I. *Semigroup Forum* **43**, 305–330 (1991)
- [5] E.I. Kleiman, On bases of identities of Brandt semigroups. *Semigroup Forum* **13**, 209–218 (1977)
- [6] E.I. Kleiman, Bases of identities of varieties of inverse semigroups. *Sib. Math. J.* **20**, 530–543 (1979). [Translated from *Sibirskii Matematicheskii Zhurnal* **20**, 760–777 (1979)]
- [7] E.I. Kleiman, A pseudovariety generated by a finite semigroup. *Ural. Gos. Univ. Mat. Zap.* **13**(1), 40–42 (1982) (Russian)
- [8] M. Kuřil and L. Polák, On varieties of semilattice-ordered semigroups. *Semigroup Forum*, **71**, 27–48 (2005)
- [9] M.V. Lawson, *Inverse Semigroups. The Theory of Partial Symmetries*. World Scientific, Singapore (1999)
- [10] E.W.H. Lee and W.T. Zhang, Finite basis problem for semigroups of order six. *London Math. Soc. J. Comput. Math.* **18**, 1–129 (2015)
- [11] J. Leech, Inverse monoids with a natural semilattice ordering. *Proc. London Math. Soc.* **s3-70**(1), 146–182 (1995)
- [12] D.B. McAlister, Semilattice ordered inverse semigroups. In J.M. André et al (eds.), *Semigroups and Formal Languages*, pp. 205–218. World Scientific, New Jersey (2007)
- [13] P. Perkins, *Decision Problems for Equational Theories of Semigroups and General Algebras*. Ph.D. thesis, Univ. of California, Berkeley (1966)
- [14] P. Perkins, Bases for equational theories of semigroups, *J. Algebra* **11**, 298–314 (1969)
- [15] M. Petrich, *Inverse Semigroups*. John Wiley & Sons, New York (1984)
- [16] B.M. Schein, Completions, translational hulls and ideal extensions of inverse semigroups. *Czechoslovak Math. J.* **23**(4), 575–610 (1973)
- [17] L. Solomon, Representations of the rook monoid. *J. Algebra* **256**(2), 309–342 (2002)