

Sequence Analysis and Expression Patterns Divide the Maize *knotted1*-like Homeobox Genes into Two Classes

Randall Kerstetter,^a Erik Vollbrecht,^a Brenda Lowe,¹ Bruce Veit,² Judy Yamaguchi,^b and Sarah Hake^{a,b,3}

^a Department of Plant Biology, University of California, Berkeley, California 94720

^b Plant Gene Expression Center, U.S. Department of Agriculture, Agricultural Research Station, 800 Buchanan Street, Albany, California 94710

The homeobox of the *knotted1* (*kn1*) gene was used to isolate 12 related sequences in maize. The homeodomains encoded by the *kn1*-like genes are very similar, ranging from 55 to 89% amino acid identity. Differences outside the precisely conserved third helix allowed us to group the genes into two classes. The homeodomains of the seven class 1 genes share 73 to 89% identical residues with *kn1*. The four class 2 genes share 55 to 58% identical residues with *kn1*, although the conservation within the class is greater than 87%. Expression patterns were analyzed by RNA gel blot analysis. Class 1 genes were highly expressed in meristem-enriched tissues, such as the vegetative meristem and ear primordia. Expression was not detectable in leaves. The class 2 genes were expressed in all tissues, although one was abundantly expressed in roots. The genes were mapped using recombinant inbred populations. We determined that clusters of two to three linked genes are present on chromosomes 1 and 8; otherwise, the genes are distributed throughout the genome. Four pairs of genes, similar in both sequence and expression patterns, mapped within duplicated regions of the genome.

INTRODUCTION

The homeobox was first identified as a cross-hybridizing DNA sequence shared among several genes that control morphogenesis in *Drosophila* (McGinnis et al., 1984; Scott and Weiner, 1984). Comparison of the deduced amino acid sequences from these genes revealed a region with similarity to known regulatory proteins encoded by the yeast mating type loci *MAT α 2* and *MAT α 1* (Laughon and Scott, 1984; Shepherd et al., 1984). This conserved protein motif became known as the homeodomain. Subsequently, the homeobox has been found in a number of genes that control cell fate or convey positional information during the development of diverse groups of animals, fungi, and plants (for review, see Gehring, 1987; Scott et al., 1989; Kessel and Gruss, 1990).

Homeodomain proteins are thought to act as sequence-specific DNA binding proteins that can directly regulate the expression of specific groups of target genes (Affolter et al., 1990a; Hayashi and Scott, 1990). Although the primary amino acid sequences show relatively low identity, 27%, high-resolution structural studies of the homeodomains of *Engrailed* (*En*) and *Antennapedia* (*Antp*) and of α 2 complexed with DNA have revealed nearly identical structures (Kissinger et al., 1990; Otting et al., 1990; Wolberger et al., 1991). The homeodomain

of each protein contains three α -helices, although the third α -helix of ANTP is kinked and is therefore referred to as helices 3 and 4 (Qian et al., 1989). Residues in helix 3, dubbed the "recognition" helix, make sequence-specific contact in the major groove of the DNA binding site, and basic residues of the N-terminal portion of the homeodomain make contact in the minor groove forming a pincer-like grip. The remarkable conformational identity between distantly related homeodomains suggests that important structural and functional properties are shared among all members of this diverse class of genes (Kornberg, 1993).

Although structurally similar, different homeodomains are able to recognize diverse DNA binding sites. This has been clearly illustrated by experiments that show that changes of as few as one residue in the third helix can dramatically alter the specificity of DNA binding (Hanes and Brent, 1989; Treisman et al., 1989; Affolter et al., 1990b; Percival-Smith et al., 1990). Different homeodomain proteins have been grouped into separate families or classes based on either sequence identity within the homeodomain or conserved protein motifs outside of the homeodomain (Scott et al., 1989; Treisman et al., 1991). Examples of such families of homeodomain proteins in animal systems include the Antennapedia, Engrailed, Paired, POU, and LIM classes. The Antennapedia class is a large family; its members comprise the homeotic complexes (HOM-C in *Drosophila* and Hox in vertebrates; for review, see Krumlauf, 1992). The distinguishing feature of this class is an identical core sequence of 12 amino acids in the recognition helix; this

¹ Current address: Department of Biology, Yale University, New Haven, CT 06511.

² Current address: Department of Biology, University of California, San Diego, La Jolla, CA 92093.

³ To whom correspondence should be addressed.

sequence is shared by all members. The POU class of homeodomain proteins, in contrast, is distinguished by the presence of a 75-amino acid POU domain separated from the homeodomain by 20 to 30 amino acids (Herr et al., 1988). In plants, four different types of homeodomain proteins have been described thus far: homeodomain zipper proteins (HD-Zip), which are distinguished by the presence of a leucine zipper adjacent to the homeodomain (Ruberti et al., 1991; Mattsson et al., 1992; Schena and Davis, 1992, 1994); plant homeodomain finger proteins (PHD-finger) represented by ZMHOX1A in maize (Bellmann and Werr, 1992), HAT3.1 in Arabidopsis (Schindler et al., 1993), and the pathogenesis-related homeodomain proteins in parsley and Arabidopsis (PRHP and PRHA, respectively; Korfhage et al., 1994), which share a conserved cysteine-rich motif; the Arabidopsis GLABRA2 homeodomain protein (Rerie et al., 1994); and KNOTTED1-like proteins (Vollbrecht et al., 1991; Matsuoka et al., 1993; Ma et al., 1994), which are the subject of this report.

The first family of homeobox genes reported in a plant species was isolated by transposon tagging the morphological mutation *Knotted1* (*Kn1*) in maize (Hake et al., 1989; Vollbrecht et al., 1991). *Kn1* is one of a number of dominant mutations that alter the developmental fate of cells in the leaf blade toward sheath-like identities (for reviews, see Freeling, 1992; Hake, 1992). Wild-type expression of *kn1* is localized to the meristem and ground tissue of the unexpanded stem and is absent from leaves and leaf primordia. In addition, *kn1* is expressed throughout the corpus of the inflorescence meristem and in spikelet and floral meristems but is not expressed in lateral organs such as glume primordia (Smith et al., 1992; Jackson et al., 1994). In the dominant *Kn1* mutants, however, ectopic expression of *kn1* in the veins of the developing leaf blade has been correlated with the mutant phenotype (Smith et al., 1992). These gain-of-function mutations typically affect the lateral veins, causing the adjacent tissues to differentiate incompletely and continue dividing, with the result that sporadic outgrowths, or knots, are formed on the leaf blade (Gelinas et al., 1969; Freeling and Hake, 1985). Transgenic tobacco plants expressing the *kn1* cDNA driven by the cauliflower mosaic virus 35S promoter have altered leaf size and shape, and in severe cases, ectopic shoots arise from the leaf surface (Sinha et al., 1993). The results suggest that ectopic expression of *kn1* causes leaf cells to adopt different fates that range from alternative types of leaf cells to meristem cells (Smith et al., 1992; Sinha et al., 1993).

In animals, families of related homeobox genes often specify related developmental processes (Gehring, 1987; Krumlauf, 1992). By taking advantage of the conserved homeobox, we isolated a number of *kn1*-like genes from maize (Vollbrecht et al., 1991, 1993). If members of this class of genes specify related developmental processes, they may, like *kn1*, play significant roles in the development of higher plants. Here, we compare the deduced amino acid sequences of the *kn1*-like genes in the homeodomain, characterize their patterns of steady state mRNA expression, and report their chromosomal locations on the maize genetic map.

RESULTS

Isolation and Sequencing of *knox* Genes

To assess the approximate number of *kn1*-like homeobox genes present in the maize genome, a maize genomic DNA gel blot was prepared and hybridized at low stringency to a probe consisting of only the homeobox (see Methods). The blot was stripped and reprobated at high stringency with a gene-specific probe from the third intron of *kn1*. Figure 1 shows that while the gene-specific probe hybridizes to only one band (Figure 1B), the homeobox probe hybridizes to 10 to 15 sequences in the maize genome at low stringency (Figure 1A). The large number of bands suggests that the homeobox in *kn1* represents one member of a fairly large gene family. We propose the name *knox* (for *knotted*-like homeobox) to distinguish this gene family from other types of homeobox genes.

Genomic and cDNA libraries were screened with the *kn1* homeobox-specific probe at low stringency. After several new clones were isolated and characterized, the libraries were




Figure 1. Maize Genomic DNA Gel Blot Analysis.

(A) Genomic DNA from the inbred maize lines CM37 (C) and T232 (T), digested with either EcoRI or BamHI, was run on an agarose gel, blotted, and probed with the *kn1* homeobox (box) at low stringency. (B) The blot in (A) was stripped and reprobated with a gene-specific probe from the third intron of *kn1* at high stringency.

Each of the *knox* genes has an intron in an absolutely conserved position within the homeobox near the N-terminal end of the second helix (solid arrow, Figure 2). In addition, the *knox* genes in the class 2 subgroup share another small intron in the adjacent ELK region (open arrow, Figure 2). The positions of the introns interrupting the plant homeoboxes are distinct from the positions of introns reported in animal or fungal homeoboxes (for a catalog of homeobox intron positions, see Allen et al., 1991).

Analysis of *knox* Gene Expression

kn1 is highly expressed in the vegetative meristem and inflorescence primordia of maize and undetectable in more determinate organs, such as leaves, roots, and floral organs (Smith et al., 1992). To investigate *knox* gene expression in plant tissues, both total RNA and poly(A)⁺-enriched RNA were isolated from various maize tissues. RNA blots were prepared using 1 μg of poly(A)⁺-enriched RNA per lane and were hybridized with DNA probes from each of the *knox* genes for which cDNA clones were isolated. All the *knox* genes examined by hybridization to RNA blots detected a single band between 1.5 and 1.7 kb in length. The class 1 genes *kn1*, *rs1*, *knox3*, *knox4*, and *knox8* were all found to be strongly expressed in ear inflorescence primordia and shoot meristem-enriched tissue, moderately to weakly expressed in embryos, and not detected in leaves, as shown in Figure 3. A subset, *rs1* and *knox4*, were weakly expressed in roots. The expression of *knox5* was similar to *rs1* and *knox4* on total RNA blots (data not shown), and *knox10*, *knox11*, and *lg3* were not extensively tested because of the inadequacy of the probes available for use in RNA gel blot analysis.

By contrast, the class 2 *knox* genes showed greater diversity in their patterns of expression. The *knox1* gene was weakly expressed in all tissues (data not shown) and strongly expressed in roots. *knox2* appeared to be expressed at a significant level in all tissues examined, although its expression level was strongest in leaves. *knox6*, like *knox2*, was strongly expressed in all the tissues examined but showed somewhat stronger expression in meristematic tissues. The pattern of *knox7* expression was indistinguishable from that of *knox6* (data not shown). Accurate quantitative comparisons between signals on poly(A)⁺-enriched RNA from different tissues were difficult because of potential sample-to-sample variation in the efficiency of poly(A)⁺ enrichment. In practice, the patterns of expression seen with 1 μg of poly(A)⁺-enriched RNA shown in Figure 3 closely mirrored the patterns seen with 10 μg of total RNA (data not shown).

RNA gel blot hybridizations were carried out at high stringency to prevent cross-hybridization of related mRNAs. Each probe was shown to be specific by preparing slot blots containing RNA synthesized with T3 polymerase from each of the other KNOX clones. Duplicate slot blots were then hybridized with probes derived from each KNOX clone. With 0.01 ng of RNA loaded per slot, self-hybridization was at least 100-fold




Figure 3. RNA Gel Blot Analysis of *knox* Genes.

RNA gel blots were prepared from 1 μg each of poly(A)⁺-enriched RNA collected from embryos 17 days postpollination (lanes E), shoot meristems including unexpanded stem and the youngest leaf primordia (less than 0.5 cm) (lanes M), ear inflorescence primordia up to 2 cm long (lanes I), primary roots from seeds germinated in air (lanes R), and the blade portion of fully expanded juvenile leaves (lanes L). Blots were probed with gene-specific probes for a subset of class 1 *knox* genes (left) and class 2 *knox* genes (right). The approximate length of each transcript is indicated below the gene designation. Reprobing with the maize ubiquitin cDNA (UBI1) allowed the quantity of RNA in each lane to be compared.

greater than hybridization to any other clone (data not shown). The KNOX4 probe is an exception. It was generated from a short genomic clone and showed only ~20-fold greater hybridization to itself than to the RS1 cDNA clone; although in the reciprocal hybridization, the RS1 cDNA probe did not hybridize significantly to the KNOX4 clone (data not shown). Differences in the patterns of expression and minor differences in the sizes of the bands revealed with different *knox* gene probes also indicate that cross-hybridization was not significant.

Chromosomal Map Locations of *knox* Sequences

The *knox* genes were mapped on recombinant inbred lines according to the method of Burr and coworkers (Burr et al., 1988). Approximate map positions are shown in Figure 4. Chromosomes 2, 3, 4, 7, and 9 have one *knox* gene, chromosome 5 has two *knox* genes, and chromosomes 1 and 8 contain small clusters of three and two linked *knox* genes, respectively. Only

chromosomes 6 and 10 remain without any mapped *knox* gene. Four pairs of closely related but unlinked *knox* genes (*knox1* and *knox2*, *knox6* and *knox7*, *knox4* and *rs1*, and *lg3* and *knox5* or *knox11*) map in regions of the genome that share a number of duplicate genes (Helentjaris et al., 1988).

At least three *knox* genes map to the long arm of chromosome 1. Within the resolution of the CO159 × Tx303 recombinant inbred mapping population, *knox3* maps to the same location as *kn1*, and *knox8* resides ~14 centimorgans distal to *kn1* and *knox3*. To examine further the linkage of *kn1* and *knox3*, we took

advantage of a deletion that completely removes the *kn1* locus but not the closely linked markers *alcohol dehydrogenase1* and *lemon-white*. The deletion, *def(kn1)-O*, is not transmitted through the male gametophyte (Hake et al., 1989; Veit et al., 1990) and results in a defective embryo when uncovered by a much larger deficiency (L. Smith and S. Hake, unpublished data). In Figure 5, a gene-specific probe for *knox3* detects a restriction fragment length polymorphism (RFLP) between the progenitor of the deletion (*Kn1-O*) and a wild-type line (+). We crossed the heterozygote carrying *Kn1-O* on one chromosome and




Figure 4. Genome Map Positions of *knox* Genes.

knox gene sequences were mapped on recombinant inbred lines according to the method of Burr et al. (1988). Segments of chromosomes or chromosome arms (S, short arm; L, long arm) are represented by solid lines, with centromeric regions indicated by a thickened line. Map distances are in centimorgans. Framework loci are shown in bold; loci in normal type on the same line are linked to the nearest framework locus by the distance indicated. RFLP markers and distances, taken from the current genome map maintained by E. Matz, F. Burr, and B. Burr (personal communication), convey the linkage relationships of the *knox* genes. The dashed lines connect two RFLP loci that hybridize to a single probe and may reflect ancient duplications of regions of the genome (Helentjaris et al., 1988). The pairs of genes *knox6* and *knox7*, *lg3* and *knox11* (or *knox5*), and *knox4* and *rs1* are similar in sequence and expression and fall within these duplicated regions. Some map segments were inverted from the conventional orientation to simplify comparisons between putative duplicated regions.




Figure 5. A Deletion of *kn1* Also Removes *knox3*.

The *knox3* probe detected a polymorphism between *Kn1-O*, the progenitor of a deletion that completely removes the *kn1* locus, and a wild-type allele (+) when DNA was digested with BamHI. The probe detected only the (+) polymorphic band in DNA of plants carrying both the deletion and the wild-type (+) allele (lane marked deletion/+). The absence of a *knox3* band associated with the deletion chromosome suggests that this locus is also missing in the deletion. Genotypes at the *kn1* locus are indicated at the top of each lane: + indicates a wild-type allele, deletion is *def(kn1)-O*, and *Kn1-O* is a dominant mutant allele and progenitor of *def(kn1)-O*.

def(kn1)-O (deletion) on the homologous chromosome by the wild-type line and analyzed the DNA of the knotted and normal progeny. Both RFLPs were clearly observed in the knotted (*Kn1-O/+*) progeny; however, the normal progeny (*def(kn1)-O/+*) showed only the wild-type RFLP. The absence of a *knox3* band associated with the deletion chromosome suggests that the locus is also missing in the deletion, although the precise extent of the deletion remains uncertain.

DISCUSSION

The *kn1* class of homeobox genes in maize (*knox* genes) comprise a family of ~13 genes that map to eight of the 10 chromosomes. The genes all contain a conserved intron position and a three-amino acid insertion within the homeodomain. The third helix of the homeodomain is invariant. The similarity extends outside the homeodomain to include an adjacent region, the ELK domain, that contains repeating hydrophobic residues. The family can be divided into two subclasses based on sequence similarity within the homeodomain and patterns of expression at the level of RNA gel blots. The first class is more like KN1, sharing 73 to 89% amino acid identity in the homeodomain and being expressed primarily in shoot and inflorescence meristems but not in leaves. The second class differs from KN1 in the extent of sequence similarity and pattern of expression. These proteins share only 55 to 58% identity with KN1 in the homeodomain, and they appear to be expressed at varying levels in all tissues, including mature leaves.

kn1-like homeobox genes have recently been reported in rice (Matsuoka et al., 1993) and identified in several dicot species, including Arabidopsis (Lincoln et al., 1994), tomato (N. Sinha and S. Hake, unpublished data), and soybean (Ma et al., 1994). The sequence similarity within the homeodomain, regardless of species, is high. Sequence conservation among *kn1*-like genes across monocot and dicot plant species suggests that there is functional significance that has been conserved over evolutionary time. As shown in Figure 6, other classes of plant homeodomains are more divergent in sequence. Most of these other proteins are also distinguished by conserved features outside of the homeodomain, such as leucine zippers in the HD-Zip proteins (Ruberti et al., 1991; Mattsson et al., 1992; Schena and Davis, 1992, 1994) and putative metal binding domains in the PHD-finger proteins (Bellmann and Werr, 1992; Schindler et al., 1993; Korfhage et al., 1994). The recognition helix of the GLABRA2 protein in Arabidopsis is somewhat similar to the HD-Zip class of homeodomains, but the protein lacks a leucine zipper, so it represents a distinct class of plant homeodomain protein.

Mapping of the *knox* genes on the maize genome reveals several interesting features. Whereas two or three genetically linked homeobox genes are found on chromosomes 1 and 8, there is no indication of extensive clusters of homeobox genes as has been observed with the HOM-C or Hox clusters of homeobox genes in animal genomes (Krumlauf, 1992). The closely linked maize genes encode similar homeodomains and may be the result of small duplications that have subsequently diverged.

A number of researchers have shown that the maize genome contains large numbers of duplicated genes that are distributed in a nonrandom manner such that certain pairs of chromosomes share many duplicated genes (Wendel et al., 1986; Helentjaris et al., 1988). Helentjaris and coworkers reported




Figure 6. Analysis of Plant Homeodomain Sequence Similarity.

The dendrogram reflects the degree of similarity between deduced amino acid sequences from various plant homeodomains using the unweighted pair group maximum averages method (Sneath and Sokal, 1973). Percent similarity between homeodomain sequences is indicated by the scale at the bottom. The maize *kn1*-like homeodomains are in bold type. Homeobox genes similar to *kn1* from other species include *OSH1* from rice (Matsuoka et al., 1993), *sbh1* from soybean (Ma et al., 1994), the *KNAT* genes in Arabidopsis (Lincoln et al., 1994), and *thb6* and *thb12* from tomato (N. Sinha, J. Mathern, and S. Hake, unpublished data). The two classes of KNOX homeodomains are easily distinguished and appear to have diverged prior to the separation of monocots and dicots. The other types of homeobox genes reported in plants fall into completely different classes: the HD-Zip class; *Athb-1*, *Athb-2* (Ruberti et al., 1991; these homeodomains have the same sequence as *HAT5* and *HAT4*, respectively; Schena and Davis, 1992), *Athb-3* (Mattsson et al., 1992), and *Athb-4* (Carabelli et al., 1993); the *GLABRA2* (*GL2*) class (Rerie et al., 1994); the PHD-finger class; *ZMHOX1A*, *ZMHOX1B* (Bellmann and Werr, 1992), *HAT3.1* (Schindler et al., 1993), *PRHP*, and *PRHA* (Korthage et al., 1994).

that ~29% of the 200 single-copy RFLP probes that they used detected an unlinked, duplicate sequence. Because related duplications are found in other grass species, such as sorghum (Berhan et al., 1993), the modern maize genome may be the result of an ancient duplication or allopolyploidy that occurred early in the evolution of the grass supertribe Andropogonodae. Certain pairs of *knox* genes, such as *knox4* and *rs1*, show significantly more similarity to each other than to the rest of the gene family and map within these duplicate regions (Figures 4 and 6). These results suggest that these *knox* gene pairs may have once been duplicate genes.

All of the KN1-like homeodomains examined in plants share an identical series of 13 amino acids in the third helix. This portion of the homeodomain is thought to be largely responsible for much of the sequence-specific contact with DNA (for review, see Kornberg, 1993). Identical recognition sequences may indicate that different proteins recognize the same set of downstream targets or that additional specificity is provided by differences in other parts of the homeodomain, by other domains of the proteins, or by interactions with other proteins that help to specify DNA recognition. Much attention has been given to the problem of specificity in target DNA recognition (for reviews, see Affolter et al., 1990a; Hayashi and Scott, 1990). It appears that amino acid sequences within the homeodomain, apart from the recognition helix, can determine target specificity or influence the ability of a particular protein to form a specific protein-DNA complex. In *Drosophila* for example, the homeotic genes *Antp* and *Sex combs reduced* (*Scr*) encode homeodomains that differ at only four residues in the N-terminal part of the homeodomain, yet they condition different phenotypes when expressed ectopically under the heat shock promoter (Zeng et al., 1993). Domain-swapping experiments have revealed that the N-terminal region of the homeodomain is sufficient to distinguish the functional specificity of the SCR protein from that of ANTP in vivo, although a few features are influenced by sequences outside of the homeodomain (Zeng et al., 1993). The mammalian octamer motif binding factors, Oct-1 and Oct-2, share little similarity apart from the POU and homeodomains, yet swapping experiments have demonstrated that the ability of Oct-1 to form a DNA binding complex with the herpes simplex virus *trans*-inducing factor, VP16, depends on only a few amino acid differences in helix 2 of the homeodomain (Stern et al., 1989). Clearly, the recognition helix may not be the only determinant of downstream target specificity. When combined with sequence differences that lie outside of the homeodomain that may mediate differential interactions with the transcriptional machinery or with specific *trans*-acting factors, there exists ample opportunity for proteins with quite similar homeodomains to regulate different sets of target genes specifically.

Expression of the five class 1 *knox* genes examined on RNA gel blots is high in meristem-enriched tissues and inflorescence primordia but not detectable in leaves. Because the meristem-enriched region includes the youngest leaf primordia, RNA gel blot analysis does not address when expression disappears during leaf development. In situ hybridization analysis has shown that expression in the meristem of *rs1*, *knox3*, and *knox8*,

like *kn1*, is exclusive of leaf primordia (Jackson et al., 1994) even at the very earliest stages of leaf initiation. The staining patterns indicate overlapping but distinct domains of expression within the shoot apical meristem. *kn1* and *knox8* are expressed throughout the central region of the shoot apical meristem in a similar pattern, although the staining of *kn1* is stronger than that of *knox8*. By contrast, *rs1* and *knox3* expression is limited to a subset of meristem cells (Jackson et al., 1994). All four genes are also expressed in the unexpanded stem, and the patterns of expression are quite similar in the inflorescence primordium.

The *kn1* locus was originally defined by dominant mutations that affect leaf development (Bryan and Sass, 1941; Gelinis et al., 1969; Freeling and Hake, 1985). The mutations, which are either transposon insertions or rearrangements of 5' sequences (Hake, 1992), result in ectopic expression of *kn1* in lateral veins of developing leaves (Smith et al., 1992). Localized proliferations of cells around lateral veins result in outgrowths or knots on the leaf blade. There is also an overall reduction in height of the plant and a broadening of the leaf. Cells along the lateral veins in the blade adopt fates of cells normally found in more basal regions of the leaf (Sinha and Hake, 1994). Transgenic tobacco plants expressing the maize *kn1* cDNA under the cauliflower mosaic virus 35S promoter are also affected in plant height and leaf shape in a dosage-dependent manner (Sinha et al., 1993). In the most severely affected plants, leaf cells adopt meristem fates and form ectopic shoots.

Two of the class 1 *knox* genes have been shown to correspond to existing morphological mutations that affect the maize plant in a manner similar to *kn1*. The dominant *Rs1* mutation (P. Becraft, R. Schneeberger, S. Hake, and M. Freeling, unpublished data) and the dominant *Lg3* mutation (J. Fowler and M. Freeling, unpublished data) correspond to homeobox genes that are closely related to *kn1* (Figures 2 and 6). Several other *knox* sequences show tight genetic linkage to known mutations, including *knox11* and *knox5* to *Liguleless4* (J. Fowler and M. Freeling, unpublished data) and *knox4* to the morphological mutation *Gnarley1* (T. Foster and S. Hake, unpublished data). All of these dominant mutations alter cell fates in the maize leaf in a manner analogous to *Kn1* but appear to affect different tissues specifically. For example, whereas *Kn1* primarily affects the lateral veins, *Rs1* affects the ligule/auricle region (Becraft and Freeling, 1994) and *Lg3* affects primarily the midrib region (for review of these mutations, see Freeling, 1992). Because the *Kn1* mutations result from ectopic expression in leaves where the gene is not normally expressed, it is likely that the dominant mutations of the other *kn1*-like genes also result from ectopic expression.

Although these class 1 genes are similar in sequence, expression patterns, and dominant phenotypes and in some cases map within regions that may have arisen as part of an ancient duplication, they are not necessarily redundant. Despite many examples of duplicate homeobox genes, gene redundancy may not be common. For example, in mice there are four copies of the *Hox* gene cluster located on four separate chromosomes (Hunt and Krumlauf, 1992). Targeted gene

disruptions of the paralogous genes from two different clusters, *Hoxd-3* and *Hoxa-3*, produce different phenotypes despite the fact that the genes have identical expression patterns (Chisaka and Capecchi, 1991; Condie and Capecchi, 1993). Recent data from our laboratory suggest that *kn1* is not redundant, as we have found a recessive inflorescence phenotype (R. Kerstetter, L. Smith, and S. Hake, unpublished data). Similar experiments are underway for related *kn1*-like genes (M. Freeling, personal communication). Exactly what role each *knox* gene plays in development and how their activities are coordinated await analysis of such recessive alleles. Examination of the patterns of expression and mutant phenotypes of the *kn1*-like homeobox genes in dicot species should also serve to elucidate further the role of these genes in plant development.

METHODS

DNA Isolation and Blot Hybridization

Maize genomic DNA was isolated as described by Chen and Dellaporta (1994). DNA (7 μ g per lane) was digested with the appropriate endonucleases, separated on 0.8% agarose gels, and transferred to Magna nylon transfer membranes (Micron Separations Inc., Westboro, MA). Low-stringency hybridization was performed in $9 \times$ SSC (1 \times SSC is 0.15 M NaCl, 15 mM sodium citrate), 2 mM EDTA, 10 mM Tris, pH 7.5, 5 \times Denhardt's solution (1 \times Denhardt's solution is 0.02% Ficoll, 0.02% polyvinylpyrrolidone, 0.02% BSA), 50 μ g/mL sheared salmon sperm DNA, 20 mM sodium phosphate, pH 7, 2% SDS at 60°C, and filters were washed in $2 \times$ SSC, 0.2% SDS twice for 1 hr at the same temperature. Normal high-stringency genomic DNA gel blot hybridizations were performed in a hybridization oven (Robbin's Scientific, Sunnyvale, CA) according to the recommendations of the manufacturer and washed in $0.2 \times$ SSC, 0.2% SDS twice for 1 hr at 68°C. Probes were labeled by the random prime method as suggested by the kit manufacturer (Amersham International). The gene-specific probe for *knotted1* (*kn1*) is H2, which was derived from a HindIII fragment from the third intron of the gene (Veit et al., 1990). The homeobox-specific probe was obtained by polymerase chain reaction amplification of the homeobox region of the *kn1* cDNA using the primers E35 and E40.

Genomic and cDNA Libraries

Genomic and cDNA libraries were screened with the homeobox-specific probe at reduced stringency under conditions similar to those described above for low-stringency DNA gel blot hybridizations. The genomic libraries were prepared from size-selected SacI fragments from *Kn1-N2* DNA in λ ZAPII according to the protocol supplied by Stratagene and from DNA partially digested with Sau3A in EMBL3 according to the protocols supplied by Stratagene. We screened cDNA libraries made from poly(A)⁺-enriched mRNA from whole seedlings (provided by A. Barkan, University of Oregon, Eugene, OR) and from poly(A)⁺-enriched mRNA isolated from ear primordia (Schmidt et al., 1993).

RNA Isolation and Blot Hybridization

Total RNA from various maize tissues was prepared as described by Smith et al. (1992). Poly(A)⁺ enrichment was performed using the Poly Attract kit from Promega. Ten micrograms of total RNA or 1 μ g of poly(A)⁺-enriched RNA per lane was glyoxalated, separated on an

agarose gel, transferred to Magna nylon transfer membranes, and hybridized as previously described (Smith et al., 1992). Blots were stripped and reprobed as many as five times. Autoradiograms were exposed for different amounts of time. The clones used to probe RNA gel blots were complete or partial maize cDNAs, except for KNOX4, and are as follows; KN1, a 1.6-kb cDNA clone; KNOX8, a 0.6-kb cDNA clone; RS1, an 0.8-kb cDNA clone; KNOX3, a 0.9-kb cDNA clone; KNOX4, a polymerase chain reaction product derived from a genomic clone that includes the ELK region, homeobox, a 127-bp intron, and 70 bp of the 3' end of the gene; KNOX1, a 1.0-kb cDNA clone; KNOX2, a 1.0-kb cDNA clone; KNOX6, a 0.7-kb cDNA clone; and UBI1, a 0.65-kb PstI-SacI subclone of the maize ubiquitin cDNA clone (Christensen and Quail, 1989).

The specificity of hybridization of the KNOX probes used on RNA gel blots was checked on slot blots. Each KNOX clone was transcribed into RNA using T3 RNA polymerase and then treated with DNase to remove the DNA template (Sambrook et al., 1989). Three different quantities (100, 1.0, and 0.01 ng) of each RNA were loaded in a MinifoldII slot blotting device (Schleicher & Schuell) and transferred to a nylon transfer membrane according to Sambrook et al. (1989). The RNA and membrane were UV cross-linked and then baked for 1 hr at 80°C. The hybridization conditions were identical to those used for RNA gel blots.

Mapping Genes on Maize Recombinant Inbred Lines

Genomic DNA was isolated from leaves of two maize recombinant inbred (RI) families (CO159 \times Tx303 and T232 \times CM37) grown from seed provided by B. Burr (Brookhaven National Laboratory, Upton, NY). Survey gel blots were prepared with 7 to 10 μ g of DNA per lane from each of the inbred parents restricted with various endonucleases. The blots were probed with gene-specific pieces derived from either the 3' end of cDNA clones or the intron dividing the homeobox of genomic clones to identify appropriate polymorphisms. Combinations of RI family, restriction endonuclease, and probe that yielded useful polymorphisms were then used in the preparation of DNA gel blots containing the 41 or 48 RI individuals. The allelic distribution was then scored and sent to B. Burr for comparison with the distribution of all previously mapped loci (Burr et al., 1988). The linkage information returned was used to position the genes on the maize genome map, portions of which are shown in Figure 6.

Oligonucleotide Synthesis

Oligonucleotides were synthesized on a DNA synthesizer (model 380B; Applied Biosystems, Foster City, CA). Primers for amplification of the *kn1* homeobox are as follows: E35, 5'-CTCGCTCAAGCAAGAAGCTGTC-3'; E40, 5'-TCCATCATCAGGTGGTGCAT-3'. Degenerate oligonucleotides for sequencing are as follows: E44, 5'-CARATKAAYAAYTGTTTATY-AAYCARMG-3'; E61, 5'-CAYNNNAAGTGGCCNTAYCC-3'; E62, 5'-CTTCKYTGRTRATRAACCA-3', where R = A or G, Y = C or T, K = G or T, M = A or C, and N = A or G or C or T.

DNA Sequencing and Analysis

Double-stranded DNA sequencing was performed using the Sequenase Version 2 kit (U.S. Biochemical Corp.) according to the manufacturer's protocols. DNA and putative protein sequences were analyzed using the PCGENE program and the IG Molecular Biology Software System Release 5.4 (IntelliGenetics, Mountain View, CA). The dendrogram in Figure 6 was generated using the CLUSTAL program in PCGENE.

ACKNOWLEDGMENTS

We would like to thank Alice Barkan for sharing her maize seedling cDNA library; Lauren Hubbard for mapping *knox4*; Neelima Sinha, Cindy Lincoln, Kyle Serikawa, Toshi Foster, Richard Schneeberger, Philip Becraft, John Fowler, and Michael Freeling for sharing unpublished results; and Eileen Matz, Francis Burr, and Ben Burr (Department of Biology, Brookhaven National Lab, Upton, NY) for permitting us to use portions of the maize recombinant inbred molecular map. We would also like to thank Peter Quail for critical comments on the manuscript. The work was supported by the U.S. Department of Agriculture Grant No. NRICGP9103192.

Received June 8, 1994; accepted October 4, 1994.

REFERENCES

- Affolter, M., Scheir, A., and Gehring, W.J.** (1990a). Homeodomain proteins and the regulation of gene expression. *Curr. Opin. Cell Biol.* **2**, 485–495.
- Affolter, M., Percival-Smith, A., Muller, M., Leupin, W., and Gehring, W.J.** (1990b). DNA binding properties of the purified Antennapedia homeodomain. *Proc. Natl. Acad. Sci. USA* **87**, 4093–4097.
- Alberti, S., Oehler, S., von Wilcken-Bergmann, B., and Muller-Hill, B.** (1993). Genetic analysis of the leucine heptad repeats of the *Lac* repressor: Evidence for a 4-helical bundle. *EMBO J.* **12**, 3227–3236.
- Allen, J.D., Lints, T., Jenkins, N.A., Copeland, N.G., Strasser, A., Harvey, R.P., and Adams, J.M.** (1991). Novel murine homeobox gene on chromosome 1 expressed in specific hematopoietic lineages and during embryogenesis. *Genes Dev.* **5**, 509–520.
- Becraft, P.W., and Freeling, M.** (1994). Genetic analysis of *Rough sheath1* developmental mutants of maize. *Genetics* **136**, 295–311.
- Bellmann, R., and Werr, W.** (1992). *Zmhox1a*, the product of a novel maize homeobox gene, interacts with the *Shrunken* 26-bp *feedback* control element. *EMBO J.* **11**, 3367–3374.
- Berhan, A.M., Hulbert, S.H., Butler, L.G., and Bennetzen, J.L.** (1993). Structure and evolution of the genomes of *Sorghum bicolor* and *Zea mays*. *Theor. Appl. Genet.* **86**, 598–604.
- Bryan, A.A., and Sass, J.E.** (1941). Heritable characters in maize. *J. Hered.* **32**, 343–346.
- Burr, B., Burr, F.A., Thompson, K.H., Albertson, M.C., and Stuber, C.W.** (1988). Gene mapping with recombinant inbreds in maize. *Genetics* **118**, 519–526.
- Carabelli, M., Sessa, G., Baima, S., Morelli, G., and Ruberti, I.** (1993). The *Arabidopsis Athb-2* and *-4* genes are strongly induced by far-red-rich light. *Plant J.* **4**, 469–479.
- Chen, J., and Dellaporta, S.** (1994). Urea-based plant DNA miniprep. In *The Maize Handbook*, M. Freeling and V. Walbot, eds (New York: Springer-Verlag), pp. 526–538.
- Chisaka, O., and Capecchi, M.R.** (1991). Regionally restricted developmental defects resulting from targeted disruption of the mouse homeobox gene *hox-1.5*. *Nature* **350**, 473–479.
- Christensen, A.H., and Quail, P.H.** (1989). Sequence analysis and transcriptional regulation by heat shock of polyubiquitin transcripts from maize. *Plant Mol. Biol.* **12**, 619–632.
- Condie, B.G., and Capecchi, M.R.** (1993). Mice homozygous for a targeted disruption of *Hoxd-3* (*Hox-4.1*) exhibit anterior transformations of the first and second cervical vertebrae, the atlas and the axis. *Development* **119**, 579–595.
- Freeling, M.** (1992). A conceptual framework for maize leaf development. *Dev. Biol.* **153**, 44–58.
- Freeling, M., and Hake, S.** (1985). Developmental genetics of mutants that specify *Knotted* leaves in maize. *Genetics* **111**, 617–634.
- Gehring, W.J.** (1987). Homeo boxes in the study of development. *Science* **236**, 1245–1252.
- Gelinas, D., Postlethwait, S.N., and Nelson, O.E.** (1969). Characterization of development in maize through the use of mutants. II. The abnormal growth conditioned by the *Knotted* mutant. *Am. J. Bot.* **56**, 671–678.
- Hake, S.** (1992). Unraveling the knots in plant development. *Trends Genet.* **8**, 109–114.
- Hake, S., Vollbrecht, E., and Freeling, M.** (1989). Cloning *Knotted*, the dominant morphological mutant in maize using *Ds2* as a transposon tag. *EMBO J.* **8**, 15–22.
- Hanes, S.D., and Brent, R.** (1989). DNA specificity of the bicoid activator protein is determined by homeodomain recognition helix residue 9. *Cell* **57**, 1275–1283.
- Hayashi, S., and Scott, M.P.** (1990). What determines the specificity of action of Drosophila homeodomain proteins? *Cell* **63**, 883–894.
- Helentjaris, T., Weber, D., and Wright, S.** (1988). Identification of the genomic locations of duplicate nucleotide sequences in maize by analysis of restriction fragment length polymorphisms. *Genetics* **118**, 353–363.
- Herr, W., Sturm, R.A., Clerc, R.G., Corcoran, L.M., Baltimore, D., Sharp, P.A., Ingraham, H.A., Rosenfeld, M.G., Finney, M., Ruvkun, G., and Horvitz, H.R.** (1988). The POU domain: A large conserved region in the mammalian *pit-1*, *oct-1*, *oct-2*, and *Caenorhabditis elegans unc-86* gene products. *Genes Dev.* **2**, 1513–1516.
- Hunt, P., and Krumlauf, R.** (1992). Hox codes and positional specification in vertebrate embryonic axes. *Annu. Rev. Cell Biol.* **8**, 227–256.
- Jackson, D., Veit, B., and Hake, S.** (1994). Expression of maize *KNOTTED-1* related homeobox genes in the shoot apical meristem predicts patterns of morphogenesis in the vegetative shoot. *Development* **120**, 405–413.
- Kessel, M., and Gruss, P.** (1990). Murine developmental control genes. *Science* **249**, 374–379.
- Kissinger, C.R., Liu, B., Martin-Bianco, E., Kornberg, T. B., and Pabo, C.O.** (1990). Crystal structure of an engrailed homeodomain-DNA complex at 2.8 Å resolution: A framework for understanding homeodomain-DNA interactions. *Cell* **63**, 579–590.
- Korfhage, U., Trezzini, G.F., Meier, I., Hahlbrock, K., and Somssich, I.E.** (1994). Plant homeodomain protein involved in transcriptional regulation of a pathogen defense-related gene. *Plant Cell* **6**, 695–708.
- Kornberg, T.B.** (1993). Understanding the homeodomain. *J. Biol. Chem.* **268**, 26813–26816.
- Krumlauf, R.** (1992). Evolution of the vertebrate Hox homeobox genes. *BioEssays* **14**, 245–252.
- Landschulz, W.H., Johnson, P.F., and McKnight, S.L.** (1988). The leucine zipper: A hypothetical structure common to a new class of DNA binding proteins. *Science* **240**, 1759–1764.
- Laughon, A., and Scott, M.P.** (1984). Sequence of a *Drosophila* segmentation gene: Protein structure homology with DNA-binding proteins. *Nature* **310**, 25–31.

- Lincoln, C., Long, J., Yamaguchi, J., Serikawa, K., and Hake, S.** (1994). A *knotted1*-like homeobox gene in *Arabidopsis* is expressed in the vegetative meristem and dramatically alters leaf morphology when overexpressed in transgenic plants. *Plant Cell* **6**, 1859–1876.
- Ma, H., McMullen, M.D., and Finer, J.J.** (1994). Identification of a homeobox-containing gene with enhanced expression during soybean (*Glycine max* L.) somatic embryo development. *Plant Mol. Biol.* **24**, 465–473.
- Matsuoka, M., Ichikawa, H., Saito, A., Tada, Y., Fujimura, T., and Kano-Murakami, Y.** (1993). Expression of a rice homeobox gene causes altered morphology of transgenic plants. *Plant Cell* **5**, 1039–1048.
- Mattsson, J., Soderman, E., Svenson, M., Borkind, C., and Engstrom, R.** (1992). A new homeobox-leucine zipper gene from *Arabidopsis thaliana*. *Plant Mol. Biol.* **18**, 1019–1022.
- McGinnis, W., Levine, M.S., Hafen, E., Kuroiwa, A., and Gehring, W.J.** (1984). A conserved DNA sequence in homeotic genes of the *Drosophila* Antennapedia and bithorax complexes. *Nature* **308**, 428–433.
- Murre, C., McCaw, P.S., and Baltimore, D.** (1989). A new DNA binding and dimerization motif in immunoglobulin enhancer binding, *daughterless*, *MyoD*, and *myc* proteins. *Cell* **56**, 777–783.
- Otting, G., Qian, Y.Q., Billeter, M., Muller, M., Affolter, M., Gehring, W.J., and Wuthrich, K.** (1990). Protein-DNA contacts in the structure of a homeodomain-DNA complex determined by nuclear magnetic resonance spectroscopy in solution. *EMBO J.* **9**, 3085–3092.
- Percival-Smith, A., Muller, M., Affolter, M., and Gehring, W.J.** (1990). The interaction with DNA of wild-type and mutant *fushi tarazu* homeodomains. *EMBO J.* **9**, 3967–3974.
- Qian, Y.Q., Billeter, M., Otting, G., Muller, M., Gehring, W.J., and Wuthrich, K.** (1989). The structure of the Antennapedia homeodomain determined by NMR spectroscopy in solution: Comparison with prokaryotic repressors. *Cell* **59**, 573–580.
- Rerie, G.W., Feldmann, K.A., and Marks, M.D.** (1994). The *GLABRA2* gene encodes a homeodomain protein required for normal trichome development in *Arabidopsis*. *Genes Dev.* **8**, 1388–1399.
- Ruberti, I., Sessa, G., Lucchetti, S., and Morelli, G.** (1991). A novel class of plant proteins containing a homeodomain with a closely linked leucine zipper motif. *EMBO J.* **10**, 1787–1791.
- Sambrook, J., Fritsch, E.F., and Maniatis, T.** (1989). *Molecular Cloning: A Laboratory Manual*. 2nd edition. (Cold Spring Harbor, NY: Cold Spring Harbor Press).
- Schena, M., and Davis, R.W.** (1992). HD-Zip proteins: Members of an *Arabidopsis* homeodomain protein superfamily. *Proc. Natl. Acad. Sci. USA* **89**, 3894–3898.
- Schena, M., and Davis, R.W.** (1994). Structure of homeobox-leucine zipper genes suggests a model for the evolution of gene families. *Proc. Natl. Acad. Sci. USA* **91**, 8393–8397.
- Schindler, U., Beckmann, H., and Cashmore, A.R.** (1993). HAT3.1, a novel *Arabidopsis* homeodomain protein containing a conserved cysteine-rich region. *Plant J.* **4**, 137–150.
- Schmidt, R.J., Veit, B., Mandel, M.A., Mena, M., Hake, S., and Yanofsky, M.F.** (1993). Identification and molecular characterization of *ZAG1*, the maize homolog of the *Arabidopsis* floral homeotic gene *AGAMOUS*. *Plant Cell* **5**, 729–737.
- Scott, M.P., and Weiner, A.J.** (1984). Structural relationships among genes that control development: Sequence homology between the *Antennapedia*, *Ultrabithorax*, and *fushi tarazu* loci of *Drosophila*. *Proc. Natl. Acad. Sci. USA* **81**, 4115–4119.
- Scott, M.P., Tamkun, J.W., and Hartzell, G.W., III.** (1989). The structure and function of the homeodomain. *Biochim. Biophys. Acta* **989**, 25–48.
- Shepherd, J.C.W., McGinnis, W., Carrasco, A.E., De Robertis, E.M., and Gehring, W.J.** (1984). Fly and frog homeo domains show homologies with yeast mating type regulatory genes. *Nature* **310**, 70–71.
- Sinha, N., and Hake, S.** (1994). The *Knotted1* leaf blade is a mosaic of blade, sheath and auricle identities. *Dev. Genet.* **15**, 401–414.
- Sinha, N., Williams, R., and Hake, S.** (1993). Overexpression of the maize homeobox gene, *KNOTTED-1*, causes a switch from determinate to indeterminate cell fates. *Genes Dev.* **7**, 787–795.
- Smith, L., Greene, B., Veit, B., and Hake, S.** (1992). A dominant mutation in the maize homeobox gene, *Knotted-1*, causes its ectopic expression in leaf cells with altered fates. *Development* **116**, 21–30.
- Sneath, P.H.A., and Sokal, R.R.** (1973). *Numerical Taxonomy: The Principles and Practice of Numerical Classification*. (San Francisco: W.H. Freeman).
- Stern, S., Tanaka, M., and Herr, W.** (1989). The Oct-1 homeodomain directs formation of a multiprotein-DNA complex with the HSV transactivator VP16. *Nature* **341**, 624–630.
- Treisman, J., Gonczyk, P., Vashishtha, M., Harris, E., and Desplan, C.** (1989). A single amino acid can determine the DNA binding specificity of homeodomain proteins. *Cell* **59**, 553–562.
- Treisman, J., Harris, E., and Desplan, C.** (1991). The paired box encodes a second DNA-binding domain in the paired homeodomain protein. *Genes Dev.* **5**, 594–604.
- Veit, B., Vollbrecht, E., Mathern, J., and Hake, S.** (1990). A tandem duplication causes the *Kn1-O* allele of *Knotted*, a dominant morphological mutant of maize. *Genetics* **125**, 623–631.
- Vollbrecht, E., Veit, B., Sinha, N., and Hake, S.** (1991). The developmental gene *Knotted-1* is a member of a maize homeobox gene family. *Nature* **350**, 241–243.
- Vollbrecht, E., Kerstetter, R., Lowe, B., Veit, B., and Hake, S.** (1993). Homeobox genes in plant development: Mutational and molecular analysis. In *Evolutionary Conservation of Developmental Mechanisms*, A.C. Spadling, ed (New York: Wiley-Liss), pp. 111–123.
- Wendel, J.F., Stuber, C.W., Edwards, M.D., and Goodman, M.M.** (1986). Duplicated chromosome segments in maize (*Zea mays* L.): Further evidence from hexokinase isozymes. *Theor. Appl. Genet.* **72**, 178–185.
- Wolberger, C., Vershon, A.K., Liu, B., Johnson, A.D., and Pabo, C.O.** (1991). Crystal structure of a *MATa2* homeodomain-operator complex suggests a general model for homeodomain-DNA interactions. *Cell* **67**, 517–528.
- Zeng, W., Andrew, D.J., Mathies, L.D., Horner, M.A., and Scott, M.P.** (1993). Ectopic expression and function of the *Antp* and *Scr* homeotic genes: The N terminus of the homeodomain is critical to functional specificity. *Development* **118**, 339–352.