

Short communication

SEX HORMONE INFLUENCES ON YAWNING BEHAVIOR

Björn HOLMGREN, Ruth URBÁ-HOLMGREN, Margarita AGUIAR
and Renzo RODRIGUEZ

Department of Neurophysiology, National Center for Scientific Research
Havana, Cuba

Abstract. Young male albino rats yawn significantly more than females or castrated males when injected with physostigmine (0.10 mg/kg). Treatment with testosterone (100 µg daily) during seven days restores cholinomimetically induced yawning in castrated males, and increases yawning in normal and androgenized females. Treatment with estradiol (200 µg daily) during one week does not modify physostigmine induced yawning behavior in castrated males nor in androgenized females.

Although yawning is a simple behavioral item not obviously related to the sexual sphere, several reports have appeared pointing towards a curious association between yawning and some elements of sexual behavior. Injections of ACTH, MSH, or peptides possessing similar hormonal activity, into the cerebrospinal fluid or the brain, induce a peculiar stretching and yawning syndrome (SYS) in several animal species (6, 7). The same treatment also produces a display of sexual excitement, both in male (2, 3) and female rabbits (1). Associated yawning and penile erections have been observed during and between waves of cortical spreading depression in male rats (4). Jakobartl and Huston (16) have recently suggested that both the SYS and the penile erections observed after intracranial injections of ACTH₁₋₂₄ (and other peptides) could be after-effects of spreading depression produced by these substances. With low doses of apomorphine and other dopaminergic agonists injected systemically, Mogilnicka and Klimek (18) have also

observed associated yawning with penile erections in adult rats, observation which we have confirmed (unpublished results).

Even if little or no sex differences exist in the SYS induced by ACTH₁₋₂₄ (1), the release of yawning behavior in monkeys in response to their mirror images seems to be sex-bound: Hall (10) mentioned that females in this experimental situation rarely "yawned". Cholinomimetically-induced yawning in adult rats also seems to be sex dependent: males injected with physostigmine yawn significantly more than females (21). In the present report we describe some experiments which suggest that cholinergic yawning is subject to distinct sex hormone influences.

Young male albino rats, 30-day-old, injected intraperitoneally with physostigmine salicylate (0.10 mg/kg, as base), and observed during one hour as formerly described (13, 22), yawn significantly more than female littermates, or males castrated when 11 day old (Fig. 1). This age for testes ablation was selected so as not to interfere with the perinatal critical period of hormonal masculinizing effects on the CNS, and distant enough from the first test with physostigmine for the effects of




Fig. 1. Influence of castration and hormonal treatment on yawning. Each bar corresponds to a mean value obtained from 20–22 rats injected with physostigmine (0.10 mg/kg). ♂ males castrated when 11-day-old; T and E, same as above but treated with testosterone (T) or estradiol (E). Comparisons between normal males or castrated males treated with testosterone, against the other groups are significantly different at the level of $P < 0.001$ (Mann-Whitney U Test). The difference between normal males and testosterone-treated castrates is not significant.




Fig. 2. Hormonal influences on cholinomimetically induced yawning in female rats. Each bar represents the mean number of yawns after physostigmine administration. Treatment with testosterone (T) during one week (100 μ g daily) increases yawning significantly ($P < 0.001$) both in normal and androgenized females. The difference between testosterone treated normal and androgenized females is also significant ($P < 0.01$). One week treatment with estradiol benzoate (E) has no significant effect. 30, 37, age of animals in days.

castration to become evident. Castrated males treated with daily injections of testosterone (100 μg) from the 30th to the 37th day, are undistinguishable from intact males in their yawning response to the anticholinesterastic drug. Such restoration of yawning is not obtainable by treatment with estradiol benzoate (200 μg daily).

A single androgenizing (19) injection of testosterone (250 μg) to one-day-old female rats ($n = 16$) does not modify the physostigmine-induced yawning response tested on the 30th postnatal day, when compared to that of normal females ($n = 16$) (Fig. 2). A second test, performed on the 37th day, after one week treatment with testosterone (100 μg daily), demonstrated highly significantly increased yawning responses in both groups, in comparison to their responses in the first test ($P < 0.001$, Mann-Whitney U Test) (20). The difference in yawning frequency between normal and androgenized females after this treatment is also statistically significant ($P < 0.01$), the average number of yawns of the latter group being more than 100% higher than in the former, during one hour after physostigmine injection.

It is unlikely that so important quantitative differences in cholinomimetically induced yawning between male and female rats are due to simple pharmacokinetic effects, as have been described for other drugs. These are generally interpreted as the result of sex hormone influences upon microsomal enzymes participating in drug metabolism, androgens stimulating, while estrogens and progesterone inhibit microsomal enzymes activity (5). Therefore, the greater yawn promoting effect of physostigmine in male rats, compared to females, in testosterone treated castrated males compared to the untreated castrates, and in testosterone treated females compared to normal females, could hardly be understood as due to pharmacokinetic effects of testosterone on the metabolism of physostigmine. These would rather tend to reduce than to increase the anticholinesterasic action of physostigmine.

The greater yawning-inducing effect in testosterone treated neonatally androgenized females is also consistent with the idea that the differences observed are the result of true androgenic hormonal effects exerted upon the central nervous system, both at an early (perinatal) period of development, and later on in life. In the particular effect analyzed by us, testosterone treatment of one-day-old females seems to have increased their sensitivity to testosterone administered one month later. The level of the central nervous system at which testosterone might interact with the neural structures responsible for cholinomimetically induced yawning is unknown, and only broad hypothetical suggestions may be advanced. Because of the early ontogenetic maturation of cholinergic yawning (13, 21), it has been suggested that

the neural circuitry responsible for yawning behavior seems to be localized at the lower brain stem region. The observation of yawning in Gamper's mesencephalic human being (17) also points in that direction. This does not necessarily mean that the influence of testosterone is exerted at such a caudal level, because it might also be mediated by descending pathways from higher placed nervous sites, i.e. in the limbic system, well known as steroid hormone targets.

Although the described effect of testosterone is upon a behavioral pattern including important cholinergic links, the hormonal action might as well take place in cholinergic or cholin-sensitive neurons, or in neurons of some of the specific monoamine pathways, which could exert some modulating influences on yawning. Such alternatives should be considered when reports have appeared describing yawning in adult rats evoked by dopaminergic agonists (18), and a strong facilitation of physostigmine-induced yawning by Lu 10-171 (22), a potent and selective serotonin uptake inhibitor (15). Sex differences and the effect of gonadal steroids on brain serotonin levels, 5-HT synthesis or turnover have been the subject of research by several groups in recent years (4, 8, 9, 11, 12) but the results remain controversial.

Even if the functional significance of the above described hormonal effects on yawning are obscure, they do confirm the possibility envisaged by Young et al (23) that "gonadal substances may affect behavior beyond that which is primarily sexual in the sense of being directed solely towards the attainment of sexual aims".

1. BALDWIN, D. B., HAUN, C. K. and SAWYER, C. H. 1974. Effects of intraventricular infusion of ACTH⁴⁻¹⁰ on LH release, ovulation and behavior in the rat. *Brain Res.* 80: 291-301.
2. BERTOLINI, A., GESSA, G. L., VERGONI, W. and FERRARI, W. 1968. Induction of sexual excitement with intraventricular ACTH; permissive role of testosterone in the male rabbit. *Life Sci.* 7: 1203-1206.
3. BERTOLINI, A., VERGONI, W., GESSA, G. L. and FERRARI, W. 1969. Induction of sexual excitement by the action of adrenocorticotrophic hormone in brain. *Nature* 221: 667-669.
4. BOURGOIN, S., ARTAUD, F., ADRIEN, J., HERY, F., GLOWINSKI, J. and HAMON, F. 1977. 5-hydroxytryptamine catabolism in the rat brain during ontogenesis. *J. Neurochem.* 22: 951-956.
5. BOWMANN, W. C., RAND, M. J. and WEST, G. B. 1970. *Farmacología*. Editorial JIMS, Barcelona, España.
6. FERRARI, W., GESSA, G. L. and VARGIU, L. 1963. Behavioral effects induced by intracysternally injected ACTH and MSH. *Ann. N. Y. Acad. Sci.* 104: 330-343.
7. GESSA, G. L., PISANO, M., VARGIU, L., GRABAI, F. and FERRARI, W. 1969. Stretching and yawning movements after intracerebral injection of ACTH. *Rev. Canad. Biol.* 26: 229-236.

8. GIULIAN, A., POHORECKY, L. A. and MCEWEN, B. S. 1973. Effects of gonadal steroids upon brain 5-hydroxytryptamine levels in the neonatal rat. *Endocrinology* 93: 1329-1335.
9. GLADUE, B. A., HUMPHRYS, R. R., DEBOLD, J. F. and CLEMENS, L. G. 1977. Ontogeny of biogenic amine systems and modification of indole levels upon adult sexual behavior in the rat. *Pharm. Biochem. Behav.* 7: 253-258.
10. HALL, K. R. 1969. Behaviour of monkeys towards mirror-images. *Nature* 196: 1258-1261.
11. HAMON, M., BOURGOIN, S. and ENJALBERT, A. 1977. Biochemical regulatory mechanisms of serotonergic neurons in the rat brain. *Prog. Reprod. Biol.* 2: 25-40.
12. HARDIN, C. M. 1973. Sex differences in serotonin synthesis from 5-hydroxytryptophan in neonatal rat brain. *Brain Res.* 59: 437-439.
13. HOLMGREN, B. and URBA-HOLMGREN, R. 1978. Ontogeny of two cholinergically mediated central effects: stereotyped yawning and potentiation of head-shaking. *Acta Neurobiol. Exp.* 38: 11-23.
14. HUSTON, J. P. 1971. Yawning and penile erections induced in rats by cortical spreading depression. *Nature* 232: 274-275.
15. HYTTTEL, J. 1977. Neurochemical characterization of a new potent and selective serotonin uptake inhibitor: Lu 10-171. *Psychopharmacology* 51: 225-233.
16. JAKOBARTL, L. and HUSTON, J. P. 1977. Spreading depression in hippocampus and neocortex of rats induced by ACTH₁₋₂₄. *Neurosci. Lett.* 5: 180-192.
17. JUNG, R. and HASSLER, R. 1960. The extrapyramidal system. In J. Field (ed.), *Handbook of physiology. Section I. Neurophysiology. Vol. II.* Am. Physiol. Soc., Washington, p. 833-927.
19. MOGILNICKA, E. and KLIMEK, V. 1977. Drugs affecting dopamine neurons and yawning behavior. *Pharmacol. Biochem. Behav.* 7: 303-305.
19. PFAFF, D. W., DIAKOW, C., ZIGMOND, R. E. and KOW, L. W. 1974. Neural and hormonal determinants of female mating behavior in rats. In F. Schmitt and F. G. Worden (ed.), *The Neurosciences. Third Study Program.* The MIT Press, Cambridge, Mass., p. 621-646.
20. SIEGEL, S. 1956. *Nonparametric statistics for the behavioral sciences.* McGraw-Hill Co., New York, 312 p.
21. URBA-HOLMGREN, R., GONZALEZ, R. M. and HOLMGREN, B. 1977. Is yawning a cholinergic response? *Nature* 267: 261-262.
22. URBA-HOLMGREN, R., HOLMGREN, B., RODRIGUEZ, R. and GONZALEZ, R. M. 1979. Serotonergic modulation of yawning. *Pharmacol. Biochem. Behav.* 11: 371-372.
23. YOUNG, W. C., GOY, R. W. and PHOENIX, C. H. 1964. Hormones and sexual behavior. *Science* 143: 212-218.

Accepted 27 November 1979

Björn HOLMGREN, Ruth URBA-HOLMGREN, Margarita AGUIAR and Renzo RODRIGUEZ, Department of Neurophysiology, National Center for Scientific Research, Apartado 6990, Havana, Cuba.