



Shape and weight concern as a moderator of program outcomes from the Prevention Across the Spectrum RCT

Citation

Wilksch, Simon, Tracey Wade, Susan Paxton, Susan Byrne, and S Bryn Austin. 2015. "Shape and weight concern as a moderator of program outcomes from the Prevention Across the Spectrum RCT." *Journal of Eating Disorders* 3 [Suppl 1]: 028. doi:10.1186/2050-2974-3-S1-028. <http://dx.doi.org/10.1186/2050-2974-3-S1-028>.

Published Version

doi:10.1186/2050-2974-3-S1-028

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:23993482>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

ORAL PRESENTATION

Open Access

Shape and weight concern as a moderator of program outcomes from the Prevention Across the Spectrum RCT

Simon Wilksch^{1*}, Tracey Wade¹, Susan Paxton², Susan Byrne³, S Bryn Austin^{4,5}

From 2015 ANZAED Conference: Riding the Waves to Recovery
Surfers Paradise, Australia. 21-22 August 2015

Objective

To investigate if baseline shape and weight concern moderated outcomes in the Prevention Across the Spectrum trial, a randomised-controlled trial (RCT) of 3 school-based programs aimed at reducing eating disorder and obesity risk factors.

Method

N = 1,316 Grade 7 and 8 girls and boys (M age = 13.21 years) across three Australian states were randomly allocated to: Media Smart; Life Smart; Helping, Encouraging, Listening and Protecting Peers Initiative (HELPP) or control (usual school class). Risk factors were measured at baseline, post-program (5-weeks later), and 6- and 12-month follow-up.

Results

Moderation was indicated by significant interaction effects for group (Media Smart; Life Smart; HELPP; Control) X moderator (high shape and weight concern; low shape and weight concern) X time (post-program; 6-month follow-up; 12-month follow-up), with baseline entered as a covariate. Such effects were found for shape concern, weight concern, eating concern, regular eating, body dissatisfaction, and physical activity. Post-hoc testing found Media Smart participants with high baseline shape and weight concern experienced a reduction in risk at 12-month follow-up for 4 of the 6 variables.

Discussion

This study shows it is possible for a school-based program to reduce eating disorder risk factors in participants with high baseline risk of an eating disorder.

Authors' details

¹School of Psychology, Flinders University, Australia. ²School of Psychological Science, La Trobe University, Australia. ³School of Psychology, University of Western Australia, Australia. ⁴Boston Children's Hospital, Australia. ⁵Harvard Medical School & Harvard School of Public Health, Australia.

Published: 23 November 2015

doi:10.1186/2050-2974-3-S1-O28

Cite this article as: Wilksch et al.: Shape and weight concern as a moderator of program outcomes from the Prevention Across the Spectrum RCT. *Journal of Eating Disorders* 2015 **3**(Suppl 1):O28.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



* Correspondence: simon.wilksch@flinders.edu.au

¹School of Psychology, Flinders University, Australia

Full list of author information is available at the end of the article