

## SHARP MAXIMAL FUNCTION ESTIMATES AND BOUNDEDNESS FOR COMMUTATORS ASSOCIATED WITH GENERAL INTEGRAL OPERATOR

Lanzhe Liu

### Abstract

In this paper, we establish the sharp maximal function estimates for the commutator associated with some integral operator with general kernel and the weighted Lipschitz functions. As an application, we obtain the boundedness of the commutator on weighted Lebesgue, Morrey and Triebel-Lizorkin space. The operator includes Littlewood-Paley operators, Marcinkiewicz operators and Bochner-Riesz operator.

## 1 Introduction and Preliminaries

As the development of singular integral operators(see [12][28][29]), their commutators have been well studied(see [8][26][27]). In [8][26][27], the authors prove that the commutators generated by the singular integral operators and  $BMO$  functions are bounded on  $L^p(R^n)$  for  $1 < p < \infty$ . Chanillo (see [4]) proves a similar result when singular integral operators are replaced by the fractional integral operators. In [5][14][23], the boundedness for the commutators generated by the singular integral operators and Lipschitz functions on Triebel-Lizorkin and  $L^p(R^n)$ ( $1 < p < \infty$ ) spaces are obtained. In [1][13], the boundedness for the commutators generated by the singular integral operators and the weighted  $BMO$  and Lipschitz functions on  $L^p(R^n)$ ( $1 < p < \infty$ ) spaces are obtained. In [3], some singular integral operators with general kernel are introduced, and the boundedness for the operators and their commutators generated by  $BMO$  and Lipschitz functions are obtained(see [3][15]). The purpose of this paper is to prove the sharp maximal function estimates for the commutator associated with some integral operator with general kernel and the weighted Lipschitz functions. As an application, we obtain the boundedness of the commutator on weighted Lebesgue, Morrey and Triebel-Lizorkin space. The operator includes Littlewood-Paley operators, Marcinkiewicz operators and Bochner-Riesz operator.

---

2010 *Mathematics Subject Classifications*: 42B20, 42B25.

*Key words and Phrases*: Commutator; Littlewood-Paley operator; Marcinkiewicz operator; Bochner-Riesz operator; Sharp maximal function; Morrey space; Triebel-Lizorkin space.

Received: May 7, 2011

Communicated by Dragan S. Djordjević

First, let us introduce some notations. Throughout this paper,  $Q$  will denote a cube of  $R^n$  with sides parallel to the axes. For any locally integrable function  $f$ , the sharp maximal function of  $f$  is defined by

$$M^\#(f)(x) = \sup_{Q \ni x} \frac{1}{|Q|} \int_Q |f(y) - f_Q| dy,$$

where, and in what follows,  $f_Q = |Q|^{-1} \int_Q f(x) dx$ . It is well-known that (see [12][28])

$$M^\#(f)(x) \approx \sup_{Q \ni x} \inf_{c \in C} \frac{1}{|Q|} \int_Q |f(y) - c| dy.$$

Let

$$M(f)(x) = \sup_{Q \ni x} \frac{1}{|Q|} \int_Q |f(y)| dy.$$

For  $\eta > 0$ , let  $M_\eta(f)(x) = M(|f|^\eta)^{1/\eta}(x)$ .

For  $0 < \eta < 1$  and  $1 \leq r < \infty$ , set

$$M_{\eta,r}(f)(x) = \sup_{Q \ni x} \left( \frac{1}{|Q|^{1-r\eta/n}} \int_Q |f(y)|^r dy \right)^{1/r}.$$

The  $A_p$  weight is defined by (see [12]), for  $1 < p < \infty$ ,

$$A_p = \left\{ w \in L^1_{loc}(R^n) : \sup_Q \left( \frac{1}{|Q|} \int_Q w(x) dx \right) \left( \frac{1}{|Q|} \int_Q w(x)^{-1/(p-1)} dx \right)^{p-1} < \infty \right\}$$

and

$$A_1 = \{ w \in L^1_{loc}(R^n) : M(w)(x) \leq Cw(x), a.e. \}.$$

The  $A(p, r)$  weight is defined by (see [22]), for  $1 < p, r < \infty$ ,

$$A(p, r) = \left\{ w > 0 : \sup_Q \left( \frac{1}{|Q|} \int_Q w(x)^r dx \right)^{1/r} \left( \frac{1}{|Q|} \int_Q w(x)^{-p/(p-1)} dx \right)^{(p-1)/p} < \infty \right\}.$$

Given a non-negative weight function  $w$ . For  $1 \leq p < \infty$ , the weighted Lebesgue space  $L^p(w)$  is the space of functions  $f$  such that

$$\|f\|_{L^p(w)} = \left( \int_{R^n} |f(x)|^p w(x) dx \right)^{1/p} < \infty.$$

For  $\beta > 0$ ,  $p > 1$  and the non-negative weight function  $w$ , let  $\dot{F}_p^{\beta, \infty}(w)$  be the weighted homogeneous Triebel-Lizorkin space (see [2][23]).

For  $0 < \beta < 1$  and the non-negative weight function  $w$ , the weighted Lipschitz space  $Lip_\beta(w)$  is the space of functions  $b$  such that

$$\|b\|_{Lip_\beta(w)} = \sup_Q \frac{1}{w(Q)^{1+\beta/n}} \int_Q |b(y) - b_Q| dy < \infty.$$

**Remark.** (1). It has been known that, for  $b \in Lip_\beta(w)$ ,  $w \in A_1$  and  $x \in Q$ ,

$$|b_Q - b_{2^k Q}|_{Lip_\beta(w)} \leq Ck \|b\|_{Lip_\beta(w)} w(x) w(2^k Q)^{\beta/n}.$$

(2). Let  $b \in Lip_\beta(w)$  and  $w \in A_1$ . By [11], we know that spaces  $Lip_\beta(w)$  coincide and the norms  $\|b\|_{Lip_\beta(w)}$  are equivalent with respect to different values  $1 \leq p \leq \infty$ .

In this paper, we will study some integral operators as following (see [3]).

**Definition 1.** Let  $F_t(x, y)$  be defined on  $R^n \times R^n \times [0, +\infty)$  and  $b$  be a locally integrable function on  $R^n$ , set

$$F_t(f)(x) = \int_{R^n} F_t(x, y) f(y) dy$$

and

$$F_t^b(f)(x) = \int_{R^n} (b(x) - b(y)) F_t(x, y) f(y) dy$$

for every bounded and compactly supported function  $f$ .

Let  $H$  be the Banach space  $H = \{h : \|h\| < \infty\}$ . For each fixed  $x \in R^n$ , we view  $F_t(f)(x)$  and  $F_t^b(f)(x)$  as the mappings from  $[0, +\infty)$  to  $H$ . Set

$$T(f)(x) = \|F_t(f)(x)\|,$$

which  $T$  is bounded on  $L^2(R^n)$ . The commutator related to  $F_t^b$  is defined by

$$T^b(f)(x) = \|F_t^b(f)(x)\|$$

and  $F_t$  satisfies: there is a sequence of positive constant numbers  $\{C_k\}$  such that for any  $k \geq 1$ ,

$$\int_{2|y-z| < |x-y|} ( \|F_t(x, y) - F_t(x, z)\| + \|F_t(y, x) - F_t(z, x)\| ) dx \leq C,$$

and

$$\begin{aligned} & \left( \int_{2^k|z-y| \leq |x-y| < 2^{k+1}|z-y|} ( \|F_t(x, y) - F_t(x, z)\| + \|F_t(y, x) - F_t(z, x)\| )^q dy \right)^{1/q} \\ & \leq C_k (2^k|z-y|)^{-n/q'}, \end{aligned}$$

where  $1 < q' < 2$  and  $1/q + 1/q' = 1$ .

**Definition 2.** Let  $\varphi$  be a positive, increasing function on  $R^+$  and there exists a constant  $D > 0$  such that

$$\varphi(2t) \leq D\varphi(t) \quad \text{for } t \geq 0.$$

Let  $w$  be a non-negative weight function on  $R^n$  and  $f$  be a locally integrable function on  $R^n$ . Set, for  $1 \leq p < \infty$ ,

$$\|f\|_{L^{p,\varphi}(w)} = \sup_{x \in R^n, d > 0} \left( \frac{1}{\varphi(d)} \int_{Q(x,d)} |f(y)|^p w(y) dy \right)^{1/p},$$

where  $Q(x, d) = \{y \in R^n : |x - y| < d\}$ . The generalized weighted Morrey space is defined by

$$L^{p,\varphi}(R^n, w) = \{f \in L^1_{loc}(R^n) : \|f\|_{L^{p,\varphi}(w)} < \infty\}.$$

If  $\varphi(d) = d^\delta$ ,  $\delta > 0$ , then  $L^{p,\varphi}(R^n, w) = L^{p,\delta}(R^n, w)$ , which is the classical Morrey spaces (see [24][25]). If  $\varphi(d) = 1$ , then  $L^{p,\varphi}(R^n, w) = L^p(R^n, w)$ , which is the weighted Lebesgue spaces (see [6]).

As the Morrey space may be considered as an extension of the Lebesgue space, it is natural and important to study the boundedness of the operator on the Morrey spaces (see [6][9][10][16][21]).

It is well known that commutators are of great interest in harmonic analysis and have been widely studied by many authors (see [26][27]). In [27], Pérez and Trujillo-Gonzalez prove a sharp estimate for the multilinear commutator. The main purpose of this paper is to prove the sharp maximal inequalities for the commutator. As the application, we obtain the weighted  $L^p$ -norm inequality, Morrey and Triebel-Lizorkin spaces boundedness for the commutator.

## 2 Theorems

We shall prove the following theorems.

**Theorem 1.** Let  $T$  be the integral operator as **Definition 1**, the sequence  $\{kC_k\} \in l^1$ ,  $w \in A_1$ ,  $0 < \beta < 1$ ,  $q' \leq s < \infty$  and  $b \in Lip_\beta(w)$ . Then there exists a constant  $C > 0$  such that, for any  $f \in C_0^\infty(R^n)$  and  $\tilde{x} \in R^n$ ,

$$M^\#(T^b(f))(\tilde{x}) \leq C \|b\|_{Lip_\beta(w)} w(\tilde{x})^{1+\beta/n} (M_{\beta,s}(f)(\tilde{x}) + M_{\beta,s}(T(f))(\tilde{x})).$$

**Theorem 2.** Let  $T$  be the integral operator as **Definition 1**, the sequence  $\{k2^{\beta k}C_k\} \in l^1$ ,  $w \in A_1$ ,  $0 < \beta < 1$ ,  $q' \leq s < \infty$  and  $b \in Lip_\beta(w)$ . Then there exists a constant  $C > 0$  such that, for any  $f \in C_0^\infty(R^n)$  and  $\tilde{x} \in R^n$ ,

$$\begin{aligned} & \sup_{Q \ni \tilde{x}} \frac{1}{|Q|^{1+\beta/n}} \int_Q |T^b(f)(x) - C_0| dx \\ & \leq C \|b\|_{Lip_\beta(w)} w(\tilde{x})^{1+\beta/n} (M_s(f)(\tilde{x}) + M_s(T(f))(\tilde{x})). \end{aligned}$$

**Theorem 3.** Let  $T$  be the integral operator as **Definition 1**, the sequence  $\{kC_k\} \in l^1$ ,  $w \in A_1$ ,  $0 < \beta < \min(1, n/q')$ ,  $q' < p < n/\beta$ ,  $1/r = 1/p - \beta/n$  and  $b \in Lip_\beta(w)$ . Then  $T^b$  is bounded from  $L^p(w)$  to  $L^r(w^{r/p-r(1+\beta/n)})$ .

**Theorem 4.** Let  $T$  be the integral operator as **Definition 1**, the sequence  $\{kC_k\} \in l^1$ ,  $w \in A_1$ ,  $0 < D < 2^n$ ,  $0 < \beta < \min(1, n/q')$ ,  $q' < p < n/\beta$ ,  $1/r = 1/p - \beta/n$  and  $b \in Lip_\beta(w)$ . Then  $T^b$  is bounded from  $L^{p,\varphi}(w)$  to  $L^{r,\varphi}(w^{r/p-r(1+\beta/n)})$ .

**Theorem 5.** Let  $T$  be the integral operator as **Definition 1**, the sequence  $\{k2^{\beta k}C_k\} \in l^1$ ,  $w \in A_1$ ,  $0 < \beta < \min(1, n/q')$ ,  $q' < p < n/\beta$ ,  $1/r = 1/p - \beta/n$  and  $b \in Lip_\beta(w)$ . Then  $T^b$  is bounded from  $L^p(w)$  to  $\dot{F}_r^{\beta,\infty}(w^{r/p-r(1+\beta/n)})$ .

### 3 Proofs of Theorems

To prove the theorems, we need the following lemma.

**Lemma 1.**(see [3]) Let  $T$  be the integral operator as **Definition 1**. Then  $T$  is bounded on  $L^p(w)$  for  $w \in A_\infty$  with  $1 < p < \infty$ .

**Lemma 2.**(see [11][13]) For any cube  $Q$ ,  $b \in Lip_\beta(w)$ ,  $0 < \beta < 1$  and  $w \in A_1$ , we have

$$\sup_{x \in Q} |b(x) - b_Q| \leq C \|b\|_{Lip_\beta(w)} w(Q)^{1+\beta/n} |Q|^{-1}.$$

**Lemma 3.**(see [2][23]). For  $0 < \beta < 1$ ,  $1 < p < \infty$  and  $w \in A_\infty$ , we have

$$\begin{aligned} \|f\|_{\dot{F}_p^{\beta, \infty}(w)} &\approx \left\| \sup_{Q \ni \cdot} \frac{1}{|Q|^{1+\beta/n}} \int_Q |f(x) - f_Q| dx \right\|_{L^p(w)} \\ &\approx \left\| \sup_{Q \ni \cdot} \inf_c \frac{1}{|Q|^{1+\beta/n}} \int_Q |f(x) - c| dx \right\|_{L^p(w)}. \end{aligned}$$

**Lemma 4.**(see [12]). Let  $0 < p < \infty$  and  $w \in \cup_{1 \leq r < \infty} A_r$ . Then, for any smooth function  $f$  for which the left-hand side is finite,

$$\int_{R^n} M(f)(x)^p w(x) dx \leq C \int_{R^n} M^\#(f)(x)^p w(x) dx.$$

**Lemma 5.**(see [22]). Suppose that  $0 \leq \eta < n$ ,  $1 < s < p < n/\eta$ ,  $1/r = 1/p - \eta/n$  and  $w \in A(p, r)$ . Then

$$\|M_{\eta, s}(f)\|_{L^r(w^r)} \leq C \|f\|_{L^p(w^p)}.$$

**Lemma 6.**(see [7][12]) If  $w \in A_p$ , then  $w\chi_Q \in A_p$  for  $1 \leq p \leq \infty$  and any cube  $Q$ .

**Lemma 7.** Let  $1 < r < \infty$ ,  $0 < \eta < \infty$ ,  $0 < D < 2^n$ ,  $w \in A_\infty$  and  $L^{r, \varphi}(R^n, w)$  be the weighted Morrey space as **Definition 2**. Then, for any smooth function  $f$  for which the left-hand side is finite,

$$\|M(f)\|_{L^{r, \varphi}(w)} \leq C \|M^\#(f)\|_{L^{r, \varphi}(w)}.$$

**Proof.** Notice that  $w\chi_Q \in A_\infty$  for any cube  $Q = Q(x, d)$  by [8] and Lemma 6, thus, for  $f \in L^{r, \varphi}(R^n, w)$  and any cube  $Q$ , we have, by Lemma 4,

$$\begin{aligned} &\int_Q M(f)(x)^r w(x) dx = \int_{R^n} M(f)(x)^r w(x) \chi_Q(x) dx \\ &\leq C \int_{R^n} M^\#(f)(x)^r w(x) \chi_Q(x) dx \\ &= C \int_Q M^\#(f)(x)^r w(x) dx, \end{aligned}$$

thus

$$\left( \frac{1}{\varphi(d)} \int_{Q(x,d)} M(f)(x)^r w(x) dx \right)^{1/r} \leq C \left( \frac{1}{\varphi(d)} \int_{Q(x,d)} M^\#(f)(x)^r w(x) dx \right)^{1/r}$$

and

$$\|M(f)\|_{L^{r,\varphi}(w)} \leq C \|M^\#(f)\|_{L^{r,\varphi}(w)}.$$

This finishes the proof.

**Lemma 8.** Let  $1 < p < \infty$ ,  $0 < D < 2^n$ ,  $w \in A_1$ ,  $T$  be the integral operator as **Definition 1** and  $L^{p,\varphi}(R^n, w)$  be the weighted Morrey space as **Definition 2**. Then

$$\|T(f)\|_{L^{p,\varphi}(w)} \leq C \|f\|_{L^{p,\varphi}(w)}.$$

**Lemma 9.** Let  $0 < D < 2^n$ ,  $1 \leq s < p < n/\eta$ ,  $1/r = 1/p - \eta/n$ ,  $w \in A(p, r)$  and  $L^{p,\varphi}(R^n, w)$  be the weighted Morrey space as **Definition 2**. Then

$$\|M_{\eta,s}(f)\|_{L^{p,\varphi}(w^r)} \leq C \|f\|_{L^{p,\varphi}(w^p)}.$$

The proofs of two Lemmas are similar to that of Lemma 7 by Lemma 1 and 5, we omit the details.

**Proof of Theorem 1.** It suffices to prove for  $f \in C_0^\infty(R^n)$  and some constant  $C_0$ , the following inequality holds:

$$\frac{1}{|Q|} \int_Q |T^b(f)(x) - C_0| dx \leq C \|b\|_{Lip_\beta(w)} w(\tilde{x})^{1+\beta/n} (M_{\beta,s}(f)(\tilde{x}) + M_{\beta,s}(T(f))(\tilde{x})).$$

Fix a cube  $Q = Q(x_0, d)$  and  $\tilde{x} \in Q$ . Write, for  $f_1 = f\chi_{2Q}$  and  $f_2 = f\chi_{(2Q)^c}$ ,

$$F_t^b(f)(x) = (b(x) - b_{2Q})F_t(f)(x) - F_t((b - b_{2Q})f_1)(x) - F_t((b - b_{2Q})f_2)(x).$$

Then

$$\begin{aligned} & \frac{1}{|Q|} \int_Q \|F_t^b(f)(x) - F_t((b_{2Q} - b)f_2)(x_0)\| dx \\ & \leq \frac{1}{|Q|} \int_Q \|(b(x) - b_{2Q})F_t(f)(x)\| dx + \frac{1}{|Q|} \int_Q \|F_t((b - b_{2Q})f_1)(x)\| dx \\ & \quad + \frac{1}{|Q|} \int_Q \|F_t((b - b_{2Q})f_2)(x) - F_t((b - b_{2Q})f_2)(x_0)\| dx \\ & = I_1 + I_2 + I_3. \end{aligned}$$

For  $I_1$ , by Hölder's inequality and Lemma 2, we obtain

$$\begin{aligned} I_1 & \leq \frac{C}{|Q|} \sup_{x \in 2Q} |b(x) - b_{2Q}| |Q|^{1-1/s} \left( \int_Q |T(f)(x)|^s dx \right)^{1/s} \\ & \leq C \|b\|_{Lip_\beta(w)} \frac{w(2Q)^{1+\beta/n}}{|2Q|} |Q|^{-1/s} |Q|^{1/s-\beta/n} \left( \frac{1}{|Q|^{1-s\beta/n}} \int_Q |T(f)(x)|^s dx \right)^{1/s} \\ & \leq C \|b\|_{Lip_\beta(w)} \left( \frac{w(Q)}{|Q|} \right)^{1+\beta/n} M_{\beta,s}(T(f))(\tilde{x}) \\ & \leq C \|b\|_{Lip_\beta(w)} w(\tilde{x})^{1+\beta/n} M_{\beta,s}(T(f))(\tilde{x}). \end{aligned}$$

For  $I_2$ , by the boundedness of  $T$ , we get

$$\begin{aligned}
I_2 &\leq \left( \frac{1}{|Q|} \int_{R^n} |T((b - b_{2Q})f_1)(x)|^s dx \right)^{1/s} \\
&\leq C \left( \frac{1}{|Q|} \int_{R^n} |(b(x) - b_{2Q})f_1(x)|^s dx \right)^{1/s} \\
&\leq C|Q|^{-1/s} \sup_{x \in 2Q} |b(x) - b_{2Q}| |2Q|^{1/s-\beta/n} \left( \frac{1}{|2Q|^{1-s\beta/n}} \int_{2Q} |f(x)|^s dx \right)^{1/s} \\
&\leq C \|b\|_{Lip_\beta(w)} \left( \frac{w(Q)}{|Q|} \right)^{1+\beta/n} M_{\beta,s}(f)(\tilde{x}) \\
&\leq C \|b\|_{Lip_\beta(w)} w(\tilde{x})^{1+\beta/n} M_{\beta,s}(f)(\tilde{x}).
\end{aligned}$$

For  $I_3$ , recalling that  $s > q'$ , we have

$$\begin{aligned}
I_3 &\leq \frac{1}{|Q|} \int_Q \int_{(2Q)^c} |b(y) - b_{2Q}| |f(y)| |F_t(x, y) - F_t(x_0, y)| dy dx \\
&\leq \frac{1}{|Q|} \int_Q \sum_{k=1}^{\infty} \int_{2^k d \leq |y-x_0| < 2^{k+1} d} \|F_t(x, y) - F_t(x_0, y)\| |b(y) - b_{2^{k+1}Q}| |f(y)| dy dx \\
&\quad + \frac{1}{|Q|} \int_Q \sum_{k=1}^{\infty} \int_{2^k d \leq |y-x_0| < 2^{k+1} d} \|F_t(x, y) - F_t(x_0, y)\| |b_{2^{k+1}Q} - b_{2Q}| |f(y)| dy dx \\
&\leq \frac{C}{|Q|} \int_Q \sum_{k=1}^{\infty} \left( \int_{2^k d \leq |y-x_0| < 2^{k+1} d} \|F_t(x, y) - F_t(x_0, y)\|^q dy \right)^{1/q} \\
&\quad \times \sup_{y \in 2^{k+1}Q} |b(y) - b_{2^{k+1}Q}| \left( \int_{2^{k+1}Q} |f(y)|^{q'} dy \right)^{1/q'} dx \\
&\quad + \frac{C}{|Q|} \int_Q \sum_{k=1}^{\infty} |b_{2^{k+1}Q} - b_{2Q}| \left( \int_{2^k d \leq |y-x_0| < 2^{k+1} d} \|F_t(x, y) - F_t(x_0, y)\|^q dy \right)^{1/q} \\
&\quad \times \left( \int_{2^{k+1}Q} |f(y)|^{q'} dy \right)^{1/q'} dx \\
&\leq C \sum_{k=1}^{\infty} C_k (2^k d)^{-n/q'} \frac{w(2^{k+1}Q)^{1+\beta/n}}{|2^{k+1}Q|} \|b\|_{Lip_\beta(w)} |2^{k+1}Q|^{1/q'-1/s} |2^{k+1}Q|^{1/s-\beta/n} \\
&\quad \times \left( \frac{1}{|2^{k+1}Q|^{1-s\beta/n}} \int_{2^{k+1}Q} |f(y)|^s dy \right)^{1/s} \\
&\quad + C \sum_{k=1}^{\infty} k \|b\|_{Lip_\beta(w)} w(\tilde{x}) w(2^k Q)^{\beta/n} C_k (2^k d)^{-n/q'} |2^{k+1}Q|^{1/q'-1/s} |2^{k+1}Q|^{1/s-\beta/n} \\
&\quad \times \left( \frac{1}{|2^{k+1}Q|^{1-s\beta/n}} \int_{2^{k+1}Q} |f(y)|^s dy \right)^{1/s}
\end{aligned}$$

$$\begin{aligned}
&\leq C \|b\|_{Lip_\beta(w)} \sum_{k=1}^{\infty} C_k \left( \frac{w(2^{k+1}Q)}{|2^{k+1}Q|} \right)^{1+\beta/n} M_{\beta,s}(f)(\tilde{x}) \\
&\quad + C \|b\|_{Lip_\beta(w)} w(\tilde{x}) \sum_{k=1}^{\infty} k C_k \left( \frac{w(2^{k+1}Q)}{|2^{k+1}Q|} \right)^{\beta/n} M_{\beta,s}(f)(\tilde{x}) \\
&\leq C \|b\|_{Lip_\beta(w)} w(\tilde{x})^{1+\beta/n} M_{\beta,s}(f)(\tilde{x}) \sum_{k=1}^{\infty} (k+1) C_k \\
&\leq C \|b\|_{Lip_\beta(w)} w(\tilde{x})^{1+\beta/n} M_{\beta,s}(f)(\tilde{x}).
\end{aligned}$$

These complete the proof of Theorem 1.

**Proof of Theorem 2.** It suffices to prove for  $f \in C_0^\infty(R^n)$  and some constant  $C_0$ , the following inequality holds:

$$\frac{1}{|Q|^{1+\beta/n}} \int_Q |T^b(f)(x) - C_0| dx \leq C \|b\|_{Lip_\beta(w)} w(\tilde{x}) (M_s(f)(\tilde{x}) + M_s(T(f))(\tilde{x})).$$

Fix a cube  $Q = Q(x_0, d)$  and  $\tilde{x} \in Q$ . Write, for  $f_1 = f\chi_{2Q}$  and  $f_2 = f\chi_{(2Q)^c}$ ,

$$F_t^b(f)(x) = (b(x) - b_{2Q})F_t(f)(x) - F_t((b - b_{2Q})f_1)(x) - F_t((b - b_{2Q})f_2)(x).$$

Then

$$\begin{aligned}
&\frac{1}{|Q|^{1+\beta/n}} \int_Q \|F_t^b(f)(x) - F_t((b_{2Q} - b)f_2)(x_0)\| dx \\
&\leq \frac{1}{|Q|^{1+\beta/n}} \int_Q \|(b(x) - b_{2Q})F_t(f)(x)\| dx + \frac{1}{|Q|^{1+\beta/n}} \int_Q \|F_t((b - b_{2Q})f_1)(x)\| dx \\
&\quad + \frac{1}{|Q|^{1+\beta/n}} \int_Q \|F_t((b - b_{2Q})f_2)(x) - F_t((b - b_{2Q})f_2)(x_0)\| dx \\
&= I_1 + I_2 + I_3.
\end{aligned}$$

By using the same argument as in the proof of Theorem 1, we get

$$\begin{aligned}
I_1 &\leq \frac{C}{|Q|^{1+\beta/n}} \sup_{x \in 2Q} |b(x) - b_{2Q}| |Q|^{1-1/s} \left( \int_Q |T(f)(x)|^s dx \right)^{1/s} \\
&\leq C \|b\|_{Lip_\beta(w)} \frac{w(2Q)^{1+\beta/n}}{|2Q|} |Q|^{-1/s} |Q|^{1/s-\beta/n} \left( \frac{1}{|Q|} \int_Q |T(f)(x)|^s dx \right)^{1/s} \\
&\leq C \|b\|_{Lip_\beta(w)} \left( \frac{w(Q)}{|Q|} \right)^{1+\beta/n} M_s(T(f))(\tilde{x}) \\
&\leq C \|b\|_{Lip_\beta(w)} w(\tilde{x})^{1+\beta/n} M_s(T(f))(\tilde{x}),
\end{aligned}$$



$$\begin{aligned}
I_2 &\leq \frac{1}{|Q|^{1+\beta/n}} |Q|^{1-1/s} \left( \int_{R^n} |T((b-b_{2Q})f_1)(x)|^s dx \right)^{1/s} \\
&\leq C \frac{1}{|Q|^{1+\beta/n}} |Q|^{1-1/s} \left( \int_{R^n} |(b(x)-b_{2Q})f_1(x)|^s dx \right)^{1/s} \\
&\leq C \frac{1}{|Q|^{1+\beta/n}} |Q|^{1-1/s} \sup_{x \in 2Q} |b(x)-b_{2Q}| |2Q|^{1/s} \left( \frac{1}{|2Q|} \int_{2Q} |f(x)|^s dx \right)^{1/s} \\
&\leq C \|b\|_{Lip_\beta(w)} \left( \frac{w(Q)}{|Q|} \right)^{1+\beta/n} M_s(f)(\tilde{x}) \\
&\leq C \|b\|_{Lip_\beta(w)} w(\tilde{x})^{1+\beta/n} M_s(f)(\tilde{x}), \\
I_3 &\leq \frac{1}{|Q|^{1+\beta/n}} \int_Q \int_{(2Q)^c} |b(y)-b_{2Q}| |f(y)| \|F_t(x,y)-F_t(x_0,y)\| dy dx \\
&\leq \frac{1}{|Q|^{1+\beta/n}} \int_Q \sum_{k=1}^{\infty} \int_{2^k d \leq |y-x_0| < 2^{k+1} d} \|F_t(x,y)-F_t(x_0,y)\| |b(y)-b_{2^{k+1}Q}| |f(y)| dy dx \\
&\quad + \frac{1}{|Q|^{1+\beta/n}} \int_Q \sum_{k=1}^{\infty} \int_{2^k d \leq |y-x_0| < 2^{k+1} d} \|F_t(x,y)-F_t(x_0,y)\| |b_{2^{k+1}Q}-b_{2Q}| |f(y)| dy dx \\
&\leq \frac{C}{|Q|^{1+\beta/n}} \int_Q \sum_{k=1}^{\infty} \left( \int_{2^k d \leq |y-x_0| < 2^{k+1} d} \|F_t(x,y)-F_t(x_0,y)\|^q dy \right)^{1/q} \\
&\quad \times \sup_{y \in 2^{k+1}Q} |b(y)-b_{2^{k+1}Q}| \left( \int_{2^{k+1}Q} |f(y)|^{q'} dy \right)^{1/q'} dx \\
&\quad + \frac{C}{|Q|^{1+\beta/n}} \int_Q \sum_{k=1}^{\infty} |b_{2^{k+1}Q}-b_{2Q}| \left( \int_{2^k d \leq |y-x_0| < 2^{k+1} d} \|F_t(x,y)-F_t(x_0,y)\|^q dy \right)^{1/q} \\
&\quad \times \left( \int_{2^{k+1}Q} |f(y)|^{q'} dy \right)^{1/q'} dx \\
&\leq C |Q|^{-\beta/n} \sum_{k=1}^{\infty} C_k (2^k d)^{-n/q'} \frac{w(2^{k+1}Q)^{1+\beta/n}}{|2^{k+1}Q|} \|b\|_{Lip_\beta(w)} |2^{k+1}Q|^{1/q'} \\
&\quad \times \left( \frac{1}{|2^{k+1}Q|} \int_{2^{k+1}Q} |f(y)|^s dy \right)^{1/s} \\
&\quad + C |Q|^{-\beta/n} \sum_{k=1}^{\infty} k \|b\|_{Lip_\beta(w)} w(\tilde{x}) w(2^k Q)^{\beta/n} C_k (2^k d)^{-n/q'} |2^{k+1}Q|^{1/q'} \\
&\quad \times \left( \frac{1}{|2^{k+1}Q|} \int_{2^{k+1}Q} |f(x)|^s dx \right)^{1/s} \\
&\leq C \|b\|_{Lip_\beta(w)} \sum_{k=1}^{\infty} 2^{\beta k} C_k \left( \frac{w(2^{k+1}Q)}{|2^{k+1}Q|} \right)^{1+\beta/n} M_s(f)(\tilde{x}) \\
&\quad + C \|b\|_{Lip_\beta(w)} w(\tilde{x}) \sum_{k=1}^{\infty} k 2^{\beta k} C_k \left( \frac{w(2^{k+1}Q)}{|2^{k+1}Q|} \right)^{\beta/n} M_s(f)(\tilde{x}) \\
&\leq C \|b\|_{Lip_\beta(w)} w(\tilde{x})^{1+\beta/n} M_s(f)(\tilde{x}) \sum_{k=1}^{\infty} (k+1) 2^{\beta k} C_k \\
&\leq C \|b\|_{Lip_\beta(w)} w(\tilde{x})^{1+\beta/n} M_s(f)(\tilde{x}).
\end{aligned}$$

This completes the proof of Theorem 2.

**Proof of Theorem 3.** Choose  $q' < s < p$  in Theorem 1, notice  $w^{r/p-r(1+\beta/n)} \in A_\infty$  and  $w^{1/p} \in A(p, r)$ , we have, by Lemma 1, 4 and 5,

$$\begin{aligned}
& \|T^b(f)\|_{L^r(w^{r/p-r(1+\beta/n)})} \\
& \leq \|M(T^b(f))\|_{L^r(w^{r/p-r(1+\beta/n)})} \\
& \leq C\|M^\#(T^b(f))^\#\|_{L^r(w^{r/p-r(1+\beta/n)})} \\
& \leq C\|b\|_{Lip_\beta(w)}(\|M_{\beta,s}(T(f))w^{1+\beta/n}\|_{L^r(w^{r/p-r(1+\beta/n)})} \\
& \quad + \|M_{\beta,s}(f)w^{1+\beta/n}\|_{L^r(w^{r/p-r(1+\beta/n)})}) \\
& = C\|b\|_{Lip_\beta(w)}(\|M_{\beta,s}(T(f))\|_{L^r(w^{r/p})} + \|M_{\beta,s}(f)\|_{L^r(w^{r/p})}) \\
& \leq C\|b\|_{Lip_\beta(w)}(\|T(f)\|_{L^p(w)} + \|f\|_{L^p(w)}) \\
& \leq C\|b\|_{Lip_\beta(w)}\|f\|_{L^p(w)}.
\end{aligned}$$

This completes the proof of Theorem 3.

**Proof of Theorem 4.** Choose  $q' < s < p$  in Theorem 1, notice  $w^{r/p-r(1+\beta/n)} \in A_\infty$  and  $w^{1/p} \in A(p, r)$ , we have, by Lemma 7-9,

$$\begin{aligned}
& \|T^b(f)\|_{L^{r,\varphi}(w^{r/p-r(1+\beta/n)})} \\
& \leq \|M(T^b(f))\|_{L^{r,\varphi}(w^{r/p-r(1+\beta/n)})} \\
& \leq C\|M^\#(T^b(f))^\#\|_{L^{r,\varphi}(w^{r/p-r(1+\beta/n)})} \\
& \leq C\|b\|_{Lip_\beta(w)}(\|M_{\beta,s}(T(f))w^{1+\beta/n}\|_{L^{s,\varphi}(w^{r/p-r(1+\beta/n)})} \\
& \quad + \|M_{\beta,s}(f)w^{1+\beta/n}\|_{L^{r,\varphi}(w^{r/p-r(1+\beta/n)})}) \\
& = C\|b\|_{Lip_\beta(w)}(\|M_{\beta,s}(T(f))\|_{L^{r,\varphi}(w^{r/p})} + \|M_{\beta,s}(f)\|_{L^{r,\varphi}(w^{r/p})}) \\
& \leq C\|b\|_{Lip_\beta(w)}(\|T(f)\|_{L^{p,\varphi}(w)} + \|f\|_{L^{p,\varphi}(w)}) \\
& \leq C\|b\|_{Lip_\beta(w)}\|f\|_{L^{p,\varphi}(w)}.
\end{aligned}$$

This completes the proof of Theorem 4.

**Proof Theorem 5.** Choose  $q' < s < p$  in Theorem 2, notice that  $w^{r/p-r(1+\beta/n)} \in A_\infty$  and  $w^{1/p} \in A(p, r)$ . By using Lemma 3, we obtain

$$\begin{aligned}
& \|T^b(f)\|_{\dot{F}_r^{\beta,\infty}(w^{r/p-r(1+\beta/n)})} \\
& \leq C \left\| \sup_{Q \in \mathfrak{D}} \frac{1}{|Q|^{1+\beta/n}} \int_Q |T^b(f)(x) - T((b_{2Q} - b)f_2)(x_0)| dx \right\|_{L^r(w^{r/p-r(1+\beta/n)})} \\
& \leq C\|b\|_{Lip_\beta(w)}(\|M_s(T(f))w^{1+\beta/n}\|_{L^r(w^{r/p-r(1+\beta/n)})} \\
& \quad + \|M_s(f)w^{1+\beta/n}\|_{L^r(w^{r/p-r(1+\beta/n)})}) \\
& = C\|b\|_{Lip_\beta(w)}(\|M_s(T(f))\|_{L^r(w^{r/p})} + \|M_s(f)\|_{L^r(w^{r/p})}) \\
& \leq C\|b\|_{Lip_\beta(w)}(\|T(f)\|_{L^p(w)} + \|f\|_{L^p(w)}) \\
& \leq C\|b\|_{Lip_\beta(w)}\|f\|_{L^p(w)}.
\end{aligned}$$

This completes the proof of the theorem.

## 4 Applications

In this section we shall apply the Theorems 1-5 of the paper to some particular operators such as the Littlewood-Paley operators, Marcinkiewicz operator and Bochner-Riesz operator.

**Application 1.** Littlewood-Paley operators.

Fixed  $\varepsilon > 0$  and  $\mu > (3n + 2)/n$ . Let  $\psi$  be a fixed function which satisfies:

- (1)  $\int_{\mathbb{R}^n} \psi(x) dx = 0$ ,
- (2)  $|\psi(x)| \leq C(1 + |x|)^{-(n+1)}$ ,
- (3)  $|\psi(x + y) - \psi(x)| \leq C|y|^\varepsilon(1 + |x|)^{-(n+1+\varepsilon)}$  when  $2|y| < |x|$ ;

We denote  $\Gamma(x) = \{(y, t) \in \mathbb{R}_+^{n+1} : |x - y| < t\}$  and the characteristic function of  $\Gamma(x)$  by  $\chi_{\Gamma(x)}$ . The Littlewood-Paley commutators are defined by

$$g_\psi^b(f)(x) = \left( \int_0^\infty |F_t^b(f)(x)|^2 \frac{dt}{t} \right)^{1/2},$$

$$S_\psi^b(f)(x) = \left[ \int \int_{\Gamma(x)} |F_t^b(f)(x, y)|^2 \frac{dy dt}{t^{n+1}} \right]^{1/2}$$

and

$$g_\mu^b(f)(x) = \left[ \int \int_{\mathbb{R}_+^{n+1}} \left( \frac{t}{t + |x - y|} \right)^{n\mu} |F_t^b(f)(x, y)|^2 \frac{dy dt}{t^{n+1}} \right]^{1/2},$$

where

$$F_t^b(f)(x) = \int_{\mathbb{R}^n} (b(x) - b(y)) \psi_t(x - y) f(y) dy,$$

$$F_t^b(f)(x, y) = \int_{\mathbb{R}^n} (b(x) - b(z)) f(z) \psi_t(y - z) dz$$

and  $\psi_t(x) = t^{-n} \psi(x/t)$  for  $t > 0$ . Set  $F_t(f)(y) = f * \psi_t(y)$ . We also define

$$g_\psi(f)(x) = \left( \int_0^\infty |F_t(f)(x)|^2 \frac{dt}{t} \right)^{1/2},$$

$$S_\psi(f)(x) = \left( \int \int_{\Gamma(x)} |F_t(f)(y)|^2 \frac{dy dt}{t^{n+1}} \right)^{1/2}$$

and

$$g_\mu(f)(x) = \left( \int \int_{\mathbb{R}_+^{n+1}} \left( \frac{t}{t + |x - y|} \right)^{n\mu} |F_t(f)(y)|^2 \frac{dy dt}{t^{n+1}} \right)^{1/2},$$

which are the Littlewood-Paley operators (see [29]). Let  $H$  be the space

$$H = \left\{ h : \|h\| = \left( \int_0^\infty |h(t)|^2 dt/t \right)^{1/2} < \infty \right\}$$

or

$$H = \left\{ h : \|h\| = \left( \int \int_{R_+^{n+1}} |h(y, t)|^2 dy dt / t^{n+1} \right)^{1/2} < \infty \right\},$$

then, for each fixed  $x \in R^n$ ,  $F_t^b(f)(x)$  and  $F_t^b(f)(x, y)$  may be viewed as the mapping from  $[0, +\infty)$  to  $H$ , and it is clear that

$$g_\psi^b(f)(x) = \|F_t^b(f)(x)\|, \quad g_\mu(f)(x) = \|F_t(f)(x)\|,$$

$$S_\psi^b(f)(x) = \|\chi_{\Gamma(x)} F_t^b(f)(x, y)\|, \quad S_\psi(f)(x) = \|\chi_{\Gamma(x)} F_t(f)(y)\|$$

and

$$g_\mu^b(f)(x) = \left\| \left( \frac{t}{t + |x - y|} \right)^{n\mu/2} F_t^b(f)(x, y) \right\|, \quad g_\mu(f)(x) = \left\| \left( \frac{t}{t + |x - y|} \right)^{n\mu/2} F_t(f)(y) \right\|.$$

It is easily to see that  $g_\psi^b$ ,  $S_\psi^b$  and  $g_\mu^b$  satisfy the conditions of Theorem 1-5 (see [17-19]), thus theorem 1-5 hold for  $g_\psi^b$ ,  $S_\psi^b$  and  $g_\mu^b$ .

**Application 2.** Marcinkiewicz operators.

Fixed  $\lambda > \max(1, 2n/(n+2))$  and  $0 < \gamma \leq 1$ . Let  $\Omega$  be homogeneous of degree zero on  $R^n$  with  $\int_{S^{n-1}} \Omega(x') d\sigma(x') = 0$ . Assume that  $\Omega \in Lip_\gamma(S^{n-1})$ . The Marcinkiewicz commutators are defined by

$$\mu_\Omega^b(f)(x) = \left( \int_0^\infty |F_t^b(f)(x)|^2 \frac{dt}{t^3} \right)^{1/2},$$

$$\mu_S^b(f)(x) = \left[ \int \int_{\Gamma(x)} |F_t^b(f)(x, y)|^2 \frac{dy dt}{t^{n+3}} \right]^{1/2}$$

and

$$\mu_\lambda^b(f)(x) = \left[ \int \int_{R_+^{n+1}} \left( \frac{t}{t + |x - y|} \right)^{n\lambda} |F_t^b(f)(x, y)|^2 \frac{dy dt}{t^{n+3}} \right]^{1/2},$$

where

$$F_t^b(f)(x) = \int_{|x-y|\leq t} (b(x) - b(y)) \frac{\Omega(x-y)}{|x-y|^{n-1}} f(y) dy$$

and

$$F_t^b(f)(x, y) = \int_{|y-z|\leq t} (b(x) - b(z)) \frac{\Omega(y-z)}{|y-z|^{n-1}} f(z) dz.$$

Set

$$F_t(f)(x) = \int_{|x-y|\leq t} \frac{\Omega(x-y)}{|x-y|^{n-1}} f(y) dy.$$

We also define

$$\mu_\Omega(f)(x) = \left( \int_0^\infty |F_t(f)(x)|^2 \frac{dt}{t^3} \right)^{1/2},$$

$$\mu_S(f)(x) = \left( \int \int_{\Gamma(x)} |F_t(f)(y)|^2 \frac{dydt}{t^{n+3}} \right)^{1/2}$$

and

$$\mu_\lambda(f)(x) = \left( \int \int_{R_+^{n+1}} \left( \frac{t}{t+|x-y|} \right)^{n\lambda} |F_t(f)(y)|^2 \frac{dydt}{t^{n+3}} \right)^{1/2},$$

which are the Marcinkiewicz operators(see [30]). Let  $H$  be the space

$$H = \left\{ h : \|h\| = \left( \int_0^\infty |h(t)|^2 dt/t^3 \right)^{1/2} < \infty \right\}$$

or

$$H = \left\{ h : \|h\| = \left( \int \int_{R_+^{n+1}} |h(y, t)|^2 dydt/t^{n+3} \right)^{1/2} < \infty \right\}.$$

Then, it is clear that

$$\begin{aligned} \mu_\Omega^b(f)(x) &= \|F_t^b(f)(x)\|, & \mu_\Omega(f)(x) &= \|F_t(f)(x)\|, \\ \mu_S^b(f)(x) &= \|\chi_{\Gamma(x)} F_t^b(f)(x, y)\|, & \mu_S(f)(x) &= \|\chi_{\Gamma(x)} F_t(f)(y)\| \end{aligned}$$

and

$$\mu_\lambda^b(f)(x) = \left\| \left( \frac{t}{t+|x-y|} \right)^{n\lambda/2} F_t^b(f)(x, y) \right\|, \quad \mu_\lambda(f)(x) = \left\| \left( \frac{t}{t+|x-y|} \right)^{n\lambda/2} F_t(f)(y) \right\|.$$

It is easily to see that  $\mu_\Omega^b$ ,  $\mu_S^b$  and  $\mu_\lambda^b$  satisfy the conditions of Theorem 1-5 (see [17-18][30]), thus Theorem 1-5 hold for  $\mu_\Omega^b$ ,  $\mu_S^b$  and  $\mu_\lambda^b$ .

**Application 3.** Bochner-Riesz operator.

Let  $\delta > (n-1)/2$ ,  $B_t^\delta(f)(\xi) = (1-t^2|\xi|^2)_+^\delta \hat{f}(\xi)$  and  $B_t^\delta(z) = t^{-n} B^\delta(z/t)$  for  $t > 0$ . Set

$$F_{\delta,t}^b(f)(x) = \int_{R^n} (b(x) - b(y)) B_t^\delta(x-y) f(y) dy,$$

The maximal Bochner-Riesz commutator is defined by

$$B_{\delta,*}^b(f)(x) = \sup_{t>0} |B_{\delta,t}^b(f)(x)|.$$

We also define that

$$B_{\delta,*}^\delta(f)(x) = \sup_{t>0} |B_t^\delta(f)(x)|$$

which is the maximal Bochner-Riesz operator(see [20]). Let  $H$  be the space  $H = \{h : \|h\| = \sup_{t>0} |h(t)| < \infty\}$ , then

$$B_{\delta,*}^b(f)(x) = \|B_{\delta,t}^b(f)(x)\|, \quad B_*^\delta(f)(x) = \|B_t^\delta(f)(x)\|.$$

It is easily to see that  $B_{\delta,*}^b$  satisfies the conditions of Theorem 1-5(see [17]), thus Theorem 1-5 hold for  $B_{\delta,*}^b$ .

## References

- [1] S. Bloom, *A commutator theorem and weighted BMO*, Trans. Amer. Math. Soc., 292(1985), 103–122.
- [2] H. Q. Bui, *Characterizations of weighted Besov and Triebel-Lizorkin spaces via temperatures*, J. Func. Anal., 55(1984), 39–62.
- [3] D. C. Chang, J. F. Li and J. Xiao, *Weighted scale estimates for Calderón-Zygmund type operators*, Contemporary Mathematics, 446(2007), 61–70.
- [4] S. Chanillo, *A note on commutators*, Indiana Univ. Math. J., 31(1982), 7–16.
- [5] W. G. Chen, *Besov estimates for a class of multilinear singular integrals*, Acta Math. Sinica, 16(2000), 613–626.
- [6] F. Chiarenza and M. Frasca, *Morrey spaces and Hardy-Littlewood maximal function*, Rend. Mat., 7(1987), 273–279.
- [7] R. Coifman and R. Rochberg, *Another characterization of BMO*, Proc. Amer. Math. Soc., 79(1980), 249–254.
- [8] R. R. Coifman, R. Rochberg and G. Weiss, *Fractorization theorems for Hardy spaces in several variables*, Ann. of Math., 103(1976), 611–635.
- [9] G. Di FaZio and M. A. Ragusa, *Commutators and Morrey spaces*, Boll. Un. Mat. Ital., 5-A(7)(1991), 323–332.
- [10] G. Di Fazio and M. A. Ragusa, *Interior estimates in Morrey spaces for strong solutions to nondivergence form equations with discontinuous coefficients*, J. Func. Anal., 112(1993), 241–256.
- [11] J. Garcia-Cuerva, *Weighted  $H^p$  spaces*, Dissert. Math., 162(1979).
- [12] J. Garcia-Cuerva and J. L. Rubio de Francia, *Weighted norm inequalities and related topics*, North-Holland Math.,16, Amsterdam, 1985.
- [13] B. Hu and J. Gu, *Necessary and sufficient conditions for boundedness of some commutators with weighted Lipschitz spaces*, J. of Math. Anal. and Appl., 340(2008), 598–605.
- [14] S. Janson, *Mean oscillation and commutators of singular integral operators*, Ark. Math., 16(1978), 263–270.
- [15] Y. Lin, *Sharp maximal function estimates for Calderón-Zygmund type operators and commutators*, Acta Math. Scientia, to appear.
- [16] L. Z. Liu, *Interior estimates in Morrey spaces for solutions of elliptic equations and weighted boundedness for commutators of singular integral operators*, Acta Math. Scientia, 25(B)(1)(2005), 89–94.

- [17] L. Z. Liu, *The continuity of commutators on Triebel-Lizorkin spaces*, Integral Equations and Operator Theory, 49(2004), 65–76.
- [18] L. Z. Liu, *Triebel-Lizorkin space estimates for multilinear operators of sublinear operators*, Proc. Indian Acad. Sci. (Math. Sci), 113(2003), 379–393.
- [19] L. Z. Liu, *Boundedness for multilinear Littlewood-Paley operators on Triebel-Lizorkin spaces*, Methods and Applications of Analysis, 10(4)(2004), 603–614.
- [20] S. Z. Lu, *Four lectures on real  $H^p$  spaces*, World Scientific, River Edge, NJ, 1995.
- [21] T. Mizuhara, *Boundedness of some classical operators on generalized Morrey spaces*, in "Harmonic Analysis", Proceedings of a conference held in Sendai, Japan, 1990, 183–189.
- [22] B. Muckenhoupt and R. L. Wheeden, *Weighted norm inequalities for fractional integral*, Trans. Amer. Math. Soc., 192(1974), 261–274.
- [23] M. Paluszynski, *Characterization of the Besov spaces via the commutator operator of Coifman, Rochberg and Weiss*, Indiana Univ. Math. J., 44(1995), 1–17.
- [24] J. Peetre, *On convolution operators leaving  $L^{p,\lambda}$ -spaces invariant*, Ann. Mat. Pura. Appl., 72(1966), 295–304.
- [25] J. Peetre, *On the theory of  $L^{p,\lambda}$ -spaces*, J. Func. Anal., 4(1969), 71–87.
- [26] C. Pérez, *Endpoint estimate for commutators of singular integral operators*, J. Func. Anal., 128(1995), 163–185.
- [27] C. Pérez and R. Trujillo-Gonzalez, *Sharp weighted estimates for multilinear commutators*, J. London Math. Soc., 65(2002), 672–692.
- [28] E. M. Stein, *Harmonic analysis: real variable methods, orthogonality and oscillatory integrals*, Princeton Univ. Press, Princeton NJ, 1993.
- [29] A. Torchinsky, *Real variable methods in harmonic analysis*, Pure and Applied Math. 123, Academic Press, New York, 1986.
- [30] A. Torchinsky and S. Wang, *A note on the Marcinkiewicz integral*, Colloq. Math., 60/61(1990), 235–243.

Lanzhe Liu  
Department of Mathematics, Hunan University, Changsha 410082, P. R. of China  
E-mail: lanzheliu@163.com