

1 Ritchie, P.D., Smith, G., Davis, K.J., Fezzi, C., Halleck-Vega, S., Harper, A., Boulton, C.A.,
2 Binner, A.R., Day, B.H., Gallego-Sala, A., Mecking, J.V., Sitch, S., Lenton, T.M. and
3 Bateman, I.J. (2019) **Shifts in national land use and food production in Great Britain after**
4 **a climate tipping point**, *Nature Food*, 1:76–83 doi:10.1038/s43016-019-0011-3.
5

6 **Authors:** Ritchie, Paul D.L.^{1,2,*} & Smith, Greg S.^{1,3,4*}; Davis, Katrina J.³; Fezzi, Carlo^{3,5};
7 Halleck-Vega, Solmaria⁶; Harper, Anna B.^{1,7}; Boulton, Chris A.^{1,2}; Binner, Amy R.³; Day,
8 Brett H.³; Gallego-Sala, Angela V.²; Mecking, Jennifer V.⁸; Sitch, Stephen A.²; Lenton,
9 Timothy M.^{1,2,†} & Bateman, Ian J.^{1,3,†}

10 * joint lead authors, † joint corresponding authors

11 ¹Global Systems Institute, University of Exeter, Exeter, EX4 4QE, United Kingdom

12 ²College of Life and Environmental Sciences, Laver Building, University of Exeter, Exeter,
13 EX4 4QE, United Kingdom

14 ³Land, Environment, Economics and Policy Institute, University of Exeter Business School,
15 Xfi Building, Rennes Drive, Exeter, EX4 4PU, United Kingdom

16 ⁴CSIRO Land and Water, Hobart, 7001, Australia

17 ⁵Department of Economics and Management, University of Trento, via Vigilio Inama 5,
18 38122 Trento, Italy

19 ⁶Wageningen University and Research, Leeuwenborch Building, Hollandseweg 1, 6706KN,
20 Wageningen, Netherlands

21 ⁷College of Engineering, Mathematics and Physical Sciences, Laver Building, University of
22 Exeter, Exeter, EX4 4QE, United Kingdom

23 ⁸Ocean and Earth Science, National Oceanography Centre Southampton, University of
24 Southampton, Southampton, SO14 3ZH, United Kingdom

25

27 **Climate change is expected to impact agricultural land use. Steadily accumulating**
28 **changes in temperature and water availability can alter the relative profitability of**
29 **different farming activities and promote land use changes. There is also potential for**
30 **high-impact ‘climate tipping points’ where abrupt, non-linear change in climate occurs**
31 **- such as the potential collapse of the Atlantic Meridional Overturning Circulation**
32 **(AMOC). Here, using data from Great Britain, we develop a methodology to analyse the**
33 **impacts of a climate tipping point on land use and economic outcomes for agriculture.**
34 **We show that economic/land use impacts of such a tipping point are likely to include**
35 **widespread cessation of arable farming with losses of agricultural output, an order of**
36 **magnitude larger than the impacts of climate change without an AMOC collapse. The**
37 **agricultural effects of AMOC collapse could be ameliorated by technological**
38 **adaptations such as widespread irrigation, but the amount of water required and the**
39 **costs appear prohibitive in this instance.**

40


41 Tipping points can occur in elements of the climate system¹, in ecosystems², and in coupled
42 social-ecological systems³ where, often because of prior cumulative effects, a small change in
43 drivers generates an abrupt response in a system - qualitatively changing its future state. The
44 potential difficulties of reversing changes caused by tipping points⁴ means there is a pressing
45 need to understand their potential impacts and the extent to which such impacts can be
46 ameliorated. However, economic assessments of the impacts of large-scale climate tipping
47 points are rare⁴⁻⁶, typically of low resolution⁷, and often contested^{8,9}.

48 To address these issues, we consider a well-studied tipping point; collapse of the Atlantic
49 Meridional Overturning Circulation (AMOC)^{10,11}. The AMOC includes surface ocean
50 currents that transport heat from the tropics to the northeast Atlantic region benefiting
51 Western Europe, including the agricultural system of Great Britain (GB). We contrast the

52 impacts of conventional (hereafter ‘smooth’) climate change with that of a climate tipping
53 point involving AMOC collapse on agricultural land use and its economic value in GB, with
54 or without a technological response. Our climate projections span 2020 to 2080 and use a
55 mid-range climate change scenario as a baseline (Figure 1a-f; see Methods, subsequent
56 discussion of uncertainties such as weather variability, and sensitivity analysis in Extended
57 Data; results reported in the main paper are mean effects). We take an existing simulation of
58 the effects of AMOC collapse^{12,13} and treat it as a set of anomalies that can be linearly
59 combined with the baseline (smooth) climate change scenario. We nominally assume AMOC
60 collapse occurs over the time period 2030 to 2050 (Figure 1g-l; see Methods). This is a low
61 probability fast and early collapse of the AMOC compared to current expectations¹⁴,
62 emphasising the idealised nature of our study and our focus on assessing impacts. That said,
63 the AMOC has recently weakened by ~15%¹⁵ and models may be biased to favour a stable
64 AMOC relative to observations¹⁶.

65 We predict the production decisions of individual farms at 2 km x 2 km grid resolution
66 building upon an econometric land-use model¹⁷ and the detailed dataset¹⁸ employed by the
67 Natural Environment Valuation (NEV) model, which underpinned the UK National
68 Ecosystem Assessment¹⁹. Smooth changes in climate (Figure 1a-f) alter the relative
69 profitability of agricultural products generating changes in land-use. For example, arable
70 production is generally more profitable than grassland meat production in GB (see Extended
71 Data Figure 1) but is limited by physical restrictions, such as topography or low temperatures.
72 Climate change can raise temperatures, extending the area where cropping is economically
73 viable provided that rainfall is sufficient¹⁸. Relative to ‘smooth’ climate change, a climate
74 tipping point is likely to induce more abrupt land-use changes. For example, an AMOC
75 collapse (Figure 1g-l) is expected to induce significant reductions in rainfall²⁰, which could
76 rapidly shift land out of arable production²¹. A technological response to rainfall reductions in

77 the agriculturally productive lowlands of the south and east might be to irrigate them. These
 78 climate and technological responses lead to four scenario combinations of land-use change
 79 under climate change; with or without AMOC collapse and with or without a technological
 80 (irrigation) response²².




81
 82 **Figure 1. Temperature and rainfall for the growing season (April to September) in 2020**
 83 **and 2080.** *a) - c) Temperature in °C under smooth climate change. g) - i) Temperature in °C*
 84 *under abrupt climate change. d) - f) Rainfall in mm/growing season under smooth climate*
 85 *change. j) - l) Rainfall in mm/growing season under abrupt climate change. a), d), g), j)*
 86 *Climate data for 2020. b), e), h), k) Climate data for 2080. c), f), i), l) Difference between*
 87 *2020 and 2080 climate variables; a positive (negative) value represents an increase*
 88 *(decrease) in 2080 compared to 2020.*

89 **Land use change under smooth climate change**

90 Figure 2a maps land-use in 2020 as predicted by the agricultural model based on a spatially
91 explicit analysis of physical environment, climate, economic, and policy data from the 1960s
92 to the present day, allowing for climate trends over that period. Here physical constraints and
93 cool temperatures are expected to constrain high value arable production mainly to the
94 lowlands of south and east GB.

95 Our smooth climate change scenario results in a substantial 1.9°C mean warming in the
96 growing season in 2080 relative to 2020 (from an average of 12.6°C, Figure 1a, c, see
97 Methods) together with a modest 20 mm mean decline in growing season rainfall (from an
98 average of 445 mm, Figure 1d,f). Assuming that the AMOC is maintained then climate
99 change is likely to induce a significant and profitable increase in the intensity of arable
100 production across most lowland areas (Figure 2b, c, contrast with Figure 2a). These results
101 indicate a modest increase in overall arable area, but in parts of eastern England, high
102 temperatures and declining rainfall result in a reduction in arable production (Figure 2b).
103 Taking these differing effects into account, overall, GB arable area rises from 32% to 36% of
104 total agricultural area (see Extended Data Figure 2, Extended Data Figure 3), increasing
105 agricultural output value by approximately £40million per annum by 2080 (assuming
106 2017/18 agricultural prices). This value may increase further if, as best estimates suggest^{22,23},
107 real (inflation adjusted) agricultural prices increase somewhat over the period as a result of
108 climate change²³⁻²⁶ and other factors^{27,28}.



Figure

109

110

111


112

113

114

2. Impact of smooth and abrupt climate and economic change on the share of arable farmland in 2020 and 2080. *a) Arable farmland for 2020. b), d), f), h), arable farmland for 2080 under the four scenarios considered. c), e), g), i) Time series (England only) for mean climate and economic measures from 2020 to 2080 under the four scenarios considered. Water supply refers to the combination of rainfall and irrigation (if applicable).*

115 Under smooth climate change, approximately 14% of GB is likely to be rainfall-limited by
 116 2080 (Figure 4). If this proportion was irrigated from 2050, this would lead to an even greater
 117 rise in arable area—up from 32% to 42% of total agricultural land (Figure 2d, e, Extended
 118 Data Figure 3). This generates an increase in agricultural production value of £125million per
 119 annum by 2080. The overall water requirements for such an intervention are relatively
 120 modest, with average demand across irrigated areas equivalent to approximately 18 mm of
 121 extra rainfall during the growing season. Nevertheless, recent estimates of the costs of



122 **Figure 3. Limiting factors from an AMOC collapse on the share of arable land.** *a)*
 123 *Arable farmland for 2020. b) Arable farmland for 2080 with temperature based on an AMOC*
 124 *collapse and rainfall under smooth climate change (no AMOC collapse). c) Arable farmland*
 125 *for 2080 with rainfall based on an AMOC collapse and temperature under smooth climate*
 126 *change (no AMOC collapse).*

127 irrigating GB wheat production²⁹ show that these costs exceed the value of additional
128 production; in short, from an economic perspective, unless future arable crop prices rose
129 sufficiently, such investment may not be worthwhile.

130 **Land use change under a climate tipping point**

131 Our remaining scenarios impose a collapse of the AMOC over the period 2030-2050 overlaid
132 on the smooth climate change trend. A previous study that combined a rapid AMOC collapse
133 with future climate projections demonstrated that temperatures will continue to rise globally,
134 but with a delay of 15 years, while GB temperatures will be dependent upon the AMOC^{12,30-}
135 ³². In the present study, the AMOC collapse reverses the warming seen in the smooth climate
136 change scenarios, generating an average fall in temperature of 3.4°C by 2080 accompanied by
137 a substantial reduction in rainfall, falling by 123 mm during the growing season (Extended
138 Data Figure 2 and Extended Data Figure 4).

139 Holding real prices constant, then in the absence of a technological response (i.e. irrigation),
140 rainfall (and to a lesser extent temperature) limitation due to AMOC collapse is predicted to
141 affect arable farming in many areas (Figure 2f, g). The expected overall area of arable
142 production is predicted to fall dramatically from 32% to 7% of land area (Extended Data
143 Figure 2, Extended Data Figure 3). This in turn generates a major reduction in the value of
144 agricultural output, falling by £346million per annum (Table 1), representing a ~10%
145 reduction in total income from GB farming³³. The key driver of the arable loss seen across
146 GB is climate drying due to AMOC collapse, rather than cooling (Figure 3b, c). This adds
147 considerably to the part of Eastern England that is already vulnerable to arable loss due to
148 drying under baseline climate change (green band in Figures 2b, 3b). Part of eastern Scotland
149 has a potential gain in arable production suppressed by the cooling effects of an AMOC
150 collapse (contrast Figures 2f and 3c), but the loss of potential arable production due to
151 cooling is small compared to the impacts of drying. However, the assumption of constant real

152 prices is less plausible under the major global food system dislocation caused by a collapse of
 153 the AMOC. While firm estimates are not available, substantial food price increases are
 154 thought likely^{22,34}. With the physical limits imposed by AMOC collapse constraining farm
 155 production, such price increases mean that wellbeing losses may be significantly higher than
 156 those calculated here, implying that our results should be viewed as lower bound,
 157 conservative estimates of the impacts of such a scenario.


	Smooth climate change, no technological change	Smooth climate change, with technological change	Abrupt climate change, no technological change	Abrupt climate change, with technological change
AMOC	Maintained	Maintained	Collapse	Collapse
Irrigation	No	Yes	No	Yes
Agricultural change value (£M p.a.)	40	125	-346	79
Irrigation cost (£M p.a.)	0	-284	0	-807
Net value change (£M p.a.)	40	-159	-346	-728

158

159 **Table 1. Net impact on GB agriculture of smooth versus tipping point (AMOC collapse)**
 160 **climate change, with and without ameliorative measures (technological response).**

161 With a change in technology to implement sufficient irrigation from 2050, the drying effects
 162 of the AMOC collapse on arable production could be substantially offset (Figure 2h, i). In
 163 this scenario, land area under arable production still rises from 32% to 38% by 2080 with an
 164 accompanying increase in output value of £79million per annum (Table 1, Extended Data
 165 Figure 3). Nevertheless, this increase in extent and value are lower than under the second
 166 scenario where the AMOC is maintained, due to lower temperatures (contrast Figure 2h with
 167 2b). Furthermore, the more extreme reduction in rainfall caused by the AMOC collapse
 168 means that water required for adequate irrigation is much greater than under the scenario

169 where the AMOC is maintained. Under the AMOC collapse scenario, 54% of GB grid cells
170 now require irrigation, with demand exceeding 150 mm in the growing season for some areas
171 in the south and east of England (and an average demand across irrigated areas of 70 mm of
172 extra rainfall) (Figure 4). This would require water storage (across seasons) or spatial
173 redistribution across the country from areas of higher rainfall in the north and western
174 uplands of GB. Irrigation costs incurred in this scenario are estimated at over £800million per
175 year, more than 10 times the value of the arable production it would support (see Methods).
176 So, again, irrigation costs outweigh amelioration benefits under climate change; a difference
177 which is massively inflated by the climate tipping point of AMOC collapse. Our analysis also
178 indicates the level of food cost increase (nearly three-quarters of a billion pounds) necessary
179 to justify such irrigation expenditure costs.



180

181 **Figure 4. GB water balance in 2080 during the growing season with irrigation available**

182 **under the climate scenarios of the AMOC either maintained or collapsed. Water deficits**

183 *(< 280 mm) during the growing season (April-September) where irrigation occurs (red) and*

184 *areas with excess water (> 280 mm) (blue) during the growing season when a) AMOC is*

185 *maintained or b) AMOC collapsed.*

186 **Future agriculture in Great Britain**

187 Table 1 summarises results from our analysis of the impacts of both smooth and abrupt
188 climate change upon agriculture in GB. In the absence of a climate tipping point, smooth
189 climate change results in an elevation of temperature with modest falls in water availability.
190 Given the cool, moist present-day conditions of GB this results in a relatively small increase
191 in agricultural net profits (smooth climate change, no technological change). A few areas,
192 notably in Eastern England, experience rainfall limitations but the costs of irrigation
193 outweigh the benefits of addressing these constraints (smooth climate change, with
194 technological change). However, the introduction of a climate tipping point in the form of an
195 AMOC collapse removes the possibility of any positive outcome for GB agriculture.
196 Reductions in temperature, and especially rainfall, result in major losses in the value of
197 agricultural production (abrupt climate change, no technological change). While
198 technological change in terms of widespread irrigation can ameliorate reductions in arable
199 output (abrupt climate change, with technological change), in the absence of major price
200 increases (which are plausible but uncertain) the costs of such investments dwarf the benefits
201 they would provide.

202 Alongside economic uncertainties, agricultural land use, production and its value will also
203 respond to a number of other variables including changes in farming systems⁴¹,
204 technology^{35,36}, national and international policy^{37,38}. Even holding all of these factors
205 constant, climate futures may themselves bring increased variability including more frequent
206 weather extremes which may not be well reflected in mean temperature and rainfall
207 trends^{26,39}. A sensitivity analysis is therefore discussed in Methods with findings presented in
208 Extended Data. This reveals substantial variability in results, however the key findings and
209 relative comparison across our four scenarios remain. There are a number of reasons for
210 expecting such relativities to be robust. First, while there is uncertainty between models

211 regarding the net effect of global warming and AMOC collapse on GB temperatures, this is
212 not the major control on arable fraction. Instead, predicted drying due to AMOC collapse is
213 the key control and this is robust across climate models (see Extended Data Figure 5). The
214 climate model we use is conservative in its predicted drying, but nevertheless arable
215 production is still largely eliminated under AMOC collapse. Hence using another climate
216 model with greater predicted drying has relatively little scope to alter this key result. The
217 major source of uncertainty in the economic analysis concerns future prices. Under smooth
218 climate change real prices are generally expected to increase although only modestly. For
219 example, IPCC²³ estimate a median increase of 7.6% (range of 1 to 23%) in cereal prices by
220 mid-century under smooth climate change. Previous analyses using the same agricultural land
221 use model show that such price increases, if sustained, could yield similar scale effects to
222 those induced by smooth climate change⁴⁰. Given that potentially transformational
223 improvements in food production technology²⁸ and diets could dampen these effects, overall
224 this suggests that the estimates reported in the present paper, which assume constant real
225 prices, should be seen as lower bound but of appropriate magnitude. There are several other
226 expected impacts of AMOC collapse on GB that are not considered. These include harsher
227 winters, with greater storminess, and shortening of the growing season^{20,41}. These would
228 further tend to suppress arable production and challenge farming more generally. Weather
229 variability is expected to increase under AMOC collapse and could lead to farmers
230 diversifying their activity. Thus, whilst we already predict a nearly complete cessation of
231 arable farming, the overall impact of AMOC collapse on farming activity and associated
232 income could be considerably greater than we predict.

233 **Conclusion**

234 We have presented the first detailed case study of the national impacts of a climate tipping
235 point on land-use, agricultural production and its economic value, together with an

236 assessment of the potential for technological change to ameliorate impacts. While smooth
237 climate change can result in major changes in land-use and accompanying economic values,
238 we show that passing a climate tipping point has the potential to generate order-of-magnitude
239 greater economic impacts and that even these may be lower bound estimates. Our case study
240 concerns just one sector in one country, within which we only examine one impact of the
241 substantial land-use changes predicted. While agricultural production is obviously important,
242 changes in land-use generate multiple impacts; the need to understand these changes, and
243 their impacts on further sectors and countries, underlines the importance of many more such
244 analyses.

245

246 **Methods**

247 *Climate data*

248 Observational temperature and rainfall data from 1981-2010⁴² were used to estimate the land-
249 use model on agricultural census data (June Agricultural Census panel from EDINA).
250 Specifically, the surface observations, provided at 5 km x 5 km resolution, are averaged over
251 the growing seasons (April to September) and bilinearly interpolated (ignoring topography)
252 onto the 2 km x 2 km grid cell resolution used in the agricultural census.

253 The projected future climate data used in the agricultural model is supplied by the Met Office
254 Hadley Centre Regional Model Perturbed Physics Ensemble simulations for the 21st Century
255 for the UK domain (HadRM3-PPE-UK)⁴³. The runs consist of daily data that spans 1950-
256 2100 at 25 km x 25 km resolution over the UK and forms part of the UK Climate Projections,
257 UKCP09⁴⁴. The ensemble is designed to simulate the regional climate over the UK for the
258 historical and medium emissions scenario SRES-A1B⁴⁵. In this paper, we chose the standard
259 run, where parameters are kept at their unperturbed values, corresponding to a 3.5K global
260 climate sensitivity and again we bilinearly interpolate the data onto the 2 km grid used for the

261 agricultural model. The climate projections used in the agricultural model for any given year
262 consist of the mean temperature and rainfall for the growing seasons (April to September) of
263 the preceding 30 years. To correct for any systematic bias in the modelled climate projections
264 the climate projections are bias corrected. The bias correction was performed by shifting the
265 future projections by the mean bias between the modelled and observed data for 1960-1989
266 (the mean temperature and rainfall for 1960-1989 during the growing season is shown in
267 Extended Data Figure 6).

268 For simulation of an AMOC collapse, we use data from an experiment that used the
269 HadGEM3 model with the global configuration 2 (GC2), N216 atmospheric (~60 km) and
270 ORCA025 ocean (~25 km)⁴⁶. The coupled climate model simulations are a present-day
271 control simulation and a simulation where the AMOC is collapsed using freshwater hosing
272 after which the model is allowed to run freely^{13,20}. Both runs contain seasonal mean averages
273 for a 30-year period (again consistent with the time span used for estimation of the
274 agricultural model) for temperature and rainfall once the model has reached steady state.
275 Specifically, the data period 50 to 80 years after freshwater perturbations had ended were
276 used for temperature and rainfall seasonal averages. Note the results of Mecking, et al.¹³
277 suggest that the reduction of rainfall over the North Atlantic following the collapse reduces
278 with time, however, this effect is believed to be negligible at GB latitudes. Extended Data
279 Figure 4 shows the temperature and rainfall for the spring and summer (effectively
280 exchanging September for March in the growing season) for the AMOC maintained and
281 AMOC collapse scenarios.

282 Combining the difference between the HadGEM3 runs and the difference between the
283 transient runs with the observation data we were able to simulate an idealised AMOC
284 collapse. This is consistent with findings from Drijfhout¹², where a freshwater hosing run and
285 a control run showed that the difference in surface air temperature after an AMOC collapse

286 between the two runs remains approximately constant. A progressive (not instantaneous)
287 collapse of the AMOC was simulated by applying a linear weighting function to the AMOC
288 difference data during the prescribed years the AMOC is weakening, namely 2030-2050. It
289 should be noted that the speed of collapse is relatively fast and the linearity assumption
290 idealised compared to what is predicted in some models.

291 The subsequent cooling and drying observed following an AMOC collapse is consistent
292 amongst models (see Extended Data Figure 5). Furthermore, the spatial pattern of greatest
293 cooling in north west GB and least cooling in south east GB is prominent in an ensemble of
294 freshwater hosing experiments in different climate models⁴⁸.

295 *Agricultural model*

296 The agricultural land-use model builds on the data and the econometric methodology
297 developed by Fezzi and Bateman¹⁷, subsequently forming an essential component of the UK
298 National Ecosystem Assessment (e.g., Bateman, et al.⁴⁷, NEA¹⁹). This approach is also
299 recently used by Fezzi and Bateman¹⁸ to appraise the environmental impact of climate change
300 adaptation on land-use and water quality. We use a simpler version of the model that focuses
301 on understanding the determinants of agricultural land-use allocation between arable and
302 grassland. While agricultural revenues change greatly with output prices, arable land is
303 typically the highest-value agricultural activity in GB (exceptions are some very intensive
304 dairy farms located in the South West of the country), and therefore provides a proxy for
305 understanding the effects of climate change on the 72% of UK land area under agricultural
306 production³³.

307 The land-use data are derived from the June Agricultural Census (JAC) panel from EDINA
308 (www.edina.ac.uk), which are collected on a 2 km x 2 km grid (400 Ha) basis covering the
309 entirety of GB for eleven unevenly spaced years from 1972 to 2010. This generates around
310 55,000 grid-square records per year.

311 The model integrates germane environmental determinants of land-use among which are
312 climate, soil characteristics and land gradient. Crop yield is not fixed but rather is allowed to
313 depend on climate, soils, input levels, etc. and can therefore change across space and time.
314 So crop productivity is allowed to alter as climate changes and farmers are allowed to adapt
315 by changing crop varieties, fertilization methods etc. What we are not changing is the bundle
316 of crop possibilities available to farmers. So, for example, no new genetically modified crops
317 are brought into the analysis. The approach taken, not modelling yield directly but focusing
318 on land use via a discrete choice model, is the most established statistical land use model
319 approach, with contributions going back to Wu and Segerson⁴⁸ and more recently Lubowski,
320 Plantinga and Stavins⁴⁹ as well as our own exposition of the approach given in Fezzi and
321 Bateman⁵. Recent research⁵⁰ also shows that such an approach implies underlying and
322 theoretically consistent profit and yield functions.

323 To account for non-linear effects, rainfall and temperature in the growing season (April to
324 September) are modelled using piecewise linear functions. This approach allows us to capture
325 changes in the proportion of land allocated to arable cropping resulting from different growth
326 factors over a range of values (cf.^{18,51}). An interaction term is also included to allow the effect
327 of rainfall to depend on the effect of temperature and vice versa^{18,52}. Soil characteristics
328 include shares of peat, (s_peat), gravel (s_gravel), stones (s_stoney), or fragipan soil
329 (s_fragipan) and three dummy variables representing soil texture, namely share of fine,
330 medium and coarse soils (s_fine, s_medium, s_coarse). We used data from the Harmonised
331 World Soil Database (HWSD): a 30 arc-second (approximately 1 km resolution) raster
332 (regular gridded) database with over 16,000 different soil mapping units⁵³. Finally, we
333 include mean altitude (elev) and slope represented as mean slope (slope), both derived from
334 the 50 m resolution Integrated Hydrological Digital Terrain Model (IHDTM) licensed from
335 the Centre for Ecology and Hydrology⁵⁴.

336 In order to address potential spatial autocorrelation, the approach in Fezzi and Bateman⁵ is
337 followed and a cell every four along both the horizontal and vertical axis is sampled. We
338 define grassland as the sum of rough grazing, permanent grassland and temporary grassland,
339 and arable land as the sum of cereals, oilseed rape, root crops, and all other agricultural lands.
340 The only significant agricultural land-use category excluded from the agricultural model is
341 rural woodland, whose expansion and contractions are mainly driven by governmental
342 subsidies which we assume remain constant across our climate change scenarios. As
343 described on the source data website (www.edina.ac.uk), grid square land-use estimates can
344 sometimes overestimate or underestimate the amount of agricultural land within an area,
345 since their collection is based on the location of the main farm house. This feature is
346 corrected by rescaling the sum of the different agricultural land-use areas assigned to each
347 grid square to match with the total agricultural land derived using satellite land cover data
348 and ancillary spatial data⁵⁵ (Meridian Developed Land Use Areas, OS roads, OS railways; the
349 National Inventory for Woodland and Trees) to locate areas that are used for agricultural
350 production, urban activities, etc.

351 For policy determinants of land-use decisions the share of each grid square designated as
352 National Park (*npark*), Environmentally Sensitive Area (*esa*) and Greenbelts (*greenbelt*) are
353 included. Environmentally Sensitive Areas, introduced in 1987 and extended in subsequent
354 years, were launched to conserve and enhance areas of particular landscape and wildlife
355 significance. Digital boundary data were downloaded from Natural England⁵⁶ and the
356 Scottish Government⁵⁷. Spatial data for English greenbelts were licensed by Defra from the
357 Ordnance Survey⁵⁵. Presently, there is no national digital spatial boundary dataset for
358 Scottish greenbelts. Each council provided information and PDF maps or ESRI shapefiles.
359 For Wales, there is currently only one area of greenbelt (Newport and Cardiff), and its
360 boundaries were derived from local development plans.

361 The dependent variable of the model is the share of agricultural land devoted to arable. We
362 model this variable as a function of all the determinants of land-use in a reduced-form
363 specification. After applying a logit transformation, this model can be estimated via quasi-
364 maximum likelihood (QML)^{58,59}. The estimation results are reported in Extended Data Figure
365 7. It can be observed that favourable environmental and topographical features (e.g. soil
366 quality and less elevated areas), significantly increase the share of arable. It is also apparent
367 that policy factors are in line with expectations, in this case reducing the share of arable as
368 these reflect a greater amount of protected areas: such as for national parks. Almost all of the
369 parameter estimates of the rainfall and temperature effects are also highly statistically
370 significant. These non-linear impacts can also be observed in Extended Data Figure 8.

371 Similarly, it emerges from Extended Data Figure 8 that warmer temperatures are beneficial
372 for arable as this promotes plant growth with the trend increasing quite rapidly at first, and
373 then more gradually. In the full sample, higher temperature extremes can have adverse
374 impacts, but this is based on a small number of observations with average growing season
375 temperatures above 14°C. For this reason, a subsample is taken as the non-linear climate
376 effects are sensitive to the inclusion of these few observations. The estimates of all other
377 variables are very similar regardless of basing the estimations on the full or subsample. A
378 simple quadratic specification shows increases in predicted arable share with increasing
379 temperature; this provides further evidence of the robustness of the study's results to the
380 model specification.

381 It is also evident that higher accumulated rainfall over the growing season negatively affects
382 arable share (e.g. from flooding or waterlogging) (Extended Data Figure 8). When all
383 observations are used, the estimates also corroborate a downward trend of arable with respect
384 to average rainfall of less than 300 mm but few observations exist below 290 mm. The few
385 observations with lower rainfall levels are also those with observed higher average

386 temperatures. However, under the smooth and abrupt (AMOC collapse) climate change
387 scenarios we consider in this study there is a growing shift towards less rainfall in the
388 summer and therefore the functional form requires extending below 290 mm. We apply a
389 conservative approach by applying a linear extrapolation to the downward trend (Extended
390 Data Figure 8). Using land cover data from the European Space Agency Climate Change
391 Initiative⁶⁰ and average growing season rainfall values from 1988-2017 (CRU TS4.02⁶¹), we
392 have provided arable share for rainfall values that go outside the range of GB data. We used
393 the CCI-Land Cover Tools (v. 3.14) to regrid the land cover data from the original 300 m
394 spatial resolution to the half-degree resolution of the CRU data. Two regions were selected
395 based on comparable agricultural extent and climate with GB: US Great Plains (87W to
396 113W; 35N to 49N) and an area covering northern Eurasia (10W to 50E; 43N to 60N). We
397 also include data from over the UK, which shows a similar increasing trend in arable share
398 with lower rainfall values (above 300 mm). We define arable as rain-fed crops, including land
399 with herbaceous, tree or shrub cover, and pasture is defined as mosaic herbaceous and
400 grassland. The turning point estimated for GB is similar to that observed for the US Great
401 Plains and a little lower for EurAsia (the latter might reflect differences in crop types used).
402 In both cases the fall in arable share for rainfall below the turning point is sharper than our
403 estimation, suggesting that we apply a conservative approach. In addition to complex rainfall
404 patterns being more difficult to predict, there is also the issue of predicting how evenly
405 distributed the rainfall is over the growing season. This would be interesting to explore in
406 another study, as well as crop variations.

407 Our agricultural model does not explicitly account for the introduction of technological
408 advances in the form of new crops, etc., which could also help to attenuate the negative
409 impacts of the AMOC collapse. Effects other than temperature and rainfall, in particular CO₂
410 fertilization are not accounted for, and CO₂ fertilization has the potential to increase the

411 water-use efficiency of C3 crop plants and thus reduce the corresponding irrigation demand⁶².
412 Any agricultural model should be sensitive to prices and subsidies, and ours is no exception.
413 Arable farm profit margins are typically higher than for beef and sheep livestocking. While
414 dairy farms currently enjoy high per hectare margins (see the statistics in Fezzi, et al.⁶³), the
415 capital costs of moving into such production are prohibitive for most livestock farms and
416 many small dairy farms are uneconomic⁶⁴.

417 *Economic analysis*

418 Estimates of changes in farm profitability for the four scenarios are calculated using country
419 estimates of arable and grassland profitability. Profitability figures are taken from the Farm
420 Business Survey (FBS)⁶⁵ for England and Wales and the Farm Business Income (FBI) survey
421 for Scotland⁵⁷. Arable profitability is calculated as the average profitability per hectare from
422 cereal and general-cropping farming for a medium sized farm. Grassland profitability is
423 dependent on whether the land is classified as being in Less Favoured Areas (LFAs). LFAs
424 were introduced by the European Union to support farming where production conditions are
425 difficult and are defined according to the different physical and socio-economic
426 characteristics across the regions. LFAs are available for England in
427 https://magic.defra.gov.uk/Dataset_Download_Summary.htm, Scotland in
428 <https://data.gov.uk/dataset/a1ba43dd-569c-47e9-9623-21664aaf49ff/less-favoured-areas>. For
429 Wales we estimate LFAs by taking the lowland areas classified in LandMap
430 (<http://lle.gov.wales/catalogue/item/LandmapVisualSensory/?lang=en>). Extended Data
431 Figure 1 shows the changes in farm profitability for farms in England, Scotland and Wales
432 under the four scenarios. Agricultural prices and irrigation costs are fixed throughout the
433 economic analyses assuming 2017/18 prices.

434 In principle, the irrigation water demands considered in our analyses could be met from either
435 storage of water during the wetter, non-growing season, or spatial redistribution from those

436 areas of GB with surplus rainfall. Irrigation costs are estimated using values from a recent
437 study on the costs of irrigating wheat production in the East of England³⁵ which estimates
438 total system costs for irrigation at £163.60 per hectare. Under the scenario with smooth
439 climate and technological change, areas in GB with insufficient rainfall for arable production
440 (14% of GB grid cells) require, on average, an additional 18 mm of rainfall in the growing
441 season. Under a scenario with abrupt climate and technological change, areas in GB that
442 require irrigation (54% of grid cells), require an additional 70 mm in the growing season. To
443 meet this latter shortfall, water could be redistributed across the country from areas that do
444 not require irrigation—there is an average excess (after use) of 167 mm of rainfall in the
445 growing season in these areas. This equates to a positive difference of 39 mm across GB: in
446 other words, there is sufficient rainfall within GB to meet all irrigation needs. However, as
447 discussed in the main text, the costs of these technological interventions dwarf the benefits
448 they would provide (Table 1).

449 *Sensitivity analysis*

450 We performed a sensitivity analysis to assess the impact the climate variables (temperature
451 and rainfall) have on arable share. Extended Data Figure 2 provides the lower and upper
452 quartiles of the temperature and rainfall for selected years, over the previous 30 years (as
453 used in the agricultural model). Using the different combinations of the lower and upper
454 quartiles of the temperature and rainfall, together with the means used in the original analysis,
455 we generate eight additional arable fraction values. The ranges of these outputs are displayed
456 in Extended Data Figure 2 and Extended Data Figure 9 for the different scenarios.

457 The ranges of arable fractions suggest that the ranking of the scenarios is consistent when
458 compared to the ranking obtained using the means. The worst scenario for the arable fraction
459 remains the abrupt climate with no technological change which drops from a range of 19% -
460 34% in 2020 to 3% - 16% by 2080. The best scenario remains the smooth climate with

461 technological change which increases from 19% - 34% in 2020 to 28% - 52% by 2080. The
462 results show that climate projection variance is important in determining land use outputs.
463 The arable fraction ranges presented in Extended Data Figure 2 are wide, reflecting the
464 uncertainty in the climate projections. This uncertainty also translates into uncertainty in the
465 economic analysis, the economic value ranges from the sensitivity analysis are displayed in
466 Extended Data Figure 10 for the different scenarios. Despite the wide ranges around the
467 economic values, the patterns are still consistent with those reported in the main text, abrupt
468 climate change generates a major reduction in the value of agricultural output, falling by
469 £218 to £393million per annum, representing a substantial reduction in total income from GB
470 farming. The ranges on the costs of irrigation become very wide as the upper quartile for
471 rainfall results in lower demand for irrigation while the lower quartile results in higher
472 demand leading to wider uncertainty about the costs of scenarios 2 and 4.

473

474 **Data Availability**

475 The modelled output data that support the findings of this study are openly available from
476 Smith and Ritchie⁶⁶.

477 **References**

- 478 1 Lenton, T. M. *et al.* Tipping elements in the Earth's climate system. *Proceedings of*
479 *the national Academy of Sciences* **105**, 1786-1793 (2008).
- 480 2 Scheffer, M., Carpenter, S., Foley, J. A., Folke, C. & Walker, B. Catastrophic shifts in
481 ecosystems. *Nature* **413**, 591 (2001).
- 482 3 Milkoreit, M. *et al.* Defining tipping points for social-ecological systems
483 scholarship—an interdisciplinary literature review. *Environmental Research Letters*
484 **13**, 033005 (2018).
- 485 4 Lenton, T. M. & Ciscar, J.-C. Integrating tipping points into climate impact
486 assessments. *Climatic Change* **117**, 585-597 (2013).
- 487 5 Kopp, R. E., Shwom, R. L., Wagner, G. & Yuan, J. Tipping elements and climate–
488 economic shocks: Pathways toward integrated assessment. *Earth's Future* **4**, 346-372
489 (2016).
- 490 6 Vaughan, D. G. & Spouge, J. R. Risk estimation of collapse of the West Antarctic Ice
491 Sheet. *Climatic Change* **52**, 65-91 (2002).

492 7 Boulton, C. A., Allison, L. C. & Lenton, T. M. Early warning signals of Atlantic
493 Meridional Overturning Circulation collapse in a fully coupled climate model. *Nature*
494 *communications* **5**, 5752 (2014).

495 8 Link, P. M. & Tol, R. S. Estimation of the economic impact of temperature changes
496 induced by a shutdown of the thermohaline circulation: an application of FUND.
497 *Climatic Change* **104**, 287-304 (2011).

498 9 Tol, R. S. The economic effects of climate change. *Journal of economic perspectives*
499 **23**, 29-51 (2009).

500 10 Hofmann, M. & Rahmstorf, S. On the stability of the Atlantic meridional overturning
501 circulation. *Proceedings of the National Academy of Sciences* **106**, 20584-20589
502 (2009).

503 11 Rahmstorf, S. *et al.* Exceptional twentieth-century slowdown in Atlantic Ocean
504 overturning circulation. *Nat Clim Change* **5**, 475 (2015).

505 12 Drijfhout, S. Competition between global warming and an abrupt collapse of the
506 AMOC in Earth's energy imbalance. *Sci Reports* **5**, 14877 (2015).

507 13 Mecking, J., Drijfhout, S., Jackson, L. & Graham, T. Stable AMOC off state in an
508 eddy-permitting coupled climate model. *Climate dynamics* **47**, 2455-2470 (2016).

509 14 Stocker, T. F. *et al.* (Cambridge University Press, 2013).

510 15 Caesar, L., Rahmstorf, S., Robinson, A., Feulner, G. & Saba, V. Observed fingerprint
511 of a weakening Atlantic Ocean overturning circulation. *Nature* **556**, 191-196,
512 doi:10.1038/s41586-018-0006-5 (2018).

513 16 Liu, W., Xie, S.-P., Liu, Z. & Zhu, J. Overlooked possibility of a collapsed Atlantic
514 Meridional Overturning Circulation in warming climate. *Science Advances* **3**,
515 e1601666, doi:10.1126/sciadv.1601666 (2017).

516 17 Fezzi, C. & Bateman, I. J. Structural agricultural land use modeling for spatial agro-
517 environmental policy analysis. *American Journal of Agricultural Economics* **93**,
518 1168-1188 (2011).

519 18 Fezzi, C. & Bateman, I. The impact of climate change on agriculture: Nonlinear
520 effects and aggregation bias in Ricardian models of farmland values. *Journal of the*
521 *Association of Environmental and Resource Economists* **2**, 57-92 (2015).

522 19 NEA. UK National Ecosystem Assessment: Technical Report [United Nations
523 Environmental Programme–World Conservation Monitoring Centre (UNEP-WCMC).
524 (Cambridge, 2011).

525 20 Jackson, L. *et al.* Global and European climate impacts of a slowdown of the AMOC
526 in a high resolution GCM. *Climate dynamics* **45**, 3299-3316 (2015).

527 21 Cook, B. I., Ault, T. R. & Smerdon, J. E. Unprecedented 21st century drought risk in
528 the American Southwest and Central Plains. *Science Advances* **1**, e1400082 (2015).

529 22 Benton, T. *et al.* Environmental tipping points and food system dynamics: Main
530 Report. (The Global Food Security Programme, UK, 2017).

531 23 IPCC. Climate Change and Land. (Intergovernmental Panel on Climate Change,
532 Geneva, Switzerland, 2019).

533 24 Porter, J. R. *et al.* in *Climate Change 2014: Impacts, Adaptation, and Vulnerability.*
534 *Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth*
535 *Assessment Report of the Intergovernmental Panel on Climate Change* (eds C.B.
536 Field *et al.*) 485-533 (Cambridge University Press, 2014).

537 25 Mbow, C. *et al.* Food Security in Climate Change and Land, Intergovernmental Panel
538 on Climate Change. (Geneva, Switzerland, 2019).

539 26 (GFS), G. F. S. Extreme weather and resilience of the global food system. (The
540 Global Food Security Programme, UK, 2015).

541 27 (WEF), W. E. F. Shaping the Future of Global Food Systems: A Scenarios Analysis.
542 (World Economic Forum, Geneva, Switzerland, 2017).

543 28 Defence, M. o. Global Strategic Trends: The future starts today (sixth edition).
544 (Development, Concepts and Doctrine Centre, Shrivenham, 2018).

545 29 El Chami, D., Knox, J., Daccache, A. & Weatherhead, E. The economics of irrigating
546 wheat in a humid climate—A study in the East of England. *Agricultural Systems* **133**,
547 97-108 (2015).

548 30 Swingedouw, D. *et al.* Impact of Freshwater Release in the North Atlantic under
549 Different Climate Conditions in an OAGCM. *Journal of Climate* **22**, 6377-6403,
550 doi:10.1175/2009jcli3028.1 (2009).

551 31 Vellinga, M. & Wood, R. A. Global Climatic Impacts of a Collapse of the Atlantic
552 Thermohaline Circulation. *Climatic Change* **54**, 251-267,
553 doi:10.1023/a:1016168827653 (2002).

554 32 Jacob, D. *et al.* Slowdown of the thermohaline circulation causes enhanced maritime
555 climate influence and snow cover over Europe. *Geophysical Research Letters* **32**,
556 doi:10.1029/2005gl023286 (2005).

557 33 National Statistics. Agriculture in the United Kingdom 2017. (The Department for
558 Environment, Food and Rural Affairs; Department of Agriculture, Environment and
559 Rural Affairs (Northern Ireland); Welsh Assembly Government, The Department for
560 Rural Affairs and Heritage; The Scottish Government, Rural and Environment
561 Science and Analytical Services, 2018).

562 34 Nordhaus, W. & Boyer, J. (Cambridge, MA: MIT Press, 2000).

563 35 Dinesh, D., Campbell, B., Bonilla-Findji, O. & Richards, M. Vol. CCAFS Working
564 Paper No. 215 (CGIAR Research Program on Climate Change, Agriculture and
565 Food Security (CAAFS), Wageningen, The Netherlands, 2017).

566 36 Madramootoo, C. *Emerging Technologies for Promoting Food Security: Overcoming
567 the World Food Crisis*. (Woodhead Publishing, 2015).

568 37 Benton, T. G., Froggatt, A., Wright, G., Thompson, C. E. & King, R. *Food Politics
569 and Policies in Post-Brexit Britain* (Chatham House, London, 2019).

570 38 Challinor, A. J. *et al.* Transmission of climate risks across sectors and borders.
571 *Philosophical Transactions of the Royal Society A: Mathematical, Physical and
572 Engineering Sciences* **376**, 20170301, doi:doi:10.1098/rsta.2017.0301 (2018).

573 39 Benton, T. G., Gallani, B., Jones, C., Lewis, K. & Tiffin, R. Severe weather and UK
574 food chain resilience. (Government Office for Science (GO-Science), London, UK,
575 2012).

576 40 Fezzi, C. *et al.* Valuing provisioning ecosystem services in agriculture: the impact of
577 climate change on food production in the United Kingdom. *Environmental and
578 Resource Economics* **57**, 197-214 (2014).

579 41 Brayshaw, D. J., Woollings, T. & Vellinga, M. Tropical and Extratropical Responses
580 of the North Atlantic Atmospheric Circulation to a Sustained Weakening of the MOC.
581 *Journal of Climate* **22**, 3146-3155, doi:10.1175/2008jcli2594.1 (2009).

582 42 Met Office. UKCP09: Met Office gridded land surface climate observations - long
583 term averages at 5km resolution. (Centre for Environmental Data Analysis, date of
584 citation., 2017).

585 43 Hadley Centre for Climate Prediction and Research. UKCP09: Met Office HadRM3-
586 PPE UK model runs. (NCAS British Atmospheric Data Centre, date of citation.,
587 2014).

588 44 Jenkins, G. UK climate projections: briefing report. (Met Office Hadley Centre,
589 2009).

590 45 Nakicenovic, N. *et al.* *Special report on emissions scenarios (SRES), a special report*
591 *of Working Group III of the intergovernmental panel on climate change.* (Cambridge
592 University Press, 2000).

593 46 Safta, C. *et al.* Global sensitivity analysis, probabilistic calibration, and predictive
594 assessment for the data assimilation linked ecosystem carbon model. *Geoscientific*
595 *Model Development (Online)*, Medium: ED; Size: p. 1899-1918 (2015).

596 47 Bateman, I. J. *et al.* Bringing Ecosystem Services into Economic Decision-Making:
597 Land Use in the United Kingdom. *Science* **341**, 45-50, doi:10.1126/science.1234379
598 (2013).

599 48 Wu, J. & Segerson, K. The impact of policies and land characteristics on potential
600 groundwater pollution in Wisconsin. *American Journal of Agricultural Economics* **77**,
601 1033-1047 (1995).

602 49 Lubowski, R. N., Plantinga, A. J. & Stavins, R. N. Land-use change and carbon sinks:
603 econometric estimation of the carbon sequestration supply function. *Journal of*
604 *Environmental Economics and Management* **51**, 135-152 (2006).

605 50 Carpentier, A. & Letort, E. Multicrop production models with Multinomial Logit
606 acreage shares. *Environmental and Resource Economics* **59**, 537-559 (2014).

607 51 Schlenker, W. & Roberts, M. J. Nonlinear temperature effects indicate severe
608 damages to US crop yields under climate change. *Proceedings of the National*
609 *Academy of sciences* **106**, 15594-15598 (2009).

610 52 Morison, J. & Morecroft, M. (Blackwell Publishing, Oxford, 2006).

611 53 Van Liedekerke, M., Jones, A. & Panagos, P. ESDBv2 Raster Library—A Set of
612 Rasters Derived from the European Soil Database Distribution v2. 0. *European*
613 *Commission and the European Soil Bureau Network, CDROM, EUR 19945* (2006).

614 54 IHDTM. Integrated Hydrological Digital Terrain Model. (Centre for Ecology and
615 Hydrology, 2002).

616 55 Ordnance Survey. Meridian 2 Developed Land Use Area. (Ordnance Survey, 2013).

617 56 Natural England. Digital map boundaries download. (2012).

618 57 Scottish Government. Scottish Government Spatial Data File Download. (2012).

619 58 Papke, L. E. & Wooldridge, J. M. Econometric methods for fractional response
620 variables with an application to 401 (k) plan participation rates. *Journal of applied*
621 *econometrics* **11**, 619-632 (1996).

622 59 Papke, L. E. & Wooldridge, J. M. Panel data methods for fractional response
623 variables with an application to test pass rates. *Journal of Econometrics* **145**, 121-133
624 (2008).

625 60 ESA. Vol. Version 2.0 (2017).

626 61 Harris, I., Jones, P. D., Osborn, T. J. & Lister, D. H. Updated high-resolution grids of
627 monthly climatic observations – the CRU TS3.10 Dataset. *International Journal of*
628 *Climatology* **34**, 623-642, doi:10.1002/joc.3711 (2014).

629 62 Long, S. P., Ainsworth, E. A., Leakey, A. D. B., Nösberger, J. & Ort, D. R. Food for
630 Thought: Lower-Than-Expected Crop Yield Stimulation with Rising
631 CO₂ Concentrations. *Science* **312**, 1918-1921,
632 doi:10.1126/science.1114722 (2006).

633 63 Fezzi, C., Rigby, D., Bateman, I. J., Hadley, D. & Posen, P. Estimating the range of
634 economic impacts on farms of nutrient leaching reduction policies. *Agricultural*
635 *Economics* **39**, 197-205, doi:10.1111/j.1574-0862.2008.00323.x (2008).

636 64 MacDonald, J. M. *et al.* Profits, costs, and the changing structure of dairy farming.
637 *USDA-ERS Economic Research Report* (2007).

638 65 DEFRA. Farm Business Survey. (Department for Environment, Food & Rural Affairs,
639 UK, 2018).

640 66 Smith, G. S. & Ritchie, P. D. L. (NERC Environmental Information Data Centre:
641 doi.org/10.5285/e1c1dbcf-2f37-429b-af19-a730f98600f6, 2019).

642 **Corresponding author**

643 Correspondence to Ian Bateman (I.Bateman@exeter.ac.uk) and/or Tim Lenton
644 (T.M.Lenton@exeter.ac.uk)

645 **Acknowledgements**

646 This work was supported by the NERC Valuing Nature Programme (NE/P007880/1). We are
647 grateful for comments from Tim Benton and anonymous Referees.

648 **Author contribution statement**

649 I.J.B. and T.M.L. designed and directed the research and P.D.L.R. and G.S.S. helped shape
650 the research. P.D.L.R., G.S.S., K.J.D., I.J.B. and T.M.L. wrote the manuscript with C.F.,
651 C.A.B., A.B.H., A.V.G.S., J.V.M., S.H.V. and S.A.S. providing support and revisions.
652 P.D.L.R., G.S.S. and K.J.D. planned and conducted simulations for all analyses. C.F.
653 designed and ran the original agriculture land use model with A.R.B., B.H.D. and I.J.B.
654 providing support. C.F. and S.H.V. further developed the agricultural land use model from a
655 global analysis of agricultural land use by A.B.H. and A.V.G.S. The climate data was sourced
656 and corrected for modelled bias by P.D.L.R., and J.V.M. designed and ran the AMOC climate
657 simulations.

658 **Competing interest**

659 The authors declare no competing interests.

660
661

662 **Extended Data Figures**

Panel a: Changes in farm profitability for England, Scotland and Wales

England	Change in Agricultural profit 2020 to 2060	Change in Agricultural profit 2020 to 2080
	(£ Million)	(£ Million)
Smooth climate, no technological change	+47	+29
Smooth climate with technological change	+82	+114
Abrupt climate, no technological change	-313	-315
Abrupt climate, with technological change	+61	+90
Scotland		
Smooth climate, no technological change	-10	+3
Smooth climate with technological change	-10	+3
Abrupt climate, no technological change	-40	-35
Abrupt climate, with technological change	-35	-26
Wales		
Smooth climate, no technological change	+6	+8
Smooth climate with technological change	+6	+8
Abrupt climate, no technological change	-1	+4
Abrupt climate, with technological change	+9	+15
Total		
Smooth climate, no technological change	+43	+40
Smooth climate with technological change	+78	+125
Abrupt climate, no technological change	-354	-346
Abrupt climate, with technological change	+35	+79

Panel b: Estimates of average Farm Profitability for England, Scotland and Wales

	Arable (£ per Ha)	Lowland grassland (Lowland Grazing Livestock) (£ per Ha)	Upland grassland (Less Favoured Areas Grazing Livestock) (£ per Ha)
England ^a	351.30	262.30	222.50
Scotland ^b	195.00	141.50	82.60
Wales ^a	351.30 ^c	306.50	225.50

Notes: ^a England and Wales farm profitability is reported as the net profits from the Farm Business Survey (FBS) 2017/2018⁶⁵. ^b Scottish farm profitability is calculated from the Scottish farm business income (FBI); annual estimates 2016-2017⁵⁷. ^c Farm Business Survey values are not available for arable profit in Wales, for which values from England are used. Note that this comparison excludes dairy production as this tends to be limited by the availability of high levels of capital input which in turn is heavily influenced by historic access to milk quota subsidies that have now been abandoned.

663
664 **Extended Data Figure 1. Changes in farm profitability between 2020 and 2060 and**
665 **between 2020 and 2080.**


Panel a: Mean temperature and rainfall for previous 30-year growing seasons (April-September) when the Atlantic meridional overturning circulation (AMOC) is maintained or collapses.

Year	AMOC maintained				AMOC collapse			
	Mean arable area (percent)		Mean arable area (percent)		Temp (°C)	Rain(mm)	Mean arable area (percent)	
	Smooth climate, no technological change	Smooth climate, technological change	Abrupt climate, no technological change	Abrupt climate, technological change				
2020	12.6	445	32%	32%	12.6	445	32%	32%
2030	12.9	446	33%	33%	12.9	446	33%	33%
2040	13.2	447	34%	34%	10.6	396	25%	25%
2050	13.5	453	34%	35%	8.3	351	7%	33%
2060	14.0	448	36%	39%	8.7	345	6%	35%
2070	14.3	442	35%	40%	9.1	339	7%	37%
2080	14.5	425	36%	42%	9.2	322	7%	38%

Panel b: Combinations of lower and upper quartiles of temperature and rainfall for previous 30-year growing seasons (April-September) when the Atlantic meridional overturning circulation (AMOC) is maintained or collapses.


Year	AMOC maintained				AMOC collapse			
	Mean arable area ranges (percent)		Mean arable area ranges (percent)		Temp (°C)	Rain(mm)	Mean arable area ranges (percent)	
	Smooth climate, no technological change	Smooth climate, technological change	Abrupt climate, no technological change	Abrupt climate, technological change				
2020	11.9 - 13.1	369 - 517	19% - 34%	19% - 34%	11.9 - 13.1	369 - 517	19% - 34%	19% - 34%
2030	12.2 - 13.3	367 - 526	18% - 34%	18% - 34%	12.2 - 13.3	367 - 526	18% - 34%	18% - 34%
2040	12.7 - 13.7	372 - 522	19% - 35%	19% - 35%	10.1 - 11.1	320 - 471	9% - 26%	9% - 26%
2050	13.1 - 14.0	377 - 531	19% - 35%	19% - 47%	7.8 - 8.8	275 - 428	3% - 23%	24% - 37%
2060	13.4 - 14.5	372 - 526	18% - 37%	21% - 49%	8.2 - 9.2	270 - 423	3% - 23%	27% - 40%
2070	13.7 - 14.7	361 - 523	17% - 36%	23% - 50%	8.5 - 9.5	258 - 421	3% - 23%	30% - 41%
2080	13.9 - 14.8	355 - 494	19% - 36%	28% - 52%	8.6 - 9.6	252 - 391	3% - 16%	32% - 42%

666
667 **Extended Data Figure 2. Predicted farm allocation to arable land for individual years**
668 **between 2020 and 2080 per 2 km grid cell.**



669
 670 **Extended Data Figure 3. Time series of mean temperature, total rainfall for the growing**
 671 **season and arable share for the four scenarios considered. a) Temperature and rainfall in**
 672 *Great Britain with AMOC maintained and collapsed over 2020 to 2080. b) Mean arable*
 673 *fraction of agricultural land in Great Britain with AMOC maintained or collapsed and*
 674 *irrigation on or off, over the period 2020 to 2080.*

675



676
677 **Extended Data Figure 4. Mean temperature and total rainfall for spring and summer**
678 **(March-August) in steady state runs of the AMOC maintained and collapsed. a) - c)**
679 *Mean temperature and d) – f) mean total rainfall for a), d) a maintained AMOC and b), e)*
680 *collapsed AMOC^{13,20}. c), f) Plots the difference between the means of the AMOC maintained*
681 *and collapsed; a positive (negative) value represents an increase (decrease) for an AMOC*
682 *collapse compared to the AMOC maintained.*

Reference	Model	Temperature (Cooling)	Rainfall (Drying)	Notes
Jackson et al., 2015	HadGEM3 GC2	5.0°C growing season	85 mm/growing season (21%)	Model used in this study, 1980's CO ₂ levels (difference between AMOC maintained and collapsed in 2080, see Extended Data Table 1)
Drijfhout, 2015	ECHAM5/MPI-OM	2-4°C	Not provided	Global atmosphere-ocean general circulation model, 5member ensemble, SRES-A1B, 15 years after onset
Jacob et al., 2005	ECHAM5/MPI-OM & REMO	2-3°C	~20%	REMO is a regional atmospheric model, summer values
Vellinga & Wood, 2002	HadCM3	2-3°C	100-150 mm/growing season	Pre-industrial GHG emissions, 20-30 years after collapse
Vellinga & Wood, 2008	HadCM3	2-5°C	90 mm/growing season	IS92a emissions scenario
Swingedouw et al., 2009	IPSL CM4	~2°C	90 mm/growing season	Ocean-atmosphere-sea ice-land coupled GCM, 5 sets of experiments over different epochs, largest weakening – Last Glacial Maximum (LGM) – 12Sv circulation decline

Note: The last three entries of the change in rainfall (drying) have been converted (assuming rainfall is evenly distributed throughout the year) to mm/growing season for consistency.

683
684


Extended Data Figure 5. Impact of an AMOC collapse on temperature and rainfall

685

across various climate model freshwater hosing experiments. First row, model used in

686

this study.




687
 688 **Extended Data Figure 6. Surface observations of the mean temperature and total**
 689 **rainfall for the growing season for 1960-1989. a) Mean temperature and b) mean total**
 690 *rainfall for the growing season (April-September) from surface observations for the period*
 691 *1960-1989.*

	Estimate	Std. Error	Z-test	P-value	
rain	0.146	0.087	1.672	0.094	.
rain >= 290	-0.313	0.128	-2.442	0.015	*
rain >= 300	0.147	0.041	3.559	<2e-16	***
rain >= 400	0.009	0.001	6.754	<2e-16	***
rain >= 600	0.010	0.001	9.970	<2e-16	***
temp	0.738	0.332	2.224	0.026	*
temp >= 10	-0.542	0.312	-1.740	0.082	.
temp >= 12	-0.243	0.128	-1.898	0.058	.
temp >= 13	0.147	0.140	1.048	0.295	
rain*temp	0.000	0.000	0.301	0.764	
elev	-0.003	0.000	-7.710	<2e-16	***
slope	-0.060	0.011	-5.546	<2e-16	***
npark	-0.004	0.001	-2.881	0.004	**
csa	-0.002	0.001	-2.750	0.006	**
greenbelt	-0.002	0.001	-2.947	0.003	**
dist300	-0.001	0.000	-3.455	0.001	***
s_peat	-0.587	0.157	-3.738	<2e-16	***
s_gravel	-0.613	0.125	-4.883	<2e-16	***
s_stoney	-0.077	0.076	-1.012	0.312	
s_fragipan	-1.278	0.173	-7.376	<2e-16	***
s_coarse	0.238	0.069	3.463	0.001	***
s_fine	-0.345	0.063	-5.487	<2e-16	***
constant	-47.352	25.079	-1.888	0.059	.
pseudo-R ²	0.76				


Notes: . *, ** and *** indicate 10% 5% 1% and 0.1% significance levels respectively. Model estimated via QML. N = 22,220. The dependent variable is arable land share. The high pseudo-R² provides an indication of good model fit. Details of variable definitions are presented in the methods section. The model includes a time fixed effect to account for potential time-varying unobserved determinants such as commodity prices. As these are not relevant to the focus of this study, they are omitted from the table but are available from the authors.

692
693

Extended Data Figure 7. Model estimates of land-use (arable land share).



694
 695 **Extended Data Figure 8. Estimated impact of temperature and rainfall on arable land**
 696 **share in Great Britain from the agricultural model. Estimated fraction of arable share in**
 697 *Great Britain based on a) temperature and b) rainfall. For b) only: arable shares based on*
 698 *land cover data from Northern Eurasia (Eurasia), United Kingdom (UK), and the US Great*
 699 *Plains (USGP).*



700
 701 **Extended Data Figure 9. Impact sensitivity analysis of climate variables has on arable**
 702 **land share for 2020. a) GB map of arable farmland for using the lower quartile temperature**
 703 *and rainfall. b) GB map of arable farmland for using the upper quartile temperature and*
 704 *lower quartile rainfall. c) GB map of arable farmland for using the mean temperature and*
 705 *rainfall. d) GB map of arable farmland for using the lower quartile temperature and upper*

706 *quartile rainfall. e) GB map of arable farmland for using the upper quartile temperature and*
 707 *rainfall.*

708

	Smooth climate change, no technological change	Smooth climate change, with technological change	Abrupt climate change, no technological change	Abrupt climate change, with technological change
AMOC	Maintained	Maintained	Collapse	Collapse
Irrigation	No	Yes	No	Yes
Agricultural change value (£M p.a.)	-169 to +48	-63 to +271	-393 to -218	-7 to +139
Irrigation cost (£M p.a.)	0	-1 to -882	0	-527 to -952
Net value change (£M p.a.)	-169 to +48	-945 to +270	-393 to -218	-959 to -388

709

710 **Extended Data Figure 10. Net impact range on GB agriculture of smooth versus tipping**
 711 **point (AMOC collapse) climate change, with and without ameliorative measures**
 712 **(technological response) using lower and upper quartile of temperature and rainfall for**
 713 **previous 30-year growing seasons (April-September).**