

# Signaling cross-talk between TGF- $\beta$ /BMP and other pathways

Xing Guo<sup>1,2</sup>, Xiao-Fan Wang<sup>1</sup>

<sup>1</sup>Department of Pharmacology and Cancer Biology, Duke University Medical Center, Durham, NC 27710, USA

**Transforming growth factor-beta (TGF- $\beta$ )/bone morphogenic protein (BMP) signaling is involved in the vast majority of cellular processes and is fundamentally important during the entire life of all metazoans. Deregulation of TGF- $\beta$ /BMP activity almost invariably leads to developmental defects and/or diseases, including cancer. The proper functioning of the TGF- $\beta$ /BMP pathway depends on its constitutive and extensive communication with other signaling pathways, leading to synergistic or antagonistic effects and eventually desirable biological outcomes. The nature of such signaling cross-talk is overwhelmingly complex and highly context-dependent. Here we review the different modes of cross-talk between TGF- $\beta$ /BMP and the signaling pathways of Mitogen-activated protein kinase, phosphatidylinositol-3 kinase/Akt, Wnt, Hedgehog, Notch, and the interleukin/interferon-gamma/tumor necrosis factor-alpha cytokines, with an emphasis on the underlying molecular mechanisms.**

**Keywords:** TGF- $\beta$ , Smad, cross-talk, signaling pathway

*Cell Research* (2009) 19:71-88. doi: 10.1038/cr.2008.302; published online 11 November 2008

## Introduction

The transforming growth factor-beta (TGF- $\beta$ ) family of cytokines, including TGF- $\beta$ , bone morphogenic proteins (BMPs), and activin/inhibin, plays crucial roles in embryonic development, adult tissue homeostasis and the pathogenesis of a variety of diseases. The highly conserved core of the canonical TGF- $\beta$ /BMP signaling is a simple linear cascade that involves the TGF- $\beta$ /BMP ligands, two types of receptors (type I and II) and the signal transducers, Smads. On activation, the receptor complex phosphorylates the carboxy-terminus of receptor-regulated Smad proteins (R-Smads), including Smad1, 5 and 8 for BMP signaling and Smad2 and 3 for TGF- $\beta$  signaling. Activated R-Smads interact with the common partner Smad, Smad4, and accumulate in the nucleus, where the Smad complex directly binds defined elements on the DNA and regulates target gene expression together with numerous other factors [1-3]. Simple as it is, the TGF- $\beta$ /BMP pathway controls a myriad of events, including cell proliferation, differentiation, apoptosis, migration, extracellular matrix (ECM) remodeling,

immune functions, and tumor invasion/metastasis [4-8]. On the other hand, the activity and the signaling outcomes of this pathway are also influenced by many intracellular and extracellular signals [1, 3, 9]. This interplay between TGF- $\beta$ /BMP and other pathways, which is tightly regulated both spatially and temporally, gives rise to the remarkable complexity, diversity, flexibility, and delicacy of TGF- $\beta$ /BMP functions that have been exemplified by a great number of studies.

Signaling cross-talk is a perennial theme of TGF- $\beta$  research. In retrospect, the entire field of TGF- $\beta$  research sprang from a few groundbreaking observations, which we now know as a typical form of signaling cross-talk (reviewed by [10]). In the early 1980s, Roberts and colleagues isolated two fractions from murine sarcoma cell extracts that could synergistically induce remarkable growth of normal fibroblasts (NRK cells) on soft agar, a hallmark of cellular transformation. These two components were therefore named transforming growth factor  $\alpha$  and  $\beta$  (TGF- $\alpha$  and TGF- $\beta$ ) [11-13]. TGF- $\alpha$  alone had only limited transforming activity, and it was soon proven to be a ligand for the epithelial growth factor receptor (EGFR) and equivalent to EGF in promoting newborn mouse eyelid opening [14]. The other polypeptide, TGF- $\beta$ , was shown to be a potent inducer of NRK cell transformation, but only in the presence of TGF- $\alpha$  or EGF. This was undoubtedly the first classic example of functional interaction between TGF- $\beta$  and other

Correspondence: Xiao-Fan Wang

Tel: +1-919-681-4861

E-mail: wang0011@mc.duke.edu

<sup>2</sup>Current address: Department of Pharmacology, University of California, San Diego, La Jolla, CA 92093, USA

signaling pathways, although the underlying mechanisms remain not completely understood even today.

At the organism level, TGF- $\beta$ /BMP talks with other pathways at every stage of the life of a metazoan from birth to death. During embryonic development, the complex but delicate interactions between the TGF- $\beta$ /BMP, Wnt/Wg, Hedgehog (Hh), Notch, mitogen-activated protein kinase (MAPK), and other pathways are crucial for stem cell maintenance, body patterning, cell fate determination, organogenesis, and so on. These signals are also instrumental for the proper growth and functioning of cells and tissues in adult animals (homeostasis), whereas concurrent alterations of these pathways are commonly found in aged or diseased animals, as repetitively seen during the development of cancer [7, 15-20].


In the past three decades since its discovery, the downstream signaling cascade of TGF- $\beta$  has been clearly delineated. The biochemical basis for the extensive cross-talk between TGF- $\beta$  and other pathways is essentially two-fold: (1) multiple components of the TGF- $\beta$  pathway (mainly Smads) make direct and dynamic contacts with numerous other proteins; and (2) TGF- $\beta$  has a great many targets, both transcriptional and non-transcriptional (Figure 1). Many of these binding partners and target molecules are essential constituents of other pathways, naturally integrating TGF- $\beta$  with other signals to produce highly regulated cellular

responses. In this review, we focus on these features and discuss the different modes of TGF- $\beta$  signaling cross-talk as well as the ensuing context-dependent outcomes.

### TGF- $\beta$ and the MAPK pathway

MAPKs, including Erk1/2, JNK1/2/3, and p38/MAPKs, are evolutionarily conserved regulators essential for a variety of cellular events. Multiple extracellular stimuli can initiate a cascade of serial phosphorylation activation from MAP kinase kinase kinase (MAPKKK) to MAP kinase kinase (MAPKK) and finally MAPK [21]. Activated MAPKs phosphorylate a battery of proteins (primarily nuclear transcription factors) with diverse functions in regulating proliferation, survival, migration, and so on. One of the best characterized trigger for this MAPK pathway is Ras activation, which propagates signals from a number of ligand- or self-activated receptor tyrosine kinases (RTKs), such as EGFR (including HER2/Neu/ErbB2), FGFR, IGFR, PDGFR, and insulin receptor [21-23]. In addition, MAPKs can also be regulated by TGF- $\beta$ /BMP stimulation [24, 25], which represents an important mechanism for non-Smad TGF- $\beta$  signaling and is discussed in a separate review. Here, we mainly focus on how MAPK activity modulates the function of TGF- $\beta$ /BMP.

A series of studies have shown that HER2/Neu/ErbB2



**Figure 1** Basic modes of signaling cross-talk. A cross-talk exists between pathways A and B when both of the following criteria are met. Functionally, the combinatorial signal from A and B must produce a different response than that triggered by A or B alone. Mechanistically, the A and B pathways must be connected in at least one of the three depicted ways: **(a)** components of the two pathways physically interact; **(b)** components of one pathway are enzymatic or transcriptional targets of the other; and **(c)** one signal modulates or competes for a key modulator or mediator (“M”) of the other. In this scheme, A and B are interchangeable, and the arrows may represent either positive or negative regulations. Note that an altered response can arise from independent (non-cross-talk) inputs **(d)**.

signaling, which activates both the MAPK and phosphatidylinositol-3 kinase (PI3K)/Akt pathways, communicates intimately with TGF- $\beta$ /Smad in controlling mammary epithelial cell biology and breast cancer development [6,26-31]. A general notion emerging from these studies is that HER2/Ras can antagonize TGF- $\beta$ -induced apoptosis and cell cycle arrest, while allowing for the pro-migratory and pro-invasive functions of TGF- $\beta$ . Therefore, both positive and negative regulations exist between the two pathways.


The synergy between the TGF- $\beta$  and HER2/Ras/MAPK pathways often leads to the secretion of additional growth factors and cytokines, including TGF- $\beta$  itself, which in turn promote epithelial-to-mesenchymal transition (EMT) and cell invasion [32-34], whereas JNK kinases seem to negatively regulate the autocrine expression of TGF- $\beta$ 1 [35]. MEK/Erk has been reported to positively regulate *SMAD3* gene transcription in epithelial and smooth muscle cells [36]. On the other hand, TGF- $\beta$ /Smad induces the expression of platelet-derived growth factor (PDGF) in liver cancer and glioma, which is required for fulfilling the pro-oncogenic and pro-metastatic functions of TGF- $\beta$  [37, 38].

Numerous studies have revealed that the linker region of Smad proteins is a critical platform for integrating RTK/MAPK signals with the TGF- $\beta$ /BMP pathway. The Smad linker region is loosely organized and highly flexible in structure, rendering it readily accessible for a number of kinases. This region is also rich in serine, threonine as well as proline residues, favoring phosphorylation by proline-directed kinases such as MAPKs and glycogen synthase kinase 3- $\beta$  (GSK3- $\beta$ ). Human cancer cells harboring oncogenic Ras are often resistant to TGF- $\beta$ -induced cytostasis, which was thought to result from Erk-mediated Smad2/3 linker phosphorylation and Smad nuclear exclusion [39]. However, this effect is debatable and may be cell context-dependent [40, 41]. Several reports showed that Erk or JNK activation by RTKs leads to strong phosphorylation of endogenous Smad2/3 in mammalian cells without affecting their nuclear accumulation and transcriptional activity [42-44]. More recently, three residues in the linker region of Smad3 (Thr178, Ser203, and Ser207) were identified as Erk1/2 phosphorylation sites both *in vitro* and *in vivo*. Erk-mediated phosphorylation of these sites inhibits Smad3 transcriptional activity but does not prevent Smad3 from entering the nucleus [45], suggesting the existence of a yet unknown mechanism for Smad3 inhibition by the linker phosphorylation. Adding more complexity, the same serine residues (203 and 207) of Smad3 are also targets for other kinases. In human breast cancer MCF10CA1h cells, both Rho-dependent kinase (ROCK) and p38 MAPK phosphorylate Ser203/207 and facilitate, rather than inhibit,

TGF- $\beta$ -induced growth inhibition [46]. Our recent study indicates that GSK3- $\beta$ , which is structurally similar to MAPKs, selectively phosphorylates Ser203 of Smad3 *in vivo* [47]. We also noticed that, unlike Erk, which primarily phosphorylates Smad3 linker in the nucleus ([45] and our unpublished result), GSK3- $\beta$  mainly phosphorylates the cytoplasmic Smad3 (our unpublished result). The linker phosphorylation by GSK3- $\beta$  does not seem to affect Smad3 localization or activity, and its functional role in TGF- $\beta$  signaling is unknown. Together, these findings suggest that linker phosphorylation of Smad2/3 can yield distinct outcomes depending on the identity of the kinase, the specific intracellular localization where the phosphorylation occurs, the collateral events caused by MAPK activation, and other cell type-specific factors (Figure 2).

MAPKs (especially Erk1/2) also phosphorylate the linker of Smad1/5, which almost always blocks Smad1/5 nuclear translocation. As a result, BMP function can be suppressed by several signals that activate RTK/MAPK, including EGF, fibroblast growth factor (FGF) and insulin-like growth factor (IGF) [48-50]. Multiple Ser/Thr residues in Smad1 linker can be sequentially phosphorylated by Erk and then GSK3- $\beta$ , creating a docking site for the Smad1/5-specific E3 ubiquitin ligase, Smurf1. Smurf1 binding not only causes ubiquitination and degradation of the Smads but also occludes their interaction with the nuclear pore complex, thereby preventing Smad nuclear translocation [50]. As a functional consequence, FGF/MAPK relieves BMP-mediated repression to induce neural differentiation of *Xenopus* embryonic cells and rat neural precursor cells [51, 52]. Importantly, Wnt signaling, which is known to inactivate GSK3- $\beta$ , reduces Smad1 ubiquitination and stabilizes the protein [53]. Together, these studies have provided a compelling molecular mechanism for the long-known Wnt—| FGF —| BMP axis during embryonic patterning and cell differentiation [54]. It is interesting to note that MAPK- and GSK3- $\beta$ -mediated linker phosphorylation has not been shown to regulate the protein stability of Smad2/3. Such difference between Smad1/5 and Smad2/3 could be because of the variation in amino-acid sequences of their respective linker regions. A different mode of regulation has been observed in prostate cells, in which Erk-mediated linker phosphorylation allows BMP-activated Smad1 to physically interact with the androgen receptor (AR) and act as a co-repressor. This induced binding of Smad1 and AR culminates in an antagonism of androgen-stimulated prostate cell growth by BMP [55]. Additionally, MH1 domain phosphorylation of Mad (*Drosophila* homolog of Smad1/5) by the MAPK-like kinase Nemo also leads to Mad nuclear exclusion [56].

In addition to R-Smads, MAPKs also phosphorylate and regulate the Co-Smad, Smad4, and the inhibitory Smad,



**Figure 2** TGF- $\beta$ /BMP and RTK/Ras-activated MAPK and PI3K/Akt pathways. The MAPK and PI3K/Akt pathways impinge on TGF- $\beta$ /BMP signaling primarily by modulating Smad functions. MAPKs and Akt bind and/or phosphorylate R-Smads to control their intracellular distribution and transcriptional activity. MAPKs and Akt also phosphorylate and regulate a variety of Smad binding partners in the nucleus, indirectly affecting the Smads.

Smad7. For example, oncogenic Ras decreases Smad4 protein stability in an MEK/Erk-dependent manner [57]. JNK and p38 seem to preferentially phosphorylate tumor-derived mutant Smad4 and promote its proteasomal degradation [58]. Erk, JNK, and p38 have all been implicated in the transcriptional regulation of Smad7, therefore indirectly regulating TGF- $\beta$  signaling [59-61].

Although FGF often suppresses BMP activity as described earlier, these pathways can have synergistic functions. In chicken liver, hepatoblast differentiation into biliary epithelial cells is induced by BMP-4 and also requires an active FGF/MAPK pathway. This collaboration of BMP and FGF occurs specifically at certain stages of liver development and results in the expression of several cell lineage-restricted genes [62]. On the other hand, FGF/MAPK activity can also be opposed by BMPs. Mice lacking the receptor BMPR1a and/or BMPR1b exhibit defects in cartilage development, partly because of an elevation in FGF signaling that suppresses chondrogenesis. In the

growth plate of these mutant mice, both FGFR1 protein level and Erk1/2 activity are higher than in wild-type animals, suggesting inhibition of the FGF/MAPK pathway by BMP [63]. In early mouse embryos, limb-bud outgrowth is promoted by Shh and FGFs, whereas termination of this growth requires BMP-mediated inhibition of FGFs. Interestingly, high-level FGF downregulates Gremlin1, an antagonist of BMP. Thus, a negative feed-back loop is established to tune down FGF signals by indirectly activating BMP, which is an ideal way to prevent limb-bud overgrowth ([64] and references therein).

MAPKs phosphorylate a number of nuclear transcription factors, many of which can physically interact with Smads and regulate TGF- $\beta$ /BMP responses. The best-characterized ones in this category are the AP-1 proteins, including members of the Jun, Fos, Maf, and ATF sub-families [65]. Functional interaction between Smad and the Jun/Fos family proteins has been widely studied, and their relationship can be synergistic or antagonistic depending



on their target genes and other binding partners [66-72]. Recently, inhibition of MafA-dependent transcription by a TGF- $\beta$ -activated Smad complex was reported [73]. The ATF proteins (ATF1, ATF2, and ATF3), which are activated by p38 MAPK, have also been shown to bind Smads and participate in a variety of TGF- $\beta$ -regulated activities [74-78]. It is noteworthy that several of the Jun and ATF sub-family members are themselves Smad target genes, thus establishing the so-called “self-enabling” or “self-disabling” TGF- $\beta$  responses [75,79-81].

### TGF- $\beta$ and the PI3K/Akt pathway

The most widely studied PI3K, p110 $\alpha$ /p85, converts phosphatidylinositol-4,5-bisphosphate (PIP<sub>2</sub>) to phosphatidylinositol-3,4,5-triphosphate (PIP<sub>3</sub>). PIP<sub>3</sub> is a potent signaling molecule that recruits and regulates a number of downstream effectors, the most important one being the serine/threonine kinase Akt (also known as PKB). RTK/Ras, integrin, and several other signals can activate the PI3K/Akt pathway, which usually promotes cell survival, growth, and motility through Akt-mediated phosphorylation of a slew of relevant proteins. Consistent with these physiological functions, oncogenic mutations and protein overproduction of PI3K and Akt are commonly found in human cancers. The activity of PI3K is counteracted by the tumor suppressor protein PTEN (phosphatase and tensin homolog deleted on chromosome 10), which is a lipid phosphatase that removes the phosphate group from the 3' position of the inositol ring of PIP<sub>3</sub>, thereby blocking Akt activation. Loss-of-function mutations of *PTEN* also occur at a high frequency in human cancers (reviewed by [23, 82-84]).

The PI3K/Akt activity is known to alleviate TGF- $\beta$ -induced apoptosis and/or cell cycle arrest in multiple types of cells in response to insulin, IGF, interleukin (IL)-6, and viral proteins [85-89]. Interestingly, Smad3, but not Smad2, seems to be the primary target of inhibition by PI3K/Akt, consistent with the indispensable function of Smad3 in mediating the pro-apoptotic effects of TGF- $\beta$  [6].

Several mechanisms have been proposed by which PI3K/Akt restricts Smad3 activity (Figure 2). Studies using pharmacological inhibitors of PI3K or the downstream kinase mammalian target of rapamycin (mTOR) have implicated PI3K/Akt in regulating Smad3 activation by the TGF- $\beta$  receptors, although conflicting results have been observed such that PI3K/Akt could either enhance or attenuate TGF- $\beta$  responses [90, 91]. Exactly how PI3K/Akt modulates Smad3 activation remains a question unanswered. Other reports have suggested a “Smad3-trap” model for the anti-TGF- $\beta$  effects of Akt in protecting liver cancer cells from cell death [92, 93]. PI3K-activated, plasma

membrane-anchored Akt can physically sequester Smad3 and block its nuclear translocation in a kinase-independent manner, without affecting the C-terminal phosphorylation (activation) of Smad3. However, this mechanism may be cell type-specific and dependent on the stoichiometry between the Akt and Smad3 proteins, as Akt can also facilitate Smad3 function (see below), and PI3K/Akt has been shown to be required for the nuclear accumulation of BMP-activated Smad1 [94]. A third way for PI3K/Akt to debilitate Smad3 is through the inactivation of certain nuclear factors that are necessary for Smad3 function. For instance, activated Smad3 directly interacts with the FoxO family of transcription factors and they are simultaneously recruited to the promoter of p21 as an integral step of the TGF- $\beta$ -induced cytostatic program [95]. However, both extracellular (e.g. IGF) and intracellular (e.g. Bcr-Abl) signals can defy this activity of TGF- $\beta$  by enhancing Akt-mediated phosphorylation of FoxO, which forces FoxO out of the nucleus and foils p21 induction [95,96].

On the flip side, the PI3K/Akt pathway is also subjected to TGF- $\beta$ /BMP regulation. Akt activity increases in response to TGF- $\beta$  treatment, which seems to be required for a variety of TGF- $\beta$ -induced activities, such as cell migration of HER2-expressing breast cancer cells [97], EMT of normal mammary epithelial cells [98, 99], cell survival of mouse hippocampal neurons and mesenchymal cells [100, 101], as well as growth stimulation of certain fibroblasts [102]. BMP also activates Akt to induce osteoblast differentiation [94]. It is to be noted that Akt activation by TGF- $\beta$ /BMP is cell type-dependent and very likely indirect, often requiring either MAPKs or autocrine actions of secreted molecules.

Alteration of PTEN function represents another route for TGF- $\beta$ /BMP to influence Akt activity. TGF- $\beta$  has been shown to transcriptionally downregulate PTEN in Smad4 null pancreatic cancer cells, which, again, seems to rely on the function of the Ras/MAPK pathway [103]. In the same cells, TGF- $\beta$  elicits EMT by dislodging  $\beta$ -catenin from the adherence junctions, a process that involves TGF- $\beta$ -dependent PTEN dissociation from  $\beta$ -catenin and Akt activation [104]. On the other hand, TGF- $\beta$ /Smad can reduce Akt activity in hematopoietic cells by inducing the expression of SHIP (SH2 domain-containing 5' inositol phosphatase), a lipid phosphatase that removes the 5 position phosphate from PIP<sub>3</sub> [105].

BMP also regulates PTEN activity. In one report, *BM-PR1a* deletion in mouse intestine caused elevated PTEN phosphorylation (indicative of PTEN inactivation) and therefore Akt hyperactivation, suggesting a positive correlation between BMP signaling and PTEN activity. As a result, those mutant mice developed polyposis that was reminiscent of human diseases with perturbed functions

of BMPR1a and PTEN [106,107]. In addition, BMP-2 treatment of MCF7 human breast cancer cells slightly increased the protein level and stability of PTEN [108]. In contrast, Beck and Carethers [109] noticed that long-term BMP-2 treatment of Smad4-null SW480 colon cancer cells could decrease the mRNA level of *PTEN*, which requires Ras/Erk activity. It is yet to be shown whether the above observations are cell type- and tissue-specific; and more work is needed to understand how TGF- $\beta$ /BMP and PTEN are connected at the molecular level.

### TGF- $\beta$ and the Wnt pathway

The Wnt proteins are secreted, lipid-modified signaling molecules that have diverse roles in regulating cell proliferation, differentiation, migration, survival, and so on. The canonical Wnt signaling is mediated by the transcription co-factor  $\beta$ -catenin, which undergoes nucleocytoplasmic shuttling and is also essential for the formation of adherence junctions between cells through its interaction with the cadherins. In the absence of Wnt, the level of cytosolic  $\beta$ -catenin is controlled by the so-called “ $\beta$ -catenin destruction complex”, comprising Axin, adenomatous polyposis coli (APC), GSK3 and casein kinase 1 alpha (CK1 $\alpha$ ). In this complex, Axin scaffolds all the other members by direct binding to facilitate CK1 $\alpha$ - and GSK3-mediated serial phosphorylation and subsequent polyubiquitination/degradation of  $\beta$ -catenin. This constitutive turnover of  $\beta$ -catenin keeps the Wnt pathway in an “OFF” state. When the Wnt ligand binds to its receptor Frizzled (Fz) and co-receptor LRP5/6, the signal is transduced to the  $\beta$ -catenin destruction complex through an intracellular protein Dishevelled (Dvl), leading to Axin downregulation, GSK3 inactivation and  $\beta$ -catenin stabilization. The cytosolic accumulation of  $\beta$ -catenin favors its translocation into the nucleus, where it binds the Lef/TCF (lymphocyte enhancer factor/T-cell transcription factor) family of transcription factors and turns the Wnt pathway on. Given the power of Wnt signaling to stimulate cell proliferation, hyperactivation of this pathway often contributes to carcinogenesis [110-113].

The cross-talk between TGF- $\beta$ /BMP and Wnt pathways has been known for a long time and is probably the most extensively studied. The two pathways are intertwined throughout the life of an animal, and molecularly they interact at multiple levels. First, TGF- $\beta$ /BMP and Wnt reciprocally regulate their ligand production, which is critical for establishing extracellular gradients of these morphogens during embryonic development. Second, the best-defined venue of TGF- $\beta$ /Wnt cross-talk is in the nucleus, where the Smad/ $\beta$ -catenin/Lef protein complex regulates a host of shared target genes, often in a synergistic manner. Third, recent research has identified cytoplasmic

interactions between components of these pathways as novel mechanisms for fine-tuning their respective signaling. In fact, this multi-level paradigm also holds true for the cross-talk between TGF- $\beta$  and other pathways such as Hh, Notch, IL, and interferon-gamma (IFN $\gamma$ ), which will be discussed later (Figures 3 and 4). Moreover, the molecular details of such cross-talk are often conserved across species, further highlighting the biological relevance of integrated signaling.


Mutual regulations between TGF- $\beta$ /BMP and Wnt ligands are prevalent and important during early development, but are also seen in adult tissues. In chicken embryos, Wnt-8c induces the expression of Nodal (a TGF- $\beta$  family member) in a  $\beta$ -catenin-dependent manner, which is required for the establishment of the left-right body axis [114]. In *Xenopus*, BMP2/4 regulate Wnt-8 expression and they cooperatively pattern the mesoderm [115], whereas BMP-2 downregulates Wnt-7a and  $\beta$ -catenin in chicken embryonic mesenchymal cells in a p38-dependent manner, leading to enhanced chondrogenesis [75]. Oncogenic  $\beta$ -catenin has been implicated in promoting BMP-4 production in human colon cancer cells [116]. However, Kosinski *et al.* [117] showed that Wnt activity overlaps with the expression of a number of BMP antagonists at the bottom of colon crypts, which is believed to be the niche for intestinal stem cells. This study suggests that one of the mechanisms by which Wnt signaling maintains the intestinal stem cell population is through inhibition of the opposing activity of BMP. In *Xenopus* embryos, Wnt and TGF- $\beta$  co-regulate the expression of connective tissue growth factor (CTGF). Interestingly, CTGF directly interacts with the BMP-4 and TGF- $\beta$ 1 ligands in the extracellular space, which blocks the ligand-receptor binding of BMP but enhances that of TGF- $\beta$  [118]. CTGF is also co-regulated by Wnt and BMP in mesenchymal stem cells, and its induction stunts the osteoblastic differentiation driven by BMP [119]. Another Wnt target, Cripto, is a TGF- $\beta$ /Nodal co-receptor that plays important roles, both positive and negative, in transmitting TGF- $\beta$ /Nodal signals during development and in cellular transformation as well as tumorigenesis (reviewed in [120]).

Many development-relevant genes in several model systems have been documented to be co-regulated by TGF- $\beta$ /BMP and Wnt pathways at the transcription level, including, but not limited to, *Ultrabithorax*, *Gooseoid*, *Siamois*, *Xnr*, *Chordin*, *Cerberus*, *Crescent*, and *Noggin* [121-125]. During the early stage of vertebrate development, TGF- $\beta$ (Nodal)/BMP and Wnt signaling play critical roles in the formation of the Spemann's organizer, an embryonic dorsal signaling center that controls the movement and fate of neighboring cells [126]. In *Xenopus*, the organizer genes *Xtwn* and *Xsia* are synergistically controlled by both

activin/Vg1-like (members of the TGF- $\beta$  subfamily) and Wnt activity, and inactivation of either pathway leads to significant reduction in *Xtwn* transcription [127]. The two pathways were found to converge at the promoter of *Xtwn*, where Smad4,  $\beta$ -catenin, and Lef1 form a complex and bind to adjacent regulatory elements to co-regulate *Xtwn* expression. The interaction between Smad4 and  $\beta$ -catenin is mediated by Lef1, and their cooperation was soon corroborated in mammalian cells [128]. Interestingly, not all Wnt target genes in the same organizer cells are regulated by Smad4 [127], and neither *Xtwn* nor *Xsia* is affected by BMP, which also signals through Smad4 [129]. These findings suggest that cross-talk between the transcription (co)factors of different pathways is target gene-specific and is dependent on the context of target gene promoters as well as other relevant co-factors (Figure 3).

Besides *Xtwn* and *Xsia*, several other genes are jointly regulated by Smads and  $\beta$ -catenin/Lef in a similar manner

as described above. During mouse embryonic development, Wnt and BMP collaborate to upregulate the *Emx2* and *Msx2* genes that play important roles in neural development, and critical elements in the promoters of both genes were found co-occupied by the Smad/ $\beta$ -catenin/Lef1 complex [130, 131]. In human embryonic carcinoma cells, *Msx1*, *Msx2*, and *Id2* were synergistically induced by Wnt3-A and BMP-4 [132]. In mouse gastric cancer cells, the pro-tumorigenic protein gastrin is activated by both Wnt and TGF- $\beta$  [133]. Interestingly, although the promoter of *gastrin* contains both Smad and Lef/Tcf binding sites, either site alone is sufficient to recruit the Smads/ $\beta$ -catenin/Tcf4 complex. Smads and  $\beta$ -catenin/Tcf4 can function as mutual co-factors, and the interaction between these proteins is thought to be stabilized by the p300 co-activator protein [133]. The implication of cooperative TGF- $\beta$  and Wnt signaling in tumor progression has recently been examined by Labbé *et al.* [134], who identified several shared target



**Figure 3** TGF- $\beta$ /BMP and the Wnt pathway. The most common format of TGF- $\beta$ /Wnt cross-talk occurs in the nucleus, where the Smad and Lef/ $\beta$ -catenin synergistically regulate a set of shared target genes. TGF- $\beta$ /BMP and Wnt can determine the ligand production of each other (see text and Figure 4). In addition, protein interactions in the cytoplasm (such as Smad7-Axin binding) also link the two pathways in various settings.

genes of the two pathways in normal mouse epithelial cells, such as *Ctgf*, *Robo1*, *Gpc1*, and *Inhba*. Importantly, when analyzed in transgenic mouse models, many of these genes were found to be overexpressed in breast and colon tumors with active TGF- $\beta$  and Wnt signaling. Controlled inactivation of the TGF- $\beta$  pathway in these animals resulted in weakened expression of some of the above genes as well as delayed tumor formation, indicating that TGF- $\beta$  and Wnt can synergistically promote tumorigenesis.

An unusual way of Smad3- $\beta$ -catenin cross-talk has been observed in human mesenchymal stem cells (hMSCs). TGF- $\beta$  and Wnt cooperatively stimulate the proliferation of these cells and inhibit their differentiation into the osteocytic and adipocytic lineages, thereby supporting hMSC self-renewal [135]. TGF- $\beta$  stimulation of hMSCs leads to a rapid co-translocation of Smad3 and  $\beta$ -catenin into the nucleus, which is a unique feature of this cell type and is required for the above functions of TGF- $\beta$ . Moreover, Smad3 and  $\beta$ -catenin co-regulate a cohort of genes in these cells that are otherwise not known to be TGF- $\beta$  or Wnt targets, such as the Src family tyrosine kinase BLK. The functions of these genes in TGF- $\beta$ /Wnt-regulated hMSC self-renewal remain to be determined [135].

In contrast to TGF- $\beta$ , BMP impedes Wnt-induced  $\beta$ -catenin translocation and cell proliferation in mouse MSCs as a result of Smad1 interaction with Dvl-1 in the cytoplasm [136]. Axin also interacts with Smad proteins in the cytoplasm. Overexpression studies suggest that Axin can facilitate TGF- $\beta$  signaling by presenting Smad3 to the type I TGF- $\beta$  receptor [137]. However, our findings indicate that endogenous Axin negatively regulates TGF- $\beta$  function by promoting Smad3 basal degradation in a GSK3- $\beta$ -dependent and Wnt-independent fashion [47].

Several lines of evidence have suggested perplexing roles of Smad7 in connecting the TGF- $\beta$  and Wnt pathways. As an inhibitory Smad, Smad7 primarily functions to downregulate the TGF- $\beta$ /BMP receptors and hence R-Smad activation through the recruitment of E3 ubiquitin ligases, Smurf1 and Smurf2 [3]. Smad7 itself is induced by TGF- $\beta$ /BMP and undergoes nucleocytoplasmic shuttling. Smad7 can directly bind  $\beta$ -catenin and promote  $\beta$ -catenin degradation by Smurf2-mediated ubiquitination, thereby reducing Wnt activity. As a result, mice carrying a Smad7 transgene experience abnormal epidermal development, which may be owing to inefficient Wnt signaling in the epidermal stem cells [138]. However, a recent report described a conflicting scenario that Smad7 could bind to Axin, disassemble the  $\beta$ -catenin destruction complex, prevent Smurf2 recruitment, and stabilize  $\beta$ -catenin as well as the adherence junctions [139]. Moreover, Smad7-Axin interaction was also reported to cause Smad7 ubiquitination and degradation through the Axin-assisted action of an E3 ubiquitin ligase

called Arkadia [140]. Finally, Edlund *et al.* [141] observed that TGF- $\beta$  could trigger  $\beta$ -catenin nuclear translocation in a Smad7-dependent manner in human prostate cancer PC-3U cells. The physical interaction between Smad7 and  $\beta$ -catenin was shown to be important for TGF- $\beta$ -induced,  $\beta$ -catenin-regulated apoptotic responses in those cells. The most probable explanation for these seemingly opposite results is that they represent different aspects of Smad7, Axin, and  $\beta$ -catenin functions that are only visible in certain cell types under specific experimental conditions.

## TGF- $\beta$ and the Hh pathway

Like the Wnt proteins, Hedgehog (Hh) is a family of lipid-modified, secreted molecules that participates in a variety of cellular functions, functioning as a potent mitogen and morphogen. In flies, Hh binds the membrane-bound receptor Patched (Ptc) and relieves its suppression on another membrane protein called Smoothed (Smo). This leads to, through a series of intracellular events, activation of the signal transducer, Ci, which is otherwise processed to a repressor form (Ci<sup>rep</sup>) by proteolysis in the cytoplasm of resting cells. Activated Ci (Ci<sup>act</sup>) joins with the co-activator protein CBP in the nucleus and controls Hh target genes. Three Hh proteins are produced in mammals: sonic, desert, and Indian hedgehog (Shh, Dhh, and Ihh), with Shh being the best characterized. The mammalian homologs of Ci are the Gli proteins (Gli1, Gli2, and Gli3). The Hh signal transduction is well conserved during evolution, and aberrant Hh signaling leads to developmental defects as well as tumorigenesis [142-144].

During development and oncogenesis, Hh and TGF- $\beta$ /BMP pathways can directly regulate key components of each other. In the wing imaginal disc of flies, Ci can either suppress or induce the transcription of *Dpp* (Decapentaplegic, the fly version of BMP) depending on the availability of Hh. The different forms of Ci (activator or repressor) bind to the same regulatory elements of the *Dpp* promoter, yet with probably distinct co-factors [145-147]. Luciferase reporter analysis also revealed potential binding sites for the Gli proteins in the promoters of human BMP-4 and -7 genes [148]. Shh provokes an invasive phenotype of cultured gastric cancer cells, which is thought to be mediated by Shh-induced TGF- $\beta$  ligand production as well as T $\beta$ RI expression [149]. During bone development, Shh upregulates TGF- $\beta$ 2 to inhibit hypertrophic chondrocyte differentiation [150]; whereas Shh/Gli2-induced BMP-2 expression is responsible for osteoblast differentiation [151]. On the other hand, notochord-derived activin- $\beta$ B (as well as FGF) represses Shh expression and permits proper development of the pancreas in chicken embryos [152]. Dorsalin-1, a BMP family member, has been shown



to compete with Hh and repress the development of muscle pioneer cells in zebrafish [153].

Smads regulate the *Gli* genes and modulate Hh activity. In several types of cells, TGF- $\beta$ /Smad3 directly induces Gli2 transcription, which in turn upregulates Gli1. Of particular relevance, in pancreatic cancer cells that are resistant to Hh inhibition, blocking TGF- $\beta$  function could attenuate Gli-mediated Hh signaling and reduce cell growth [154]. In developing cerebellum, Shh stimulates the proliferation of granule cell precursors (GCPs) [155], whereas BMPs have the opposite functions [156]. BMP-2 and -4, but not BMP-7, are expressed in the same group of cells and antagonize the proliferative function of Shh specifically through Smad5. Interestingly, Shh pathway components such as Smo and Gli1 can be downregulated by BMP treatment, probably through direct or indirect transcriptional repression [157, 158]. In addition, the BMP-2 target gene TIEG-1 (TGF- $\beta$  inducible early gene-1) blocks Gli-mediated transcription of *N-myc*, an oncogene and essential target of Shh in GCPs, thereby inhibiting cell proliferation and promoting cell differentiation [158].

Several Smad proteins were shown to bind to a C-terminally truncated form of Gli3, which is produced in human diseases and may resemble the endogenously expressed Gli<sup>rep</sup>. Despite the unknown function of this Smad-Gli3 complex, TGF- $\beta$  or BMP treatment could dissociate Smads from the truncated Gli3, which may allow the Gli repressor to antagonize Shh signaling [159].

Our knowledge about the molecular nature of the TGF- $\beta$ /Hh interaction is still limited. However, in view of the similarity between the Hh and Wnt pathways [143, 160], more sophisticated cross-talk between Hh and TGF- $\beta$  is expected.

### TGF- $\beta$ and the Notch pathway

The Notch signaling plays integral roles in cell fate determination and is activated in a cell contact-dependent manner. The Notch protein (receptor) of one cell binds the transmembrane ligand, Jagged or Delta-like, that is expressed on the surface of an adjacent cell. Such ligand engagement triggers the shedding of the ectodomain of Notch proteins and further proteolytic cleavage that releases the Notch intracellular domain (NICD). This NICD fragment translocates into the nucleus and activates the CSL family of transcription factors. This NICD-CSL complex, in conjunction with other co-factor proteins, then drives the expression of Notch target genes, including *Hes*, *Herp*, and *Hey*, which encode the basic helix-loop-helix (bHLH) transcription factors necessary for mediating the downstream effects of Notch [161].

TGF- $\beta$  can induce the expression of Notch ligands. In

Ciona embryos, Nodal induces the local expression of Delta2 to specify the fate of notochord cells [162, 163]. Jagged has been shown to be a TGF- $\beta$  target gene in multiple types of mammalian cells. Smad3-dependent expression of Jagged1 and Hey1 seems to be critical for TGF- $\beta$ -induced EMT in cells derived from several organs [164]. Jagged1 upregulation also contributes to TGF- $\beta$ -stimulated p21 expression and cytoostasis in epithelial cells [165]. Induction of Jagged and Hes1 by TGF- $\beta$  seems particularly evident in diabetic patients with nephropathy, which might be relevant to the pathogenic process [166].

TGF- $\beta$ /BMP and Notch synergistically regulate their common target genes in many cell types. In chicken embryos and in mouse myofibroblasts with active TGF- $\beta$  and Notch signaling, Smad3 and NICD directly interact and form a complex with CSL that binds to specific DNA sequences as found in the promoter of *Hes-1* [167]. A similar cooperation is seen in mouse regulatory T cells (Treg) in that the Notch1 ICD not only interacts with activated Smad3 and facilitates its nuclear translocation [168], but also remains bound with pSmad3 in the nucleus, where they jointly upregulate the transcription factor Foxp3 [169]. Moreover, the membrane-bound form of TGF- $\beta$  expressed in Treg cells activates the Notch pathway in the target cells, an event necessary for Treg/TGF- $\beta$ -regulated immunosuppression [170]. BMP inhibits myogenic differentiation of C2C12 cells, and this function requires an intact Notch pathway as well as Smad1-regulated induction of Notch target genes *Hes-1* and *Hey-1* [171]. In mouse neuroepithelial cells, BMP-2-activated Smad1 joins with NICD with the aid of p300 to regulate the expression of *Hes-5* and *Hesr-1* [172]. Similarly, the Notch target gene *Herp2* is positively regulated by BMP in endothelial cells. However, *Herp2* in turn suppresses the expression of the BMP target gene, *Id1*, which is required for BMP-stimulated endothelial cell migration. Consequently, the transcriptional synergy of the two pathways in fact leads to functional antagonism [173].

Notch can also act against TGF- $\beta$ /BMP. Activated Notch1 has been shown to inhibit the anti-proliferative function of TGF- $\beta$  by sequestering the p300 co-factor from Smads [174]. Notch4 ICD can interact with Smad2/3/4 and inhibit TGF- $\beta$ -regulated cytoostasis, although it does not interfere with Smad2/3 activation [175]. Very recently, Carlson *et al.* [176] reported an interesting antagonism between TGF- $\beta$  and Notch in the muscles. In old muscles with failing ability to regenerate, high levels of TGF- $\beta$  ligand and activated Smad3 were observed in association with decreased activity of Notch. Re-activation of Notch could rejuvenate the muscle cells by preventing Smad3 binding to the promoters of p15, p16, p21, and p27, key regulators of cell cycle and senescence that contribute to

the aging of muscle [176]. A mutual inhibition of Notch3 and TGF- $\beta$  was also seen during the differentiation of 10T1/2 fibroblasts into smooth muscle-like cells [177]. BMP and Notch signaling can also have opposing effects on prostate development in mice. Deletion of BMP7 in mouse urogenital cells led to excessive branching morphogenesis and elevated Notch activity *in vivo*, whereas BMP7 treatment decreased *Hes1* expression *in vitro*. Therefore, BMP and Notch are antagonistic in this setting, although the underlying mechanism is not clear [178].

### TGF- $\beta$ and the IL, TNF $\beta$ , and IFN- $\gamma$ pathways

ILs, tumor necrosis factor-alpha (TNF $\alpha$ ), and IFN $\gamma$  are key regulators of immune functions, inflammatory responses, and many other physiological/pathological activities. These cytokines collectively signal through the Jak/STAT (signal transducers and activators of transcription) pathway and the NF- $\kappa$ B pathway to regulate multiple aspects of cell survival, proliferation, and differentiation. When ILs and IFN $\gamma$  bind their cognate receptors, the Jak kinase becomes activated and phosphorylates the STAT proteins, allowing for STAT dimerization and nuclear translocation. The NF- $\kappa$ B proteins, including p50, p52, RelA/p65, and RelB, exist as dimers and are normally kept inactive by the I $\kappa$ B protein. Extracellular stimuli such as TNF $\alpha$  can trigger the degradation of I $\kappa$ B thereby freeing the NF- $\kappa$ B complexes. Activated NF- $\kappa$ B proteins enter the nucleus and, like the STATs, function as transcription factors to regulate a wide range of target genes involved in the aforementioned processes [179-183]. TGF- $\beta$ , with its diverse and crucial functions in the immune system, inevitably overlaps with ILs, TNF $\alpha$ , and IFN- $\gamma$  in many signaling events (reviewed by [8, 184]). TGF- $\beta$  regulates the bioavailability of these cytokines as well as their signal transduction. In turn, TGF- $\beta$  activity is modulated by these factors in various ways.

TGF- $\beta$  potently suppresses T cell proliferation by inhibiting IL-2 production via Smad3 [185]. TGF- $\beta$ /Smad3 also selectively blocks the expression of IL-2 target genes involved in cell proliferation, without affecting other IL-2 targets that support cell survival [186]. However, TGF- $\beta$ /Smad3 also induces the expression of IL-2 receptor [187]. In addition, TGF- $\beta$  probably synergizes with IL-2/STAT5 signals to upregulate Foxp3, an essential transcription factor for the differentiation of induced Treg (iTreg) cells [188-191]. These results therefore suggest a two-faced relationship between TGF- $\beta$  and IL-2 in T cells, depending on the specific developmental and functional status of the cells.

IL-11 has been identified as a TGF- $\beta$ /Smad target gene, and it plays a unique part in mediating the facilitatory ef-

fects of TGF- $\beta$  during breast cancer bone metastasis [2, 192, 193]. A related finding is that osteoblast-derived TGF- $\beta$ , probably in conjunction with AP-1 and NF- $\kappa$ B, stimulates the expression of IL-8 in human cancer cells [194].

TGF- $\beta$  and IL-6 signaling cooperate in promoting the differentiation of Th17 cells, a new type of T helper cell, although the molecular mechanisms for such cooperation are still elusive ([8] and references therein). IL-6 treatment of human renal proximal tubular epithelial cells promotes TGF- $\beta$  receptor internalization via the endosomal (signaling) pathway as opposed to the caveolar (degradation) pathway, thereby stabilizing the receptor and augmenting Smad3 activity [195]. In contrast, Smad2-mediated inhibition of STAT3 activation by IL-6 has been observed in human intestinal epithelial cells [196].

In mice with T-cell-specific deletion of T $\beta$ RII, a higher amount of IFN $\gamma$  is produced by the CD4<sup>+</sup> and CD8<sup>+</sup> T cells, which correlates with an altered pattern of T-cell differentiation and suggests an inhibition of IFN $\gamma$  expression by TGF- $\beta$  [197, 198]. In line with this, TGF- $\beta$  secreted from tumor cells directly inhibits the production of IFN $\gamma$  and several other factors by cytotoxic T cells, providing an important mechanism for the escape of tumor cells from immune surveillance [77].

Conversely, IFN- $\gamma$  antagonizes TGF- $\beta$  activity in many cellular responses, for which Smad7 induction by Jak1/STAT1 seems to be a major mechanism [199]. On skin injury, loss of IFN- $\gamma$  correlates with a reduction in the Smad7 level and an increase in TGF- $\beta$  ligand production and Smad2 activation, which favors the wound-healing process [200]. Smad7 is also a target of ILs and TNF $\alpha$  (see below). In mouse gastric epithelial cells, hyperactive gp130 (the receptor for IL-6) upregulates Smad7 through activation of STAT3, and subsequent inhibition of TGF- $\beta$  by Smad7 leads to gastric adenoma growth in the animals [201]. IL-7 signaling in pulmonary fibrosis fibroblasts also induces Smad7 expression via Jak1/STAT1 and inhibits the pro-fibrotic functions of TGF- $\beta$  [202].

TGF- $\beta$ /Smad and NF- $\kappa$ B antagonize each other in inflammatory and adaptive immune responses [203]. NF- $\kappa$ B/RelA-mediated Smad7 induction inhibits Smad2/3 activation and allows cells to respond to a variety of pro-inflammatory stimuli, including TNF- $\alpha$  and IL-1 $\beta$  [204]. IL-1 $\beta$  has also been shown to alleviate TGF- $\beta$ -induced growth arrest and transcriptional responses in epithelial cancer cells and hematopoietic cells without conscripting Smad7. Instead, IL-1 $\beta$  activates TAK1, an MAPKKK known to be activated by TGF- $\beta$  and hence the name. TAK1 directly interacts with Smad3 in response to IL-1 $\beta$  and modulates Smad3 activity (probably by phosphorylation) without affecting its C-terminal phosphorylation or nuclear translocation [205].


NF- $\kappa$ B is also subject to TGF- $\beta$  regulation. For example, TGF- $\beta$  treatment of mouse intestinal epithelial cells decreases NF- $\kappa$ B activity by downregulation of the Toll-like receptor 2 (TLR2) protein and subsequently lowers IL-6 production. In IL-10 null mice, owing to severely impaired TGF- $\beta$  signaling, bacterial infection leads to persistent NF- $\kappa$ B activation, which does not involve Smad7 induction [206,207]. On the other hand, an additive effect of Smad and NF- $\kappa$ B/RelA on the expression of type VII collagen (COL7A1) has been reported [208]. Smads have also been shown to bind and cooperate with NF- $\kappa$ B/p52 to upregulate JunB [209].

In *Xenopus* embryos, the leukemia inhibitory factor (LIF), which signals through gp130 and STAT3, inhibits activin/Smad2 activity [210]. On the other hand, BMP2 and LIF (or IL-6) synergistically induce astrocyte formation

from mouse neural progenitor cells. In response to these ligands, Smad1 and STAT3 form a complex with p300 on the promoter of astrocyte-specific genes such as *GFAP* and co-direct cell differentiation [211]. In addition, BMP-induced Id expression provides a permissive environment for LIF to support self-renewal of the ES cells [212].

### Concluding remarks

With a growing list of new regulatory factors and targets being identified, the TGF- $\beta$  pathway has been interwoven into the vast network of cell signaling. Countless experiments performed in distinct systems and/or with different approaches have provided all sorts of answers as to how the TGF- $\beta$ /BMP pathway interacts with the rest of this signaling network. Sometimes these studies show contra-



**Figure 4** A simplified and unified view of the multi-level cross-talk. TGF- $\beta$ /BMP communicates with other signals at several levels. Physical interactions between pathway components, both in the cytoplasm and in the nucleus, can lead to protein redistribution and/or post-translational modifications, and eventually changes in target gene expression. Often the ligands themselves, as well as certain extracellular regulators of the ligands, are transcriptional targets of other pathways. All these events are context-dependent.

dictory results that are not easily reconcilable. Although such discrepancies could have arisen from the variations of experimental conditions, they may also reflect the true adaptability that an organism must possess to survive in a constantly changing environment. In this sense, signaling cross-talk between different pathways creates a comprehensive view of the outside world, so that the cell can orchestrate these pieces of information and respond in an accurate, efficient, and balanced manner.

The core concept of TGF- $\beta$  signaling cross-talk is the context dependency. It is clear from all the above discussions that no simple rule can be easily generalized to describe how TGF- $\beta$  interacts with any other signaling cascade. All experimental data should be interpreted with specific confinement parameters, including cell type, developmental stage, physiological/pathological status, protein intracellular localization, nature of modifying enzymes, co-factors, identity of targets, and so forth.

It should also be noted that cross-talk between pathways can be direct or indirect, unidirectional or bidirectional, and often occurs as part of a feed-back loop. On the other hand, synergy or antagonism can also result from independent inputs that do not cross-talk. Therefore, in-depth mechanistic studies are necessary to distinguish the cause from the consequence and to identify the specific convergence point of the pathways (Figure 4).

The outcome of any signaling cross-talk is an integrated and quantitative reflection of all individual input signals, which should be kept within a physiologically relevant range. Otherwise, distorted artifacts may occur. With the availability of techniques such as RNAi and gene targeting, some of the earlier conclusions drawn from experiments solely dependent on overexpression and dominant-negative strategies warrant re-evaluation.

Owing to space limit, several other pathways that functionally interact with TGF- $\beta$  signaling are not covered in this review, including the nuclear receptor and apoptosis pathways. On the other hand, recent literatures have suggested some new areas that are interconnected with TGF- $\beta$  activity, such as energy metabolism (glucose uptake/consumption, AMPK and mTOR signaling) and NO (nitric oxide) signaling. In addition, the recent discovery that TGF- $\beta$ /BMP/Smad signaling also regulates microRNA expression has pointed out a path toward yet another unexplored territory of TGF- $\beta$  research [213]. It can be envisioned that macro-scale screening/modeling in combination with current techniques in molecular biology, biochemistry, genetics, structural biology, and bioinformatics will reveal a good number of new signaling partners of TGF- $\beta$  and greatly assist the elucidation of many fundamental mechanisms regarding TGF- $\beta$  function and regulation.

## Acknowledgments

We apologize to the researchers whose work was not included in this review owing to space limit. Xiao-Fan Wang was supported by NIH grants DK064113 and GM083000.

## References

- 1 Massagué J. How cells read TGF-beta signals. *Nat Rev Mol Cell Biol* 2000; **1**:169-178.
- 2 Gomis RR, Alarcon C, Nadal C, Van Poznak C, Massague J. C/EBPbeta at the core of the TGFbeta cyostatic response and its evasion in metastatic breast cancer cells. *Cancer Cell* 2006; **10**:203-214.
- 3 Shi Y, Massagué J. Mechanisms of TGF-beta signaling from cell membrane to the nucleus. *Cell* 2003; **113**:685-700.
- 4 Bierie B, Moses HL. Tumour microenvironment: TGFbeta: the molecular Jekyll and Hyde of cancer. *Nat Rev Cancer* 2006; **6**:506-520.
- 5 Massagué J, Blain SW, Lo RS. TGF $\beta$  signaling in growth control, cancer, and heritable disorders. *Cell* 2000; **103**:295-309.
- 6 Siegel PM, Shu W, Cardiff RD, Muller WJ, Massagué J. Transforming growth factor  $\beta$  signaling impairs Neu-induced mammary tumorigenesis while promoting pulmonary metastasis. *Proc Natl Acad Sci USA* 2003; **100**:8430-8435.
- 7 Massagué J. TGF[beta] in cancer. *Cell* 2008; **134**:215-230.
- 8 Li MO, Flavell RA. TGF-[beta]: a master of all T cell trades. *Cell* 2008; **134**:392-404.
- 9 Heldin C-H, Miyazono K, Dijke Pt. TGF- $\beta$  signalling from cell membrane to nucleus through SMAD proteins. *Nature* 1997; **390**:465.
- 10 Sporn MB. The early history of TGF-beta, and a brief glimpse of its future. *Cytokine Growth Factor Rev* 2006; **17**:3-7.
- 11 Anzano MA, Roberts AB, Meyers CA, *et al.* Communication: synergistic interaction of two classes of transforming growth factors from murine sarcoma cells. *Cancer Res* 1982; **42**:4776-4778.
- 12 Roberts AB, Anzano MA, Lamb LC, Smith JM, Sporn MB. New class of transforming growth factors potentiated by epidermal growth factor: isolation from non-neoplastic tissues. *Proc Natl Acad Sci USA* 1981; **78**:5339-5343.
- 13 Roberts AB, Lamb LC, Newton DL, Sporn MB, De Larco JE, Todaro GJ. Transforming growth factors: isolation of polypeptides from virally and chemically transformed cells by acid/ethanol extraction. *Proc Natl Acad Sci USA* 1980; **77**:3494-3498.
- 14 Smith JM, Sporn MB, Roberts AB, Derynck R, Winkler ME, Gregory H. Human transforming growth factor-[alpha] causes precocious eyelid opening in newborn mice. *Nature* 1985; **315**:515-516.
- 15 Mishra L, Shetty K, Tang Y, Stuart A, Byers SW. The role of TGF-beta and Wnt signaling in gastrointestinal stem cells and cancer. *Oncogene* 2005; **24**:5775-5789.
- 16 Pires-daSilva A, Sommer RJ. The evolution of signalling pathways in animal development. *Nat Rev Genet* 2003; **4**:39-49.
- 17 Attisano L, Labbe E. TGFbeta and Wnt pathway cross-talk. *Cancer Metastasis Rev* 2004; **23**:53-61.
- 18 Gerhart J. 1998 Warkany lecture: signaling pathways in develop-



- ment. *Teratology* 1999; **60**:226-239.
- 19 Sumi T, Tsuneyoshi N, Nakatsuji N, Suemori H. Defining early lineage specification of human embryonic stem cells by the orchestrated balance of canonical Wnt/ $\beta$ -catenin, Activin/Nodal and BMP signaling. *Development* 2008; **135**:2969-2979.
  - 20 Rao M. Conserved and divergent paths that regulate self-renewal in mouse and human embryonic stem cells. *Dev Biol* 2004; **275**:269-286.
  - 21 Chang L, Karin M. Mammalian MAP kinase signalling cascades. *Nature* 2001; **410**:37-40.
  - 22 Karnoub AE, Weinberg RA. Ras oncogenes: split personalities. *Nat Rev Mol Cell Biol* 2008; **9**:517-531.
  - 23 Hanahan D, Weinberg RA. The hallmarks of cancer. *Cell* 2000; **100**:57-70.
  - 24 Moustakas A, Heldin CH. Non-Smad TGF- $\beta$  signals. *J Cell Sci* 2005; **118**:3573-3584.
  - 25 Derynck R, Zhang YE. Smad-dependent and Smad-independent pathways in TGF- $\beta$  family signalling. *Nature* 2003; **425**:577-584.
  - 26 Muraoka RS, Koh Y, Roebuck LR, *et al.* Increased malignancy of Neu-induced mammary tumors overexpressing active transforming growth factor  $\beta$ 1. *Mol Cell Biol* 2003; **23**:8691-8703.
  - 27 Ueda Y, Wang S, Dumont N, Yi JY, Koh Y, Arteaga CL. Overexpression of HER2 (erbB2) in human breast epithelial cells unmasks transforming growth factor  $\beta$ -induced cell motility. *J Biol Chem* 2004; **279**:24505-24513.
  - 28 Janda E, Lehmann K, Killisch I, *et al.* Ras and TGF $\beta$  cooperatively regulate epithelial cell plasticity and metastasis: dissection of Ras signaling pathways. *J Cell Biol* 2002; **156**:299-313.
  - 29 Wang SE, Xiang B, Guix M, *et al.* Transforming growth factor  $\beta$  engages TACE and ErbB3 to activate PI3K/Akt in ErbB2-overexpressing breast cancer and desensitizes cells to Trastuzumab. *Mol Cell Biol* 2008; **28**: 5605-5620.
  - 30 Seton-Rogers SE, Brugge JS. ErbB2 and TGF- $\beta$ : a cooperative role in mammary tumor progression? *Cell Cycle* 2004; **3**:597-600.
  - 31 Seton-Rogers SE, Lu Y, Hines LM, *et al.* Cooperation of the ErbB2 receptor and transforming growth factor  $\beta$  in induction of migration and invasion in mammary epithelial cells. *Proc Natl Acad Sci USA* 2004; **101**:1257-1262.
  - 32 Lehmann K, Janda E, Pierreux CE, *et al.* Raf induces TGF- $\beta$  production while blocking its apoptotic but not invasive responses: a mechanism leading to increased malignancy in epithelial cells. *Genes Dev* 2000; **14**:2610-2622.
  - 33 Oft M, Peli J, Rudaz C, Schwarz H, Beug H, Reichmann E. TGF- $\beta$ 1 and Ha-Ras collaborate in modulating the phenotypic plasticity and invasiveness of epithelial tumor cells. *Genes Dev* 1996; **10**:2462-2477.
  - 34 Yue J, Mulder KM. Requirement of Ras/MAPK pathway activation by transforming growth factor  $\beta$  for transforming growth factor  $\beta$ 1 production in a Smad-dependent pathway. *J Biol Chem* 2000; **275**:30765-30773.
  - 35 Ventura JJ, Kennedy NJ, Flavell RA, Davis RJ. JNK regulates autocrine expression of TGF- $\beta$ 1. *Mol Cell* 2004; **15**:269-278.
  - 36 Ross KR, Corey DA, Dunn JM, Kelley TJ. SMAD3 expression is regulated by mitogen-activated protein kinase kinase-1 in epithelial and smooth muscle cells. *Cell Signal* 2007; **19**:923-931.
  - 37 Gotzmann J, Fischer ANM, Zojer M, *et al.* A crucial function of PDGF in TGF- $\beta$ -mediated cancer progression of hepatocytes. *Oncogene* 2006; **25**:3170-3185.
  - 38 Bruna A, Darken RS, Rojo F, *et al.* High TGF $\beta$ -Smad activity confers poor prognosis in glioma patients and promotes cell proliferation depending on the methylation of the PDGF-B gene. *Cancer Cell* 2007; **11**:147-160.
  - 39 Kretzschmar M, Doody J, Timokhina I, Massague J. A mechanism of repression of TGF $\beta$ /Smad signaling by oncogenic Ras. *Genes Dev* 1999; **13**:804-816.
  - 40 Hu PP-c, Shen X, Huang D, Liu Y, Counter C, Wang XF. The MEK pathway is required for stimulation of p21WAF1/CIP1 by transforming growth factor- $\beta$ . *J Biol Chem* 1999; **274**:35381-35387.
  - 41 Dunfield LD, Nachtigal MW. Inhibition of the antiproliferative effect of TGF $\beta$  by EGF in primary human ovarian cancer cells. *Oncogene* 2003; **22**:4745-4751.
  - 42 de Caestecker MP, Parks WT, Frank CJ, *et al.* Smad2 transduces common signals from receptor serine-threonine and tyrosine kinases. *Genes Dev* 1998; **12**:1587-1592.
  - 43 Brown JD, DiChiara MR, Anderson KR, Gimbrone Jr MA, Topper JN. MEKK-1, a component of the stress (stress-activated protein kinase/c-Jun N-terminal kinase) pathway, can selectively activate Smad2-mediated transcriptional activation in endothelial cells. *J Biol Chem* 1999; **274**:8797-8805.
  - 44 Mori S, Matsuzaki K, Yoshida K, *et al.* TGF- $\beta$  and HGF transmit the signals through JNK-dependent Smad2/3 phosphorylation at the linker regions. *Oncogene* 2004; **23**:7416-7429.
  - 45 Matsuura I, Wang G, He D, Liu F. Identification and characterization of ERK MAP kinase phosphorylation sites in Smad3. *Biochemistry* 2005; **44**:12546-12553.
  - 46 Kamaraju AK, Roberts AB. Role of Rho/ROCK and p38 MAP kinase pathways in transforming growth factor- $\beta$ -mediated Smad-dependent growth inhibition of human breast carcinoma cells *in vivo*. *J Biol Chem* 2005; **280**:1024-1036.
  - 47 Guo X, Ramirez A, Waddell DS, Li Z, Liu X, Wang XF. Axin and GSK3- $\beta$  control Smad3 protein stability and modulate TGF- $\beta$  signaling. *Genes Dev* 2008; **22**:106-120.
  - 48 Eivers E, Fuentealba LC, De Robertis EM. Integrating positional information at the level of Smad1/5/8. *Curr Opin Genet Dev* 2008; **18**:304-310.
  - 49 Kretzschmar M, Doody J, Massagu J. Opposing BMP and EGF signalling pathways converge on the TGF- $\beta$  family mediator Smad1. *Nature* 1997; **389**:618-622.
  - 50 Sapkota G, Alarcon C, Spagnoli FM, Brivanlou AH, Massague J. Balancing BMP signaling through integrated inputs into the Smad1 linker. *Mol Cell* 2007; **25**:441-454.
  - 51 Kuroda H, Fuentealba L, Ikeda A, Reversade B, De Robertis EM. Default neural induction: neuralization of dissociated *Xenopus* cells is mediated by Ras/MAPK activation. *Genes Dev* 2005; **19**:1022-1027.
  - 52 Bilican B, Fiore-Herich C, Compston A, Allen ND, Chandran S. Induction of Olig2 precursors by FGF involves BMP signalling blockade at the Smad level. *PLoS ONE* 2008; **3**:e2863.
  - 53 Fuentealba LC, Eivers E, Ikeda A, *et al.* Integrating patterning signals: Wnt/GSK3 regulates the duration of the BMP/Smad1 signal. *Cell* 2007; **131**:980-993.
  - 54 Wilson S, Rydstrom A, Trimborn T, *et al.* The status of Wnt signalling regulates neural and epidermal fates in the chick embryo. *Nature* 2001; **411**:325-330.

- 55 Qiu T, Grizzle WE, Oelschlagel DK, Shen X, Cao X. Control of prostate cell growth: BMP antagonizes androgen mitogenic activity with incorporation of MAPK signals in Smad1. *EMBO J* 2007; **26**:346-357.
- 56 Zeng YA, Rahnama M, Wang S, Sosu-Sedzorme W, Verheyen EM. *Drosophila Nemo* antagonizes BMP signaling by phosphorylation of Mad and inhibition of its nuclear accumulation. *Development* 2007; **134**:2061-2071.
- 57 Saha D, Datta PK, Beauchamp RD. Oncogenic Ras represses transforming growth factor-beta/Smad signaling by degrading tumor suppressor Smad4. *J Biol Chem* 2001; **276**:29531-29537.
- 58 Liang M, Liang YY, Wrighton K, et al. Ubiquitination and proteolysis of cancer-derived Smad4 mutants by SCFSkp2. *Mol Cell Biol* 2004; **24**:7524-7537.
- 59 Brodin G, Ahgren A, ten Dijke P, Heldin C-H, Heuchel R. Efficient TGF-beta induction of the Smad7 gene requires cooperation between AP-1, Sp1, and Smad proteins on the mouse Smad7 promoter. *J Biol Chem* 2000; **275**:29023-29030.
- 60 Dowdy SC, Mariani A, Janknecht R. HER2/Neu- and TAK1-mediated up-regulation of the transforming growth factor {beta} inhibitor Smad7 via the ETS protein ER81. *J Biol Chem* 2003; **278**:44377-44384.
- 61 Uchida K, Suzuki H, Ohashi T, Nitta K, Yumura W, Nihei H. Involvement of MAP kinase cascades in Smad7 transcriptional regulation. *Biochem Biophys Res Commun* 2001; **289**:376-381.
- 62 Yanai M, Tatsumi N, Hasunuma N, Katsu K, Endo F, Yokouchi Y. FGF signaling segregates biliary cell-lineage from chick hepatoblasts cooperatively with BMP4 and ECM components *in vitro*. *Dev Dyn* 2008; **237**:1268-1283.
- 63 Yoon BS, Pogue R, Ovchinnikov DA, et al. BMPs regulate multiple aspects of growth-plate chondrogenesis through opposing actions on FGF pathways. *Development* 2006; **133**:4667-4678.
- 64 Verheyden JM, Sun X. An Fgf/Gremlin inhibitory feedback loop triggers termination of limb bud outgrowth. *Nature* 2008; **454**:638-641.
- 65 Shaulian E, Karin M. AP-1 as a regulator of cell life and death. *Nat Cell Biol* 2002; **4**:E131-E136.
- 66 Zhang Y, Feng XH, Derynck R. Smad3 and Smad4 cooperate with c-Jun/c-Fos to mediate TGF-[beta]-induced transcription. *Nature* 1998; **394**:909-913.
- 67 Wong C, Rougier-Chapman EM, Frederick JP, et al. Smad3-Smad4 and AP-1 complexes synergize in transcriptional activation of the c-Jun promoter by transforming growth factor beta. *Mol Cell Biol* 1999; **19**:1821-1830.
- 68 Yamamura Y, Hua X, Bergelson S, Lodish HF. Critical role of Smads and AP-1 complex in transforming growth factor-beta-dependent apoptosis. *J Biol Chem* 2000; **275**:36295-36302.
- 69 Dennler S, Prunier C, Ferrand N, Gauthier JM, Atfi A. c-Jun inhibits transforming growth factor beta-mediated transcription by repressing Smad3 transcriptional activity. *J Biol Chem* 2000; **275**:28858-28865.
- 70 Attisano L, Wrana JL. Smads as transcriptional co-modulators. *Curr Opin Cell Biol* 2000; **12**:235-243.
- 71 Javelaud D, Mauviel A. Crosstalk mechanisms between the mitogen-activated protein kinase pathways and Smad signaling downstream of TGF-[beta]: implications for carcinogenesis. *Oncogene* 2005; **24**:5742-5750.
- 72 Verrecchia F, Vindevoghel L, Lechleider RJ, Uitto J, Roberts AB, Mauviel A. Smad3/AP-1 interactions control transcriptional responses to TGF-beta in a promoter-specific manner. *Oncogene* 2001; **20**:3332-3340.
- 73 Matsumura H, Kudo T, Harada A, et al. Suppression of MafA-dependent transcription by transforming growth factor-beta signaling. *Biochem Biophys Res Commun* 2007; **364**:151-156.
- 74 Hanafusa H, Ninomiya-Tsuji J, Masuyama N, et al. Involvement of the p38 mitogen-activated protein kinase pathway in transforming growth factor-beta-induced gene expression. *J Biol Chem* 1999; **274**:27161-27167.
- 75 Jin EJ, Lee SY, Choi YA, Jung JC, Bang OS, Kang SS. BMP-2-enhanced chondrogenesis involves p38 MAPK-mediated down-regulation of Wnt-7a pathway. *Mol Cells* 2006; **22**:353-359.
- 76 Sano Y, Harada J, Tashiro S, Gotoh-Mandeville R, Maekawa T, Ishii S. ATF-2 is a common nuclear target of Smad and TAK1 pathways in transforming growth factor-beta signaling. *J Biol Chem* 1999; **274**:8949-8957.
- 77 Thomas DA, Massagué J. TGF-[beta] directly targets cytotoxic T cell functions during tumor evasion of immune surveillance. *Cancer Cell* 2005; **8**:369-380.
- 78 Monzen K, Hiroi Y, Kudoh S, et al. Smads, TAK1, and their common target ATF-2 play a critical role in cardiomyocyte differentiation. *J Cell Biol* 2001; **153**:687-698.
- 79 Verrecchia F, Tacheau C, Schorpp-Kistner M, Angel P, Mauviel A. Induction of the AP-1 members c-Jun and JunB by TGF-beta/Smad suppresses early Smad-driven gene activation. *Oncogene* 2001; **20**:2205-2211.
- 80 Jonk LJ, Itoh S, Heldin CH, ten Dijke P, Kruijer W. Identification and functional characterization of a Smad binding element (SBE) in the JunB promoter that acts as a transforming growth factor-beta, activin, and bone morphogenetic protein-inducible enhancer. *J Biol Chem* 1998; **273**:21145-21152.
- 81 Kang Y, Chen CR, Massagué J. A self-enabling TGF $\beta$  response coupled to stress signaling: Smad engages stress response factor ATF3 for Id1 repression in epithelial cells. *Mol Cell* 2003; **11**:915-926.
- 82 Cantley LC. The phosphoinositide 3-kinase pathway. *Science* 2002; **296**:1655-1657.
- 83 Bader AG, Kang S, Zhao L, Vogt PK. Oncogenic PI3K deregulates transcription and translation. *Nat Rev Cancer* 2005; **5**:921-999.
- 84 Salmena L, Carracedo A, Pandolfi PP. Tenets of PTEN tumor suppression. *Cell* 2008; **133**:403-14.
- 85 Chen RH, Su YH, Chuang RL, Chang TY. Suppression of transforming growth factor-beta-induced apoptosis through a phosphatidylinositol 3-kinase/Akt-dependent pathway. *Oncogene* 1998; **17**:1959-1968.
- 86 Chen RH, Chang MC, Su YH, Tsai YT, Kuo ML. Interleukin-6 inhibits transforming growth factor-beta-induced apoptosis through the phosphatidylinositol 3-kinase/Akt and signal transducers and activators of transcription 3 pathways. *J Biol Chem* 1999; **274**:23013-23019.
- 87 Fukuda M, Longnecker R. Latent membrane protein 2A inhibits transforming growth factor-{beta}1-induced apoptosis through the phosphatidylinositol 3-kinase/Akt pathway. *J Virol* 2004; **78**:1697-1705.
- 88 Shih WL, Kuo ML, Chuang SE, Cheng AL, Doong SL. Hepatitis B virus X protein inhibits transforming growth factor-beta-induced apoptosis through the activation of phosphatidylinositol 3-kinase pathway. *J Biol Chem* 2000; **275**:25858-25864.
- 89 Song K, Cornelius SC, Reiss M, Danielpour D. Insulin-like growth factor-I inhibits transcriptional responses of transform-

- ing growth factor- $\{\beta\}$  by phosphatidylinositol 3-kinase/Akt-dependent suppression of the activation of Smad3 but not Smad2. *J Biol Chem* 2003; **278**:38342-38351.
- 90 Asano Y, Ihn H, Yamane K, Jinnin M, Mimura Y, Tamaki K. Phosphatidylinositol 3-kinase is involved in  $\{\alpha\}2(I)$  collagen gene expression in normal and scleroderma fibroblasts. *J Immunol* 2004; **172**:7123-7135.
- 91 Song K, Wang H, Krebs TL, Danielpour D. Novel roles of Akt and mTOR in suppressing TGF- $\beta$ /ALK5-mediated Smad3 activation. *EMBO J* 2006; **25**:58-69.
- 92 Conery AR, Cao Y, Thompson EA, Townsend Jr CM, Ko TC, Luo K. Akt interacts directly with Smad3 to regulate the sensitivity to TGF- $\beta$  induced apoptosis. *Nat Cell Biol* 2004; **6**:366-372.
- 93 Remy I, Montmarquette A, Michnick SW. PKB/Akt modulates TGF- $\beta$  signalling through a direct interaction with Smad3. *Nat Cell Biol* 2004; **6**:358-365.
- 94 Ghosh-Choudhury N, Abboud SL, Nishimura R, Celeste A, Mahimainathan L, Choudhury GG. Requirement of BMP-2-induced phosphatidylinositol 3-kinase and Akt serine/threonine kinase in osteoblast differentiation and Smad-dependent BMP-2 gene transcription. *J Biol Chem* 2002; **277**:33361-33368.
- 95 Seoane J, Le HV, Shen L, Anderson SA, Massague J. Integration of Smad and forkhead pathways in the control of neuroepithelial and glioblastoma cell proliferation. *Cell* 2004; **117**:211-223.
- 96 Atfi A, Abecassis L, Bourgeade MF. Bcr-Abl activates the AKT/Fox O3 signalling pathway to restrict transforming growth factor- $\beta$ -mediated cytostatic signals. *EMBO Rep* 2005; **6**:985-991.
- 97 Wang SE, Shin I, Wu FY, Friedman DB, Arteaga CL. HER2/Neu (ErbB2) signaling to Rac1-Pak1 is temporally and spatially modulated by transforming growth factor  $\{\beta\}$ . *Cancer Res* 2006; **66**:9591-9600.
- 98 Bakin AV, Tomlinson AK, Bhowmick NA, Moses HL, Arteaga CL. Phosphatidylinositol 3-kinase function is required for transforming growth factor  $\beta$ -mediated epithelial to mesenchymal transition and cell migration. *J Biol Chem* 2000; **275**:36803-36810.
- 99 Lamouille S, Derynck R. Cell size and invasion in TGF- $\{\beta\}$  induced epithelial to mesenchymal transition is regulated by activation of the mTOR pathway. *J Cell Biol* 2007; **178**:437-451.
- 100 Zhu Y, Culmsee C, Klump S, Kriegelstein J. Neuroprotection by transforming growth factor- $\{\beta\}1$  involves activation of nuclear factor- $\{\kappa\}B$  through phosphatidylinositol-3-OH kinase/Akt and mitogen-activated protein kinase-extracellular-signal regulated kinase1, 2 signaling pathways. *Neuroscience* 2004; **123**:897-906.
- 101 Horowitz JC, Lee DY, Waghay M, *et al.* Activation of the pro-survival phosphatidylinositol 3-kinase/AKT pathway by transforming growth factor- $\{\beta\}1$  in mesenchymal cells is mediated by p38 MAPK-dependent induction of an autocrine growth factor. *J Biol Chem* 2004; **279**:1359-1367.
- 102 Wilkes MC, Mitchell H, Penheiter SG, *et al.* Transforming growth factor- $\{\beta\}$  activation of phosphatidylinositol 3-kinase is independent of Smad2 and Smad3 and regulates fibroblast responses via p21-activated kinase-2. *Cancer Res* 2005; **65**:10431-10440.
- 103 Chow JY, Quach KT, Cabrera BL, Cabral JA, Beck SE, Carethers JM. RAS/ERK modulates TGF $\beta$ -regulated PTEN expression in human pancreatic adenocarcinoma cells. *Carcinogenesis* 2007; **28**:2321-2327.
- 104 Vogelmann R, Nguyen-Tat MD, Giehl K, Adler G, Wedlich D, Menke A. TGF $\beta$ -induced downregulation of E-cadherin-based cell-cell adhesion depends on PI3-kinase and PTEN. *J Cell Sci* 2005; **118**:4901-4912.
- 105 Valderrama-Carvajal H, Cocolakis E, Lacerte A, *et al.* Activin/TGF- $\beta$  induce apoptosis through Smad-dependent expression of the lipid phosphatase SHIP. *Nat Cell Biol* 2002; **4**:963-969.
- 106 He XC, Zhang J, Tong WG, *et al.* BMP signaling inhibits intestinal stem cell self-renewal through suppression of Wnt- $\beta$ -catenin signaling. *Nat Genet* 2004; **36**:1117-1121.
- 107 Tian Q, He XC, Hood L, Li L. Bridging the BMP and Wnt pathways by PI3 kinase/Akt and 14-3-3 $\zeta$ . *Cell Cycle* 2005; **4**:215-216.
- 108 Waite KA, Eng C. BMP2 exposure results in decreased PTEN protein degradation and increased PTEN levels. *Hum Mol Genet* 2003; **12**:679-684.
- 109 Beck SE, Carethers JM. BMP suppresses PTEN expression via RAS/ERK signaling. *Cancer Biol Ther* 2007; **6**:1313-1317.
- 110 Haramis A-PG, Begthel H, van den Born M, *et al.* De novo crypt formation and juvenile polyposis on BMP inhibition in mouse intestine. *Science* 2004; **303**:1684-1686.
- 111 Cadigan KM, Nusse R. Wnt signaling: a common theme in animal development. *Genes Dev* 1997; **11**:3286-3305.
- 112 Peifer M, Polakis P. Wnt signaling in oncogenesis and embryogenesis—a look outside the nucleus. *Science* 2000; **287**:1606-1609.
- 113 Polakis P. The many ways of Wnt in cancer. *Curr Opin Genet Dev* 2007; **17**:45-51.
- 114 Rodriguez-Esteban C, Capdevila J, Kawakami Y, Belmonte JCI. Wnt signaling and PKA control nodal expression and left-right determination in the chick embryo. *Development* 2001; **128**:3189-3195.
- 115 Hoppler S, Moon RT. BMP-2/-4 and Wnt-8 cooperatively pattern the *Xenopus* mesoderm. *Mech Dev* 1998; **71**:119-129.
- 116 Kim J-S, Crooks H, Dracheva T, *et al.* Oncogenic  $\{\beta\}$ -catenin is required for bone morphogenetic protein 4 expression in human cancer cells. *Cancer Res* 2002; **62**:2744-2748.
- 117 Kosinski C, Li VSW, Chan ASY, *et al.* Gene expression patterns of human colon tops and basal crypts and BMP antagonists as intestinal stem cell niche factors. *Proc Natl Acad Sci* 2007; **104**:15418-15423.
- 118 Abreu JG, Ketpura NI, Reversade B, De Robertis EM. Connective-tissue growth factor (CTGF) modulates cell signalling by BMP and TGF- $\beta$ . *Nat Cell Biol* 2002; **4**:599-604.
- 119 Luo Q, Kang Q, Si W, *et al.* Connective tissue growth factor (CTGF) is regulated by Wnt and bone morphogenetic proteins signaling in osteoblast differentiation of mesenchymal stem cells. *J Biol Chem* 2004; **279**:55958-55968.
- 120 Strizzi L, Bianco C, Normanno N, Salomon D. Cripto-1: a multifunctional modulator during embryogenesis and oncogenesis. *Oncogene* 2005; **24**:5731-5741.
- 121 Crease DJ, Dyson S, Gurdon JB. Cooperation between the activin and Wnt pathways in the spatial control of organizer gene expression. *Proc Natl Acad Sci USA* 1998; **95**:4398-4403.
- 122 Watabe T, Kim S, Candia A, *et al.* Molecular mechanisms of Spemann's organizer formation: conserved growth factor synergy between *Xenopus* and mouse. *Genes Dev* 1995; **9**:3038-3050.
- 123 Agius E, Oelgeschlager M, Wessely O, Kemp C, De Robertis EM. Endodermal Nodal-related signals and mesoderm induction in *Xenopus*. *Development* 2000; **127**:1173-1183.



- 124 Xanthos JB, Kofron M, Tao Q, Schaible K, Wylie C, Heasman J. The roles of three signaling pathways in the formation and function of the Spemann organizer. *Development* 2002; **129**:4027-4043.
- 125 Riese J, Yu X, Munneryn A, *et al.* LEF-1, a nuclear factor coordinating signaling inputs from wingless and decapentaplegic. *Cell* 1997; **88**:777-787.
- 126 Niehrs C. Regionally specific induction by the Spemann-Mangold organizer. *Nat Rev Genet* 2004; **5**:425-434.
- 127 Nishita M, Hashimoto MK, Ogata S, *et al.* Interaction between Wnt and TGF-beta signalling pathways during formation of Spemann's organizer. *Nature* 2000; **403**:781-785.
- 128 Labbé E, Letamendia A, Attisano L. Association of Smads with lymphoid enhancer binding factor 1/T cell-specific factor mediates cooperative signaling by the transforming growth factor-beta and wnt pathways. *Proc Natl Acad Sci USA* 2000; **97**:8358-8363.
- 129 Laurent MN, Cho KWY. Bone morphogenetic protein antagonism of Spemann's organizer is independent of Wnt signaling. *Dev Biol* 1999; **206**:157-162.
- 130 Theil T, Aydin S, Koch S, Grotewold L, Ruther U. Wnt and Bmp signalling cooperatively regulate graded Emx2 expression in the dorsal telencephalon. *Development* 2002; **129**:3045-3054.
- 131 Hussein SM, Duff EK, Sirard C. Smad4 and {beta}-catenin co-activators functionally interact with lymphoid-enhancing factor to regulate graded expression of Msx2. *J Biol Chem* 2003; **278**:48805-48814.
- 132 Willert J, Epping M, Pollack J, Brown P, Nusse R. A transcriptional response to Wnt protein in human embryonic carcinoma cells. *BMC Dev Biol* 2002; **2**:8.
- 133 Lei S, Dubeykovskiy A, Chakladar A, Wojtukiewicz L, Wang TC. The murine gastrin promoter is synergistically activated by transforming growth factor-{beta}/Smad and Wnt signaling pathways. *J Biol Chem* 2004; **279**:42492-42502.
- 134 Labbé E, Lock L, Letamendia A, *et al.* Transcriptional cooperation between the transforming growth factor-{beta} and Wnt pathways in mammary and intestinal tumorigenesis. *Cancer Res* 2007; **67**:75-84.
- 135 Jian H, Shen X, Liu I, Semenov M, He X, Wang XF. Smad3-dependent nuclear translocation of beta-catenin is required for TGF-beta1-induced proliferation of bone marrow-derived adult human mesenchymal stem cells. *Genes Dev* 2006; **20**:666-674.
- 136 Liu Z, Tang Y, Qiu T, Cao X, Clemens TL. A Dishevelled-1/Smad1 interaction couples WNT and bone morphogenetic protein signaling pathways in uncommitted bone marrow stromal cells. *J Biol Chem* 2006; **281**:17156-17163.
- 137 Furuhashi M, Yagi K, Yamamoto H, *et al.* Axin facilitates Smad3 activation in the transforming growth factor- $\beta$  signaling pathway. *Mol Cell Biol* 2001; **21**:5132-5141.
- 138 Han G, Li AG, Liang Y-Y, *et al.* Smad7-induced [beta]-catenin degradation alters epidermal appendage development. *Dev Cell* 2006; **11**:301-312.
- 139 Tang Y, Liu Z, Zhao L, Clemens TL, Cao X. Smad7 stabilizes beta-catenin binding to E-cadherin complex and promotes cell-cell adhesion. *J Biol Chem* 2008; **283**:23956-23963.
- 140 Liu W, Rui H, Wang J, *et al.* Axin is a scaffold protein in TGF-beta signaling that promotes degradation of Smad7 by Arkadia. *EMBO J* 2006; **25**:1646-1658.
- 141 Edlund S, Lee SY, Grimsby S, *et al.* Interaction between Smad7 and {beta}-catenin: importance for transforming growth factor {beta}-induced apoptosis. *Mol Cell Biol* 2005; **25**:1475-1488.
- 142 Hooper JE, Scott MP. Communicating with Hedgehogs. *Nat Rev Mol Cell Biol* 2005; **6**:306-317.
- 143 Taipale J, Beachy PA. The Hedgehog and Wnt signalling pathways in cancer. *Nature* 2001; **411**:349-354.
- 144 Lum L, Beachy PA. The Hedgehog response network: sensors, switches, and routers. *Science* 2004; **304**:1755-1759.
- 145 Aza-Blanc P, Kornberg TB. Ci: a complex transducer of the Hedgehog signal. *Trends Genet* 1999; **15**:458-462.
- 146 Hepker J, Blackman RK, Holmgren R. Cubitus interruptus is necessary but not sufficient for direct activation of a wing-specific decapentaplegic enhancer. *Development* 1999; **126**:3669-3677.
- 147 Muller B, Basler K. The repressor and activator forms of Cubitus interruptus control Hedgehog target genes through common generic gli-binding sites. *Development* 2000; **127**:2999-3007.
- 148 Kawai S, Sugiura T. Characterization of human bone morphogenetic protein (BMP)-4 and -7 gene promoters: activation of BMP promoters by Gli, a sonic hedgehog mediator. *Bone* 2001; **29**:54-61.
- 149 Yoo YA, Kang MH, Kim JS, Oh SC. Sonic hedgehog signaling promotes motility and invasiveness of gastric cancer cells through TGF-beta-mediated activation of the ALK5-Smad 3 pathway. *Carcinogenesis* 2008; **29**:480-490.
- 150 Alvarez J, Sohn P, Zeng X, Doetschman T, Robbins DJ, Serra R. TGFbeta2 mediates the effects of hedgehog on hypertrophic differentiation and PTHrP expression. *Development* 2002; **129**:1913-1924.
- 151 Zhao M, Qiao M, Harris SE, Chen D, Oyajobi BO, Mundy GR. The zinc finger transcription factor Gli2 mediates bone morphogenetic protein 2 expression in osteoblasts in response to hedgehog signaling. *Mol Cell Biol* 2006; **26**:6197-6208.
- 152 Hebrok M, Kim SK, Melton DA. Notochord repression of endodermal Sonic hedgehog permits pancreas development. *Genes Dev* 1998; **12**:1705-1713.
- 153 Du SJ, Devoto SH, Westerfield M, Moon RT. Positive and negative regulation of muscle cell identity by members of the hedgehog and TGF-beta gene families. *J Cell Biol* 1997; **139**:145-156.
- 154 Dennler S, Andre J, Alexaki I, *et al.* Induction of sonic hedgehog mediators by transforming growth factor-beta: Smad3-dependent activation of Gli2 and Gli1 expression *in vitro* and *in vivo*. *Cancer Res* 2007; **67**:6981-6986.
- 155 Wechsler-Reya RJ, Scott MP. Control of neuronal precursor proliferation in the cerebellum by Sonic Hedgehog. *Neuron* 1999; **22**:103-114.
- 156 Zhu G, Mehler MF, Zhao J, Yu Yung S, Kessler JA. Sonic hedgehog and BMP2 exert opposing actions on proliferation and differentiation of embryonic neural progenitor cells. *Dev Biol* 1999; **215**:118-129.
- 157 Rios I, Alvarez-Rodriguez R, Marti E, Pons S. Bmp2 antagonizes sonic hedgehog-mediated proliferation of cerebellar granule neurons through Smad5 signalling. *Development* 2004; **131**:3159-3168.
- 158 Alvarez-Rodriguez R, Barzi M, Berenguer J, Pons S. Bone morphogenetic protein 2 opposes Shh-mediated proliferation in cerebellar granule cells through a TIEG-1-based regulation of Nmyc. *J Biol Chem* 2007; **282**:37170-37180.
- 159 Liu F, Massagué J, Ruiz i Altaba A. Carboxy-terminally truncated Gli3 proteins associate with Smads. *Nat Genet* 1998; **20**:325-326.



- 160 Kalderon D. Similarities between the Hedgehog and Wnt signaling pathways. *Trends Cell Biol* 2002; **12**:523-531.
- 161 Lai EC. Notch signaling: control of cell communication and cell fate. *Development* 2004; **131**:965-973.
- 162 Hudson C, Yasuo H. A signalling relay involving Nodal and Delta ligands acts during secondary notochord induction in Ciona embryos. *Development* 2006; **133**:2855-2864.
- 163 Hudson C, Lotito S, Yasuo H. Sequential and combinatorial inputs from Nodal, Delta2/Notch and FGF/MEK/ERK signalling pathways establish a grid-like organisation of distinct cell identities in the ascidian neural plate. *Development* 2007; **134**:3527-3537.
- 164 Zavadil J, Cermak L, Soto-Nieves N, Bottinger EP. Integration of TGF-beta/Smad and Jagged1/Notch signalling in epithelial-to-mesenchymal transition. *EMBO J* 2004; **23**:1155-1165.
- 165 Niimi H, Pardali K, Vanlandewijck M, Heldin CH, Moustakas A. Notch signaling is necessary for epithelial growth arrest by TGF- $\beta$ . *J Cell Biol* 2007; **176**:695-707.
- 166 Walsh DW, Roxburgh SA, McGettigan P, et al. Co-regulation of Gremlin and Notch signalling in diabetic nephropathy. *Biochim Biophys Acta* 2008; **1782**:10-21.
- 167 Blokzijl A, Dahlqvist C, Reissmann E, et al. Cross-talk between the Notch and TGF- $\beta$  signaling pathways mediated by interaction of the Notch intracellular domain with Smad3. *J Cell Biol* 2003; **163**:723-728.
- 168 Asano N, Watanabe T, Kitani A, Fuss IJ, Strober W. Notch1 signaling and regulatory T cell function. *J Immunol* 2008; **180**:2796-2804.
- 169 Samon JB, Champhekar A, Minter LM, et al. Notch1 and TGF-beta1 cooperatively regulate Foxp3 expression and the maintenance of peripheral regulatory T cells. *Blood* 2008; **112**:1813-1821.
- 170 Ostroukhova M, Qi Z, Oriss TB, Dixon-McCarthy B, Ray P, Ray A. Treg-mediated immunosuppression involves activation of the Notch-HES1 axis by membrane-bound TGF-beta. *J Clin Invest* 2006; **116**:996-1004.
- 171 Dahlqvist C, Blokzijl A, Chapman G, et al. Functional Notch signaling is required for BMP4-induced inhibition of myogenic differentiation. *Development* 2003; **130**:6089-6099.
- 172 Takizawa T, Ochiai W, Nakashima K, Taga T. Enhanced gene activation by Notch and BMP signaling cross-talk. *Nucleic Acids Res* 2003; **31**:5723-5731.
- 173 Itoh F, Itoh S, Goumans MJ, et al. Synergy and antagonism between Notch and BMP receptor signaling pathways in endothelial cells. *EMBO J* 2004; **23**:541-551.
- 174 Masuda S, Kumano K, Shimizu K, et al. Notch1 oncoprotein antagonizes TGF- $\beta$ /Smad-mediated cell growth suppression via sequestration of coactivator p300. *Cancer Sci* 2005; **96**:274-282.
- 175 Sun Y, Lowther W, Kato K, et al. Notch4 intracellular domain binding to Smad3 and inhibition of the TGF-beta signaling. *Oncogene* 2005; **24**:5365-5374.
- 176 Carlson ME, Hsu M, Conboy IM. Imbalance between pSmad3 and Notch induces CDK inhibitors in old muscle stem cells. *Nature* 2008; **454**:528-532.
- 177 Kennard S, Liu H, Lilly B. Transforming growth factor-beta (TGF-1) down-regulates Notch3 in fibroblasts to promote smooth muscle gene expression. *J Biol Chem* 2008; **283**:1324-1333.
- 178 Grishina IB, Kim SY, Ferrara C, Makarenkova HP, Walden PD. BMP7 inhibits branching morphogenesis in the prostate gland and interferes with Notch signaling. *Dev Biol* 2005; **288**:334-347.
- 179 Haan C, Kreis S, Margue C, Behrmann I. Jaks and cytokine receptors—an intimate relationship. *Biochem Pharmacol* 2006; **72**:1538-1546.
- 180 Yoshimura A. Signal transduction of inflammatory cytokines and tumor development. *Cancer Sci* 2006; **97**:439-447.
- 181 Hayden MS, Ghosh S. Signaling to NF-kappaB. *Genes Dev* 2004; **18**:2195-2224.
- 182 Karin M, Cao Y, Greten FR, Li ZW. NF-kappaB in cancer: from innocent bystander to major culprit. *Nat Rev Cancer* 2002; **2**:301-310.
- 183 Li Q, Verma IM. NF-kappaB regulation in the immune system. *Nat Rev Immunol* 2002; **2**:725-734.
- 184 Gorelik L, Flavell RA. Transforming growth factor-beta in T-cell biology. *Nat Rev Immunol* 2002; **2**:46-53.
- 185 McKarns SC, Schwartz RH, Kaminski NE. Smad3 Is Essential for TGF- $\beta$ 1 to suppress IL-2 production and TCR-induced proliferation, but not IL-2-induced proliferation. *J Immunol* 2004; **172**:4275-4284.
- 186 Nelson BH, Martyak TP, Thompson LJ, Moon JJ, Wang T. Uncoupling of promitogenic and antiapoptotic functions of IL-2 by Smad-dependent TGF-beta signaling. *J Immunol* 2003; **170**:5563-5570.
- 187 Kim HP, Kim BG, Letterio J, Leonard WJ. Smad-dependent cooperative regulation of interleukin 2 receptor alpha chain gene expression by T cell receptor and transforming growth factor-beta. *J Biol Chem* 2005; **280**:34042-34047.
- 188 Burchill MA, Yang J, Vogtenhuber C, Blazar BR, Farrar MA. IL-2 receptor beta-dependent STAT5 activation is required for the development of Foxp3+ regulatory T cells. *J Immunol* 2007; **178**:280-290.
- 189 Davidson TS, DiPaolo RJ, Andersson J, Shevach EM. Cutting edge: IL-2 is essential for TGF-beta-mediated induction of Foxp3+ T regulatory cells. *J Immunol* 2007; **178**:4022-4026.
- 190 Tone Y, Furuuchi K, Kojima Y, Tykocinski ML, Greene MI, Tone M. Smad3 and NFAT cooperate to induce Foxp3 expression through its enhancer. *Nat Immunol* 2008; **9**:194-202.
- 191 Zheng SG, Wang J, Wang P, Gray JD, Horwitz DA. IL-2 Is Essential for TGF-beta to convert naive CD4+CD25- cells to CD25+Foxp3+ regulatory T cells and for expansion of these cells. *J Immunol* 2007; **178**:2018-2027.
- 192 Kang Y, Siegel PM, Shu W, et al. A multigenic program mediating breast cancer metastasis to bone. *Cancer Cell* 2003; **3**:537-549.
- 193 Kang Y, He W, Tulley S, et al. Breast cancer bone metastasis mediated by the Smad tumor suppressor pathway. *Proc Natl Acad Sci USA* 2005; **102**:13909-13914.
- 194 Fong Y-C, Maa M-C, Tsai F-J, et al. Osteoblast-derived TGF- $\beta$  stimulates IL-8 release through AP-1 and NF- $\kappa$ B in human cancer cells. *J Bone Miner Res* 2008; **23**:961-970.
- 195 Zhang XL, Topley N, Ito T, Phillips A. Interleukin-6 regulation of transforming growth factor (TGF)-beta receptor compartmentalization and turnover enhances TGF-beta1 signaling. *J Biol Chem* 2005; **280**:12239-12245.
- 196 Walia B, Wang L, Merlin D, Sitaraman SV. TGF-beta down-regulates IL-6 signaling in intestinal epithelial cells: critical role of SMAD-2. *FASEB J* 2003; **17**:2130-2132.
- 197 Li MO, Sanjabi S, Flavell R. Transforming growth factor- $\beta$  controls development, homeostasis, and tolerance of T cells by

- regulatory T cell-dependent and -independent mechanisms. *Immunity* 2006; **25**:455-471.
- 198 Marie JC, Liggitt D, Rudensky AY. Cellular mechanisms of fatal early-onset autoimmunity in mice with the T cell-specific targeting of transforming growth factor- $\beta$  receptor. *Immunity* 2006; **25**:441-454.
- 199 Ulloa L, Doody J, Massague J. Inhibition of transforming growth factor- $\beta$ /SMAD signalling by the interferon- $\gamma$ /STAT pathway. *Nature* 1999; **397**:710-713.
- 200 Ishida Y, Kondo T, Takayasu T, Iwakura Y, Mukaida N. The essential involvement of cross-talk between IFN- $\gamma$  and TGF- $\beta$  in the skin wound-healing process. *J Immunol* 2004; **172**:1848-1855.
- 201 Jenkins BJ, Grail D, Nheu T, *et al.* Hyperactivation of Stat3 in gp130 mutant mice promotes gastric hyperproliferation and desensitizes TGF- $\beta$  signaling. *Nat Med* 2005; **11**:845-852.
- 202 Huang M, Sharma S, Zhu LX, *et al.* IL-7 inhibits fibroblast TGF- $\beta$  production and signaling in pulmonary fibrosis. *J Clin Invest* 2002; **109**:931-937.
- 203 Letterio JJ, Roberts AB. Regulation of immune responses by TGF- $\beta$ . *Annu Rev Immunol* 1998; **16**:137-161.
- 204 Bitzer M, von Gersdorff G, Liang D, *et al.* A mechanism of suppression of TGF- $\beta$ /SMAD signaling by NF- $\kappa$ B/RelA. *Genes Dev* 2000; **14**:187-197.
- 205 Benus GFJD, Wierenga ATJ, de Gorter DJJ, *et al.* Inhibition of the transforming growth factor  $\beta$  (TGF $\beta$ ) pathway by interleukin-1 $\beta$  is mediated through TGF $\beta$ -activated kinase 1 phosphorylation of SMAD3. *Mol Biol Cell* 2005; **16**:3501-3510.
- 206 Ruiz PA, Shkoda A, Kim SC, Sartor RB, Haller D. IL-10 gene-deficient mice lack TGF- $\beta$ /Smad signaling and fail to inhibit proinflammatory gene expression in intestinal epithelial cells after the colonization with colitogenic *Enterococcus faecalis*. *J Immunol* 2005; **174**:2990-2999.
- 207 Ruiz PA, Shkoda A, Kim SC, Sartor RB, Haller D. IL-10 gene-deficient mice lack TGF- $\beta$ /Smad-mediated TLR2 degradation and fail to inhibit proinflammatory gene expression in intestinal epithelial cells under conditions of chronic inflammation. *Ann NY Acad Sci* 2006; **1072**:389-394.
- 208 Kon A, Vindevoghel L, Kouba DJ, Fujimura Y, Uitto J, Mauviel A. Cooperation between SMAD and NF- $\kappa$ B in growth factor regulated type VII collagen gene expression. *Oncogene* 1999; **18**:1837-1844.
- 209 Lopez-Rovira T, Chalaux E, Rosa JL, Bartrons R, Ventura F. Interaction and functional cooperation of NF- $\kappa$ B with Smads. Transcriptional regulation of the junB promoter. *J Biol Chem* 2000; **275**:28937-28946.
- 210 Nishinakamura R, Matsumoto Y, Matsuda T, *et al.* Activation of Stat3 by cytokine receptor gp130 ventralizes *Xenopus* embryos independent of BMP-4. *Dev Biol* 1999; **216**:481-490.
- 211 Nakashima K, Yanagisawa M, Arakawa H, *et al.* Synergistic signaling in fetal brain by STAT3-Smad1 complex bridged by p300. *Science* 1999; **284**:479-482.
- 212 Ying QL, Nichols J, Chambers I, Smith A. BMP induction of Id proteins suppresses differentiation and sustains embryonic stem cell self-renewal in collaboration with STAT3. *Cell* 2003; **115**:281-292.
- 213 Davis BN, Hilyard AC, Lagna G, Hata A. SMAD proteins control DROSHA-mediated microRNA maturation. *Nature* 2008; **454**:56-61.