

## ORIGINAL ARTICLE

# Significant N<sub>2</sub> fixation by heterotrophs, photoheterotrophs and heterocystous cyanobacteria in two temperate estuaries

Mikkel Bentzon-Tilia<sup>1</sup>, Sachia J Traving<sup>1</sup>, Mustafa Mantikci<sup>2</sup>, Helle Knudsen-Leerbeck<sup>2</sup>, Jørgen LS Hansen<sup>2</sup>, Stiig Markager<sup>2</sup> and Lasse Riemann<sup>1</sup>

<sup>1</sup>Marine Biological Section, Department of Biology, University of Copenhagen, Helsingør, Denmark and

<sup>2</sup>Department of Bioscience, Aarhus University, Roskilde, Denmark

**Nitrogen (N) fixation is fueling planktonic production in a multitude of aquatic environments. In meso- and poly-haline estuaries, however, the contribution of N by pelagic N<sub>2</sub> fixation is believed to be insignificant due to the high input of N from land and the presumed absence of active N<sub>2</sub>-fixing organisms. Here we report N<sub>2</sub> fixation rates, *nifH* gene composition and *nifH* gene transcript abundance for key diazotrophic groups over 1 year in two contrasting, temperate, estuarine systems: Roskilde Fjord (RF) and the Great Belt (GB) strait. Annual pelagic N<sub>2</sub> fixation rates averaged 17 and 61 mmol N m<sup>-2</sup> per year at the two sites, respectively. In RF, N<sub>2</sub> fixation was mainly accompanied by transcripts related to heterotrophic (for example, *Pseudomonas* sp.) and photoheterotrophic bacteria (for example, unicellular diazotrophic cyanobacteria group A). In the GB, the first of two N<sub>2</sub> fixation peaks coincided with a similar *nifH*-expressing community as in RF, whereas the second peak was synchronous with increased *nifH* expression by an array of diazotrophs, including heterotrophic organisms as well as the heterocystous cyanobacterium *Anabaena*. Thus, we show for the first time that significant planktonic N<sub>2</sub> fixation takes place in mesohaline, temperate estuaries and that the importance of heterotrophic, photoheterotrophic and photosynthetic diazotrophs is clearly variable in space and time.**

*The ISME Journal* (2015) 9, 273–285; doi:10.1038/ismej.2014.119; published online 15 July 2014

## Introduction

Assimilation of reactive nitrogen (N) compounds and subsequent sedimentation of organic matter in aquatic environments may lead to N limitation of primary production (Howarth, 1988a). Fixation of atmospheric N<sub>2</sub> accounts for more than 80% of the N input in some of these environments, thereby sustaining a significant fraction of the new production (Granhall and Lundgren, 1971; Horne and Galat, 1985). However, other N-limited aquatic systems do not display significant N<sub>2</sub> fixation rates (Howarth *et al.*, 1988b). In temperate estuaries, N is often limiting in summer due to the assimilation of reactive N by phytoplankton. As a consequence, the N:P ratio is below the Redfield ratio favoring N<sub>2</sub> fixation (Boynton *et al.*, 1982; Howarth *et al.*, 1988b). Nonetheless, planktonic N<sub>2</sub> fixation in temperate estuaries seems very low and is believed to be absent in estuaries exhibiting salinities >10

(Marino *et al.*, 2002; Howarth and Marino, 2006; Conley *et al.*, 2009). As an exception, the low-salinity (2–8) estuary of the Baltic Sea accommodates extensive summer blooms of heterocystous cyanobacteria (Stal *et al.*, 2003). Yet, the high-sulfate concentrations that accompany more saline water may inhibit growth and N<sub>2</sub> fixation by heterocystous cyanobacteria entering meso–polyhaline estuaries (Marino *et al.*, 2002, 2003). The diazotrophic cyanobacteria currently believed to dominate oceanic N<sub>2</sub> fixation, that is, members of the *Trichodesmium* genus, various cyanobacterial endosymbionts of diatoms, and unicellular diazotrophic cyanobacteria group A (UCYN-A) and B (*Crocospheera watsonii*), have been found almost exclusively in tropical and subtropical waters, with growth and N<sub>2</sub> fixation optima at temperatures >24 °C (Breitbarth *et al.*, 2007; Church *et al.*, 2008; Webb *et al.*, 2009; Moisaner *et al.*, 2010). The UCYN-A group has, however, been observed in *nifH* cDNA clone libraries from waters as cold as 12 °C (Short and Zehr, 2007) and it has been associated with N<sub>2</sub> fixation in temperate coastal waters (Rees *et al.*, 2009; Mulholland *et al.*, 2012). Yet, neither freshwater cyanobacterial diazotrophs nor oceanic cyanobacterial diazotrophs are thought to fix significant amounts of N in temperate estuaries.

Correspondence: L Riemann, Marine Biological Section, Department of Biology, University of Copenhagen, Strandpromenaden 5, Helsingør DK-3000, Denmark.

E-mail: [riemann@bio.ku.dk](mailto:riemann@bio.ku.dk)

Received 15 March 2014; revised 3 June 2014; accepted 5 June 2014; published online 15 July 2014

New data on nitrogenase reductase (*nifH*) gene composition continue to expand the potential biome of marine N<sub>2</sub> fixation both in terms of latitude and depth (Mehta *et al.*, 2003; Farnelid *et al.*, 2011; Fernandez *et al.*, 2011; Hamersley *et al.*, 2011) and diverse putative heterotrophic diazotrophs have been reported from temperate estuaries (for example, Affourtit *et al.*, 2001; Jenkins *et al.*, 2004; Farnelid *et al.*, 2009), yet these organisms may thrive due to other traits than diazotrophy (Short and Zehr, 2007). They do, however, seem to fix N in estuarine sediments (Fulweiler *et al.*, 2013) and heterotrophic *nifH* gene expression accompanies N<sub>2</sub> fixation in the mesohaline deep waters of the Baltic Sea (Farnelid *et al.*, 2013), indicating that these organisms could potentially be important diazotrophs in estuarine pelagic waters as well.


In this study, we investigated the temporal dynamics of N<sub>2</sub> fixation in surface waters of two contrasting, temperate estuarine systems through monthly samples taken over a 1-year period. We determined the composition of present and active diazotrophs by Illumina amplicon sequencing of *nifH* genes and their RNA transcripts. Finally, we compared N<sub>2</sub> fixation with a suite of environmental parameters and to the temporal dynamics of abundances of key groups of diazotrophs and their *nifH* transcription by quantitative PCR (qPCR) and reverse-transcription quantitative PCR (RT-qPCR).

## Materials and methods

### *Sampling and environmental parameters*

The two contrasting estuaries Roskilde Fjord (RF) and the Great Belt (GB) were sampled monthly throughout 2012. RF is a shallow (3-m mean depth), semi-enclosed and nutrient-rich estuary of 122 km<sup>2</sup>, which receives high-nutrient loads from its 1127 km<sup>2</sup> drainage area (Flindt *et al.*, 1997). The GB is part of the 37 333 km<sup>2</sup> Kattegat and Belt Sea area (17-m mean depth), and is the main connection between the Baltic Sea and the North Sea and is therefore almost permanently stratified (Jørgensen *et al.*, 2013). In GB, strong currents occur, changing between inflow of saline Baltic Sea water from the north and inflow of less saline Baltic Sea water from the south.

Using 5-l free-flow bottles, surface water (1 m) was collected monthly from the stations in RF (55°42.00'N, 12°04.46'E, 4.8-m depth) and GB (55°30.27'N, 10°51.43'E, 35-m depth, Figure 1) in collaboration with the Danish Marine Monitoring Program. Samples were always obtained around noon (1000 to 1300 hours). In January, RF was inaccessible due to ice coverage and water was collected from a pier 1 km southeast of the station. Once on shore, aliquots for bacterial enumeration were fixed, bacterial production incubation initiated, and water for DNA and RNA extractions were filtered and preserved. The remaining analyses were conducted within 1–3 h of sampling.



**Figure 1** Map depicting the sampling locations. One sampling site was located in the shallow RF estuary and the other in the GB strait.

Nutrient concentrations were analyzed using a San ++ Continuous Flow Analyzer (Skalar, Breda, The Netherlands) as previously described (Grasshoff, 1976). Detection limits were 0.3, 0.04 and 0.1  $\mu\text{mol N l}^{-1}$  for  $\text{NH}_4^+$ ,  $\text{NO}_2^-$  and  $\text{NO}_3^-$ , respectively. Chlorophyll a (Chl a) was measured fluorometrically (Holm-Hansen *et al.*, 1965) and dissolved organic carbon (DOC) was measured using a TOC-V<sub>cph</sub> analyzer (Shimadzu, Japan) according to Cauwet (1999). Bacterial production was estimated by <sup>3</sup>H-thymidine incorporation (20 nmol l<sup>-1</sup> final concentration, PerkinElmer, Waltham, MA, USA; Fuhrman and Azam, 1982) using microcentrifugation (Smith and Azam, 1992). Bacterial abundance was determined from samples fixed with glutaraldehyde (1% final concentration) using a FACSCanto II flow cytometer (BD Biosciences, Franklin Lakes, NJ, USA) according to Gasol and Del Giorgio (2000).

### *N<sub>2</sub> fixation rate measurements*

N<sub>2</sub> fixation was measured by <sup>15</sup>N-incorporation (Montoya *et al.*, 1996) on all sampling dates, except for May where <sup>15</sup>N<sub>2</sub> gas was commercially unavailable. Enriched artificial sea water was used as a source of <sup>15</sup>N rather than injecting <sup>15</sup>N<sub>2</sub> tracer gas directly into the samples, as this may underestimate N<sub>2</sub> fixation (Mohr *et al.*, 2010). The artificial sea water was prepared as previously described (Boström *et al.*, 2007a). Following autoclaving, the hot artificial sea water was distributed into 50-ml borosilicate serum vials, which were sealed with butyl rubber stoppers and crimp sealed. In total, 1 ml <sup>15</sup>N<sub>2</sub> tracer gas (Campro Scientific, Veenendaal, The Netherlands;  $\geq 98\%$ ) was introduced into each vial using a gas-tight syringe and incubated

horizontally for 24 h, at 150 r.p.m. rotation, which should ensure 90–100% tracer equilibration (Mohr *et al.*, 2010). Sea water was transferred into 1.2-l polycarbonate bottles; filling the bottles three-fourth of the way up before adding 50 ml <sup>15</sup>N<sub>2</sub>-enriched artificial sea water. Bottles were then filled with sea water until the formation of a meniscus and capped. Six replicate bottles were incubated for 24 h at *in situ* temperature  $\pm 1.5$  °C; three replicates under *in situ* light conditions (Supplementary Table S1) and three replicates in darkness. Following incubation, 600 ml from each replicate was filtered onto pre-combusted (450 °C, 8 h) GF/F filters (Whatman, GE Healthcare, Little Chalfont, UK) and the remaining 600 ml from each replicate was filtered through a 10- $\mu$ m polycarbonate filter (GE Water & Process Technologies, Trevose, PA, USA) and onto pre-combusted GF/F filters for differentiation of N<sub>2</sub> fixation rates in the respective size fractions. Six controls were done at each sampling time point: two background  $\delta^{15}$ N controls, two un-amended controls filtered at the end of incubation and two amended controls filtered immediately after tracer addition. Filters were stored at  $-20$  °C and dried at 60 °C before analysis at the Laboratory of Applied Physical Chemistry, Gent, Belgium on an isotope ratio mass spectrometer (PDZ Europa, Northwich, UK).

#### Extraction of nucleic acids and synthesis of cDNA

Water for DNA extraction was separated into size fractions. In total, 2 l of sea water was filtered onto a 10- $\mu$ m polycarbonate filter (GE Water & Process Technologies) and 1 l of the filtrate was filtered onto a 0.22- $\mu$ m Supor membrane filter (PALL Corporation, Port Washington, NY, USA). The filters were stored at  $-20$  °C in 2-ml cryotubes containing 1 ml sucrose lysis buffer (400 mM NaCl, 750 mM sucrose, 20 mM EDTA, 50 mM Tris-HCL, pH 8). DNA was extracted using a phenol/chloroform-based protocol (Boström *et al.*, 2004) and quantified (PicoGreen, Molecular Probes, Invitrogen, Eugene, OR, USA).

Water samples for RNA extraction were fixed immediately after sampling using stop solution (5% water-saturated phenol in absolute ethanol, Feike *et al.*, 2012). Size fractionation was done as described for DNA samples, and filters were stored at  $-80$  °C in 2-ml cryotubes containing 0.5 ml RNAlater (Ambion, Carlsbad, CA, USA). RNA was extracted (RNeasy kit; Qiagen, Hilden, Germany) using a slightly modified protocol (Boström *et al.*, 2007b) and quantified (NanoDrop, 2000; Thermo Scientific, Waltham, MA, USA). cDNA was synthesized using TaqMan reverse-transcription reagents (Applied Biosystems, Foster City, CA, USA) and the *nifH3* reverse primer (Zehr and McReynolds, 1989). Reverse transcriptase-free control reactions were included for the verification of complete DNA digestion during RNA extraction. Nested *nifH* PCR reactions with these were always negative.

#### *nifH* amplicon sequencing and sequence analysis

*nifH* amplicons were obtained from 56 samples, 28 from each station representing DNA and cDNA from the two size fractions in each of the 7 months: February, March, April, June, July, August and October. In total, 10 ng of DNA or cDNA corresponding to 4 ng RNA was added to Pure Taq Ready-To-Go PCR Beads (GE Healthcare) along with the *nifH3* and *nifH4* primers in a nested PCR approach (Zehr and Turner, 2001). Illumina indices (Supplementary Table S2) were added to amplicons in the second PCR round, which were done in triplicates for each sample. The triplicate PCR products were pooled, purified (AmPure, Agencourt Bioscience Corporation, Beverly, MA, USA), quantified (PicoGreen), and 30 ng from each sample were pooled and subjected to paired-end Illumina sequencing (MiSeq, Berlin Center for Genomics in Biodiversity Research, Berlin, Germany).

Sequence reads were de-multiplexed using CASAVA 1.8.2, assembled in Mothur 1.32.0 (Schloss *et al.*, 2009), trimmed to 216 nucleotides and screened for ambiguous base calls. Poorly assembled sequences and sequences with frame shifts, that is, exhibiting poor alignment to the *nifH* seed sequences in the functional gene pipeline and repository database (<http://fungene.cme.msu.edu/>), were excluded. Chimeras were removed using UCHIME through Mothur. The sequences were clustered at 97% nucleotide similarity and representative sequences from the 84 most abundant operational taxonomic units (OTUs) were aligned in MEGA 5.05 (Tamura *et al.*, 2011). The closest cultivated representatives were identified using the BLAST resource on NCBI and added to the alignment.

Inverse Simpson indices were used as a measure of alpha diversity, and subsampling down to 482 sequences per sample was chosen to accommodate the sample with the lowest number of sequences. Yue & Clayton measures of dissimilarity between samples and non-metrical multidimensional scaling coordinates were calculated in Mothur. Subsampling to 1200 sequences was applied and samples containing less sequences were excluded from the analysis. Non-metrical multidimensional scaling coordinates were plotted using KiNG through the `make_3d_plots.py` script in Qiime (Caporaso *et al.*, 2010). The significance of the divergence between samples explained by size fraction, sampling station, N<sub>2</sub> fixation rates or sample type (DNA or cDNA) was done using analysis of molecular variance. OTUs occurring in more than one cDNA sample from time points exhibiting high N<sub>2</sub> fixation rates ( $> 10$  nmol N l<sup>-1</sup> per day) were identified for each station using the Venn function in Mothur. Log<sub>10</sub> + 1-transformed abundances of these OTUs were used to produce a heatmap using the CIMminer resource (<http://discover.nci.nih.gov/cimminer/>).

Sequences were uploaded to the Sequencing Read Archive database on NCBI (Accession number SRP038933).

### Quantification of *nifH* genes and their expression

Five abundant OTUs (OTU0001, OTU0002, OTU0005, OTU0009 and OTU0014) were selected for quantification in DNA and RNA samples using qPCR and RT-qPCR. Primers and TaqMan probes were designed using Primer3 (<http://bioinfo.ut.ee/primer3-0.4.0/>, Table 1). Specificities of the assays were verified using Primer-BLAST and by PCR tests against six available non-target DNA templates; the four non-target standards above, a *Rhodospseudomonas nifH* template (Farnelid *et al.*, 2014), and a *Clostridium*-like Cluster III template (Bentzon-Tilia *et al.*, 2014). The qPCR standards were synthesized and cloned into the pUC57 vector at GenScript (Piscataway, NJ, USA) and linearized by *Hind*III digestion. Reaction mixtures, thermal cycling settings, standard preparations and inhibition tests were done as previously described (Farnelid *et al.*, 2013), supplementing the reaction mixtures with 1 mM MgCl<sub>2</sub> and 0.2 vol/w % BSA. DNA and cDNA from the two size fractions were pooled according to the concentrations of DNA and RNA observed in the two size fractions after extraction. A total of 5 ng of DNA and cDNA corresponding to 4 ng RNA was added to the respective reactions. The obtained gene and transcript abundances were corrected for the PCR efficiency (Table 1).

## Results

### Environmental parameters

In RF, salinity and water temperature ranged from 12–15 and from –0.6 to 20 °C, respectively. In GB, salinities ranged from 11–24 with reduced values of 11 in February, 13–17 in June–August and 15 in October, suggesting extensive Baltic Sea water outflow in these periods. Water temperatures ranged from 0.4–17 °C in GB. In winter, the concentration of inorganic N was almost 10-fold higher in RF than in GB with combined NO<sub>2</sub><sup>–</sup> and NO<sub>3</sub><sup>–</sup> concentrations reaching values of 45.9 μmol N l<sup>–1</sup> and 5.55 μmol N l<sup>–1</sup>, respectively (Figures 2a and b). NH<sub>4</sub><sup>+</sup> levels were 0.47–7.2 μmol N l<sup>–1</sup> in RF and 0.34–1.9 μmol N l<sup>–1</sup> in GB over the year (Figures 2a and b). Spring blooms appeared during February and March with Chl a concentrations reaching 7.2 μg l<sup>–1</sup> in RF and 4.7 μg l<sup>–1</sup> in GB (Figures 2c and d). Blooms lead to depletion of inorganic N, which increased again from September. PO<sub>4</sub><sup>3–</sup> concentrations displayed the same overall patterns as inorganic N (Figures 2a and b).

The molecular N:P ratios exceeded the Redfield ratio in winter in RF, but decreased to <1 after the spring bloom with dissolved inorganic N concentrations <1 μmol l<sup>–1</sup> in April, June and July. In GB, a N:P ratio of 25 was observed in June, but ratios were otherwise in the range of 2–10 and dissolved inorganic N concentrations were <1 μmol l<sup>–1</sup> most of the year, collectively suggesting that N availability limits primary production in summer at both sites. The level of DOC was higher in RF (478–626 μmol l<sup>–1</sup>) than in GB (175–410 μmol l<sup>–1</sup>, Figure 2a and b). In RF, bacterial production increased from February to April and from May to June where it peaked at 18 μg C l<sup>–1</sup> per day (Figure 2c). In GB, peaks were observed in March (14 μg C l<sup>–1</sup> per day) and in August (16 μg C l<sup>–1</sup> per day; Figure 2d). Bacterial abundances were highest in summer with RF exhibiting abundances about 10-fold higher than GB (Figures 2c and d).


### N<sub>2</sub> fixation

In RF, N<sub>2</sub> fixation increased from the beginning of March with a pronounced N<sub>2</sub> fixation peak in April at 47 nmol N l<sup>–1</sup> per day (Figure 3a) coincided with the demise of the spring bloom and a 100-fold drop in inorganic N. Following the peak, N<sub>2</sub> fixation stayed at 6.0–10 nmol N l<sup>–1</sup> per day in June–August before decreasing in fall to 1.1–2.0 nmol N l<sup>–1</sup> per day and being undetectable throughout the remainder of the year. N<sub>2</sub> fixation rates in bottles incubated in darkness exhibited the same pattern as those exposed to a diurnal light cycle, but rates were lower. At the onset of the N<sub>2</sub> fixation peak in late March, 76% of the fixed N was assimilated in the <10-μm size fraction. This changed to 46% in April and 61% in July and August. The annual areal input of fixed N at this shallow (4.8 m) station was 17 mmol N m<sup>–2</sup> per year.

In GB, N<sub>2</sub> fixation in light was detectable from February to September with distinct peaks in March and July where N<sub>2</sub> fixation rates reached 63 and 83 nmol N l<sup>–1</sup> per day, respectively (Figure 3b). In the parallel dark incubations, a peak was also observed in March, but like for RF, the rates were lower than in the diurnal light cycle incubations. No peak was observed in the parallel dark incubations from July. The proportion of N fixed in the smaller size fraction was higher during the N<sub>2</sub> fixation peak in March (33%) than in July (8%). The upper mixed

**Table 1** Primers and probes designed in this study for quantitative PCR and reverse-transcription quantitative PCR where R<sup>2</sup> is the correlation coefficient of the standard curve and E is PCR efficiency

Target	Forward primer (5'-3')	Reverse primer (5'-3')	Probe (5'-3')	R <sup>2</sup>	E (%)
OTU0001 ( <i>Pseudomonas</i> )	CAAGGCCAGAACACCATCA	TTCGAGGAAGTTGATCGCGG	AAATGGCGGCGGAGGCCG	0.932–0.989	97–104
OTU0002 (UCYN-A)	AGCTGCAGAAAGAGGAACGTG	GGCTCAGGACCACCAGATTC	TCCTGGTTACAACAAGTTTATGTG	0.968–0.971	90–93
OTU0005 ( <i>Anabaena</i> )	TCCAAGGCTCAAACAACCGT	CAACCTACACCGGTTCTGG	CCGCGACGTTAAATCGGTGGAATC	0.926–0.935	95–106
OTU0009 ( <i>Klebsiella</i> )	ACCATTATGGAGATGGCCGC	GCCAATTTGCAGCACGTCTT	TCCGCTCGGTGCGAGGACCTCGA	0.971–0.972	92–100
OTU0014 ( <i>Bradyrhizobium</i> )	GATGCTTGGTGGATTGGCAC	ACAGAGTTCGGCTAAAGCCC	TCCGAGAAGAAGCGGAGGACGTCGAG	0.949–0.995	92–93



**Figure 2** Seasonal changes in environmental data collected in RF (left panels) and the GB (right panels). Concentrations of inorganic nutrients and DOC for RF (a) and GB (b). (c) and (d) show concentrations of Chl a, bacterial production and bacterial abundance in RF and in GB, respectively. Note the difference in bacterial abundance units between stations. Error bars represent standard deviations.


layer was 13-m deep, and the integrated areal input of fixed N in this layer was  $61 \text{ mmol N m}^{-2}$  per year.

A negative correlation was observed between N<sub>2</sub> fixation and  $\text{NH}_4^+$  concentrations in RF ( $P=0.009$ ,  $n=11$ , Supplementary Table S3), but not in GB. Here, N<sub>2</sub> fixation was significantly negatively correlated with combined  $\text{NO}_2^-$  and  $\text{NO}_3^-$  ( $P=0.017$ ,  $n=11$ , Supplementary Table S3). In RF, significant positive correlations were found between N<sub>2</sub> fixation and DOC ( $P=0.005$ ,  $n=11$ ), bacterial production ( $P=0.015$ ,  $n=11$ ) and bacterial abundance ( $P=0.021$ ,  $n=11$ ; Supplementary Table S3). N<sub>2</sub> fixation was not correlated with temperature or Chl a at either of the sampling sites.

#### Composition of the diazotrophic communities and transcript abundances

A total of 455 596 *nifH* sequences were obtained after initial quality check and assembly. Further evaluation of assembly quality and the occurrence of chimeric sequences reduced the number of sequences to a total of 287 494. Clustering the sequences at 97% similarity yielded 11 534 OTUs

after elimination of singletons. The acquired sequences covered 87–99% of the *nifH* gene diversity in the samples. The 84 most abundant OTUs ( $\geq 250$  sequences each) accounted for  $>80\%$  of all sequences and were exclusively affiliated with *nifH* clusters I and III (Figure 4). The most abundant OTU, OTU0001, was related to the gammaproteobacterium *Pseudomonas stutzeri* (cluster I; 99% nucleotide sequence similarity and 99% amino-acid sequence similarity) and accounted for 22% of all sequences. It was not recovered from all time points, but was present in both size fractions at both stations in DNA and RNA samples. The relative abundance of OTU0001 was highest in RNA samples (Figure 4). Eighteen OTUs were affiliated with the *Geobacter/Pelobacter* sub-group of Cluster I. These OTUs were recovered from both sampling sites, yet they all had highest relative abundances in DNA and were almost exclusively detected here. In total, 30 cluster III OTUs were detected at both stations. The majority of these exhibited a higher relative abundance in RF and most were exclusively recovered from this station. They showed higher relative abundance in DNA compared with RNA



**Figure 3** N<sub>2</sub> fixation measured throughout the year in RF (a) and the GB (b). Total N<sub>2</sub> fixation in diurnal light cycle treatments is depicted as open circles, whereas closed circles represent the equivalent fixation in bottles incubated in complete darkness. N<sub>2</sub> fixation data from the two different size fractions (>10 μm or <10 μm) are depicted as bars where available. Error bars indicate standard deviations.

with OTU0014, OTU0090 and OTU0139 being the only three of the 30 cluster III OTUs exhibiting relative abundances of  $\geq 8\%$  in RNA (Figure 4). Seven OTUs were affiliated with cluster I cyanobacteria (Figure 4). Most of these were related to the UCYN-A (16% of all sequences; 99% nucleotide sequence similarity and 99% amino-acid sequence similarity) and recovered from both stations, yet, exhibiting a higher relative abundance in GB (23%) compared with RF (5%). Similarly, the genus *Anabaena* showed a higher relative abundance in GB (8%) than in RF (1%). The fact that a large fraction of the diazotrophic community did not express the *nifH* gene at the seven selected time points was mirrored in the alpha diversity (Inverse Simpson). In DNA from RF, values ranged from 5.6–68 compared with 1.0–3.9 for RNA (Supplementary Figure S1). In GB, *nifH* diversity was higher in the RNA and lower in the DNA compared with RF.


The molecular variance analysis showed a large discrepancy between the diazotrophic community present and the fraction of the community expressing *nifH* ( $P < 0.001$ ,  $n = 52$ , Supplementary Figure S2a) and different community compositions at the two sites ( $P = 0.004$ ,  $n = 52$ , Supplementary Figure

S2b). To identify differences in the active diazotrophic communities, a core *nifH*-expressing microbiome (CExM) was defined for each station (i.e. OTUs recovered in more than one of the four samples per station exhibiting N<sub>2</sub> fixation >10 nmol N l<sup>-1</sup> per day; Supplementary Figure S3). In RF, the *Pseudomonas*-like OTU0001 dominated the CexM at most time points (Figure 5a). Most OTUs from the CexM were related to *Pseudomonas*-like gammaproteobacteria, yet their relative expression level seemed reduced during summer. To gain a quantitative measure of the *nifH* transcription of OTU0001, it was quantified by RT-qPCR (Table 1). Indeed, it was detectable in RNA at 7 of 10 dates from February to November (Figure 6) and transcripts were quantifiable in March ( $1.5 \times 10^3$  l<sup>-1</sup>) and April ( $2.1 \times 10^4$  l<sup>-1</sup>), coinciding with the N<sub>2</sub> fixation peak (Figure 6a). Two OTUs similar to the UCYN-A group (OTU0002 and OTU0151) were also identified as belonging to the CexM and displayed roughly the same pattern as the *Pseudomonas*-like OTUs (Figure 5a). RT-qPCR quantification of the most abundant of these (OTU0002) showed highest activity in spring as well, with transcript abundances of  $1.1 \times 10^3$ – $1.6 \times 10^4$  l<sup>-1</sup> (Figure 6a). In July and August, the relative abundance of OTUs related to other proteobacteria increased in cDNA (Figure 5a). RT-qPCR data showed that transcripts of the *Klebsiella*-like OTU, OTU0009, was undetectable at all but one sampling date in July where it accounted for  $2.3 \times 10^3$  transcripts per litre.

In GB, a more complex CexM was present, representing 27 OTUs (Figure 5b, Supplementary Figure S2b). During the peak in N<sub>2</sub> fixation in July, the relative proportion of *nifH* transcripts of the OTUs belonging to the CexM increased. These OTUs were related to the cyanobacteria UCYN-A and *Anabaena*, but also to several non-cyanobacterial genera. Four of the five OTUs that were included in the RT-qPCR assay were detectable in July, including OTUs related to *Pseudomonas*, UCYN-A, *Anabaena* and *Bradyrhizobium*. UCYN-A and *Anabaena*-like OTUs were, however, the only two quantifiable OTUs with  $1.6 \times 10^3$  and  $3.4 \times 10^4$  transcripts per litre, respectively (Figure 6b). Many of the same OTUs were detected during the N<sub>2</sub> fixation peak in early spring, but a distinct change in the relative abundances of OTUs in the CexM was not observed at this time point. The most active OTUs, up to and during the spring peak, seemed to be the ones related to *Pseudomonas*, UCYN-A, *Bradyrhizobium*, *Vibrio* and *Desulfovibrio* (Figure 5b). In the RT-qPCR assay, OTUs related to *Pseudomonas* and UCYN-A were detected during spring but below the level of quantification (Figure 6b).

#### *nifH* gene abundances

In RF, the OTU related to *Pseudomonas* (OTU0001) reached highest abundance at the onset of the N<sub>2</sub> fixation peak with  $7.1 \times 10^5$  gene copies per




**Figure 4** Neighbor-joining tree of representative sequences from the 84 most abundant OTUs (each representing  $\geq 250$  sequences) resulting from the 97% nucleotide sequence similarity clustering. Each of the OTUs are designated a number chosen by Mothur. Bootstrap values  $\geq 50\%$  are represented by size-proportional gray circles in the tree. The heatmap indicates the relative abundance of sequences associated with a given OTU in each of the following types of samples: GB, RNA,  $< 10 \mu\text{m}$  (A), GB, RNA,  $> 10 \mu\text{m}$  (B), GB, DNA,  $< 10 \mu\text{m}$  (C), GB, DNA,  $> 10 \mu\text{m}$  (D), RF, RNA,  $< 10 \mu\text{m}$  (E), RF, RNA,  $> 10 \mu\text{m}$  (F), RF, DNA,  $< 10 \mu\text{m}$  (G), RF, DNA,  $> 10 \mu\text{m}$  (H). The affiliations of the OTUs with the canonical *nifH* clusters (Chien and Zinder, 1996) is indicated by roman numerals.

litre (Figure 6c). It was not quantifiable in the summer, but increased in abundance again in autumn. During the N<sub>2</sub> fixation peak in April, four of the five targeted OTUs were detectable by qPCR, but only the UCYN-A-like OTU0002 was quantifiable with  $1.1 \times 10^4$  gene copies per litre (Figure 6c).


The *Klebsiella*-like OTU (OTU0009) was detected in February and again from July–September exhibiting relatively low abundances ( $\approx 10^3$  gene copies per litre).

As in RF, the *Pseudomonas*-like OTU reached highest abundances in March in GB just before the



**Figure 5** The relative abundances of OTUs identified to be part of the core *nifH*-expressing microbiome over time in RF (a) and in the GB (b). The OTU-names are given to the left of the heatmap and the closest cultivated relative is indicated to the right of the heatmap. For reference, the volumetric N<sub>2</sub> fixation rates showed in Figure 3 are given above the heatmaps. These rates are split into size fractions where available.





**Figure 6** Results from the reverse-transcription quantitative PCR (RT-qPCR) and quantitative PCR (qPCR) assays performed on RNA and DNA samples taken from February to November in RF and from March to November in the GB. RT-qPCR results from RF (a) and the GB (b). qPCR results from RF (c) and the GB (d). The quantification limit (QL) is defined as the number of gene copies and transcripts needed *in situ* to match the lowest standard; for example, 10 target transcripts in 4 ng of RNA. QL was calculated for all samples and is depicted as a solid black line. OTUs that were detectable, but not quantifiable, are depicted as bars without error bars at the QL line. Error bars indicate standard deviations.

spring peak in N<sub>2</sub> fixation with  $4.8 \times 10^5$  gene copies per litre (Figure 6d). During this peak, all but OTU0009 was detectable (this OTU was never detected at this station). The *Bradyrhizobium*-like OTU0014 was, however, the only quantifiable OTU during the peak where it reached gene abundances of  $9.0 \times 10^3$  gene copies per litre (Figure 6d). During the summer peak in N<sub>2</sub> fixation, all OTUs, except for OTU0009, increased in abundance with OTU0001 reaching the highest abundance with  $1.4 \times 10^5$  gene copies per litre (Figure 6d). The two cyanobacterial OTUs reached their highest abundance just after the N<sub>2</sub> fixation peak in July with  $7.6 \times 10^5$  and  $8.1 \times 10^6$  gene copies per litre for OTU0002 (UCYN-A) and OTU0005 (*Anabaena*), respectively. These abundances decreased gradually following the peak (Figure 6d).

## Discussion

N<sub>2</sub> fixation in meso–polyhaline estuarine waters is believed to be negligible due to the high land runoff discharges of N and the presumed absence or inactivity of diazotrophic bacteria (Marino *et al.*, 2002; Howarth and Marino, 2006; Conley *et al.*, 2009). In contrast, we report substantial N<sub>2</sub> fixation in surface waters of two contrasting temperate estuaries. In fact, the measured rates are among the highest reported for pelagic systems, reaching  $83 \text{ nmol N}_2 \text{ l}^{-1}$  per day in GB. This is higher than, for example, measurements from oceanic *Trichodesmium* blooms (Capone *et al.*, 1997) and from productive, temperate coastal waters, for example, the North American east coast (Mulholland *et al.*, 2012) and the western English Channel (Rees

*et al.*, 2009), but are similar to those measured in the Baltic Sea (Wasmund *et al.*, 2005). Moreover, our molecular analyses revealed that the relative importance of the organisms responsible for the N<sub>2</sub> fixation, heterotrophic, photoheterotrophic and heterocystous cyanobacteria, differed in space and time. This implies that heterocystous cyanobacteria are not always a pre-requisite for pelagic N<sub>2</sub> fixation in temperate estuaries as previously assumed (Howarth *et al.*, 1988b) and that significant N<sub>2</sub> fixation takes place in these environments.

In both estuaries, the N<sub>2</sub> fixation was accompanied by diverse diazotrophic bacteria, mainly related to (photo)heterotrophic bacteria, including the supposedly tropical to subtropical group of unicellular cyanobacteria, UCYN-A (Mazard *et al.*, 2004; Moisaner *et al.*, 2010), which has recently been shown to be a metabolically reduced symbiont of a unicellular prymnesiophyte (Thompson *et al.*, 2012). Interestingly, *nifH* transcripts related to this group have been reported from Chesapeake Bay at water temperatures as low as 12 °C (Short and Zehr, 2007), yet, here we measured *nifH* transcript abundances of  $>10^4 \text{ l}^{-1}$  in RF in February at only 2.5 °C. Hence, our observations suggest that the geographic range of UCYN-A, conceivably of its eukaryotic host, and the associated N<sub>2</sub> fixation extends to temperate regions, even in winter time.

The different physical properties of the two systems were reflected in community composition, with RF exhibiting a more diverse collection of diazotrophs. The cluster III diazotrophs, putative anaerobes, were particularly prevalent in the shallow, more heterotrophic RF, whereas almost all the cyanobacterial OTUs showed highest relative abundance in GB. Hence, we speculate that the surface

community of RF was influenced by re-suspended sediment bacteria, whereas the GB community was influenced by the changing currents (see below).

Our data suggest that the genetic potential to fix N<sub>2</sub> was, in many cases, not mirrored in the diversity of the genes transcribed; especially in the shallow RF in late fall and winter where the water column is well-mixed. Likewise for the temperate Chesapeake Bay, clone libraries indicated that only few phylogenotypes expressed *nifH*, and it was suggested that the distribution of indigenous diazotrophs is not always due to the selective advantage of diazotrophy *per se* (Zehr and Paerl, 2008). Yet, by applying high temporal sampling resolution and sequencing depth, we show that a significant fraction of the diverse heterotrophic diazotrophs present in such estuarine systems indeed occasionally expresses *nifH*. Cyanobacterial diazotrophs may exhibit diel patterns in *nifH* gene expression (for example, Shi *et al.*, 2010). Sampling time can, therefore, have affected the relative proportions of transcripts from the respective functional types of diazotrophs in our data set. However, data on diurnal variation of *nifH* gene expression in heterotrophs is scarce (but see Church *et al.*, 2005; Moisander *et al.*, 2014); hence, at present, the consequences of the sampling time used here remains elusive. Future studies with increased diurnal sampling resolution would conceivably further advance our understanding of the explicit coupling between the key diazotrophs present and their *nifH* gene expression.

In RF, the CexM consisted exclusively of heterotrophic and photoheterotrophic diazotrophs. Their importance for N<sub>2</sub> fixation was substantiated by the fact that significant N<sub>2</sub> fixation was measured in dark incubations and in the <10- $\mu$ m size fraction. N<sub>2</sub> fixation was stimulated by light, but as photoheterotrophic groups like UCYN-A and *Rhodospseudomonas*-like organisms did express *nifH* at this site, light would indeed be expected to stimulate N<sub>2</sub> fixation. Due to co-variation of environmental parameters over a season, correlation analyses should be interpreted with caution. Nevertheless, we see the positive correlations of N<sub>2</sub> fixation with DOC, bacterial abundance and bacterial production as further support for the conclusion that heterotrophic organisms were the principal N<sub>2</sub>-fixing organisms in RF. In GB, the March peak in N<sub>2</sub> fixation was accompanied by a CexM relatively similar to the one in RF, yet it was more diverse and qPCR of transcripts of the five predominant OTUs were below the level of quantification. OTUs related to UCYN-A and *Pseudomonas* were, however, detectable both by qPCR and RT-qPCR at the time of the spring peak in N<sub>2</sub> fixation, and it is likely that these (photo)heterotrophic organisms played an important role at this time where currents moved polyhaline water (>18) in from the north. In fact, the *Pseudomonas*-like OTU reached the highest gene abundances (10<sup>5</sup>–10<sup>6</sup> copies per litre) in spring at both stations. During

the spring peaks, however, abundances of all quantified OTUs were generally low. Whether they were sufficient to account for the measured N<sub>2</sub> fixation rates is difficult to determine, as empirical data on the N<sub>2</sub> fixation potential of marine, heterotrophic diazotrophs and of the UCYN-A group are not currently available. However, adopting recently published assumptions about cell-specific N<sub>2</sub> fixation rates in different groups of marine diazotrophs (Turk-Kubo *et al.*, 2014), the five quantified OTUs could collectively account for N<sub>2</sub> fixation rates in the range of 5–10 nmol N<sub>2</sub>l<sup>-1</sup> per day during the N<sub>2</sub> fixation peak in RF. In contrast to the spring peaks, the summer N<sub>2</sub> fixation peak was mainly associated with the >10- $\mu$ m size fraction and a very low rate of N<sub>2</sub> fixation in dark incubations, as would be expected for heterocystous cyanobacteria (Fay, 1976). Indeed, the *nifH* gene expression of *Anabaena* peaked at this time and *nifH* gene abundances from this group of organisms increased during the N<sub>2</sub> fixation peak in summer, reaching the highest level observed in this study (8.1  $\times$  10<sup>6</sup> copies per litre) just after the peak. It should, however, be noted that cyanobacteria, like the *Anabaena* genus, host multiple chromosomes per cell (Simon, 1977), and hence this high gene abundance does not translate directly into cell numbers. It is conceivable that the *Anabaena*-like OTUs in the Belt Sea area are advected from the Baltic Sea, since the summer N<sub>2</sub> fixation peak coincided with a period of reduced salinity (13–17) presumably driven by a northerly current of Baltic Sea water. The salinity in GB should, however, swiftly constrain the activity of *Anabaena* (Marino *et al.*, 2002; Moisander *et al.*, 2002). Hence, the observation of *nifH* gene-expressing *Anabaena* in the Belt Sea area is surprising. Taken together, we conclude that heterotrophic and photoheterotrophic diazotrophs were responsible for N<sub>2</sub> fixation in both estuaries following the spring bloom, while the heterocystous *Anabaena* was responsible for the pronounced peak in N<sub>2</sub> fixation during summer in the GB.

Our study reports for the first time significant measured rates of pelagic N<sub>2</sub> fixation in Danish coastal waters, even in two contrasting systems. Assuming that the sampled station is representative for the whole RF estuary, the annual N<sub>2</sub> fixation input amounts to about 18 tons. In comparison, the combined RF/Isefjord system received 688 tons of N from land in 2011 (Pedersen *et al.*, 2013). If RF receives half of this N load, pelagic N<sub>2</sub> fixation corresponds to 5% of the N from land. It is worth noting though that part of the input from land will be bound as refractory-dissolved organic N (Jørgensen *et al.*, 2013), and that the biological impact of N from N<sub>2</sub> fixation may be relatively larger. It should be noted, though, that extrapolating N<sub>2</sub> fixation measured at a single station to the basin scale is speculative, since the spatial variability in the RF estuary is not known. It is even more

complicated to calculate the contribution from N<sub>2</sub> fixation in GB due to the dynamic nature of this system. Nonetheless, the GB station is part of the 37 333 km<sup>2</sup> Belt Sea and Kattegat area, which receives 63 000 tons N per year from land (Jørgensen *et al.*, 2013), equal to 121 mmol N m<sup>-2</sup> per year. In comparison, the mean rate of N<sub>2</sub> fixation measured in this study was 61 mmol N m<sup>-2</sup> per year, corresponding to 34 000 tons N per year for the Belt Sea and Kattegat area. Hence, it seems that in GB, pelagic N<sub>2</sub> fixation could represent a substantial input. However, the rates are probably much lower in the northern and more saline Kattegat, and as the N fixation is mainly restricted to two days, any extrapolation is very uncertain. Nevertheless, it is roughly consistent with the annual input from N<sub>2</sub> fixation of 11 000 tons N suggested from a recent modeling study (Jørgensen *et al.*, 2013).

The present study highlights the need for more N<sub>2</sub> fixation measurements to get confident measures of the contribution of pelagic N<sub>2</sub> fixation to the systemic N budgets in the estuaries examined here. Taken together with the recent reports of significant N<sub>2</sub> fixation in estuarine sediments (Fulweiler *et al.*, 2013; Brown and Jenkins, 2014), the basin-scale implications of estuarine N<sub>2</sub> fixation are likely considerable, suggesting that a reevaluation of the significance of N<sub>2</sub> fixation in temperate estuaries and coastal waters is warranted. Moreover, the N<sub>2</sub> fixation by heterotrophic, photoheterotrophic and heterocystous bacteria suggested from the present study highlights the importance of integrating all these three fundamentally different groups in future surveys of estuarine diazotrophy.

## Conflict of Interest

The authors declare no conflict of interest.

## Acknowledgements

We thank Pia Moisander, Hanna Farnelid, and three anonymous reviewers for constructive comments improving the manuscript. We thank the Danish Nature Agency for including us in the sampling program. This work was supported by grants 09-066396 and 11-105450 from The Danish Council for Independent Research, Natural Sciences to LR and grant 067259 from the Strategic Research Council of Denmark to SM and HK-L.

## References

Affourtit J, Zehr JP, Paerl HW. (2001). Distribution of nitrogen-fixing microorganisms along the Neuse River Estuary, North Carolina. *Microb Ecol* **41**: 114–123.

Bentzon-Tilia M, Farnelid H, Jürgens K, Riemann L. (2014). Cultivation and isolation of N<sub>2</sub>-fixing bacteria

from suboxic waters in the Baltic Sea. *FEMS Microbiol Ecol* **88**: 358–371.

Boström KH, Riemann L, Kühl M, Hagström Å. (2007a). Isolation and gene quantification of heterotrophic N<sub>2</sub>-fixing bacterioplankton in the Baltic Sea. *Environ Microbiol* **9**: 152–164.

Boström KH, Riemann L, Zweifel UL, Hagström Å. (2007b). *Nodularia* sp. *nifH* gene transcripts in the Baltic Sea proper. *J Plankton Res* **29**: 391–399.

Boström KH, Simu K, Hagström Å, Riemann L. (2004). Optimization of DNA extraction for quantitative marine bacterioplankton community analysis. *Limnol Oceanogr-Meth* **2**: 365–373.

Boynon WR, Kemp WM, Keefe CW. (1982). A comparative analysis of nutrients and other factors influencing estuarine phytoplankton production. In: Kennedy VS (ed) *Estuarine Comparisons*. Academic Press: New York, pp 69–90.

Breitbarth E, Oschlies A, LaRoche J. (2007). Physiological constraints on the global distribution of *Trichodesmium*—effect of temperature on diazotrophy. *Biogeosciences* **4**: 53–61.

Brown SM, Jenkins BD. (2014). Profiling gene expression to distinguish the likely active diazotrophs from a sea of genetic potential in marine sediments. *Environ Microbiol*; e-pub ahead of print 25 March 2014; doi:10.1111/1462-2920.12403.

Capone DG, Zehr JP, Paerl HW, Bergman B, Carpenter EJ. (1997). *Trichodesmium*, a globally significant marine cyanobacterium. *Science* **276**: 1221–1229.

Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK *et al.* (2010). QIIME allows analysis of high-throughput community sequencing data. *Nat Methods* **7**: 335–336.

Cauwet G. (1999). Determination of dissolved organic carbon (DOC) and nitrogen (DON) by high temperature combustion. In: Grasshoff K, Kremling K, Erhard M (eds) *Methods of Seawater Analyses*. Wiley-VCH, pp 407–420.

Chien YT, Zinder SH. (1996). Cloning, functional organization, transcript studies, and phylogenetic analysis of the complete nitrogenase structural genes (*nifHDK2*) and associated genes in the archaeon *Methanosarcina barkeri* 227. *J Bacteriol* **178**: 143–148.

Church MJ, Bjorkman KM, Karl DM, Saito MA, Zehr JP. (2008). Regional distributions of nitrogen-fixing bacteria in the Pacific Ocean. *Limnol Oceanogr* **53**: 63–77.

Church MJ, Short CM, Jenkins BD, Karl DM, Zehr JP. (2005). Temporal patterns of nitrogenase gene (*nifH*) expression in the oligotrophic North Pacific Ocean. *Appl Environ Microbiol* **71**: 5362–5370.

Conley DJ, Paerl HW, Howarth RW, Boesch DF, Seitzinger SP, Havens KE *et al.* (2009). ECOLOGY controlling eutrophication: nitrogen and phosphorus. *Science* **323**: 1014–1015.

Farnelid H, Andersson AF, Bertilsson S, Abu Al-Soud W, Hansen LH, Sørensen S *et al.* (2011). Nitrogenase gene amplicons from global marine surface waters are dominated by genes of non-cyanobacteria. *PLoS One* **6**: e19223.

Farnelid H, Bentzon-Tilia M, Andersson AF, Bertilsson S, Jost G, Labrenz M *et al.* (2013). Active nitrogen-fixing heterotrophic bacteria at and below the chemocline of the central Baltic Sea. *ISME J* **7**: 1413–1423.

Farnelid H, Harder J, Bentzon-Tilia M, Riemann L. (2014). Isolation of heterotrophic diazotrophic bacteria from

- estuarine surface waters. *Environ Microbiol*; e-pub ahead of print 16 December 2013; doi:10.1111/1462-2920.12335.
- Farnelid H, Oberg T, Riemann L. (2009). Identity and dynamics of putative N<sub>2</sub>-fixing picoplankton in the Baltic Sea proper suggest complex patterns of regulation. *Environ Microbiol Rep* **1**: 145–154.
- Fay P. (1976). Factors influencing dark nitrogen fixation in a blue-green alga. *Appl Environ Microbiol* **31**: 376–379.
- Feike J, Jürgens K, Hollibaugh JT, Kruger S, Jost G, Labrenz M. (2012). Measuring unbiased metatranscriptomics in suboxic waters of the central Baltic Sea using a new *in situ* fixation system. *ISME J* **6**: 461–470.
- Fernandez C, Farias L, Ulloa O. (2011). Nitrogen fixation in denitrified marine waters. *PLoS One* **6**: e20539.
- Flindt MR, Kamp-Nielsen L, Marques JC, Pardal MA, Bocci M, Bendoricchio G *et al.* (1997). Description of the three shallow estuaries: Mondego River (Portugal), Roskilde Fjord (Denmark) and the Lagoon of Venice (Italy). *Ecol Model* **102**: 17–31.
- Fuhrman JA, Azam F. (1982). Thymidine Incorporation as a measure of heterotrophic bacterioplankton production in marine surface waters—evaluation and field results. *Mar Biol* **66**: 109–120.
- Fulweiler RW, Brown SM, Nixon SW, Jenkins BD. (2013). Evidence and a conceptual model for the co-occurrence of nitrogen fixation and denitrification in heterotrophic marine sediments. *Mar Ecol-Prog Ser* **482**: 57–68.
- Gasol JM, Del Giorgio PA. (2000). Using flow cytometry for counting natural planktonic bacteria and understanding the structure of planktonic bacterial communities. *Sci Mar* **64**: 197–224.
- Granhall U, Lundgren A. (1971). Nitrogen fixation in Lake Erken. *Limnol Oceanogr* **16**: 711–719.
- Grasshoff K. (1976). *Methods of Seawater Analysis*. Verlag Chemie: Weinheim.
- Hamersley MR, Turk KA, Leinweber A, Gruber N, Zehr JP, Gunderson T *et al.* (2011). Nitrogen fixation within the water column associated with two hypoxic basins in the Southern California Bight. *Aquat Microb Ecol* **63**: 193–205.
- Holm-Hansen O, Lorenzen CJ, Holms RW, Strickland JDH. (1965). Fluorometric determination of chlorophyll. *J Cons Int Explor Mer* **30**: 3–15.
- Horne AJ, Galat DL. (1985). Nitrogen-fixation in an oligotrophic, saline desert lake—Pyramid Lake, Nevada. *Limnol Oceanogr* **30**: 1229–1239.
- Howarth RW. (1988a). Nutrient limitation of net primary production in marine ecosystems. *Annu Rev Ecol Syst* **19**: 89–110.
- Howarth RW, Marino R. (2006). Nitrogen as the limiting nutrient for eutrophication in coastal marine ecosystems: evolving views over three decades. *Limnol Oceanogr* **51**: 364–376.
- Howarth RW, Marino R, Lane J. (1988b). Nitrogen fixation in freshwater, estuarine, and marine ecosystems. 1. Rates and importance. *Limnol Oceanogr* **33**: 669–687.
- Jenkins BD, Steward GF, Short SM, Ward BB, Zehr JP. (2004). Fingerprinting diazotroph communities in the Chesapeake Bay by using a DNA macroarray. *Appl Environ Microbiol* **70**: 1767–1776.
- Jørgensen L, Markager S, Maar M. (2013). On the importance of quantifying bioavailable nitrogen instead of total nitrogen. *Biogeochemistry*; e-pub ahead of print 17 July 2013; doi:10.1007/s10533-013-9890-9.
- Marino R, Chan F, Howarth RW, Pace M, Likens GE. (2002). Ecological and biogeochemical interactions constrain planktonic nitrogen fixation in estuaries. *Ecosystems* **5**: 719–725.
- Marino R, Howarth RW, Chan F, Cole JJ, Likens GE. (2003). Sulfate inhibition of molybdenum-dependent nitrogen fixation by planktonic cyanobacteria under seawater conditions: a non-reversible effect. *Hydrobiologia* **500**: 277–293.
- Mazard SL, Fuller NJ, Orcutt KM, Bridle O, Scanlan DJ. (2004). PCR analysis of the distribution of unicellular cyanobacterial diazotrophs in the Arabian Sea. *Appl Environ Microbiol* **70**: 7355–7364.
- Mehta MP, Butterfield DA, Baross JA. (2003). Phylogenetic diversity of nitrogenase (*nifH*) genes in deep-sea and hydrothermal vent environments of the Juan de Fuca ridge. *Appl Environ Microbiol* **69**: 960–970.
- Mohr W, Grosskopf T, Wallace DWR, LaRoche J. (2010). Methodological underestimation of oceanic nitrogen fixation rates. *PLoS One* **5**: e12583.
- Moisander PH, Beinart RA, Hewson I, White AE, Johnson KS, Carlson CA *et al.* (2010). Unicellular cyanobacterial distributions broaden the oceanic N<sub>2</sub> fixation domain. *Science* **327**: 1512–1514.
- Moisander PH, McClinton E, Paerl HW. (2002). Salinity effects on growth, photosynthetic parameters, and nitrogenase activity in estuarine planktonic cyanobacteria. *Microb Ecol* **43**: 432–442.
- Moisander PH, Serros T, Paerl RW, Beinart RA, Zehr JP. (2014). Gammaproteobacterial diazotrophs and *nifH* gene expression in surface waters of the South Pacific Ocean. *ISME J* **8**: 1962–1973.
- Montoya JP, Voss M, Kahler P, Capone DG. (1996). A simple, high-precision, high-sensitivity tracer assay for N<sub>2</sub> fixation. *Appl Environ Microbiol* **62**: 986–993.
- Mulholland MR, Bernhardt PW, Blanco-Garcia JL, Mannino A, Hyde K, Mondragon E *et al.* (2012). Rates of dinitrogen fixation and the abundance of diazotrophs in North American coastal waters between Cape Hatteras and Georges Bank. *Limnol Oceanogr* **57**: 1067–1083.
- Pedersen TM, Sand-Jensen K, Markager S, Nielsen LS. (2013). Optical changes in a eutrophic estuary during reduced nutrient loadings. *Estuaries Coasts*; e-pub ahead of print 4 December 2013; doi:10.1007/s12237-013-9732-y.
- Rees AP, Gilbert JA, Kelly-Gerreyn BA. (2009). Nitrogen fixation in the western English Channel (NE Atlantic Ocean). *Mar Ecol-Prog Ser* **374**: 7–12.
- Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB *et al.* (2009). Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Appl Environ Microbiol* **75**: 7537–7541.
- Shi T, Ilikchyan I, Rabouille S, Zehr JP. (2010). Genome-wide analysis of diel gene expression in the unicellular N<sub>2</sub>-fixing cyanobacterium *Crocospaera watsonii* WH 8501. *ISME J* **4**: 621–632.
- Short SM, Zehr JP. (2007). Nitrogenase gene expression in the Chesapeake Bay estuary. *Environ Microbiol* **9**: 1591–1596.
- Simon R. (1977). Macromolecular composition of spores from the filamentous cyanobacterium *Anabaena cylindrica*. *J Bacteriol* **129**: 1154–1155.
- Smith DC, Azam F. (1992). A simple, economical method for measuring bacterial protein synthesis rates in

- seawater using <sup>3</sup>H-leucine. *Mar Microb Food Webs* **6**: 107–114.
- Stal LJ, Albertano P, Bergman B, von Brockel K, Gallon JR, Hayes PK *et al.* (2003). BASIC: Baltic Sea cyanobacteria. An investigation of the structure and dynamics of water blooms of cyanobacteria in the Baltic Sea—responses to a changing environment. *Cont Shelf Res* **23**: 1695–1714.
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. (2011). MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol* **28**: 2731–2739.
- Thompson AW, Foster RA, Krupke A, Carter BJ, Musat N, Vault D *et al.* (2012). Unicellular cyanobacterium symbiotic with a single-celled eukaryotic alga. *Science* **337**: 1546–1550.
- Turk-Kubo KA, Karamchandani M, Capone DG, Zehr JP. (2014). The paradox of marine heterotrophic nitrogen fixation: abundances of heterotrophic diazotrophs do not account for nitrogen fixation rates in the Eastern Tropical South Pacific. *Environ Microbiol*; e-pub ahead of print 7 January 2014; doi:10.1111/1462-2920.12346.
- Wasmund N, Nausch G, Schneider B, Nagel K, Voss M. (2005). Comparison of nitrogen fixation rates determined with different methods: a study in the Baltic Proper. *Mar Ecol-Prog Ser* **297**: 23–31.
- Webb EA, Ehrenreich IM, Brown SL, Valois FW, Waterbury JB. (2009). Phenotypic and genotypic characterization of multiple strains of the diazotrophic cyanobacterium, *Crocosphaera watsonii*, isolated from the open ocean. *Environ Microbiol* **11**: 338–348.
- Zehr JP, McReynolds LA. (1989). Use of degenerate oligonucleotides for amplification of the *nifH* gene from the marine cyanobacterium *Trichodesmium thiebautii*. *Appl Environ Microbiol* **55**: 2522–2526.
- Zehr JP, Paerl HW. (2008). Molecular ecological aspects of nitrogen fixation in the marine environment. In: Kirchman DL (ed) *Microbial Ecology of the Oceans*. John Wiley & Sons: Hoboken, pp 481–525.
- Zehr JP, Turner PJ. (2001). Nitrogen fixation: nitrogenase genes and gene expression. In: Paul JH (ed) *Methods in Microbiology*. Academic Press: New York, pp 271–285.

Supplementary Information accompanies this paper on The ISME Journal website (<http://www.nature.com/ismej>)