

1-1-2009

Simplicity of C^* -algebras associated to row-finite locally convex higher-rank graphs

David Robertson
Syddansk Universitet, Denmark

Aidan Sims
University of Wollongong, asims@uow.edu.au

Follow this and additional works at: <https://ro.uow.edu.au/infopapers>



Part of the [Physical Sciences and Mathematics Commons](#)

Recommended Citation

Robertson, David and Sims, Aidan: Simplicity of C^* -algebras associated to row-finite locally convex higher-rank graphs 2009, 171-192.
<https://ro.uow.edu.au/infopapers/783>

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library: research-pubs@uow.edu.au

Simplicity of C^* -algebras associated to row-finite locally convex higher-rank graphs

Abstract

In a previous work, the authors showed that the C^* -algebra $C^*(\Lambda)$ of a row-finite higher-rank graph Λ with no sources is simple if and only if Λ is both cofinal and aperiodic. In this paper, we generalise this result to row-finite higher-rank graphs which are locally convex (but may contain sources). Our main tool is Farthing's "removing sources" construction which embeds a row-finite locally convex higher-rank graph in a row-finite higher-rank graph with no sources in such a way that the associated C^* -algebras are Morita equivalent.

Keywords

associated, row, finite, simplicity, graphs, C^* , locally, convex, algebras, higher, rank

Disciplines

Physical Sciences and Mathematics

Publication Details

Robertson, D. & Sims, A. (2009). Simplicity of C^* -algebras associated to row-finite locally convex higher-rank graphs. *Israel Journal of Mathematics*, 172 171-192.

SIMPLICITY OF C^* -ALGEBRAS ASSOCIATED TO ROW-FINITE LOCALLY CONVEX HIGHER-RANK GRAPHS

BY

DAVID ROBERTSON

*Institut for Matematik og Datalogi (IMADA), Syddansk Universitet
Campusvej 55, DK-5230 Odense M, Danmark
e-mail: dir@imada.sdu.dk*

AND

AIDAN SIMS

*School of Mathematics and Applied Statistics, Austin Keane Building (15)
University of Wollongong
NSW, 2522, Australia
e-mail: asims@uow.edu.au*

ABSTRACT

In a previous work, the authors showed that the C^* -algebra $C^*(\Lambda)$ of a row-finite higher-rank graph Λ with no sources is simple if and only if Λ is both cofinal and aperiodic. In this paper, we generalise this result to row-finite higher-rank graphs which are locally convex (but may contain sources). Our main tool is Farthing's "removing sources" construction which embeds a row-finite locally convex higher-rank graph in a row-finite higher-rank graph with no sources in such a way that the associated C^* -algebras are Morita equivalent.

1. Introduction

A directed graph is a quadruple (E^0, E^1, r, s) : E^0 is a countable set of vertices; E^1 is a countable set of directed edges; and r, s are maps from E^1 to E^0 which encode the directions of the edges: an edge e points from the vertex $s(e)$ to the

Received August 2, 2007

vertex $r(e)$. In [4] and [8], C^* -algebras were associated to directed graphs so as to generalise the Cuntz–Krieger algebras of [3]. These graph C^* -algebras have been studied intensively over the last ten years; see [10] for a good summary of the literature.

For technical reasons related to the groupoid models used to analyse their C^* -algebras, the graphs considered in [7, 8] were assumed to be row-finite and to have no sources. This means that $r^{-1}(v)$ is finite and nonempty for every vertex v . To eliminate the “no sources” hypothesis, Bates et al. [2] introduced a construction known as **adding tails**. Adding tails to a graph E with sources produces a graph F with no sources so that the associated C^* -algebras $C^*(E)$ and $C^*(F)$ are Morita equivalent. Proving theorems about $C^*(E)$ then often becomes a question of identifying hypotheses on E which are equivalent to the hypotheses of [7] for F . In [2], many important theorems about C^* -algebras of row-finite graphs with no sources were extended to C^* -algebras of arbitrary row-finite graphs using this strategy.

Higher-rank graphs (or k -graphs) Λ and the associated C^* -algebras $C^*(\Lambda)$ were introduced by Kumjian and Pask in [6] as a common generalisation of graph C^* -algebras and the higher-rank Cuntz–Krieger algebras developed by Robertson and Steger in [15, 16]. As done in [7, 8] for graph C^* -algebras, [6] studied k -graph C^* -algebras using a groupoid model, and technical considerations associated to this model made it necessary to restrict attention to k -graphs which were row-finite and had no sources. Like graph C^* -algebras, k -graph C^* -algebras have received substantial attention in recent years. Unlike graph C^* -algebras, however, some fundamental structure-theoretic questions regarding k -graph algebras have not yet been answered, primarily due to the combinatorial complexities of k -graphs themselves.

In [14], the authors established that the **cofinality** and **aperiodicity** conditions formulated in [6] as sufficient conditions for simplicity of the C^* -algebra of a row-finite higher-rank graph with no sources are in fact also necessary. The resulting simplicity theorem is an exact analogue of the original simplicity theorem for C^* -algebras of row-finite graphs with no sources [7].

In this paper, we extend our previous simplicity result to a large class of higher-rank graphs with sources. To do this we use Farthing’s **removing sources** construction which produces from a higher-rank graph Λ a higher-rank graph $\bar{\Lambda}$ with no sources such that if Λ is row-finite, then $C^*(\Lambda)$ and $C^*(\bar{\Lambda})$ are

Morita equivalent [5]. By establishing how infinite paths in $\overline{\Lambda}$ are related to boundary paths in Λ , we use Farthing's results to generalise the simplicity result of [14] to the **locally convex** row-finite k -graphs with sources considered in [11]. We should point out that the restriction to locally convex k -graphs is not forced on us by Farthing's results, which are applicable for arbitrary row-finite graphs. Indeed, local convexity plays no role in Farthing's analysis, and initially we had no expectation that it would impinge upon our analysis here. However, it turns out, interestingly enough, that local convexity is needed to ensure that the natural projection of $\overline{\Lambda}$ onto Λ extends to a projection from the space of infinite paths of $\overline{\Lambda}$ to the space of boundary paths of Λ . We use this projection to translate aperiodicity and cofinality hypotheses on $\overline{\Lambda}$ to analogous conditions on Λ .

We begin the paper with preliminary notation and definitions in Section 2. We also outline Farthing's removing sources construction in the setting of row-finite locally convex higher-rank graphs, and explore the relationship between boundary paths in Λ and infinite paths in $\overline{\Lambda}$. In Section 3, we turn to our main objective. Using the results of the previous section, we formulate notions of cofinality and aperiodicity for a locally convex row-finite higher-rank graph Λ which are equivalent to the corresponding conditions of [6] for $\overline{\Lambda}$. Combining this with Farthing's Morita equivalence between $C^*(\Lambda)$ and $C^*(\overline{\Lambda})$ and the results of [14] yields the desired simplicity theorem. In Section 4, we apply the same methods to a number of other results of [14] concerning the relationship between graph-theoretic properties of a k -graph Λ and the ideal-structure of $C^*(\Lambda)$.

2. Preliminaries

In this section we summarise the standard notation and conventions for higher-rank graphs.

We regard $\mathbb{N} = \{0, 1, 2, \dots\}$ as a semigroup under addition. We write \mathbb{N}^k for the set of k -tuples $n = (n_1, n_2, \dots, n_k)$, $n_i \in \mathbb{N}$, which we regard as a semigroup under pointwise addition with identity $0 = (0, 0, \dots, 0)$. We denote the canonical generators of \mathbb{N}^k by e_1, e_2, \dots, e_k . Given $m, n \in \mathbb{N}^k$, we say $m \leq n$ if $m_i \leq n_i$ for $1 \leq i \leq k$. We write $m \vee n$ for the coordinate-wise maximum of m and n , and $m \wedge n$ for the coordinate-wise minimum. Unless otherwise

indicated through parentheses, \vee and \wedge always take precedence over addition and subtraction, so for example $m \vee n - n = (m \vee n) - n$.

Note that \vee and \wedge distribute over addition and subtraction:

$$(2.1) \quad a \wedge b + c = (a + c) \wedge (b + c) \quad \text{and} \quad a \vee b + c = (a + c) \vee (b + c)$$

for all $a, b, c \in \mathbb{N}^k$, and

$$(2.2) \quad a \wedge b - c = (a - c) \wedge (b - c) \quad \text{and} \quad a \vee b - c = (a - c) \vee (b - c)$$

(as elements of \mathbb{Z}^k) for all $a, b, c \in \mathbb{N}^k$.

2.1. HIGHER-RANK GRAPHS. The notion of a higher-rank graph is best phrased in terms of categories. For the basics of categories we refer the reader to Chapter 1 of [9]. We assume that all our categories are small in the sense that $\text{Obj}(\mathcal{C})$ and $\text{Mor}(\mathcal{C})$ are sets. Given a category \mathcal{C} , we identify $\text{Obj}(\mathcal{C})$ with $\{\text{id}_o : o \in \text{Obj}(\mathcal{C})\} \subset \text{Mor}(\mathcal{C})$, and we write $c \in \mathcal{C}$ to mean $c \in \text{Mor}(\mathcal{C})$. We write composition in our categories as juxtaposition; that is, for $c_1, c_2 \in \mathcal{C}$ with $\text{dom}(c_1) = \text{cod}(c_2)$, $c_1 c_2$ means $c_1 \circ c_2$. When convenient, we regard \mathbb{N}^k as a category with just one object, and composition implemented by addition. We write $\mathcal{C} \times_{\text{Obj}(\mathcal{C})} \mathcal{C}$ for the set $\{(c_1, c_2) : \text{dom}(c_1) = \text{cod}(c_2)\}$ of composable pairs in \mathcal{C} .

Fix $k \in \mathbb{N} \setminus \{0\}$. A **graph of rank k** or **k -graph** is a countable category Λ equipped with a functor $d : \Lambda \rightarrow \mathbb{N}^k$ called the **degree functor**, which satisfies the **factorisation property**: Given $\lambda \in \Lambda$ with $d(\lambda) = m + n$, there are unique paths $\mu, \nu \in \Lambda$ such that $d(\mu) = m$, $d(\nu) = n$ and $\lambda = \mu\nu$. **Higher-rank graph** is the generic term when the rank k is not specified.

Given a k -graph Λ one can check that $d^{-1}(0) = \{\text{id}_v : v \in \text{Obj}(\Lambda)\}$ using the factorisation property. We define $r, s : \Lambda \rightarrow d^{-1}(0)$ by $r(\lambda) := \text{id}_{\text{cod}(\lambda)}$ and $s(\lambda) := \text{id}_{\text{dom}(\lambda)}$, so $\lambda = r(\lambda)\lambda = \lambda s(\lambda)$ for all $\lambda \in \Lambda$. We think of $d^{-1}(0)$ as the vertices of Λ , and refer to $r(\lambda)$ as the **range** of λ and to $s(\lambda)$ as the **source** of λ .

For $n \in \mathbb{N}^k$ we write $\Lambda^n := d^{-1}(n)$, so Λ^0 is the collection of vertices. For $v \in \Lambda^0$ and $E \subset \Lambda$, we write vE for $\{\lambda \in E : r(\lambda) = v\}$ and $Ev := \{\lambda \in E : s(\lambda) = v\}$.

For $\lambda \in \Lambda$ with $d(\lambda) = l$, and $0 \leq m \leq n \leq l$, the factorisation property ensures that there are unique paths $\lambda' \in \Lambda^m$, $\lambda'' \in \Lambda^{n-m}$ and $\lambda''' \in \Lambda^{l-n}$ such that $\lambda = \lambda'\lambda''\lambda'''$. We write $\lambda(0, m)$, $\lambda(m, n)$ and $\lambda(n, l)$ for λ, λ' and λ''' , respectively.

We say a k -graph Λ is **row finite** if $|v\Lambda^n| < \infty$ for all $v \in \Lambda^0$ and $n \in \mathbb{N}^k$, and that it has **no sources** if $v\Lambda^n \neq \emptyset$ for all $v \in \Lambda^0$ and $n \in \mathbb{N}^k$. We say Λ is **locally convex** if for distinct $i, j \in \{1, \dots, k\}$, and paths $\lambda \in \Lambda^{e_i}$ and $\mu \in \Lambda^{e_j}$ such that $r(\lambda) = r(\mu)$, the sets $s(\lambda)\Lambda^{e_j}$ and $s(\mu)\Lambda^{e_i}$ are non-empty.

For $n \in \mathbb{N}^k$, we write

$$\Lambda^{\leq n} := \{\lambda \in \Lambda : d(\lambda) \leq n \text{ and } s(\lambda)\Lambda^{e_i} = \emptyset \text{ whenever } d(\lambda) + e_i \leq n\}.$$

When Λ has no sources, $\Lambda^{\leq n} = \Lambda^n$. For a locally convex k -graph Λ , $v\Lambda^{\leq n} \neq \emptyset$ for all $n \in \mathbb{N}^k$ and $v \in \Lambda^0$, but $v\Lambda^n \neq \emptyset$ for all n, v if and only if Λ has no sources.

2.2. BOUNDARY PATHS AND INFINITE PATHS. Fix $k > 0$ and $m \in (\mathbb{N} \cup \{\infty\})^k$. Then $\Omega_{k,m}$ denotes the category with objects $\text{Obj}(\Omega_{k,m}) = \{p \in \mathbb{N}^k : p \leq m\}$, morphisms $\text{Mor}(\Omega_{k,m}) = \{(p, q) : p, q \in \mathbb{N}^k, p \leq q \leq m\}$ and $\text{dom}(p, q) = q$, $\text{cod}(p, q) = p$, $\text{id}(p) = (p, p)$, $(p, q) \circ (q, t) = (p, t)$. The formula $d(p, q) = q - p$ defines a functor $d : \Omega_{k,m} \rightarrow \mathbb{N}^k$, and the pair $(\Omega_{k,m}, d)$ is a row-finite locally convex k -graph.

Let Λ be a row-finite locally convex k -graph. A degree-preserving functor $x : \Omega_{k,m} \rightarrow \Lambda$ is a **boundary path** of Λ if

$$(p \leq m \text{ and } p_i = m_i) \implies x(p)\Lambda^{e_i} = \emptyset \quad \text{for all } p \in \mathbb{N}^k, 1 \leq i \leq k.$$

We regard m as the degree of x and denote it $d(x)$, and we regard $x(0)$ as the range of x and denote it $r(x)$. If $m = (\infty, \dots, \infty)$ we call x an **infinite path**. We write $\Lambda^{\leq \infty}$ for the collection of all boundary paths and Λ^∞ for the collection of all infinite paths of Λ .

For $x \in \Lambda^{\leq \infty}$ and $n \in \mathbb{N}^k$ with $n \leq d(x)$, there is a boundary path $\sigma^n(x) : \Omega_{k,d(x)-n} \rightarrow \Lambda$ defined by $\sigma^n(x)(p, q) := x(p+n, q+n)$ for all $p \leq q \leq d(x) - n$. Given $x \in v\Lambda^{\leq \infty}$ and $\lambda \in \Lambda$ with $s(\lambda) = v$ there is a unique boundary path $\lambda x : \Omega_{k,m+d(\lambda)} \rightarrow \Lambda$ satisfying $(\lambda x)(0, d(\lambda)) = \lambda$ and $(\lambda x)(d(\lambda), d(\lambda) + p) = x(0, p)$ for all $p \leq d(x)$. For $x \in \Lambda^{\leq \infty}$, $\lambda \in \Lambda r(x)$ and $p \leq d(x)$, we have $x(0, p)\sigma^p(x) = x = \sigma^{d(\lambda)}(\lambda x)$. The set $\Lambda^{\leq \infty}$ has similar properties to the $\Lambda^{\leq n}$: if Λ has no sources, then $\Lambda^{\leq \infty} = \Lambda^\infty$; and for Λ locally convex, $v\Lambda^{\leq \infty}$ is non-empty for all $v \in \Lambda^0$, but $v\Lambda^\infty \neq \emptyset$ for all v if and only if Λ has no sources.

2.3. C^* -ALGEBRAS ASSOCIATED TO k -GRAPHS. Let Λ be a row-finite locally convex k -graph. Let A be a C^* -algebra, and let $\{t_\lambda : \lambda \in \Lambda\}$ be a collection of partial isometries in A . We call $\{t_\lambda : \lambda \in \Lambda\}$ a **Cuntz–Krieger Λ -family** if

- (i) $\{t_v : v \in \Lambda^0\}$ is a collection of mutually orthogonal projections;
- (ii) $t_\lambda t_\mu = t_{\lambda\mu}$ whenever $s(\lambda) = r(\mu)$;
- (iii) $t_\mu^* t_\mu = t_{s(\mu)}$ for all $\mu \in \Lambda$; and
- (iv) $t_v = \sum_{\lambda \in v\Lambda \leq n} t_\lambda t_\lambda^*$ for all $v \in \Lambda^0, n \in \mathbb{N}^k$.

There is a C^* -algebra $C^*(\Lambda)$ generated by a Cuntz–Krieger Λ -family $\{s_\lambda : \lambda \in \Lambda\}$ which is universal in the following sense: if $\{t_\lambda : \lambda \in \Lambda\}$ is another Cuntz–Krieger Λ -family in a C^* -algebra A , then there is a homomorphism $\pi_t : C^*(\Lambda) \rightarrow A$ satisfying $\pi_t(s_\lambda) = t_\lambda$ for all $\lambda \in \Lambda$.

2.4. REMOVING SOURCES FROM k -GRAPHS. The following construction is due to Farthing [5], and we refer the reader there for details and proofs. We have modified the formulation given in [5] slightly to streamline later proofs; lengthy calculations show that the resulting $\tilde{\Lambda}$ is isomorphic to Farthing’s $\bar{\Lambda}$.

We present Farthing’s construction only for row-finite locally convex k -graphs because that is the generality in which we will be working for the remainder of the paper. However, the construction makes sense for arbitrary k -graphs as long as the appropriate notion of a boundary path is used.

Fix a row-finite locally convex k -graph Λ . Define

$$V_\Lambda := \{(x; m) : x \in \Lambda^{\leq \infty}, m \in \mathbb{N}^k\}$$

and

$$P_\Lambda := \{(x; (m, n)) : x \in \Lambda^{\leq \infty}, m \leq n \in \mathbb{N}^k\}.$$

The relation \sim on V_Λ defined by $(x; m) \sim (y; n)$ if and only if

- (V1) $x(m \wedge d(x)) = y(n \wedge d(y))$; and
- (V2) $m - m \wedge d(x) = n - n \wedge d(y)$

is an equivalence relation. We denote the equivalence class of $(x; m)$ under \sim by $[x; m]$. The relation \approx on P_Λ defined by $(x; (m, n)) \approx (y; (p, q))$ if and only if

- (P1) $x(m \wedge d(x), n \wedge d(x)) = y(p \wedge d(y), q \wedge d(y))$;
- (P2) $m - m \wedge d(x) = p - p \wedge d(y)$; and
- (P3) $n - m = q - p$

is also an equivalence relation. We denote the equivalence class of $(x; (m, n))$ under \approx by $[x; (m, n)]$.

Theorem 3.24 of [5] implies that there is a row-finite k -graph $\tilde{\Lambda}$ with objects V_Λ / \sim , morphisms P_Λ / \approx , and structure-maps specified by the following formulae:

$$\begin{aligned} \tilde{r}([x; (m, n)]) &:= [x; m], \\ \tilde{s}([x; (m, n)]) &:= [x; n], \\ \tilde{\text{id}}([x; m]) &:= [x; (m, m)], \\ [x; (m, n)]\tilde{\sigma}[y; (p, q)] &:= [x(0, n \wedge d(x))\sigma^{p \wedge d(y)}(y); (m, n + q - p)], \quad \text{and} \\ \tilde{d}([x; (m, n)]) &:= n - m. \end{aligned}$$

Theorem 3.24 further states that $(\tilde{\Lambda}, \tilde{d})$ is a row-finite k -graph with no sources.

For $\lambda \in \Lambda$ and boundary paths $x, y \in s(\lambda)\Lambda^{\leq \infty}$, the elements $(\lambda x; (0, d(\lambda)))$ and $(\lambda y; (0, d(\lambda)))$ of P_Λ are equivalent under \sim . Hence there is a map $\iota : \Lambda \rightarrow \tilde{\Lambda}$ satisfying $\iota(\lambda) := [\lambda x; (0, d(\lambda))]$ for any $x \in s(\lambda)\Lambda^{\leq \infty}$. Indeed ι is an injective k -graph morphism, and hence an isomorphism of Λ onto $\iota(\Lambda) \subset \tilde{\Lambda}$.

Theorem 3.29 of [5] shows that $\sum_{v \in \Lambda^0} s_{\iota(v)}$ converges to a full projection P in the multiplier algebra of $C^*(\tilde{\Lambda})$, and that $PC^*(\tilde{\Lambda})P = C^*(\{s_{\iota(\lambda)} : \lambda \in \Lambda\}) \cong C^*(\Lambda)$. In particular, $C^*(\tilde{\Lambda})$ is Morita equivalent to $C^*(\Lambda)$.

In light of the preceding two paragraphs, we can — and do — regard Λ as a subset of $\tilde{\Lambda}$, dropping the inclusion map ι , and regard $C^*(\Lambda)$ as a C^* -subalgebra of $C^*(\tilde{\Lambda})$ also.

2.5. FROM BOUNDARY PATHS TO INFINITE PATHS AND BACK. Our aim in this paper is to combine Farthing’s results with the results of [14] to characterise simplicity of $C^*(\Lambda)$ in terms of the structure of $\Lambda^{\leq \infty}$. Applied directly, the results of [14] characterise the simplicity of $C^*(\Lambda)$ in terms of the structure of $\tilde{\Lambda}^\infty$. Hence we begin by establishing how elements of $\Lambda^{\leq \infty}$ correspond to elements of $\tilde{\Lambda}^\infty$.

For an element $x \in \Lambda^{\leq \infty}$ and $n \in \mathbb{N}^k$, define

$$[x; (n, \infty)] : \Omega_k \rightarrow \tilde{\Lambda} \text{ by } [x; (n, \infty)](p, q) := [x; (n + p, n + q)].$$

Our notation was chosen to suggest the relationship between $[x; (n, \infty)]$ and $[x; (n, p)]$, $p \geq n$; however, the reader should note that \approx is not defined on

boundary paths, and in particular $[x; (n, \infty)]$ is not itself an equivalence class under \approx .

Define $\pi : \tilde{\Lambda} \rightarrow \Lambda$ by $\pi([x; (m, n)]) = [x; (m \wedge d(x), n \wedge d(x))]$ for all $x \in \Lambda^{\leq \infty}$ and $m \leq n \in \mathbb{N}^k$. Note that $\pi([x; (m, n)]) \in \iota(\Lambda) \subseteq \tilde{\Lambda}$, so we regard π as a k -graph morphism from $\tilde{\Lambda}$ to Λ , and identify $\pi([x; (m, n)])$ with $x(m \wedge d(x), n \wedge d(x))$. That π is well-defined follows immediately from (P1). It is straightforward to check that π is a functor, is surjective onto Λ and is a projection in the sense that $\pi \circ \pi = \pi$.

LEMMA 2.1: *Let Λ be a locally convex row-finite k -graph. For $x \in \Lambda^{\leq \infty}$ and $m \leq n \in \mathbb{N}^k$ we have*

$$[x; (m, n)] = [\sigma^{m \wedge d(x)}(x); (m - m \wedge d(x), n - m \wedge d(x))].$$

Also $(m - m \wedge d(x)) \wedge d(\sigma^{m \wedge d(x)}(x)) = 0$. In particular, for each $\lambda \in \tilde{\Lambda}$ there exist $y \in \Lambda^{\leq \infty}$ and $p \in \mathbb{N}^k$ such that $\lambda = [y; (p, p + d(\lambda))]$ and $p \wedge d(y) = 0$.

Proof. We begin by verifying $(m - m \wedge d(x)) \wedge d(\sigma^{m \wedge d(x)}(x)) = 0$. For each $i \in \{1, \dots, k\}$, either $(m \wedge d(x))_i = m_i$ or $(m \wedge d(x))_i = d(x)_i$. Hence, either $(m - m \wedge d(x))_i = 0$ or $(d(x) - m \wedge d(x))_i = 0$ or both. Since $d(\sigma^{m \wedge d(x)}(x)) = d(x) - m \wedge d(x)$, we therefore obtain

$$(2.3) \quad (m - m \wedge d(x)) \wedge d(\sigma^{m \wedge d(x)}(x)) = 0$$

as required.

To see that $[x; (m, n)] = [\sigma^{m \wedge d(x)}(x); (m - m \wedge d(x), n - m \wedge d(x))]$, we must check that $(x; (m, n)) \approx (\sigma^{m \wedge d(x)}(x); (m - m \wedge d(x), n - m \wedge d(x)))$.

For (P1), we note that

$$(2.4) \quad x(m \wedge d(x), n \wedge d(x)) = \sigma^{m \wedge d(x)}(x)(0, n \wedge d(x) - m \wedge d(x))$$

By (2.3) we may replace the 0 in the right hand side of (2.4) with

$$(m - m \wedge d(x)) \wedge d(\sigma^{m \wedge d(x)}(x)),$$

giving

$$(2.5) \quad \begin{aligned} x(m \wedge d(x), n \wedge d(x)) &= \sigma^{m \wedge d(x)}(x)((m - m \wedge d(x)) \wedge d(\sigma^{m \wedge d(x)}(x)), n \wedge d(x) - m \wedge d(x)) \end{aligned}$$

Taking $a = n, b = d(x)$ and $c = m \wedge d(x)$ in (2.2) we have

$$\begin{aligned} n \wedge d(x) - m \wedge d(x) &= (n - m \wedge d(x)) \wedge (d(x) - m \wedge d(x)) \\ &= (n - m \wedge d(x)) \wedge d(\sigma^{m \wedge d(x)}(x)). \end{aligned}$$

Hence (2.5) implies that

$$\begin{aligned} x(m \wedge d(x), n \wedge d(x)) &= \sigma^{m \wedge d(x)}(x)((m - m \wedge d(x)) \wedge d(\sigma^{m \wedge d(x)}(x)), \\ &\quad (n - m \wedge d(x)) \wedge d(\sigma^{m \wedge d(x)}(x))) \end{aligned}$$

establishing (P1). For (P2), we calculate:

$$\begin{aligned} m - m \wedge d(x) &= (m - m \wedge d(x)) - 0 \\ &= (m - m \wedge d(x)) - (m - m \wedge d(x)) \wedge d(\sigma^{m \wedge d(x)}(x)). \end{aligned}$$

For (P3), just note that $n - m = (n - m \wedge d(x)) - (m - m \wedge d(x))$.

For the last statement of the lemma, write $\lambda = [x; (m, n)]$ for some $x \in \Lambda^{\leq \infty}$ and $m, n \in \mathbb{N}^k$, and take $y = \sigma^{m \wedge d(x)}(x)$ and $p = m - m \wedge d(x)$. ■

LEMMA 2.2: *Let Λ be a row-finite locally convex k -graph and let $v \in \Lambda^0$. Suppose $x \in v\Lambda^{\leq \infty}$ and $p \in \mathbb{N}^k$ satisfy $p \wedge d(x) = 0$. Then for any other $z \in v\Lambda^{\leq \infty}$ we have $p \wedge d(z) = 0$ and $[x; (0, p)] = [z; (0, p)]$.*

Proof. Fix $z \in v\Lambda^{\leq \infty}$. We must show that $p_i \neq 0$ implies $d(z)_i = 0$. Suppose $i \in \{1, \dots, k\}$ satisfies $p_i \neq 0$. Then $p \wedge d(x)_i = 0$ implies that $d(x)_i = 0$. Since x is a boundary path it follows that $v\Lambda^{e_i} = \emptyset$, so $d(z)_i = 0$ also. One now easily verifies (P1)–(P3) directly to see that $[x; (0, p)] = [z; (0, p)]$. ■

Before stating the next proposition, we need some notation. Fix $\mu, \nu \in \Lambda$. We say that $\lambda \in \Lambda$ is a **common extension** of μ and ν if $\lambda = \mu\mu' = \nu\nu'$ for some $\mu', \nu' \in \Lambda$. If λ is a common extension of μ and ν , then in particular, $d(\lambda) \geq d(\mu) \vee d(\nu)$. We say that a common extension λ of μ and ν is a **minimal common extension** if $d(\lambda) = d(\mu) \vee d(\nu)$. We write $\text{MCE}(\mu, \nu)$ for the collection of all minimal common extensions of μ and ν .

Now fix $v \in \Lambda^0$ and let $E \subset v\Lambda$. We say that E is **exhaustive** if for each $\lambda \in v\Lambda$ there exists $\mu \in E$ such that $\text{MCE}(\lambda, \mu) \neq \emptyset$. If E is also finite, we call E **finite exhaustive**.

PROPOSITION 2.3: *Let (Λ, d) be a row-finite, locally convex k -graph. Then for each infinite path $y \in \tilde{\Lambda}^\infty$, there exist a unique $p_y \in \mathbb{N}^k$ and $\pi(y) \in \Lambda^{\leq \infty}$ such*

that $p_y \wedge d(\pi(y)) = 0$ and $y = [\pi(y); (p_y, \infty)]$. We then have $\pi(y(m, n)) = \pi(y)(m \wedge d(\pi(y)), n \wedge d(\pi(y)))$ for all $m, n \in \mathbb{N}^k$.

Proof. We first argue existence. Fix $y \in \tilde{\Lambda}^\infty$. For each $n \in \mathbb{N}^k$ there exists an $x_n \in \Lambda^{\leq \infty}$ and $p(n) \in \mathbb{N}^k$ such that $[x_n; (p(n), p(n) + n)] = y(0, n)$. By Lemma 2.1 we may assume that $p(n) \wedge d(x_n) = 0$ for all $n \in \mathbb{N}^k$. This forces $p(n) = p(0)$ and $r(x_n) = r(x_0)$ for all $n \in \mathbb{N}^k$. We will henceforth just write p for $p(0)$. For $a, b \in \mathbb{N}^k$ with $a \leq b$, we have

$$[x_a; (p, p + a)] = y(0, a) = [x_b; (p, p + a)].$$

Since $p \wedge d(x_a) = 0 = p \wedge d(x_b)$, we therefore have

$$(2.6) \quad x_a(0, a \wedge d(x_a)) = x_b(0, a \wedge d(x_a)) \quad \text{for } a \leq b \in \mathbb{N}^k.$$

Define $m \in (\mathbb{N} \cup \{\infty\})^k$ by

$$m := \bigvee_{n \in \mathbb{N}^k} n \wedge d(x_n).$$

Fix $q \in \mathbb{N}^k$ with $q \leq m$. Suppose that $a, b \in \mathbb{N}^k$ satisfy $a \wedge d(x_a), b \wedge d(x_b) \geq n$. Since $q \leq a \wedge d(x_a)$, we may use (2.6) to calculate:

$$x_a(0, q) = (x_a(0, a \wedge d(x_a)))(0, q) = (x_{a \vee b}(0, a \wedge d(x_a)))(0, q) = x_{a \vee b}(0, q).$$

Similarly, $x_b(0, q) = x_{a \vee b}(0, q)$, and in particular, $x_a(0, q) = x_b(0, q)$. Hence there is a unique graph morphism $x : \Omega_{k,m} \rightarrow \Lambda$ such that for $q, q' \in \mathbb{N}^k$ with $q \leq q' \leq m$,

$$x(q, q') := x_a(q, q') \text{ for any } a \in \mathbb{N}^k \text{ such that } a \wedge d(x_a) \geq q'.$$

We claim that $x \in \Lambda^{\leq \infty}$. Suppose $q \in \mathbb{N}^k$ and $1 \leq i \leq k$ satisfy $q \leq m$ and $q_i = m_i$. We must show that $x(q)\Lambda^{e_i} = \emptyset$. We first claim that $q \wedge d(x_q) = q$. To see this, we argue by contradiction. Suppose that $d(x_q)_j < q_j$ for some j . Since x_q is a boundary path, $x_q(q \wedge d(x_q))\Lambda^{e_j} = \emptyset$. By (2.6), $x_{q'}(q \wedge d(x_q)) = x_q(q \wedge d(x_q))$ for all $q' \geq q$, and this forces $d(x_{q'})_j = d(x_q)_j < q_j$ for all $q' \geq q$. In particular, since $q \leq m$, this contradicts the definition of m . This proves the claim. Now, by definition of m , we also have $(q + e_i) \wedge d(x_{q+e_i}) = q_i$, and hence $x_{q+e_i}(q)\Lambda^{e_i} = \emptyset$. By (2.6), we have $x_{q+e_i}(q) = x_q(q) = x(q)$, so $x(q)\Lambda^{e_i} = \emptyset$ as claimed.

Now we aim to show that $y = [x; (p, \infty)]$. To do this we show that

$$(x; (p, p + n)) \approx (x_n; (p, p + n))$$

for all $n \in \mathbb{N}^k$. Fix $n \in \mathbb{N}^k$.

Condition (P1) follows directly from the definition of x .

For (P2), note that $p \wedge d(x_n) = 0$ for all n implies $p \wedge d(x) = 0$. So

$$p - p \wedge d(x) = p = p - p \wedge d(x_n).$$

Condition (P3) is immediate. Hence $y = [x; (p, \infty)]$. Since $p \wedge d(x) = 0$ as observed above, taking $p_y := p$ and $\pi(y) := x$ establishes existence.

For uniqueness, suppose that x' and p' have the same properties. Then $[x'; (p', p' + m)] = y(0) = [\pi(y); (p_y, p_y + m)]$ for all $m \in \mathbb{N}^k$. Since $p' \wedge d(x') = 0 = p_y \wedge d(\pi(y))$, condition (P1) forces $x'(0, m \wedge d(x')) = \pi(y)(0, m \wedge d(\pi(y)))$ for all $m \in \mathbb{N}^k$. In particular, $m \wedge d(x') = m \wedge d(\pi(y))$ for all $m \in \mathbb{N}^k$, so $d(x') = d(\pi(y))$, and since x' and $\pi(y)$ agree on all initial segments, they must be equal. Condition (P2) together with $p' \wedge d(x') = p_y \wedge d(\pi(y)) = 0$ forces $p' = p$. Thus p_y and $\pi(y)$ are unique.

For the final statement fix $m, n \in \mathbb{N}^k$. By the preceding paragraphs $y(m, n) = [\pi(y); (p_y + m, p_y + n)]$, so

$$\pi(y(m, n)) = \pi(y)((p_y + m) \wedge d(\pi(y)), (p_y + n) \wedge d(\pi(y))).$$

Since $p_y \wedge d(\pi(y)) = 0$, we have $(p_y + m) \wedge d(\pi(y)) = m \wedge d(\pi(y))$ and similarly for n , and this completes the proof. ■

3. Simplicity

Definition 3.1: Let (Λ, d) be a row-finite locally convex k -graph. We say Λ is **cofinal** if, for each $x \in \Lambda^{\leq \infty}$ and $v \in \Lambda^0$, there exists $n \in \mathbb{N}^k$ such that $n \leq d(x)$ and $v\Lambda x(n)$ is nonempty.

Definition 3.2: Let (Λ, d) be a row-finite k -graph, fix $v \in \Lambda^0$, and fix $m \neq n \in \mathbb{N}^k$. We say Λ has **local periodicity** m, n **at** v if for every $x \in v\Lambda^{\leq \infty}$, we have $m - m \wedge d(x) = n - n \wedge d(x)$ and $\sigma^{m \wedge d(x)}(x) = \sigma^{n \wedge d(x)}(x)$. We say Λ has **no local periodicity** if Λ does not have local periodicity m, n at $v \in \Lambda^0$ for any $m \neq n \in \mathbb{N}^k$ and $v \in \Lambda^0$.

Remark 3.3: Note that if Λ has no sources, then every boundary path is an infinite path. In particular, $m \wedge d(x) = m$ and $n \wedge d(x) = n$, for all $m, n \in \mathbb{N}^k$ and all $x \in \Lambda^{\leq \infty}$. Hence the definitions of cofinality, local periodicity and no local periodicity presented above reduce to the definitions of the same conditions given in [14] when Λ has no sources.

THEOREM 3.4: *Let (Λ, d) be a row-finite locally convex k -graph with no sources. Then $C^*(\Lambda)$ is simple if and only if both of the following conditions hold:*

- (1) Λ is cofinal; and
- (2) Λ has no local periodicity.

We prove Theorem 3.4 at the end of the section. To do so, we first establish two key propositions.

PROPOSITION 3.5: *Let (Λ, d) be a row-finite, locally convex k -graph. Then Λ is cofinal if and only if $\tilde{\Lambda}$ is cofinal.*

Proof. (\implies) Suppose Λ is cofinal. Fix $z \in \tilde{\Lambda}^\infty$ and $[y; p] \in \tilde{\Lambda}^0$. Proposition 2.3 implies $z = [x; (m, \infty)]$ for some $x \in \Lambda^{\leq \infty}$ and $m \in \mathbb{N}^k$. We must show that there exists $p' \in \mathbb{N}^k$ such that $[y; p]\tilde{\Lambda}[x; (m, \infty)](p') \neq \emptyset$.

Fix $q \geq p$ such that $y(q \wedge d(y))\Lambda^{e_i} = \emptyset$ whenever $d(y)_i < \infty$. Since Λ is cofinal,

$$y(q \wedge d(y))\Lambda x(m \wedge d(x) + n) \neq \emptyset \quad \text{for some } n \leq d(x) - m \wedge d(x).$$

Fix a path $\lambda \in y(q \wedge d(y))\Lambda x(m \wedge d(x) + n)$. Lemma 2.10 of [11] implies that $y' := y(0, q \wedge d(y)) \lambda \sigma^{m \wedge d(x) + n}(x) \in \Lambda^{\leq \infty}$. We claim that $q \wedge d(y) = q \wedge d(y')$. We have

$$\begin{aligned} q \wedge d(y') &= q \wedge d(y(0, q \wedge d(y)) \lambda \sigma^{m \wedge d(x) + n}(x)) \\ &= q \wedge (q \wedge d(y) + d(\lambda) + (d(x) - m \wedge d(x) - n)). \end{aligned}$$

For $i \in \{1, \dots, k\}$ such that $d(y)_i < \infty$, we have $y(q \wedge d(y))\Lambda^{e_i} = \emptyset$ by choice of q , so $d(\lambda x(m \wedge d(x) + n, d(x)))_i = 0$ and

$$(q \wedge (q \wedge d(y) + d(\lambda) + (d(x) - m \wedge d(x) - n)))_i = (q \wedge (q \wedge d(y)))_i = (q \wedge d(y))_i.$$

For $i \in \{1, \dots, k\}$ such that $d(y)_i = \infty$, we have

$$(q \wedge (q \wedge d(y) + d(\lambda) + (d(x) - m \wedge d(x) - n)))_i = q_i = (q \wedge d(y))_i.$$

Hence $q \wedge d(y') = q \wedge d(y)$ and so $[y'; q] = [y; q]$. Now, consider

$$\mu := [y; (p, q)][y'; (q, q + q \wedge d(y) + d(\lambda) + m)].$$

We have $r(\mu) = [y; p]$ and

$$\begin{aligned} s(\mu) &= [y'; q + q \wedge d(y) + d(\lambda) + m] \\ &= [\sigma^{(q \wedge d(y) + d(\lambda))}(y'); q + m] \\ &= [\sigma^{m \wedge d(x) + n}(x); q + m] \\ &= [x; m \wedge d(x) + n + q + m] \\ &= [x; (m, \infty)](m \wedge d(x) + n + q). \end{aligned}$$

Hence $\mu \in [y; p]\tilde{\Lambda}[x; (m, \infty)](m \wedge d(x) + n + q)$. As $[y; p] \in \tilde{\Lambda}^0$ and $z = [x; (m, \infty)] \in \tilde{\Lambda}^\infty$ were arbitrary, it follows that $\tilde{\Lambda}$ is cofinal.

(\Leftarrow) Suppose $\tilde{\Lambda}$ is cofinal. Fix $v \in \Lambda^0$ and $x \in \Lambda^{\leq \infty}$. Since $\tilde{\Lambda}$ is cofinal we may fix $n \in \mathbb{N}^k$ such that

$$v\tilde{\Lambda}[x; (0, \infty)](n) \neq \emptyset;$$

say $\lambda \in v\tilde{\Lambda}[x; (0, \infty)](n)$. Then

$$r(\pi(\lambda)) = \pi(r(\lambda)) = \pi(v) = v$$

and

$$s(\pi(\lambda)) = \pi(s(\lambda)) = \pi([x; n]) = [x; n \wedge d(x)].$$

Hence $\pi(\lambda) \in v\Lambda[x; n \wedge d(x)]$. Once again, since $v \in \Lambda^0$ and $x \in \Lambda^{\leq \infty}$ were arbitrary, it follows that Λ is cofinal. ■

PROPOSITION 3.6: *Let (Λ, d) be a row-finite locally convex k -graph. Fix $v \in \tilde{\Lambda}^0$ and $m \neq n \in \mathbb{N}^k$. Then $\tilde{\Lambda}$ has local periodicity m, n at v if and only if Λ has local periodicity m, n at $\pi(v)$. In particular, $\tilde{\Lambda}$ has no local periodicity if and only if Λ has no local periodicity.*

In order to prove this proposition, we require some preliminary results.

LEMMA 3.7: *Let (Λ, d) be a row-finite locally convex k -graph. Then for any $[x; (n, \infty)] \in \tilde{\Lambda}^\infty$ and $m \in \mathbb{N}^k$, we have $\sigma^m([x; (n, \infty)]) = [x; (n + m, \infty)]$. If $m \leq d(x)$, we also have $\sigma^m([x; (n, \infty)]) = [\sigma^m(x); (n, \infty)]$.*

Proof. For any $p \in \mathbb{N}^k$, we have

$$\begin{aligned} \sigma^m([x; (n, \infty)])(0, p) &= [x; (n, \infty)](m, m + p) = [x; (n + m, n + m + p)] \\ &= [x; (n + m, \infty)](0, p). \end{aligned}$$

Since $\sigma^m([x; (n, \infty)])$ and $[x; (n + m, \infty)]$ agree on every initial segment we conclude that they are equal. Now, fix $m \leq d(x)$. We must show that $\sigma^m([x; (n, \infty)])(0, p) = [\sigma^m(x); (n, \infty)](0, p)$ for all $p \in \mathbb{N}^k$, that is we must check that $(x; (n + m, n + m + p)) \approx (\sigma^m(x); (n, n + p))$. For (P1), fix $p \in \mathbb{N}^k$ and calculate

$$\begin{aligned} \sigma^m(x)(n \wedge d(\sigma^m(x)), (n + p) \wedge d(\sigma^m(x))) & \\ &= x(n \wedge d(\sigma^m(x)) + m, (n + p) \wedge d(\sigma^m(x)) + m) \\ &= x((n + m) \wedge (d(\sigma^m(x)) + m), (n + m + p) \wedge (d(\sigma^m(x)) + m)) \quad \text{by 2.1} \\ &= x((n + m) \wedge d(x), (n + m + p) \wedge d(x)). \end{aligned}$$

For (P2), we have

$$\begin{aligned} n - n \wedge d(\sigma^m(x)) &= (n + m) - (n \wedge d(\sigma^m(x)) + m) \\ &= (n + m) - (n + m) \wedge (d(\sigma^m(x)) + m) \quad \text{by 2.1} \\ &= (n + m) - (n + m) \wedge d(x). \end{aligned}$$

For (P3), we have $(n + m + p) - (n + m) = p = (n + p) - n$. ■

LEMMA 3.8: *Let Λ be a row-finite locally convex k -graph. Suppose that $x \in \Lambda^{\leq \infty}$ and $p, m, n \in \mathbb{N}^k$ satisfy $\sigma^m([x; (p, \infty)]) = \sigma^n([x; (p, \infty)])$. Then $d(x)_i = \infty$ whenever $m_i \neq n_i$.*

Proof. We prove the contrapositive statement. Suppose $d(x)_i < \infty$ and fix $l \in \mathbb{N}^k$ such that $(p + m + l)_i > d(x)_i$ and $(p + n + l)_i > d(x)_i$. Then

$$\begin{aligned} \sigma^m([x; (p, \infty)]) &= \sigma^n([x; (p, \infty)]) \\ \implies [x; (p + m, \infty)] &= [x; (p + n, \infty)] \quad \text{by Lemma 3.7} \\ \implies [x; (p + m, \infty)](0, l) &= [x; (p + n, \infty)](0, l) \\ \implies [x; (p + m, p + m + l)] &= [x; (p + n, p + n + l)] \\ &\quad \text{by definition of } [x; (p, \infty)] \\ \implies ((p + m + l) - (p + m + l) \wedge d(x))_i &= ((p + n + l) - (p + n + l) \wedge d(x))_i \\ &\quad \text{by (V2)} \\ \implies ((p + m + l) - d(x))_i &= ((p + n + l) - d(x))_i \\ &\quad \text{since } (p + m + l), (p + n + l) > d(x)_i \\ \implies m_i &= n_i. \end{aligned}$$

So $d(x)_i < \infty$ implies $m_i = n_i$ as claimed. ■

LEMMA 3.9: Let Λ be a row-finite locally convex k -graph. Fix $y \in \tilde{\Lambda}^\infty$ and $m, n \in \mathbb{N}^k$. Then $\sigma^m(y) = \sigma^n(y)$ if and only if: (a) $\sigma^{m \wedge d(\pi(y))}(\pi(y)) = \sigma^{n \wedge d(\pi(y))}(\pi(y))$; and (b) $m - m \wedge d(\pi(y)) = n - n \wedge d(\pi(y))$.

Proof. First suppose that (a) and (b) hold. By Proposition 2.3,

$$y = [\pi(y); (p_y, \infty)]$$

with $p_y \wedge d(\pi(y)) = 0$. Hence

$$\sigma^{m \wedge d(\pi(y))}(y) = \sigma^{m \wedge d(\pi(y))}([\pi(y); (p_y, \infty)]) = [\sigma^{m \wedge d(\pi(y))}(\pi(y)); (p_y, \infty)]$$

by Lemma 3.7. Similarly, $\sigma^{n \wedge d(\pi(y))}(y) = [\sigma^{n \wedge d(\pi(y))}(\pi(y)); (p_y, \infty)]$, so (a) implies that

$$(3.1) \quad \sigma^{m \wedge d(\pi(y))}(y) = \sigma^{n \wedge d(\pi(y))}(y).$$

Using (b) and (3.1), we now calculate:

$$\sigma^m(y) = \sigma^{m - m \wedge d(\pi(y))}(\sigma^{m \wedge d(\pi(y))}(y)) = \sigma^{n - n \wedge d(\pi(y))}(\sigma^{n \wedge d(\pi(y))}(y)) = \sigma^n(y).$$

Now suppose that $\sigma^m(y) = \sigma^n(y)$. Using Lemma 3.7, we see that

$$\sigma^m(y)(0) = \sigma^m([\pi(y); (p_y, \infty)])(0) = [\pi(y); p_y + m]$$

and similarly, $\sigma^n(y)(0) = [\pi(y); p_y + n]$. In particular, condition (V2) implies that

$$(p_y + m) - ((p_y + m) \wedge d(\pi(y))) = (p_y + n) - ((p_y + n) \wedge d(\pi(y))).$$

Since $p_y \wedge d(\pi(y)) = 0$, this establishes (b). Note that by Lemma 3.8, $d(\pi(y))_i = \infty$ whenever $m_i \neq n_i$, and hence $d(\pi(y)) - m \wedge d(\pi(y)) = d(\pi(y)) - n \wedge d(\pi(y))$. Thus $p \leq d(\pi(y)) - m \wedge d(\pi(y))$ if and only if $p \leq d(\pi(y)) - n \wedge d(\pi(y))$, and for such p we may use the final statement of Proposition 2.3 to calculate

$$\sigma^{m \wedge d(\pi(y))}(\pi(y))(0, p) = \pi(y)(m \wedge d(\pi(y)), m \wedge d(\pi(y)) + p) = \pi(y)(m, m + p).$$

Similarly, $\sigma^{n \wedge d(\pi(y))}(\pi(y))(0, p) = \pi(y)(n, n + p)$. We have $\sigma^m(y) = \sigma^n(y)$ by hypothesis, so $y(m, m + p) = y(n, n + p)$, and hence $\pi(y(m, m + p)) = \pi(y(n, n + p))$. Since $p \leq d(\pi(y)) - m \wedge d(\pi(y)) = d(\pi(y)) - n \wedge d(\pi(y))$ was arbitrary, this completes the proof. ■

Proof of Proposition 3.6. Lemma 2.2 implies that for any $z \in w\tilde{\Lambda}^\infty$, we have $p_z \wedge d(x) = 0$ and $[x; p_z] = w$ for all $x \in \pi(w)\Lambda^{\leq \infty}$. It follows that for each $x \in \pi(w)\Lambda^{\leq \infty}$ we have $[x; (p_z, \infty)]$ in $w\tilde{\Lambda}^\infty$, and $x = \pi([x; (p_z, \infty)])$. That is, $\pi(w)\Lambda^{\leq \infty} = \{\pi(z) : z \in w\tilde{\Lambda}^\infty\}$. The result now follows from Lemma 3.9. ■

Proof of Theorem 3.4. Propositions 3.5 and 3.6 imply that Λ is cofinal and has no local periodicity if and only if $\tilde{\Lambda}$ is cofinal and has no local periodicity. Since $\tilde{\Lambda}$ is row-finite and has no sources by [5, Theorem 3.23], Theorem 3.3 of [14] implies that $C^*(\tilde{\Lambda})$ is simple if and only if Λ is cofinal and has no local periodicity. Theorem 3.29 of [5] implies that $C^*(\Lambda)$ is a full corner of $C^*(\tilde{\Lambda})$, so by [13, Theorem 3.19] $C^*(\Lambda)$ and $C^*(\tilde{\Lambda})$ are Morita equivalent. In particular, $C^*(\Lambda)$ is simple if and only if $C^*(\tilde{\Lambda})$ is simple. Bringing these implications together,

$$\begin{aligned}
 C^*(\Lambda) \text{ is simple} &\iff C^*(\tilde{\Lambda}) \text{ is simple} \\
 &\iff \tilde{\Lambda} \text{ is cofinal and has no local periodicity} \\
 &\iff \Lambda \text{ is cofinal and has no local periodicity}
 \end{aligned}$$

as required. ■

Before concluding this section, we pause to discuss briefly the local periodicity condition presented in Definition 3.2. This definition is not, perhaps, the obvious extrapolation of the condition given in [14] to the locally convex setting (though certainly Proposition 3.6 indicates that it is the right one). The more obvious definition would be to say that Λ has local periodicity p, q at w if

$$(3.2) \quad \text{for every } x \in w\Lambda^{\leq \infty} \text{ we have } p, q \leq d(x) \text{ and } \sigma^p(x) = \sigma^q(x).$$

The two are not equivalent: if Λ is the 2-graph whose skeleton appears on the left of Figure 1, then one can check that the skeleton of $\tilde{\Lambda}$ is that which appears on the right of Figure 1. One can also check that $\tilde{\Lambda}$ has local periodicity $(1, 2), (0, 2)$ at $v = \pi(v)$, but there do not exist $p, q \in \mathbb{N}^2$ satisfying (3.2) with $w = v$. That having been said, the key notion for the purposes of characterising simplicity is that of no local periodicity, and the following lemma shows that Definition 3.2 and equation (3.2) correspond to the same notion of no local periodicity.




Figure 1. Local periodicity p, q at w is not equivalent to (3.2)

LEMMA 3.10: *Let Λ be a row-finite locally convex k -graph. Then there exist $v \in \Lambda^0$ and $m \neq n \in \mathbb{N}^k$ such that Λ has local periodicity at v if and only if there exist $w \in \Lambda^0$ and $p, q \in \mathbb{N}^k$ satisfying (3.2).*

Proof. Suppose first that w, p, q satisfy (3.2). Then, in particular, for $x \in w\Lambda^{\leq \infty}$, we have $p - p \wedge d(x) = 0 = q - q \wedge d(x)$, and

$$\sigma^{p \wedge d(x)}(x) = \sigma^p(x) = \sigma^q(x) = \sigma^{q \wedge d(x)}(x).$$

Hence for $m = p, n = q$ and $v = w$, Λ has local periodicity p, q at w .

Now suppose that Λ has local periodicity m, n at v . Then Proposition 3.6 implies that $\tilde{\Lambda}$ has local periodicity m, n at v . Let $p := m - m \wedge n$ and $q := n - m \wedge n$, fix $\lambda \in v\tilde{\Lambda}^{m \wedge n}$, and let $u := s(\lambda)$. For $x \in u\tilde{\Lambda}^\infty$,

$$\sigma^p(x) = \sigma^p(\sigma^{m \wedge n}(\lambda x)) = \sigma^m(\lambda x),$$

and likewise $\sigma^q(x) = \sigma^n(\lambda x)$. Since $\lambda x \in v\tilde{\Lambda}^\infty$, it follows that $\sigma^p(x) = \sigma^q(x)$. So $\tilde{\Lambda}$ has local periodicity p, q at u , and another application of Proposition 3.6 then shows that Λ has local periodicity p, q at $\pi(u)$. Since $p \wedge q = 0$, Lemma 3.8 implies that $p, q \leq d(x)$ for all $x \in \pi(u)\Lambda^{\leq \infty}$. Hence $w := \pi(u)$, $p = m - m \wedge n$ and $q = n - m \wedge n$ satisfy (3.2). ■

4. Ideal structure

In this section, we describe the relationship between cofinality and local periodicity of Λ and the ideal structure of $C^*(\Lambda)$. In order to state the results we need some background. For details of the following, see [6].

For $z \in \mathbb{T}^k$ and $n \in \mathbb{N}^k$, we use the multi-index notation $z^n := \prod_{i=1}^k z_i^{n_i} \in \mathbb{T}$. There is a strongly continuous action γ of \mathbb{T}^k on $C^*(\Lambda)$ satisfying $\gamma_z(s_\lambda) = z^{d(\lambda)} s_\lambda$ for all $\lambda \in \Lambda$. The fixed point algebra $C^*(\Lambda)^\gamma$ is called the **core** of $C^*(\Lambda)$ and is equal to $\overline{\text{span}}\{s_\mu s_\nu^* : d(\mu) = d(\nu)\}$.

The following lemma will be useful in proving both of our main results in this section. In the statement of the lemma and what follows, $\mathcal{M}(A)$ denotes the multiplier algebra of a C^* -algebra A .

LEMMA 4.1: *Let Λ be a row-finite locally convex k -graph. Let \tilde{I} be an ideal of $C^*(\tilde{\Lambda})$ and let $P := \sum_{v \in \Lambda^0} s_v \in \mathcal{M}(C^*(\tilde{\Lambda}))$. Then*

- (1) *for $v \in \tilde{\Lambda}^0$, we have $s_v \in \tilde{I}$ if and only if $s_{\pi(v)} \in P\tilde{I}P$; and*
- (2) *$P\tilde{I}P \cap C^*(\Lambda)^\gamma \neq \{0\}$ if and only if $\tilde{I} \cap C^*(\tilde{\Lambda})^\gamma \neq \{0\}$.*

Proof. For (1), fix $v \in \tilde{\Lambda}^0$, and use Lemma 2.1 to write $v = [x; p]$ where $x \in \pi(v)\Lambda^{\leq \infty}$ and $p \wedge d(x) = 0$. Let $\lambda := [x; (0, p)] \in \pi(v)\tilde{\Lambda}v$. Lemma 2.2 implies that $\pi(v)\tilde{\Lambda}^p = \{\lambda\}$. Hence the Cuntz–Krieger relations show that

$$s_v = s_\lambda^* s_{\pi(v)} s_\lambda \quad \text{and} \quad s_{\pi(v)} = s_\lambda s_v s_\lambda^*.$$

Since $Ps_{\pi(v)}P = s_{\pi(v)}$, this proves (a).

For (2), the “only if” implication is trivial, so it suffices to establish the “if” direction. Suppose that $\tilde{I} \cap C^*(\tilde{\Lambda})^\gamma \neq \{0\}$. The argument of (ii) \implies (iii) in [14, Proposition 3.4] shows that there exists $w \in \tilde{\Lambda}^0$ such that $s_w \in \tilde{I}$. Hence (a) implies that there exists $v \in \Lambda^0$ such that $s_v \in P\tilde{I}P$, and since the vertex projections are fixed by γ , we then have $s_v \in P\tilde{I}P \cap C^*(\Lambda)^\gamma$. ■

PROPOSITION 4.2: *Let Λ be a row-finite locally convex k -graph. The following are equivalent.*

- (1) *Λ is cofinal;*
- (2) *if I is an ideal of $C^*(\Lambda)$ and $s_v \in I$ for some $v \in \Lambda^0$, then $I = C^*(\Lambda)$;*
- (3) *if I is an ideal of $C^*(\Lambda)$ and $I \cap C^*(\Lambda)^\gamma \neq \{0\}$, then $I = C^*(\Lambda)$.*

Proof. That these three statements are equivalent for $\tilde{\Lambda}$ follows from [14, Proposition 3.5] because $\tilde{\Lambda}$ has no sources. Proposition 3.5 shows that $\tilde{\Lambda}$ is cofinal if and only if Λ is cofinal. Recall that $\sum_{v \in \Lambda^0} s_v$ converges to a full projection $P \in \mathcal{M}C^*(\tilde{\Lambda})$ and that $PC^*(\tilde{\Lambda})P$ is canonically isomorphic to $C^*(\Lambda)$. Hence the map $\tilde{I} \mapsto P\tilde{I}P$ is a bijection between ideals of $C^*(\tilde{\Lambda})$ and ideals of Λ . The proposition then follows from Lemma 4.1. ■

In order to state the next result we need more background. As in [11, Theorem 3.15] let $\mathcal{H} := \ell^2(\Lambda^{\leq\infty})$ with standard basis denoted $\{u_x : x \in \Lambda^{\leq\infty}\}$. For each $\lambda \in \Lambda$ define $S_\lambda \in \mathcal{B}(\mathcal{H})$ by

$$S_\lambda u_x = \begin{cases} u_{\lambda x} & \text{if } s(\lambda) = r(x) \\ 0 & \text{otherwise.} \end{cases}$$

Then $\{S_\lambda : \lambda \in \Lambda\} \subset \mathcal{H}$ is a Cuntz–Krieger Λ -family. By the universal property of $C^*(\Lambda)$ there exists a homomorphism $\pi_S : C^*(\Lambda) \rightarrow \mathcal{B}(\mathcal{H})$ such that $\pi_S(s_\lambda) = S_\lambda$ for all $\lambda \in \Lambda$. We call π_S the **boundary-path representation**.

PROPOSITION 4.3: *Let Λ be a row-finite locally convex k -graph. The following are equivalent:*

- (1) Λ has no local periodicity;
- (2) every non-zero ideal of $C^*(\Lambda)$ contains a vertex projection;
- (3) the boundary-path representation π_S is faithful.

In order to prove this proposition we need a technical lemma.

LEMMA 4.4: *Let Λ be a row-finite locally convex k -graph and suppose that Λ has local periodicity m, n at v . Fix $x \in v\Lambda^{\leq\infty}$, let $\mu := x(0, m \wedge d(x))$, let $\alpha := x(m \wedge d(x), (m \vee n) \wedge d(x))$, and let $\nu := x(0, n \wedge d(x))$. Then $d(\mu) \neq d(\nu)$, and $\mu\alpha y = \nu\alpha y$ for all $y \in s(\alpha)\Lambda^{\leq\infty}$.*

Proof. Since $x \in v\Lambda^{\leq\infty}$ we have $\sigma^{m \wedge d(x)}(x) = \sigma^{n \wedge d(x)}(x)$ and $m - m \wedge d(x) = n - n \wedge d(x)$. Since $m \neq n$ and $m - m \wedge d(x) = n - n \wedge d(x)$, we immediately have $d(\mu) \neq d(\nu)$.

Fix $y \in s(\alpha)\Lambda^{\leq\infty}$; we must show that $\mu\alpha y = \nu\alpha y$. Let $z := \mu\alpha y$. Then $z \in v\Lambda^{\leq\infty}$, and hence $\sigma^{m \wedge d(z)}(z) = \sigma^{n \wedge d(z)}(z)$.

We claim that $m \wedge d(z) = m \wedge d(x)$. To see this, fix $1 \leq i \leq k$. If $m_i \leq d(x)_i$, then $d(z)_i \geq d(\mu)_i = m_i$, so we also have $m_i \leq d(z)_i$, and $(m \wedge d(x))_i = m_i = (m \wedge d(z))_i$. If $m_i > d(x)_i$, then $x \in \Lambda^{\leq\infty}$ forces $s(\mu)\Lambda^{e_i} = \emptyset$. Hence $d(z)_i = d(\mu)_i = d(x)_i$, and $(m \wedge d(x))_i = d(x)_i = d(z)_i = (m \wedge d(z))_i$. This establishes the claim. A similar argument shows that $n \wedge d(z) = n \wedge d(x)$.

We now have

$$(4.1) \quad \alpha y = \sigma^{d(\mu)}(z) = \sigma^{m \wedge d(x)}(z) = \sigma^{m \wedge d(z)}(z) = \sigma^{n \wedge d(z)}(z) = \sigma^{n \wedge d(x)}(z).$$

We now calculate coordinate-wise to see that $(m \vee n) \wedge d(x) = (m \wedge d(x)) \vee (n \wedge d(x))$. In particular, $d(\mu\alpha) \geq d(\nu)$, so $\nu = (\mu\alpha)(0, d(\nu))$. By definition of

ν , we have $z = \nu\sigma^{n\wedge d(x)}(z)$. So $\mu\alpha y = z = \nu\sigma^{n\wedge d(x)}(z)$, and this is equal to $\nu\alpha y$ by (4.1). ■

Proof of Proposition 4.3. (1) \implies (2) Suppose Λ has no local periodicity and let I be an ideal in $C^*(\Lambda)$. Then $I = P\tilde{I}P$ for some ideal \tilde{I} of $C^*(\tilde{\Lambda})$. By Proposition 3.6 $\tilde{\Lambda}$ has no local periodicity so [14, Proposition 3.6] implies that \tilde{I} contains a vertex projection $s_{[x;m]}$. Lemma 4.1(1) then implies that $I = P\tilde{I}P$ also contains a vertex projection.

(2) \implies (3) For $v \in \Lambda^0$, $\pi_S(s_v) = S_v$ is the projection onto

$$\overline{\text{span}}\{u_x : x \in \Lambda^{\leq\infty}\}$$

and so is non-zero. So $\ker(\pi_S)$ contains no vertex projection and is trivial by (2).

(3) \implies (1) The proof of this implication runs almost exactly the same as in the proof of [14, Proposition 3.6], but we substitute Lemma 4.4 for [14, Lemma 3.4]. The broad strategy is as follows: we argue by contrapositive, supposing Λ has local periodicity m, n at v . Lemma 4.4 implies that there exist distinct elements m', n' of \mathbb{N}^k and paths $\mu \in v\Lambda^{m'}, \nu \in v\Lambda^{n'}s(\mu)$ and $\alpha \in s(\mu)\Lambda$ such that $\mu\alpha y = \nu\alpha y$ for all $y \in s(\alpha)\Lambda^{\leq\infty}$. Let

$$a := s_{\mu\alpha}s_{\mu\alpha}^* - s_{\nu\alpha}s_{\nu\alpha}^*$$

We use the gauge action to prove that $a \neq 0$, and show directly that $a \in \ker(\pi_S)$, so the boundary path representation is not faithful. ■

Before we state the next result we recall some terminology from [11, Section 5]. We say a subset $H \subseteq \Lambda^0$ is **hereditary** if $r(\lambda) \in H$ implies $s(\lambda) \in H$ for all $\lambda \in \Lambda$. We say H is **saturated** if

$$\{s(\lambda) : \lambda \in v\Lambda^{\leq e_i}\} \subseteq H \text{ for some } i \in \{1, \dots, k\} \text{ implies } v \in H.$$

If $H \subset \Lambda^0$ is saturated and hereditary, then $\Lambda \setminus \Lambda H$ is a sub- k -graph of Λ .

PROPOSITION 4.5: *Let Λ be a row-finite locally convex k -graph. Then the following are equivalent.*

- (1) every ideal of $C^*(\Lambda)$ is gauge-invariant;
- (2) for every saturated hereditary subset $H \subset \Lambda^0$, $\Lambda \setminus \Lambda H$ has no local periodicity.

Proof. (2) \implies (1) Suppose $\Lambda \setminus \Lambda H$ has no local periodicity for each saturated hereditary subset H of Λ^0 . Fix a non-zero ideal I of $C^*(\Lambda)$ and let $H := \{v \in \Lambda^0 : s_v \in I\}$. Theorem 5.2 of [11] guarantees that this set is saturated and hereditary. Let $\{t_\lambda : \lambda \in \Lambda \setminus \Lambda H\}$ be the universal generating Cuntz–Krieger family in $C^*(\Lambda \setminus \Lambda H)$. Theorem 5.2 of [11] also implies that the ideal I_H generated by $\{s_v : v \in H\}$ is a gauge invariant ideal of $C^*(\Lambda)$, and that there is an isomorphism $\varphi : C^*(\Lambda \setminus \Lambda H) \rightarrow C^*(\Lambda)/I_H$ satisfying $\varphi(t_\lambda) = s_\lambda + I_H$ for all $\lambda \in \Lambda \setminus \Lambda H$. So it suffices to show that $I = I_H$. Since $I_H \subset I$, the quotient map $q_I : C^*(\Lambda) \rightarrow C^*(\Lambda)/I$ defined by

$$q_I(s_\lambda) = s_\lambda + I$$

descends to a C^* -homomorphism $\tilde{q}_I : C^*(\Lambda)/I_H \rightarrow C^*(\Lambda)/I$ satisfying

$$\tilde{q}_I(s_\lambda + I_{H_I}) = s_\lambda + I$$

for all $\lambda \in \Lambda$. Consider the composition

$$\tilde{q}_I \circ \varphi : C^*(\Lambda \setminus \Lambda H) \rightarrow C^*(\Lambda)/I.$$

We claim $\tilde{q}_I \circ \varphi(t_v) \neq 0$ for all $v \in \Lambda^0 \setminus H = (\Lambda \setminus \Lambda H)^0$.

To see this, fix $v \in (\Lambda \setminus \Lambda H)^0$. Then

$$v \notin H \implies s_v \notin I \implies q_I(s_v) \neq 0 \implies \tilde{q}_I(s_v + I_{H_I}) \neq 0 \implies \tilde{q}_I \circ \varphi(t_v) \neq 0.$$

Hence $\ker(\tilde{q}_I \circ \varphi)$ contains no vertex projections and Proposition 4.3 implies that $\ker(\tilde{q}_I \circ \varphi) = \{0\}$. So $\tilde{q}_I \circ \varphi$ is an isomorphism, and in particular $\tilde{q}_I : C^*(\Lambda)/I_{H_I} \rightarrow C^*(\Lambda)/I$ is injective. This forces $I_H = I$ as required.

(1) \implies (2) The proof of this implication is almost identical to the proof of (1) \implies (2) in [14, Proposition 3.7]. We argue by contrapositive. Suppose there are a saturated hereditary subset $H \subset \Lambda$, a vertex $v \in \Lambda^0 \setminus H$ and elements $m \neq n \in \mathbb{N}^k$ such that $\Lambda \setminus \Lambda H$ has local periodicity m, n at v . Let $\{t_\lambda : \lambda \in \Lambda \setminus \Lambda H\}$ be the universal generating Cuntz–Krieger family for $C^*(\Lambda \setminus \Lambda H)$. Theorem 5.2 of [11] guarantees there exists an isomorphism $\varphi : C^*(\Lambda \setminus \Lambda H) \rightarrow C^*(\Lambda)/I_H$ satisfying $\varphi(t_\lambda) = s_\lambda + I_H$. Let q_{I_H} denote the quotient map from $C^*(\Lambda)$ to $C^*(\Lambda)/I_H$.

As in the proof of Proposition 4.3, we construct $\mu, \nu, \alpha \in \Lambda \setminus \Lambda H$ such that $a = t_{\mu\alpha}t_{\mu\alpha}^* - t_{\nu\alpha}t_{\nu\alpha}^* \in C^*(\Lambda \setminus \Lambda H) \setminus \{0\}$ such that $\pi_T(a) = 0$ where π_T is the boundary path representation of $C^*(\Lambda \setminus \Lambda H)$. Let $b := s_{\mu\alpha}s_{\mu\alpha}^* - s_{\nu\alpha}s_{\nu\alpha}^* \in C^*(\Lambda)$. Then $\varphi \circ q_{I_H}(b) = a \neq 0$ but $\pi_T \circ \varphi \circ q_{I_H}(b) = \pi_T(a) = 0$. Since φ is an isomorphism, the kernel of $\varphi \circ q_{I_H}$ is I_H . Since the kernel of π_T contains no

vertices of $\Lambda \setminus \Lambda H$, the ideal $J = \ker(\pi_T \circ \varphi \circ q_{I_H})$ also satisfies $J_H = H$. We have $J \neq I_H$ because $b \in J \setminus I_H$. Theorem 5.2 of [11] implies that $H \mapsto I_H$ is an isomorphism of the saturated hereditary subsets of Λ and gauge-invariant ideals of $C^*(\Lambda)$, and the implication (2) \implies (1) shows that this has inverse $I \mapsto I_H$. Since $J \neq I = I_{H,J}$ it follows that J is a non-trivial ideal of $C^*(\Lambda)$ that is not gauge invariant. ■

References

- [1] S. Allen, D. Pask and A. Sims, *A dual graph construction for higher-rank graphs, and K-theory for finite 2-graphs*, Proceedings of the American Mathematical Society **134** (2006), 455–464.
- [2] T. Bates, D. Pask, I. Raeburn and W. Szymański, *The C^* -algebras of row-finite graphs*, New York Journal of Mathematics **6** (2000), 307–324.
- [3] J. Cuntz and W. Krieger, *A class of C^* -algebras and topological Markov chains*, Inventiones Mathematicae **56** (1980), 251–268.
- [4] M. Enomoto and Y. Watatani, *A graph theory for C^* -algebras*, Mathematica Japonica **25** (1980), 435–442.
- [5] C. Farthing, *Removing sources from higher-rank graphs*, Journal of Operator Theory **60** (2008), 165–198.
- [6] A. Kumjian and D. Pask, *Higher-rank graph C^* -algebras*, New York Journal of Mathematics **6** (2000), 1–20.
- [7] A. Kumjian, D. Pask and I. Raeburn, *Cuntz–Krieger algebras of directed graphs*, Pacific Journal of Mathematics **184** (1998), 161–174.
- [8] A. Kumjian, D. Pask, I. Raeburn and J. Renault, *Graphs, groupoids and Cuntz–Krieger algebras*, Journal of Functional Analysis **144** (1997), 505–541.
- [9] S. Mac Lane, *Categories for the Working Mathematician*, 2nd ed., Graduate Texts in Mathematics, Vol. 5. Springer-Verlag, New York, 1998.
- [10] I. Raeburn, *Graph Algebras*, CBMS Regional Conference Series in Mathematics, vol. 103, American Mathematical Society, 2005.
- [11] I. Raeburn, A. Sims and T. Yeend, *Higher-rank graphs and their C^* -algebras*, Proceedings of the Edinburgh Mathematical Society. Series II **46** (2003), 99–115.
- [12] I. Raeburn, A. Sims and T. Yeend, *The C^* -algebras of finitely aligned higher-rank graphs*, Journal of Functional Analysis **213** (2004), 206–240.
- [13] I. Raeburn and D. P. Williams, *Morita Equivalence and Continuous-Trace C^* -Algebras*, Mathematical Surveys and Monographs, vol. 60, 1998.
- [14] D. I. Robertson and A. Sims, *Simplicity of C^* -algebras associated to higher-rank graphs*, Proceedings of the London Mathematical Society **39** (2007), 337–344.
- [15] G. Robertson and T. Steger, *Affine buildings, tiling systems and higher-rank Cuntz–Krieger algebras*, Journal für die reine und angewandte Mathematik **513** (1999), 115–144.
- [16] G. Robertson and T. Steger, *Asymptotic K-theory for groups acting on \tilde{A}_2 buildings*, Canadian Journal of Mathematics **53** (2001), 809–833.