

Pacific Journal of Mathematics

**SIMPLICITY OF PARTIAL AND SCHMIDT DIFFERENTIAL
OPERATOR RINGS**

DENNIS R. MALM

SIMPLICITY OF PARTIAL AND SCHMIDT DIFFERENTIAL OPERATOR RINGS

D. R. MALM

In this paper we develop necessary and sufficient conditions for certain differential operator rings to be simple. We do this for a ring with finitely many commuting derivations and for a ring with a commuting Schmidt higher derivation of finite length. Also we give a correspondence between finite sets of derivations and Schmidt higher derivations.

In this first section we deal with the simplicity of a differential operator ring over a ring with finitely many commuting derivations. Many of the results of this section overlap those of others, among them Jordan [8], Hauger [5] and Voskoglou [11]. The second section deals with the construction of a Schmidt higher derivation from a set of derivations, and vice versa. The correspondence is the same as that of Heerema [6] but we give a different construction. The third section deals with simplicity of a differential operator ring over a ring with a Schmidt higher derivation. All rings are considered to be associative with unit. This research will form part of the author's Ph.D. dissertation at the University of Utah. The author wishes to thank K. R. Goodearl for his help and suggestions.

1. Let $\delta_1, \dots, \delta_k$ be derivations on a ring R . A $(\delta_1, \dots, \delta_k)$ -ideal of R is any ideal I of R such that $\delta_i(I) \subseteq I$ for all i . The ring R is said to be $(\delta_1, \dots, \delta_k)$ -simple if R is nonzero and the only $(\delta_1, \dots, \delta_k)$ -ideals of R are 0 and R . The elements $r \in R$ such that $\delta_i(r) = 0$ for all i are called $(\delta_1, \dots, \delta_k)$ -constants and form a subring of R .

If $\delta_1, \dots, \delta_k$ are commuting derivations on R , the formal linear differential operator ring

$$T = R[\Theta_1, \dots, \Theta_k; \delta_1, \dots, \delta_k]$$

is the free left R -module generated by the symbols

$$\Theta_k^{n(k)} \dots \Theta_1^{n(1)}$$

with each $n(i)$ a nonnegative integer and given a ring structure by the relations

$$\begin{aligned}\Theta_i \Theta_j &= \Theta_j \Theta_i \\ \Theta_i r &= r \Theta_i + \delta_i(r)\end{aligned}$$

for all i, j , all $r \in R$, together with the usual multiplication in R . It is convenient to think of the elements of T as polynomials, with left coefficients in R , in the Θ 's. For a k -tuple $I = (i(k), \dots, i(1))$, with each $i(j)$ a nonnegative integer, let

$$|I| = i(k) + \dots + i(1)$$

and

$$\Theta^I = \Theta_k^{i(k)} \dots \Theta_1^{i(1)}$$

and call I the *order* of Θ^I . Let $N = (n(k), \dots, n(1))$ and $M = (m(k), \dots, m(1))$. We say the order of Θ^N is greater than that of Θ^M if $|N| > |M|$ or, if $|N| = |M|$, if there exists $j \in \{1, \dots, k\}$ such that $n(i) = m(i)$ for $i > j$ and $n(j) > m(j)$. Otherwise the order of Θ^N is less than or equal to that of Θ^M .

For nonzero $t \in T$, the *leading coefficient* of t is the coefficient (in R) of the monomial of highest order in t with nonzero coefficient. We will usually denote an element $t \in T$ by

$$t = \sum \alpha_I \Theta^I$$

with $\alpha_I \in R$. If α_J is the leading coefficient of t , the *order* of t is said to be J . One should note that for nonzero $t, s \in T$, the leading coefficient of ts is the product of the leading coefficients of t and s unless that product is zero. When $k = 1$, the order will be denoted by a lower case letter.

LEMMA 1. *Let $\delta_1, \dots, \delta_k$ be commuting derivations on a ring R and F the subring of central $(\delta_1, \dots, \delta_k)$ -constants of R . If the differential operator ring*

$$T = R[\Theta_1, \dots, \Theta_k; \delta_1, \dots, \delta_k]$$

is simple then the center of T is F and R is $(\delta_1, \dots, \delta_k)$ -simple.

Proof. Certainly F is contained in the center of T and any element of R central in T is in F . So it will be enough to show that the center of T is contained in R to show F is the center of T .

If A is a proper nonzero $(\delta_1, \dots, \delta_k)$ -ideal of R , then AT is a proper nonzero ideal of T . Hence, T simple implies R is $(\delta_1, \dots, \delta_k)$ -simple.

As R is $(\delta_1, \dots, \delta_k)$ -simple, any nonzero central $(\delta_1, \dots, \delta_k)$ -constant is invertible and therefore F is a field.

Suppose $t = \sum \alpha_I \Theta^I \neq 0$ is central in T with leading coefficient α_J and order J . Since $\Theta_i t - t \Theta_i = 0$ for all i and $rt - tr = 0$ for all $r \in R$, we get $\alpha_J \in F$. Since T is simple, t is invertible. If $|J| > 0$, and the leading coefficient of t^{-1} is β_N , then $\beta_N \alpha_J = 0$ implying $\beta_N = 0$, a contradiction. Therefore $|J| = 0$ and $t \in R$. \square

We can think of the derivations $\delta_1, \dots, \delta_k$ being defined on T such that for $t \in T$

$$\delta_i(t) = \Theta_i t - t \Theta_i.$$

Note that this implies $\delta_i(\Theta_j) = 0$ for all i, j and if $t = \sum \alpha_I \Theta^I$, then $\delta_i(t) = \sum \delta_i(\alpha_I) \Theta^I$. Also note that Leibniz' rule holds:

$$\delta_j^n(ab) = \sum_{i=0}^n \binom{n}{i} \delta_j^i(a) \delta_j^{n-i}(b)$$

and also

$$\Theta_j^n a = \sum_{i=0}^n \binom{n}{i} \delta_j^i(a) \Theta_j^{n-i}$$

for all i, j , all $a, b \in R$.

The ring R can also be considered as a left T -module with a T -module multiplication $*$ under which $\Theta_j * r = \delta_j(r)$ and $s * r = sr$ for all j , all $s, r \in R$. With this module structure, right multiplication in R by a $(\delta_1, \dots, \delta_k)$ -constant is a left T -module endomorphism of R , and these are the only left T -module endomorphisms of R .

The next theorem has been proved in [5] and [11, Theorems 3.3, 3.4]. We supply a proof to indicate the difference between the \mathbf{Q} -algebra case and the $(\mathbf{Z}/p\mathbf{Z})$ -algebra case. The case where $k = 1$ was originally proved by Amitsur [1].

THEOREM 2. *Let R be a \mathbf{Q} -algebra, $\delta_1, \dots, \delta_k$ commuting derivations on R and F the subring of central $(\delta_1, \dots, \delta_k)$ -constants of R . Then the ring*

$$T = R[\Theta_1, \dots, \Theta_k; \delta_1, \dots, \delta_k]$$

is simple if and only if R is $(\delta_1, \dots, \delta_k)$ -simple and no nontrivial F -linear combination of $\delta_1, \dots, \delta_k$ is an inner derivation by a $(\delta_1, \dots, \delta_k)$ -constant. In this case the center of T is F .

Proof. If T is simple, R is $(\delta_1, \dots, \delta_k)$ -simple and the center of T is F by Lemma 1. If $\alpha_1, \dots, \alpha_k \in F$ and

$$(\alpha_1 \delta_1 + \dots + \alpha_k \delta_k)(r) = ar - ra$$

for all $r \in R$ and some $(\delta_1, \dots, \delta_k)$ -constant a , then, for

$$s = \alpha_1 \Theta_1 + \dots + \alpha_k \Theta_k - a,$$

we have $s * r = r(-a)$. So multiplication by s is a left T -module endomorphism of R . Then for any $t \in T$, the commutator $st - ts$ is in the annihilator of ${}_T R$. Since T is simple, ${}_T R$ is faithful. This forces s to be central in T and so $s \in F$ by Lemma 1. Therefore $\alpha_i = 0$ for all i .

For the converse we will use induction on k . If $k = 1$, let A be a nonzero ideal of $T = R[\Theta; \delta]$. If n is the minimal order of nonzero elements of A , then the set

$$J = \{a \in R \mid a\Theta^n + a_{n-1}\Theta^{n-1} + \dots + a_0 \in A \text{ for some } a_{n-1}, \dots, a_0 \in R\}$$

is a nonzero δ -ideal of R . Since R is δ -simple, $1 \in J$, and so we can choose $q \in A$,

$$q = \sum_{j=0}^n a_j \Theta^j$$

with $a_n = 1$. For any $r \in R$, we have $qr - rq \in A$ of smaller order than n , so $qr - rq = 0$. Similarly $\Theta q - q\Theta = 0$, so each a_j is a δ -constant. If $n > 0$,

$$0 = qr - rq = (n\delta(r) + a_{n-1}r - ra_{n-1})\Theta^{n-1} + \text{terms of lower order,}$$

implying $n\delta(r) + a_{n-1}r - ra_{n-1} = 0$ for all $r \in R$, a contradiction. Therefore $n = 0$ and $q = 1 \in A$. Thus $A = T$ and T is simple.

Suppose $k > 1$ and the theorem is true for fewer than k derivations. Now

$$T = R[\Theta_1, \dots, \Theta_k; \delta_1, \dots, \delta_k] \cong S[\Theta_2, \dots, \Theta_k; \delta_2, \dots, \delta_k]$$

where $S = R[\Theta_1; \delta_1]$. By induction, T is simple if S is $(\delta_2, \dots, \delta_k)$ -simple and no nontrivial linear combination of $\delta_2, \dots, \delta_k$ by central $(\delta_2, \dots, \delta_k)$ -constants of S is an inner derivation on S by a $(\delta_2, \dots, \delta_k)$ -constant of S . If

$$\alpha = \sum_{i=0}^n a_i \Theta_1^i$$

is a central $(\delta_2, \dots, \delta_k)$ -constant of S with $a_n \neq 0$, then

$$\delta_j(\alpha) = \sum_{i=0}^n \delta_j(a_i) \Theta_1^i = 0$$

for $j \geq 2$ and

$$\Theta_1 \alpha - \alpha \Theta_1 = \sum_{i=0}^n \delta_1(a_i) \Theta_1^i = 0.$$

So each a_j is a $(\delta_1, \dots, \delta_k)$ -constant. If $n > 0$, then for any $r \in R$,

$$0 = \alpha r - r \alpha = (a_n r - r a_n) \Theta_1^n + (n a_n \delta_1(r) + a_{n-1} r - r a_{n-1}) \Theta_1^{n-1} \\ + \text{terms of lower order.}$$

The vanishing of the Θ_1^n term yields $a_n \in F$, and then from the vanishing of the Θ_1^{n-1} term we get a contradiction. Therefore $n = 0$ and all central $(\delta_2, \dots, \delta_k)$ -constants of S are in F .

Suppose $\alpha_2 \delta_2 + \dots + \alpha_k \delta_k = \delta_b$, the inner derivation on S by the $(\delta_2, \dots, \delta_k)$ -constant $b = b_0 + \dots + b_m \Theta_1^m$, with $b_m \neq 0$ and the $\alpha_i \in F$. Since

$$0 = (\alpha_2 \delta_2 + \dots + \alpha_k \delta_k)(\Theta_1) = b \Theta_1 - \Theta_1 b = \sum_{i=0}^m -\delta_1(b_i) \Theta_1^i$$

and b is a $(\delta_2, \dots, \delta_k)$ -constant, we have that each b_j is a $(\delta_1, \dots, \delta_k)$ -constant. Now

$$\delta_b(R) = (\alpha_2 \delta_2 + \dots + \alpha_k \delta_k)(R) \subseteq R,$$

so for all $r \in R$, the element

$$br - rb = (b_m r - r b_m) \Theta_1^m + (m b_m \delta_1(r) + b_{m-1} r - r b_{m-1}) \Theta_1^{m-1} \\ + \text{terms of lower order}$$

is in R . If $m \geq 2$, then $b_m \in F$ and we get a contradiction from the vanishing of the Θ_1^{m-1} term. Therefore $m \leq 1$ and $b = b_0 + b_1 \Theta_1$ (here b_1 may be 0). As before $b_1 \in F$ and for any $r \in R$,

$$(\alpha_2 \delta_2 + \dots + \alpha_k \delta_k)(r) = br - rb = b_1 \delta_1(r) + b_0 r - r b_0.$$

Hence each $\alpha_i = 0$. Therefore no nontrivial linear combination of $\delta_2, \dots, \delta_k$ by central $(\delta_2, \dots, \delta_k)$ -constants of S is an inner derivation on S by a $(\delta_2, \dots, \delta_k)$ -constant of S .

Suppose S has a nonzero $(\delta_2, \dots, \delta_k)$ -ideal B . Let l be the minimal order of nonzero elements of B . Then the set

$$J = \{a \in R \mid a \Theta_1^l + a_{l-1} \Theta_1^{l-1} + \dots + a_0 \in B \text{ for some } a_{l-1}, \dots, a_0 \in R\}$$

is a nonzero $(\delta_1, \dots, \delta_k)$ -ideal of R . Since R is $(\delta_1, \dots, \delta_k)$ -simple, $1 \in J$, and so there is an operator

$$s = \Theta_1^l + s_{l-1} \Theta_1^{l-1} + \dots + s_0 \in B.$$

Since $\delta_j(s) \in B$ for $j \geq 2$ and $\delta_1(s) = \Theta_1 s - s \Theta_1 \in B$, we have $\delta_j(s) = 0$ for all j since these operators have lower order than l . Therefore each s_j is a $(\delta_1, \dots, \delta_k)$ -constant. Also, if $l > 0$, for any $r \in R$, we obtain

$$0 = sr - rs = (l\delta_1(r) + s_{l-1}r - rs_{l-1})\Theta_1^{l-1} + \text{terms of lower order}$$

and the vanishing of the Θ_1^{l-1} term yields a contradiction. Therefore $l = 0$ and $1 \in B$ proving that S is $(\delta_2, \dots, \delta_k)$ -simple.

The induction hypothesis is now verified, and thus T is simple. \square

When R is a $(\mathbf{Z}/p\mathbf{Z})$ -algebra for some prime p , if δ is a derivation on R , then

$$\delta^p, \delta^{p^2}, \dots$$

are all derivations on R . This complicates the simplicity criteria for the differential operator ring in the positive characteristic case.

LEMMA 3. *Let R be a $(\mathbf{Z}/p\mathbf{Z})$ -algebra, p a prime, $\delta_1, \dots, \delta_k$ commuting derivations on R and F the subring of central $(\delta_1, \dots, \delta_k)$ -constants of R . Assume that no nontrivial F -linear combination of*

$$\delta_1, \delta_1^p, \delta_1^{p^2}, \dots$$

is an inner derivation by a $(\delta_1, \dots, \delta_k)$ -constant. Set $S = R[\Theta_1; \delta_1]$ and let $b \in S$ be a $(\delta_1, \dots, \delta_k)$ -constant.

(a) *If $br - rb \in R$ for all $r \in R$, then*

$$b = a + \sum_{j=0}^m a_j \Theta_1^{p^j}$$

for some $(\delta_1, \dots, \delta_k)$ -constant $a \in R$ and some $a_0, \dots, a_m \in F$.

(b) *If b is central in S , then $b \in F$.*

Proof. (a) Write out

$$b = \sum_{i=0}^n b_i \Theta_1^i,$$

and suppose $b_n \neq 0$. Since b is a $(\delta_1, \dots, \delta_k)$ -constant, each b_i is a $(\delta_1, \dots, \delta_k)$ -constant. From assumption (a), if $n > 0$ we get $b_n \in F$. If $n \leq 1$, we are done. If $n > 1$ and if p does not divide n , then, for all $r \in R$,

$$\begin{aligned} br - rb &= (b_n r - r b_n) \Theta_1^n + (n b_n \delta_1(r) + b_{n-1} r - r b_{n-1}) \Theta_1^{n-1} \\ &\quad + \text{terms of lower order.} \end{aligned}$$

Since $br - rb \in R$, we get a contradiction from the vanishing of the Θ_1^{n-1} term.

If $p \mid n$, set

$$t = \max\{i \mid b_i \neq 0 \text{ and } p \text{ does not divide } i\}$$

or 0 if no such i exists. Let

$$A = \{j \mid b_j \neq 0 \text{ and } n \geq j > t\}.$$

Note that $n \in A$, and that for $j \in A$, we have

$$j = p^{l(j)}m_j$$

with $(m_j, p) = 1$ and $l(j) \geq 1$. Let

$$s = \max(\{t - 1\} \cup \{p^{l(j)}(m_j - 1) \mid j \in A\})$$

and

$$C = \{j \in A \mid p^{l(j)}(m_j - 1) = s\} \cup \{t \text{ if } t - 1 = s\}.$$

If $t \in C$, let $m_t = t$ and $l(t) = 0$, so that each $j \in C$ can be written as

$$j = p^{l(j)}m_j.$$

Let

$$B = \{b_j \mid j \in C\}.$$

Since $br - rb \in R$ for all $r \in R$, we have $B \subseteq F$.

If $s > 0$, evaluating the coefficient of Θ_1^s in $br - rb$, we have

$$\sum_{j \in C} m_j b_j \delta_1^{p^{l(j)}}(r) + b_s r - r b_s = 0$$

for all $r \in R$. Since $m_j b_j \neq 0$ for all $j \in C$, this gives a contradiction. Therefore $s = 0$, and hence $t \leq 1$ and $m_j = 1$ for all $j \in A$. This proves (a).

(b) If b is central, then $br - rb = 0 \in R$ for all $r \in R$, so by part (a) we have

$$b = a + \sum_{j=0}^m a_j \Theta_1^{p^j}$$

for some $(\delta_1, \dots, \delta_k)$ -constant $a \in R$ and some $a_0, \dots, a_m \in F$. But

$$0 = br - rb = \sum_{j=0}^m a_j \delta_1^{p^j}(r) + ar - ra$$

for all $r \in R$, implying $a_j = 0$ for all j . Then a is central in R and so $b \in F$. \square

The next theorem was proved for $k = 1$ in [8, Thm. 4.1.6]. The case where $k = 1$ and R is commutative was considered in [2, Thm. 3.2a] and [3, Thm. 2.3]. There are related results in [5] and [11, Thm. 3.5]; in particular, Voskoglou proves that the given conditions are necessary for T to be simple.

THEOREM 4. *Let R be a $(\mathbf{Z}/p\mathbf{Z})$ -algebra, p a prime, $\delta_1, \dots, \delta_k$ commuting derivations on R and F the subring of central $(\delta_1, \dots, \delta_k)$ -constants of R . Then the ring*

$$T = R[\Theta_1, \dots, \Theta_k; \delta_1, \dots, \delta_k]$$

is simple if and only if R is $(\delta_1, \dots, \delta_k)$ -simple and no nontrivial F -linear combination of

$$\{\delta_i^{p^j} \mid i = 1, \dots, k, j = 0, 1, \dots\}$$

is an inner derivation by a $(\delta_1, \dots, \delta_k)$ -constant. In this case the center of T is F .

Proof. If T is simple, R is $(\delta_1, \dots, \delta_k)$ -simple and the center of T is F by Lemma 1. If there are some $\alpha_{ij} \in F$ such that

$$\sum \alpha_{ij} \delta_i^{p^j}$$

is an inner derivation by the $(\delta_1, \dots, \delta_k)$ -constant a , then

$$s = \sum \alpha_{ij} \Theta_i^{p^j} - a$$

acts on ${}_T R$ as right multiplication by $-a$. For any $t \in T$, we get that $st - ts$ is in the annihilator of ${}_T R$. Then T being simple implies s is central in T . By Lemma 1, we have $s \in F$ and so each $\alpha_{ij} = 0$.

For the converse we use induction on k . If $k = 1$ and A is a nonzero ideal of $T = R[\Theta; \delta]$ with n the minimal order of nonzero elements of A , the set

$$J = \{a \in R \mid a\Theta^n + a_{n-1}\Theta^{n-1} + \dots + a_0 \in A \text{ for some } a_{n-1}, \dots, a_0 \in R\}$$

is a nonzero δ -ideal of R . Since R is δ -simple, $1 \in J$, and we can pick $q \in A$,

$$q = \sum_{i=0}^n a_i \Theta^i$$

with $a_n = 1$. Since $\Theta q - q\Theta \in A$ and this operator is of lower order than n , we get $\Theta q - q\Theta = 0$ and so each a_i is a δ -constant. Also

$qr - rq = 0$ for all $r \in R$, and so q is central in T . By Lemma 3, we have $q \in F$. Therefore $n = 0$ and $1 \in A$, and so T is simple.

Suppose $k > 1$ and the theorem is true for fewer than k derivations. Now

$$T = R[\Theta_1, \dots, \Theta_k; \delta_1, \dots, \delta_k] \cong S[\Theta_2, \dots, \Theta_k; \delta_2, \dots, \delta_k]$$

where $S = R[\Theta_1; \delta_1]$. By induction T is simple if S is $(\delta_2, \dots, \delta_k)$ -simple and no nontrivial linear combination of

$$\{\delta_i^{p_j} \mid i = 2, \dots, k, j = 0, 1, \dots\}$$

by central $(\delta_2, \dots, \delta_k)$ -constants of S is an inner derivation by a $(\delta_2, \dots, \delta_k)$ -constant of S .

Suppose there exist α_{ij} , with $i > 1$, central $(\delta_2, \dots, \delta_k)$ -constants of S such that

$$\sum \alpha_{ij} \delta_i^{p_j}$$

is inner by the $(\delta_2, \dots, \delta_k)$ -constant

$$b = \sum_{i=0}^n b_i \Theta_1^i.$$

Since the α_{ij} are central in S , they are also δ_1 -constants and so each $\alpha_{ij} \in F$ by Lemma 3. For all $r \in R$,

$$br - rb = \sum \alpha_{ij} \delta_i^{p_j}(r) \in R.$$

So, by Lemma 3,

$$b = a + \sum_{j=0}^m a_j \Theta_1^{p_j}$$

for some $(\delta_1, \dots, \delta_k)$ -constant $a \in R$ and some $a_0, \dots, a_m \in F$. Then, for any $r \in R$,

$$\sum \alpha_{ij} \delta_i^{p_j}(r) - \sum_{j=0}^m a_j \delta_1^{p_j}(r) = ar - ra,$$

and so each $\alpha_{ij} = 0$.

If S has a nonzero $(\delta_2, \dots, \delta_k)$ -ideal B , let n be the minimal order of nonzero elements of B . Then the set

$$J = \{a \in R \mid a\Theta_1^n + a_{n-1}\Theta_1^{n-1} + \dots + a_0 \in B \text{ for some } a_{n-1}, \dots, a_0 \in R\}$$

is a nonzero $(\delta_1, \dots, \delta_k)$ -ideal of R . Therefore there is a $q \in B$ with

$$q = \sum_{i=0}^n a_i \Theta_1^i$$

and $a_n = 1$. Also the operators $\Theta_1 q - q \Theta_1 = \delta_1(q)$, as well as $\delta_i(q)$ for $i > 1$ and $qr - rq$ for all $r \in R$ are in B , have lower order than n , and so are 0. Therefore q is a central $(\delta_1, \dots, \delta_k)$ -constant of S and so $q \in F$ by Lemma 3. Then $n = 0$ and $1 \in B$, proving that S is $(\delta_2, \dots, \delta_k)$ -simple.

The induction hypothesis is now established, and therefore T is simple. \square

As an example to satisfy the hypotheses of Theorem 4, let K be a field of characteristic $p > 0$ and let $\{x_{ij} \mid i = 1, \dots, k, j = 1, 2, \dots\}$ be independent indeterminates over K . Set $R = K(\{x_{ij}\})$. Define commuting K -linear derivations $\delta_1, \dots, \delta_k$ on R by

$$\delta_i(x_{ij}) = x_{i,j+1} \quad \text{and} \quad \delta_i(x_{mj}) = 0$$

for all $m \neq i$.

First we show the set

$$\{\delta_i^j \mid i = 1, \dots, k, j = 1, 2, \dots\}$$

is linearly independent over R . Suppose there exists some $\alpha_{ij} \in R$ such that

$$\sum \alpha_{ij} \delta_i^j = 0.$$

For fixed n , pick l such that no term $x_{n,l+i}$, for $i > 0$, is involved in any of the $\alpha_{n,j}$. Then

$$0 = \sum \alpha_{ij} \delta_i^j(x_{nl}) = \sum_j \alpha_{nj} x_{n,l+j}$$

which implies each $\alpha_{nj} = 0$. Since n was arbitrary we have $\alpha_{ij} = 0$ for all i, j .

Therefore the set

$$\{\delta_i^{p^j} \mid i = 1, \dots, k, j = 0, 1, \dots\}$$

is linearly independent over the central $(\delta_1, \dots, \delta_k)$ -constants of R . As R is simple, the differential operator ring T is simple.

2. A finite *Schmidt higher derivation* (S.h.d.) on a ring R is a set of additive maps d_0, d_1, \dots, d_k on R such that d_0 is the identity map and

$$d_i(ab) = \sum_{j=0}^i d_j(a) d_{i-j}(b)$$

for all $a, b \in R$ and $i = 1, \dots, k$. Infinite S.h.d.'s were first used in [4] and [10]. For example, if δ is a derivation on a \mathbf{Q} -algebra R , then the maps

$$d_i = \delta^i / i!$$

define a S.h.d. on R . For another example, let K be any field and $K[x]$ a polynomial ring over K . Define K -linear maps d_0, d_1, \dots, d_k on $K[x]$ such that

$$d_i(x^j) = \binom{j}{i} x^{j-i}$$

for all i, j . Then d_0, d_1, \dots, d_k is a S.h.d. on $K[x]$, with $d_0 =$ identity map and $d_1 = d/dx$. If $\text{char}(K) = 2$, then

$$d_2(x^{2n}) = nx^{2n-2}$$

and

$$d_2(x^{2n+1}) = nx^{2n-1}.$$

Let d_0, d_1, \dots, d_k be a S.h.d. on a ring R and observe that $d_i(1) = 0$ for $i = 1, \dots, k$. A (d_0, d_1, \dots, d_k) -ideal of R is any ideal A of R such that $d_i(A) \subseteq A$ for all i . The ring R is said to be (d_0, d_1, \dots, d_k) -simple if R is nonzero and the only (d_0, \dots, d_k) -ideals of R are 0 and R . The elements $r \in R$ such that $d_i(r) = 0$ for all $i > 0$ are called (d_1, \dots, d_k) -constants and form a subring of R . For d_0, d_1, \dots, d_k a commuting S.h.d. on R , we can form a differential operator ring over R , analogous to the differential operator rings of the previous section, as in the following theorem. We first fix some notation.

Let

$$M = \{(l(0), \dots, l(k)) \mid l(0) = -\infty \text{ and } l(i) \text{ is a nonnegative integer for } i > 0\}$$

and observe that M is an additive semigroup with zero element $e(0) = (-\infty, 0, \dots, 0)$. For $i > 0$, let $e(i) = (l(0), \dots, l(k)) \in M$ with $l(i) = 1$ and $l(j) = 0$ for all other $j > 0$. We have for $L = (l(0), \dots, l(k)) \in M$ and a nonnegative integer j ,

$$L + je(i) = (l(0), \dots, l(i) + j, \dots, l(k))$$

for $i > 0$, while

$$L + je(0) = L.$$

THEOREM 5. Let d_0, d_1, \dots, d_k be a commuting S.h.d. on R . Then there is a differential operator ring S generated by R and elements D_0, D_1, \dots, D_k with relations

$$D_i D_j = D_j D_i, \quad D_0 = 1, \quad D_j r = \sum_{i=0}^j d_i(r) D_{j-i}$$

for all i, j and all $r \in R$, such that S is free as a left R -module with the monomials

$$D_k^{n(k)} \dots D_1^{n(1)}$$

for $n(i) \geq 0$ forming a basis.

Proof. Let U be a free left R -module with basis

$$\{u_L \mid L = (l(0), \dots, l(k)) \in M\}.$$

Let

$$E = \text{End}_{\mathbf{Z}}(U).$$

There is an injective ring map $R \rightarrow E$ under which any element $r \in R$ corresponds to left multiplication by r on U . We identify R with its image under this map so that R is now a subring of E .

Since U is the direct sum of the subgroups Ru_L , we may define maps $D_j \in E$, for $0 \leq j \leq k$, such that

$$D_j(ru_L) = \sum_{i=0}^j d_j(r) u_{L+e(j-i)}$$

for $r \in R$. Let S be the subring of E generated by D_0, D_1, \dots, D_k and R .

Given $r \in R$ and $L \in M$,

$$\begin{aligned} D_i D_j(ru_L) &= \sum_{m=0}^i \sum_{n=0}^j d_m(d_n(r)) u_{L+e(j-n)+e(i-m)} \\ &= \sum_{n=0}^j \sum_{m=0}^i d_n(d_m(r)) u_{L+e(i-m)+e(j-n)} = D_j D_i(ru_L). \end{aligned}$$

Thus $D_i D_j = D_j D_i$ for all i, j .

Given $r, s \in R$,

$$\begin{aligned} D_j(rsu_L) &= \sum_{i=0}^j d_i(rs) u_{L+e(j-i)} = \sum_{i=0}^j \left(\sum_{n=0}^i d_{i-n}(r) d_n(s) \right) u_{L+e(j-i)} \\ &= \sum_{i=0}^j \sum_{n=0}^{j-i} d_i(r) d_n(s) u_{L+e(j-i-n)} = \sum_{i=0}^j d_i(r) D_{j-i}(su_L). \end{aligned}$$

Thus

$$D_j r = \sum_{i=0}^j d_i(r) D_{j-i}.$$

It follows that if

$$D^L = D_k^{l(k)} \dots D_1^{l(1)}$$

for $L = (l(0), \dots, l(k)) \in M$, then

$$\sum_{L \in M} R D^L$$

is closed under multiplication and so

$$S = \sum_{L \in M} R D^L.$$

Thus S is generated as a left R -module by

$$\{D^L \mid L \in M\}.$$

Since $d_i(1) = 0$ for $i > 0$, we have

$$D_j u_L = u_{L+e(j)} \quad \text{and} \quad D_j^i u_L = u_{L+ie(j)}.$$

To show the D^L are left linearly independent over R , let $r_0, \dots, r_n \in R$ such that

$$r_0 + r_1 D^{L(1)} + \dots + r_n D^{L(n)} = 0$$

for distinct $L(i) \in M$. Then

$$0 = (r_0 + r_1 D^{L(1)} + \dots + r_n D^{L(n)}) u_{e(0)} = r_0 u_{e(0)} + r_1 u_{L(1)} + \dots + r_n u_{L(n)},$$

implying $r_0 = r_1 = \dots = r_n = 0$. Thus S is free as a left R -module with the monomials D^L forming a basis. \square

We denote the differential operator ring constructed in Theorem 5 by

$$S = R[D_1, \dots, D_k; d_0, d_1, \dots, d_k].$$

It is also convenient to define leading coefficients for elements of S . For $L = (l(0), \dots, l(k)) \in M$, set

$$|L| = l(1) + 2l(2) + \dots + kl(k) \quad \text{and} \quad D^L = D_k^{l(k)} \dots D_1^{l(1)}$$

and call L the *weighted order* of D^L . Let $N = (n(0), \dots, n(k)) \in M$. We say the weighted order of D^N is greater than that of D^L if $|N| > |L|$ or, if $|N| = |L|$, if there exists $j \in \{1, \dots, k\}$ such that $n(i) = l(i)$ for $i > j$ and $n(j) > l(j)$. Otherwise the weighted order of D^N is less than or equal to that of D^L .

For nonzero $s \in S$, the *leading coefficient* of s is the coefficient of the monomial of highest weighted order in s with nonzero coefficient. We will usually denote an element $s \in S$ by

$$s = \sum \alpha_L D^L$$

with $\alpha_L \in R$. If α_J is the leading coefficient of s , the *weighted order* of s is said to be J . One should note that for nonzero $t, s \in S$, the leading coefficient of ts is the product of the leading coefficients unless that product is zero. Although we could have defined order as in the last section, weighted order is useful in Theorem 11.

Before developing criteria for the simplicity of these rings, we note a case when S is noetherian and we give a construction of a S.h.d. from derivations and vice versa.

THEOREM 6. *Let d_0, d_1, \dots, d_k be a commuting S.h.d. on a ring R . If R is right (left) noetherian, the ring*

$$S = R[D_1, \dots, D_k; d_0, d_1, \dots, d_k]$$

is right (left) noetherian.

Proof. Set

$$T = R[D_1, \dots, D_{k-1}; d_0, d_1, \dots, d_{k-1}].$$

Note that for $t \in T$, we have $\delta(t) = D_k t - t D_k \in T$. Therefore, δ defines a derivation on T . Also $S \cong T[\Theta; S]$. Then, by induction on k , and by [9, Thm. 2, p. 65], we have R right (left) noetherian implies S is right(left) noetherian. \square

For the rest of this section, we fix some notation. Given derivations $\delta_1, \dots, \delta_r$ and $i = 1, \dots, r$, let

$$\delta_{(i,r)} = \sum \delta_{j(1)} \cdots \delta_{j(i)}$$

where the sum is over all formally distinct i -tuples $(j(1), \dots, j(i))$ with each $j(l)$ a positive integer and $j(1) + \cdots + j(i) = r$. For example,

$$\delta_{(2,4)} = \delta_1 \delta_3 + \delta_3 \delta_1 + \delta_2 \delta_2.$$

We define $\Theta_{(i,r)}$ similarly. Also, given a S.h.d. d_0, d_1, \dots, d_k , we define $d_{(i,r)}$ and $D_{(i,r)}$ in a similar manner.

LEMMA 7. Let $k \in \mathbf{N}$, let R be a ring in which $k!$ is invertible, let $\delta_1, \dots, \delta_k$ be derivations on R , let $R[t]$ be a polynomial ring over R , and set

$$R[s] = R[t]/(t^{k+1}).$$

Extend each δ_i to a derivation on $R[s]$ where $\delta_i(s) = 0$, and set

$$\partial = s\delta_1 + s^2\delta_2 + \dots + s^k\delta_k.$$

(a) $\partial^n = \sum_{j=0}^{k-n} s^{n+j}\delta_{(n,n+j)}$ for $1 \leq n \leq k$.

(b) $\sigma = \sum_{n=0}^k \partial^n/n!$ is a ring endomorphism of $R[s]$.

Proof. (a) If $n = 1$,

$$\sum_{j=0}^{k-1} s^{j+1}\delta_{(1,j+1)} = \sum_{j=0}^{k-1} s^{j+1}\delta_{j+1} = \partial.$$

Suppose $n > 1$ and (a) is true for $n - 1$. Then

$$\begin{aligned} \partial^n &= \partial^{n-1}\partial = \left(\sum_{j=0}^{k-n+1} s^{n-1+j}\delta_{(n-1,n-1+j)} \right) (s\delta_1 + \dots + s^k\delta_k) \\ &= \sum_{i=0}^{k-n} s^{n+i} \left(\sum_{j=0}^i \delta_{(n-1,n+j-1)}\delta_{i-j+1} \right) = \sum_{i=0}^{k-n} s^{n+i}\delta_{(n,n+i)}. \end{aligned}$$

(b) Set $d = \delta_1 + s\delta_2 + \dots + s^{k-1}\delta_k$. Then $\partial = sd$ and d is a derivation on $R[s]$. For $a, b \in R[s]$,

$$\begin{aligned} \sigma(a)\sigma(b) &= \left(\sum_{n=0}^k s^n d^n(a)/n! \right) \left(\sum_{i=0}^k s^i d^i(b)/i! \right) \\ &= \sum_{n=0}^k s^n \left(\sum_{i=0}^n d^{n-i}(a)d^i(b)/(n-i)!i! \right) \\ &= \sum_{n=0}^k s^n \left(\sum_{i=0}^n \binom{n}{i} d^{n-i}(a)d^i(b)/n! \right) \\ &= \sum_{n=0}^k s^n d^n(ab)/n! = \sigma(ab). \end{aligned}$$

Clearly σ is additive, so σ is a ring endomorphism. \square

LEMMA 8. Let $k \in \mathbf{N}$, let R be a ring in which $k!$ is invertible, let d_0, d_1, \dots, d_k be a S.h.d. on R , let $R[t]$ be a polynomial ring over R , and set

$$R[s] = R[t]/(t^{k+1}).$$

Extend d_0, d_1, \dots, d_k to a S.h.d. on $R[s]$ where $d_i(s) = 0$ for $i = 1, \dots, k$.

(a) $\phi = \sum_{i=0}^k s^i d_i$ is a ring endomorphism of $R[s]$.

(b) If $\rho = \phi - 1$, then

$$\rho^n(ab) = \sum_{i=0}^n \sum_{j=0}^{n-i} \binom{n}{i} \binom{n-i}{j} \rho^{n-i}(a) \rho^{n-j}(b)$$

for $1 \leq n \leq k$ and all $a, b \in R[s]$.

(c) $\rho^n = \sum_{j=0}^{k-n} s^{n+j} d_{(n,n+j)}$ for $1 \leq n \leq k$.

Proof. (a) Clearly ϕ is additive. For $a, b \in R$,

$$\begin{aligned} \phi(a)\phi(b) &= \left(\sum_{i=0}^k s^i d_i(a) \right) \left(\sum_{j=0}^k s^j d_j(b) \right) \\ &= \sum_{i=0}^k s^i \left(\sum_{j=0}^i d_j(a) d_{i-j}(b) \right) \\ &= \sum_{i=0}^k s^i d_i(ab) = \phi(ab). \end{aligned}$$

Hence ϕ is a ring endomorphism.

(b) If $n = 1$,

$$\begin{aligned} &\sum_{i=0}^1 \sum_{j=0}^{1-i} \binom{1}{i} \binom{1-i}{j} \rho^{1-i}(a) \rho^{1-j}(b) \\ &= \rho(a)\rho(b) + \rho(a)b + a\rho(b) \\ &= (\phi(a) - a)(\phi(b) - b) + (\phi(a) - a)b + a(\phi(b) - b) \\ &= \phi(a)\phi(b) - ab = \phi(ab) - ab = \rho(ab). \end{aligned}$$

Suppose $n > 1$ and (b) is true for $i < n$. Then

$$\begin{aligned}
\rho^n(ab) &= \rho\rho^{n-1}(ab) \\
&= \rho \left(\sum_{i=0}^{n-1} \sum_{j=0}^{n-1-i} \binom{n-1}{i} \binom{n-1-i}{j} \rho^{n-1-i}(a) \rho^{n-1-j}(b) \right) \\
&= \sum_{i=0}^{n-1} \sum_{j=0}^{n-1-i} \binom{n-1}{i} \binom{n-1-i}{j} \\
&\quad \times [\rho^{n-i}(a) \rho^{n-j}(b) + \rho^{n-i}(a) \rho^{n-j-1}(b) + \rho^{n-i-1}(a) \rho^{n-j}(b)] \\
&= \sum_{j=0}^n \left[\binom{n-1}{j} + \binom{n-1}{j-1} \right] \rho^n(a) \rho^{n-j}(b) \\
&\quad + \sum_{i=1}^n \left[\binom{n-1}{i} + \binom{n-1}{i-1} \right] \rho^{n-i}(a) \rho^n(b) \\
&\quad + \sum_{i=1}^n \sum_{j=1}^{n-i} \left[\binom{n-1}{i} \binom{n-1-i}{j} + \binom{n-1}{i} \binom{n-1-i}{j-1} \right. \\
&\quad \quad \left. + \binom{n-1}{i-1} \binom{n-i}{j} \right] \rho^{n-i}(a) \rho^{n-j}(b) \\
&= \sum_{i=0}^n \sum_{j=0}^{n-i} \binom{n}{i} \binom{n-i}{j} \rho^{n-i}(a) \rho^{n-j}(b).
\end{aligned}$$

(c) Since $\rho = \phi - 1 = sd_1 + \cdots + s^k d_k$, the proof is similar to Lemma 7(a). \square

LEMMA 9. For integers $l, m > 0$,

$$\sum_{i=l}^{l+m} (-1)^{i+1} \binom{i}{i-l} \binom{l}{i-m} / i = 0.$$

Proof.

$$\begin{aligned}
&\sum_{i=l}^{l+m} (-1)^{i+1} \binom{i}{i-l} \binom{l}{i-m} / i \\
&= \sum_{j=0}^m (-1)^{j+l+1} \binom{j+l}{j} \binom{l}{j+l-m} / (j+l) \\
&= \sum_{j=0}^m (-1)^{j+l+1} (j+l-1)! / j!(m-j)!(j+l-m)!
\end{aligned}$$

which is the coefficient for x^m in the power series expansion of fg where

$$f(x) = (-1)^{l+1}(1+x)^{-l}$$

and

$$g(x) = (1+x)^l/l.$$

Since fg is constant, this coefficient is 0 for $m > 0$. \square

THEOREM 10. *Let $k \in \mathbf{N}$ and R a ring in which $k!$ is invertible.*

(a) *If $\delta_1, \dots, \delta_k$ are (commuting) derivations on R , then the rules*

$$d_0 = \text{identity map}$$

$$d_r = \sum_{i=1}^r \delta_{(i,r)}/i!$$

define a (commuting) S.h.d. d_0, d_1, \dots, d_k on R .

(b) *If d_0, d_1, \dots, d_k is a (commuting) S.h.d. on R then the rules*

$$\partial_r = \sum_{j=1}^r (-1)^{j+1} d_{(j,r)}/j$$

define (commuting) derivations $\partial_1, \dots, \partial_k$ on R .

(c) *If $\delta_1, \dots, \delta_k$ are derivations on R and d_0, d_1, \dots, d_k the S.h.d. constructed in (a), then the derivations $\partial_1, \dots, \partial_k$ constructed in (b) equal $\delta_1, \dots, \delta_k$.*

(d) *If d'_0, d'_1, \dots, d'_k is a S.h.d. on R and $\partial_1, \dots, \partial_k$ the derivations constructed in (b), then the S.h.d. d_0, d_1, \dots, d_k constructed in (a) is equal to d'_0, d'_1, \dots, d'_k .*

Proof. (a) For the ring endomorphism σ defined in Lemma 7,

$$\begin{aligned} \sigma &= \sum_{n=0}^k \partial^n/n! = 1 + \sum_{n=1}^k \sum_{j=0}^{k-n} s^{n+j} \delta_{(n,n+j)}/n! \\ &= 1 + \sum_{n=1}^k s^n \sum_{i=1}^n \delta_{(i,n)}/i!. \end{aligned}$$

For any ring homomorphism

$$d = \sum_{i=0}^k s^i d_i: R \rightarrow R[s]$$

where $d_i: R \rightarrow R$, the maps d_0, d_1, \dots, d_k form a S.h.d. on R . Hence (a) is proved.

(b) Using the map ρ defined in Lemma 8, set

$$\delta = \sum_{n=1}^k ((-1)^{n+1}/n)\rho^n.$$

We have, for any $a, b \in R[s]$,

$$\begin{aligned} \delta(ab) &= \sum_{n=1}^k ((-1)^{n+1}/n)\rho^n(ab) \\ &= \sum_{n=1}^k \sum_{i=0}^n \sum_{j=0}^{n-1} ((-1)^{n+1}/n) \binom{n}{i} \binom{n-i}{j} \rho^{n-i}(a)\rho^{n-j}(b). \end{aligned}$$

Note that $\rho^l(a)\rho^m(b) = 0$ whenever $l + m > k$. The coefficient of a particular term $\rho^l(a)\rho^m(b)$ above for $l \geq m > 0$ and $l + m \leq k$ is

$$\sum_{i=l}^{l+m} ((-1)^{i+1}/i) \binom{i}{i-l} \binom{l}{i-m}$$

which is 0 by Lemma 9. Similarly, for $m \geq l > 0$ and $l + m \leq k$, the coefficient of $\rho^l(a)\rho^m(b)$ in the above is 0. The coefficient of a term $\rho^l(a)b$ (or $a\rho^l(b)$) is $(-1)^{l+1}/l$. Hence, $\delta(ab) = \delta(a)b + a\delta(b)$ and δ is a derivation on $R[s]$.

If β is any derivation from R to $R[s]$ and

$$\beta = \sum_{i=0}^k s^i \beta_i$$

with each $\beta_i: R \rightarrow R$, then each β_i is a derivation on R . Now, by Lemma 8,

$$\begin{aligned} \delta &= \sum_{n=1}^k ((-1)^{n+1}/n)\rho^n = \sum_{n=1}^k \sum_{j=0}^{k-n} ((-1)^{n+1}/n)s^{n+j}d_{(n,n+j)} \\ &= \sum_{n=1}^k s^n \sum_{j=1}^n ((-1)^{j+1}/j)d_{(j,n)} = \sum_{n=1}^k s^n \partial_n, \end{aligned}$$

so each ∂_n is a derivation on R .

(c) If $\partial = s\delta_1 + \cdots + s^k\delta_k$ and $\rho = sd_1 + \cdots + s^k d_k$, we have

$$\sum_{n=0}^k \partial^n/n! = \rho + 1$$

as in the proof of (a) and

$$\sum_{n=1}^k ((-1)^{n+1}/n) \rho^n = \sum_{n=1}^k s^n \partial_n$$

as in the proof of (b). Hence,

$$\sum_{n=1}^k ((-1)^{n+1}/n) \left(\sum_{i=1}^k \partial^i / i! \right)^n = \sum_{n=1}^k s^n \partial_n.$$

But

$$\partial = \sum_{n=1}^k ((-1)^{n+1}/n) \left(\sum_{i=1}^k \partial^i / i! \right)^n$$

since the coefficient of a term ∂^j for $j \leq k$ is the coefficient of x^j in the power series expansion of $\log e^x$ and $\partial^j = 0$ for $j > k$.

So $\partial = s\delta_1 + \dots + s^k \delta_k = s\partial_1 + \dots + s^k \partial_k$ and $\partial_i = \delta_i$ for all i .

(d) For $\sigma = d'_0 + sd'_1 + \dots + s^k d'_k$,

$$\sigma = \sum_{n=0}^k \left(\sum_{i=1}^k ((-1)^{i+1}/i) (\sigma - 1)^i \right)^n / n!$$

since the coefficient of σ^j for $j \leq k$ is the coefficient of x^j in the power series expansion of $e^{\log x}$ and $\sigma^j = 0$ for $j > k$.

On the other hand,

$$\sum_{i=1}^k ((-1)^{i+1}/i) (\sigma - 1)^i = \sum_{n=1}^k s^n \partial_n$$

as in the proof of (b). If we denote this map by ∂ , then as in the proof of (a),

$$\sum_{n=0}^k \partial^n / n! = 1 + \sum_{n=1}^k s^n \sum_{i=1}^n \partial_{(i,n)} / i! = 1 + sd_1 + \dots + s^k d_k.$$

Since the left hand side equals σ (as shown above) we obtain $d'_i = d_i$ for all i . \square

THEOREM 11. *Let $k \in \mathbb{N}$, and let R be a ring in which $k!$ is invertible. Let d_0, d_1, \dots, d_k be a commuting S.h.d. on R and $\delta_1, \dots, \delta_k$ the corresponding commuting derivations as in Theorem 10(b). Then the rings*

$$S = R[D_1, \dots, D_k; d_0, d_1, \dots, d_k] \quad \text{and} \quad T = R[\Theta_1, \dots, \Theta_k; \delta_1, \dots, \delta_k]$$

are isomorphic. Moreover, there is an isomorphism that is the identity on R and preserves leading coefficients.

Proof. Define $\bar{\Theta}_1, \dots, \bar{\Theta}_k$ by

$$\bar{\Theta}_n = \sum_{i=1}^n ((-1)^{i+1}/i) D_{(i,n)}.$$

Note that $\bar{\Theta}_i \bar{\Theta}_j = \bar{\Theta}_j \bar{\Theta}_i$ for all i, j . Let $R[t]$ be a polynomial ring over R and set

$$R[s] = R[t]/(t^{k+1}).$$

Set

$$D = sD_1 + \dots + s^k D_k \quad \text{and} \quad \rho = sd_1 + \dots + s^k d_k.$$

As in the proof of Lemma 7(a)

$$D^n = \sum_{j=0}^{k-n} s^{n+j} D_{(n,n+j)}$$

for $1 \leq n \leq k$. Hence

$$\begin{aligned} \sum_{n=1}^k ((-1)^{n+1}/n) D^n &= \sum_{n=1}^k \sum_{j=0}^{k-n} s^{n+j} ((-1)^{n+1}/n) D_{(n,n+j)} \\ &= \sum_{n=1}^k s^n \sum_{j=1}^n ((-1)^{j+1}/j) D_{(j,n)} = \sum_{n=1}^k s^n \bar{\Theta}_n. \end{aligned}$$

Now for $1 \leq n \leq k$ and $a \in R[s]$ we show that

$$D^n a = \sum_{i=0}^n \sum_{j=0}^{n-i} \binom{n}{i} \binom{n-i}{j} \rho^{n-i}(a) D^{n-j}.$$

To see this note that

$$Da = \rho(a)D + aD + \rho(a),$$

which is the above with $n = 1$. Then by induction,

$$\begin{aligned}
 D^n a &= D^{n-1} D a = D^{n-1} (\rho(a) D + a D + \rho(a)) \\
 &= \sum_{i=0}^{n-1} \sum_{j=0}^{n-1-i} \binom{n-1}{i} \binom{n-1-i}{j} \\
 &\quad \times (\rho^{n-i}(a) D + \rho^{n-1-i}(a) D + \rho^{n-i}(a)) D^{n-1-j} \\
 &= \sum_{i=1}^n \sum_{j=1}^{n-i} \left[\binom{n-1}{i} \binom{n-1-i}{j} + \binom{n-1}{i} \binom{n-1-i}{j-1} \right. \\
 &\quad \left. + \binom{n-1}{i-1} \binom{n-i}{j} \right] \rho^{n-i}(a) D^{n-j} \\
 &\quad + \sum_{j=0}^n \left[\binom{n-1}{j} + \binom{n-1}{j-1} \right] \rho^n(a) D^{n-j} \\
 &\quad + \sum_{i=1}^n \left[\binom{n-1}{i} + \binom{n-1}{i-1} \right] \rho^{n-i}(a) D^n \\
 &= \sum_{i=0}^n \sum_{j=0}^{n-i} \binom{n}{i} \binom{n-i}{j} \rho^{n-i}(a) D^{n-j}.
 \end{aligned}$$

Set

$$\Theta = \sum_{n=1}^k s^n \bar{\Theta}_n = \sum_{n=1}^k ((-1)^{n+1}/n) D^n.$$

Then for $a \in R$ we get

$$\begin{aligned}
 \Theta a &= \sum_{n=1}^k \sum_{i=0}^n \sum_{j=0}^{n-i} ((-1)^{n+1}/n) \binom{n}{i} \binom{n-i}{j} \rho^{n-i}(a) D^{n-j} \\
 &= a \Theta + \sum_{n=1}^k s^n \delta_n(a)
 \end{aligned}$$

as in the proof of Theorem 10(b). Hence,

$$\bar{\Theta}_n a = a \bar{\Theta}_n + \delta_n(a),$$

for $1 \leq n \leq k$ and all $a \in R$.

Consequently, there exists a unique ring homomorphism $\phi: T \rightarrow S$ such that ϕ is the identity on R and $\phi(\Theta_n) = \bar{\Theta}_n$ for all n .

If

$$0 \neq \sum r_L \Theta^L \in T$$

with leading coefficient r_J , then

$$\phi\left(\sum r_L \Theta^L\right) \in S$$

also has leading coefficient r_J . To see this, note that the highest weighted order monomial of $\bar{\Theta}_n$ is D_n . Therefore

$$\sum r_L \Theta^L \neq 0$$

implies

$$\phi\left(\sum r_L \Theta^L\right) \neq 0$$

and hence ϕ is injective.

As in the proof of Theorem 10(d), we have

$$\sum_{n=0}^k \left(\sum_{i=1}^k ((-1)^{i+1}/i) D^i \right)^n / n! = D + 1.$$

Therefore

$$\begin{aligned} D + 1 &= \sum_{n=0}^k \left(\sum_{i=1}^k s^i \bar{\Theta}_i \right)^n / n! \\ &= 1 + \sum_{n=1}^k \left(\sum_{j=0}^{k-n} s^{n+j} \bar{\Theta}_{(n,n+j)} \right) / n! = 1 + \sum_{n=1}^k s^n \sum_{i=1}^n \bar{\Theta}_{(i,n)}/i! \end{aligned}$$

and it follows that

$$D_n = \sum_{i=1}^n \bar{\Theta}_{(i,n)}/i! = \phi \left(\sum_{i=1}^n \Theta_{(i,n)}/i! \right)$$

for $1 \leq n \leq k$. Hence ϕ is also surjective and therefore an isomorphism. \square

We end this section with two lemmas that allow us to extend a S.h.d.

LEMMA 12. *Let d_0, d_1, \dots, d_k be a S.h.d. on a ring R and let C be a right denominator set in R . Then d_0, d_1, \dots, d_k can be uniquely extended to a S.h.d. on $R[C^{-1}]$.*

Proof. Let $R[t]$ be a polynomial ring over R and set

$$R[s] = R[t]/(t^{k+1}).$$

Note that $d = d_0 + sd_1 + \cdots + s^k d_k$ defines a ring homomorphism $d: R \rightarrow R[s]$. Let $\phi: R \rightarrow R[C^{-1}]$ and $\phi': R[s] \rightarrow R[C^{-1}][s]$ be the natural maps. For any $c \in C$,

$$\begin{aligned}\phi'd(c) &= \phi'(d_0(c) + sd_1(c) + \cdots + s^k d_k(c)) \\ &= \phi'(c) + s\phi'(d_1(c)) + \cdots + s^k \phi'(d_k(c)).\end{aligned}$$

Since $\phi'(c)^{-1}$ exists in $R[C^{-1}][s]$, we can inductively solve for coefficients of a right or left inverse for $\phi'(d(c))$. Therefore, there exists $\phi'(d(c))^{-1}$ in $R[C^{-1}][s]$ for all $c \in C$.

Hence, there exists a unique ring homomorphism

$$d': R[C^{-1}] \rightarrow R[C^{-1}][s]$$

such that $d'\phi = \phi'd$. If we write $d' = d'_0 + sd'_1 + \cdots + s^k d'_k$ with each $d'_i: R[C^{-1}] \rightarrow R[C^{-1}]$, d'_0, d'_1, \dots, d'_k is a S.h.d. on $R[C^{-1}]$ which uniquely extends d_0, d_1, \dots, d_k . \square

LEMMA 13. *Let d_0, d_1, \dots, d_k be a S.h.d. on a ring R . Given f_1, \dots, f_k central elements of the polynomial ring $R[x]$, there is a unique extension of d_0, d_1, \dots, d_k to a S.h.d. on $R[x]$ such that $d_i(x) = f_i$ for $1 \leq i \leq k$.*

Proof. Let $R[t]$ be a polynomial ring over R and set

$$R[s] = R[t]/(t^{k+1}).$$

Then we have a ring homomorphism

$$d = d_0 + sd_1 + \cdots + s^k d_k: R \rightarrow R[s].$$

This extends uniquely to a ring homomorphism

$$d': R[x] \rightarrow R[x][s]$$

such that

$$d'(x) = x + sf_1 + \cdots + s^k f_k,$$

because $x + sf_1 + \cdots + s^k f_k$ is central in $R[x][s]$. If

$$d' = d'_0 + sd'_1 + \cdots + s^k d'_k$$

with each $d'_i: R[x] \rightarrow R[x]$, then d'_0, d'_1, \dots, d'_k is a S.h.d. on $R[x]$ extending d_0, d_1, \dots, d_k . The uniqueness is clear. \square

3. We develop some necessary and sufficient conditions for simplicity of differential operator rings coming from a S.h.d. The proof of the next lemma is similar to that of Lemma 1.

LEMMA 14. Let d_0, d_1, \dots, d_k be a commuting S.h.d. on a ring R and let F be the subring of central (d_1, \dots, d_k) -constants of R . If the ring

$$S = R[D_1, \dots, D_k; d_0, d_1, \dots, d_k]$$

is simple then the center of S is F and R is (d_0, \dots, d_k) -simple. \square

If d_0, d_1, \dots, d_k is a commuting S.h.d. on a ring R , then R can be made into a left module over the ring

$$S = R[D_1, \dots, D_k; d_0, d_1, \dots, d_k]$$

using an S -module multiplication $*$ by which $D_i * r = d_i(r)$ and $s * r = sr$ for all i and all $s, r \in R$. With this left module structure, right multiplication in R by a (d_1, \dots, d_k) -constant is a left S -module endomorphism of R .

THEOREM 15. Let R be a \mathbf{Q} -algebra, d_0, d_1, \dots, d_k a commuting S.h.d. on R and F the subring of central (d_1, \dots, d_k) -constants of R . Then the ring

$$S = R[D_1, \dots, D_k; d_0, d_1, \dots, d_k]$$

is simple if and only if R is (d_0, d_1, \dots, d_k) -simple and no nontrivial F -linear combination of

$$\{d_k^{i(k)} \dots d_1^{i(1)} \mid i(j) \geq 0 \text{ and } 0 < i(1) + 2i(2) + \dots + ki(k) \leq k\}$$

is an inner derivation by a (d_1, \dots, d_k) -constant. In this case the center S is F .

Proof. If S is simple then the center of S is F and R is (d_0, \dots, d_k) -simple by Lemma 14. In a manner similar to that of Theorem 2, no nontrivial F -linear combination of

$$\{d_k^{i(k)} \dots d_1^{i(1)} \mid i(j) \geq 0 \text{ and } 0 < i(1) + 2i(2) + \dots + ki(k) \leq k\}$$

is an inner derivation by a (d_1, \dots, d_k) -constant.

Conversely, using the construction of Theorem 10 and Theorem 11, we have

$$S \cong T = R[\Theta_1, \dots, \Theta_k; \delta_1, \dots, \delta_k]$$

for commuting derivations $\delta_1, \dots, \delta_k$ related to d_0, d_1, \dots, d_k as in Theorem 10. Then R being (d_0, d_1, \dots, d_k) -simple implies R is $(\delta_1, \dots, \delta_k)$ -simple. No nontrivial F -linear combinations of

$$\{d_k^{i(k)} \dots d_1^{i(1)} \mid i(j) \geq 0 \text{ and } 0 < i(1) + 2i(2) + \dots + ki(k) \leq k\}$$

being inner by a d_1, \dots, d_k -constant implies no nontrivial F -linear combination of $\delta_1, \dots, \delta_k$ is inner by a $(\delta_1, \dots, \delta_k)$ -constant. Since F is also the subring of central $(\delta_1, \dots, \delta_k)$ -constants of R , by Theorem 2 the ring S is simple. \square

LEMMA 16. Let d_0, d_1, \dots, d_k be a commuting S.h.d. on a ring R and I a nonzero ideal of the ring

$$S = R[D_1, \dots, D_k; d_0, d_1, \dots, d_k].$$

If $q = \sum r_L D^L \in I$, then $d_i(q) = \sum d_i(r_L) D^L \in I$ for all i .

Proof. If $i = 1$ then $\sum d_1(r_L) D^L = D_1 q - q D_1 \in I$. If $i > 1$, suppose the lemma is true for all $j < i$. Then

$$\sum d_i(r_L) D^L = D_i q - \sum_{j=0}^{i-1} d_j(q) D_{i-j} \in I. \quad \square$$

THEOREM 17. Let d_0, d_1, \dots, d_k be a commuting S.h.d. on a ring R and F the subring of central (d_1, \dots, d_k) -constants of R . Then the ring

$$S = R[D_1, \dots, D_k; d_0, d_1, \dots, d_k]$$

is simple if and only if R is (d_0, d_1, \dots, d_k) -simple and no nontrivial F -linear combination of

$$\{d_k^{n(k)} \dots d_1^{n(1)} \mid n(i) \geq 0 \text{ and } n(k) + \dots + n(1) > 0\}$$

is an inner derivation by a (d_1, \dots, d_k) -constant. In this case the center of S is F .

Proof. If S is simple, the proof is similar to that of Theorem 15. Conversely, assume that R is (d_0, d_1, \dots, d_k) -simple and no nontrivial F -linear combination of

$$\{d_k^{n(k)} \dots d_1^{n(1)} \mid n(i) \geq 0 \text{ and } n(k) + \dots + n(1) > 0\}$$

is inner by a (d_1, \dots, d_k) -constant.

If A is a nonzero ideal of S , let J be the minimal weighted order of nonzero elements of A . Using Lemma 16, the set of 0 together with those $r \in R$ that are leading coefficient of elements of A of weighted order J forms a (d_0, d_1, \dots, d_k) -ideal of R . Hence, we have $q \in A$ of weighted order J with leading coefficient 1. Using Lemma 16 again, all other coefficients of q are (d_1, \dots, d_k) -constants. Also, $qr - rq = 0$ for any $r \in R$.

If $q = \sum r_L D^L$, let I denote the highest weighted order such that $r_I \notin F$ (or if all $r_L \in F$ let $I = e(0)$). For all $r \in R$, the coefficient of D^I in $qr - rq$ is 0. If $|J| > 0$, then from the vanishing of these coefficients we get some F -linear combination of

$$\{d_k^{n(k)} \dots d_1^{n(1)} \mid n(i) \geq 0 \text{ and } n(k) + \dots + n(1) > 0\}$$

being an inner derivation by a (d_1, \dots, d_k) -constant. Therefore $|J| = 0$, whence $q = 1$ and so $A = S$. Thus S is simple. \square

As an example to satisfy the hypotheses of Theorem 17, let K be a field of characteristic 2 and let x_1, x_2, \dots be independent indeterminates over K . By Lemma 13, there exists a unique K -linear S.h.d. d_0, d_1, d_2 on $K[\{x_i\}]$ such that

$$d_1(x_i) = x_{i+1} \quad \text{and} \quad d_2(x_i) = 0$$

for all i . Then Lemma 12 implies that d_0, d_1, d_2 extends uniquely to a S.h.d. on $R = K(\{x_i\})$.

Note that $d_1(x_i^2) = 0$ and $d_2(x_i^2) = x_{i+1}^2$ for all i . As R is a field, we only have to show that the set

$$\{d_1^{n(1)} d_2^{n(2)} \mid n(i) \geq 0 \text{ and } n(1) + n(2) > 0\}$$

is linearly independent over the subring of central (d_1, d_2) -constants of R . Actually, we show this set is linearly independent over R .

If we have some $\alpha_{ij} \in R$ such that

$$\sum_{i,j} \alpha_{ij} d_1^i d_2^j = 0,$$

find l such that x_{l+h} is not involved in any of the α_{ij} for $h \geq 0$. Then

$$\begin{aligned} 0 &= \sum_{i,j} \alpha_{ij} d_1^i d_2^j (x_l^3) = \sum_{i,j} \alpha_{ij} d_1^i d_2^j (x_l x_{l+j}^2) \\ &= \sum_{i,j} \alpha_{ij} d_1^i (x_l x_{l+j}^2) = \sum_{i,j} \alpha_{ij} x_{l+i} x_{l+j}^2. \end{aligned}$$

Therefore each $\alpha_{ij} = 0$ and the set

$$\{d_1^{n(1)} d_2^{n(2)} \mid n(i) \geq 0 \text{ and } n(1) + n(2) > 0\}$$

is linearly independent over R . Thus the ring $R[D_1, D_2; d_0, d_1, d_2]$ is simple.

REFERENCES

- [1] S. Amitsur, *Derivations in simple rings*, Proc. London Math. Soc., (3) 7 (1957), 87–112.
- [2] J. Cozzens and C. Faith, *Simple Noetherian Rings*, Cambridge University Press 1975.
- [3] K. R. Goodearl and R. Warfield, *Primitivity in differential operator rings*, Math. Z., **180** (1982), 503–523.
- [4] H. Hasse and F. K. Schmidt, *Noch eine Begründung der Theorie der höheren Differentialquotienten (Zusatz bei der Korrektur)*, J. Reine Angew. Math., **177** (1937), 223–239.
- [5] G. Hauger, *Einfache Derivationspolynomringe*, Arch. Math., **29** (1977), 491–496.
- [6] N. Heerema, *Derivations and embeddings of a field in its power series ring*, PAMS, **11** (1960), 188–194.
- [7] ———, *Derivations and Embeddings of a Field in its Power Series Ring II*, Mich. Math. J., **8** (1961), 129–134.
- [8] D. Jordan, *Ore Extensions and Jacobson Rings*, Ph.D. Thesis, Leeds, 1975.
- [9] G. Renault, *Algèbre Non Commutative*, Gauthier-Villars 1975.
- [10] O. Teichmüller, *Differentialrechnung bei Charakteristik p* , J. Reine Angew. Math., **175** (1937), 89–99.
- [11] M. Voskoglou, *Simple skew polynomial rings*, Publ. Inst. Math. (Beograd) (N.S.), **37(51)** 1985, 37–41.

Received September 4, 1986.

THE UNIVERSITY OF UTAH
SALT LAKE CITY, UT 84112

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112

R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721

RAMESH A. GANGOLLI
University of Washington
Seattle, WA 98195

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720

ROBION KIRBY
University of California
Berkeley, CA 94720

C. C. MOORE
University of California
Berkeley, CA 94720

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906-1982)

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

Scott W. Brown , Full analytic subspaces for contractions with rich spectrum . . .	1
Robert Main Burton, Jr. and Tae-Sung Kim , An invariance principle for associated random fields	11
Ana M. Viola-Prioli and Jorge Viola-Prioli , Rings whose kernel functors are linearly ordered	21
David E. Handelman , Representing polynomials by positive linear functions on compact convex polyhedra	35
Patrick Keef , On the Tor functor and some classes of abelian groups	63
Dennis R. Malm , Simplicity of partial and Schmidt differential operator rings	85
José M. Montesinos and Carmen Safont , On the Birman invariants of Heegaard splittings	113
N. P. Mukherjee and Prabir Bhattacharya , The normal index of a finite group	143
Mario Petrich and Norman R. Reilly , Semigroups generated by certain operators on varieties of completely regular semigroups	151
Robert Tijdeman and Lian Xiang Wang , Sums of products of powers of given prime numbers	177
Joel Larry Weiner , First integrals for a direction field on a simply connected plane domain	195