

# Simultaneous Equations as a Tool in the Spectrophotometric Analysis of Two Non-interacting Substances in a Binary Mixture: Senior Undergraduate Physical and Physical-Organic Chemistry Laboratory Experiment

R. Sanjeev<sup>1</sup>, V. Jagannadham<sup>2,\*</sup>, R. Ravi<sup>2</sup>, R. Veda Vrath<sup>3</sup>, Arijit Das<sup>4</sup>

<sup>1</sup>Departments of Chemistry, Avanthi Degree and PG College, Hyderabad, 500029, India

<sup>2</sup>Departments of Chemistry, Osmania University, Hyderabad, 500007, India

<sup>3</sup>Departments of Chemistry, L N Gupta Evening College, Hyderabad, 500002, India

<sup>4</sup>Departments of Chemistry, Government Degree College, Dharmanagar, Tripura, India

**Abstract** A simple and lucid spectrophotometric way of analyzing a mixture of two compounds is explained in this article for senior undergraduate level students. The essential feature of this method is that there should be at least 30-40 nm difference in the  $\lambda_{\text{max}}$  of the two compounds so that there will not be any overlap of the two individual spectra of the two compounds when recorded together in a mixture.

**Keywords** Analysis of Binary Mixtures, Spectrophotometric Analysis, Simultaneous Equations

## 1. Introduction

Interest in the analysis of drugs and some organic compounds in binary mixtures by spectrophotometric method have become seriously interesting among analytical chemists and pharmacists. As a sample some references are given at the end of the article[1-8]. In all these articles the aim of the authors was much focused on the determination of the concentration/purity of the two components in the mixture. Therefore we have tried to make the analysis much lucid and simple for a senior undergraduate laboratory experiment using the method of solving simultaneous equations[9].

## 2. Experimental

All the chemicals were of analytical grade.  $\text{KMnO}_4$  and  $\text{K}_2\text{Cr}_2\text{O}_7$  are from Aldrich (Bangalore, India). Distilled water used was from all glass still. The UV-visible spectra are recorded on UVIKON-323 (Italy) spectrophotometer. All the analytical data were stored in a personal computer and

analyzed using KaleidaGraph software (Synergy Software, Reading, PA, USA).

## 3. Theory

### 3.1. Discussion

According to Beer-Lambert law[10]

$$\text{Absorbance} = \epsilon t C \quad (1)$$

where ' $\epsilon$ ' is the molar absorptivity of any substance, ' $t$ ' is the path length in cm of the cuvette through which light passes and ' $C$ ' is the concentration of the substance taken. The conditions that need to be fulfilled in order for Beer-Lambert law to be valid are also to be valid here[10]. The conditions are: The absorption spectra of the two components must be independent of each other i.e. non-interfering, the solutions must be homogeneous, the incident radiation must be monochromatic, and the radiation must not do any photochemistry or photophysics with the substances. For all practical purposes to make the situation simple let a cuvette of 1 cm path length be chosen so that equation (1) becomes

$$\text{Absorbance} = \epsilon C \quad (2)$$

Let a mixture contain two compounds say **A** and **B**. Let the compounds **A** and **B** have their  $\lambda_{\text{max}}$  at  $\lambda_1$  and  $\lambda_2$ . Figure 1 shows the UV-Vis absorption spectrum of compound **A** with maximum absorbance at its  $\lambda_{\text{max}}$  i.e.  $\lambda_1$  and a small absorbance at  $\lambda_2$  which is the  $\lambda_{\text{max}}$  of compound **B**.

\* Corresponding author:

jagannadham1950@yahoo.com (V. Jagannadham)

Published online at <http://journal.sapub.org/jlce>

Copyright © 2013 Scientific & Academic Publishing. All Rights Reserved




Figure 1.

Let  $Abs_1$  and  $Abs_2$  be the absorbencies of compound A at  $\lambda_1$  and  $\lambda_2$  respectively. Therefore according to equation 2, the absorbencies at  $\lambda_1$  and  $\lambda_2$  are:

$$Abs_1 = \epsilon_A^1 C_A \quad \text{at } \lambda_1 \quad (3)$$

$$Abs_2 = \epsilon_A^2 C_A \quad \text{at } \lambda_2 \quad (4)$$

Similarly Figure 2 shows the UV-Vis absorption spectrum of compound B with maximum absorbance at its  $\lambda_{max}$  i.e.  $\lambda_2$  and a small absorbance at  $\lambda_1$  which is the  $\lambda_{max}$  of compound A.




Figure 2.

Let  $Abs_3$  and  $Abs_4$  be the absorbencies of compound B at  $\lambda_1$  and  $\lambda_2$  respectively. Therefore again according to equation 2, the absorbencies at  $\lambda_1$  and  $\lambda_2$  are:

$$Abs_3 = \epsilon_B^1 C_B \quad \text{at } \lambda_1 \quad (5)$$

$$Abs_4 = \epsilon_B^2 C_B \quad \text{at } \lambda_2 \quad (6)$$

Figure 3 shows the UV-Vis absorption spectrum of the mixture of compounds A and B.




Figure 3.

Let  $\text{Abs}_5$  be the total absorbance of the mixture of the compounds **A** and **B** at  $\lambda_1$  and  $\text{Abs}_6$  is the total absorbance of the mixture of the compounds **A** and **B** at  $\lambda_2$  respectively. Therefore again according to equation 2, the total absorbencies of the two components in the mixture at  $\lambda_1$  and  $\lambda_2$  are:

$$\text{Abs}_5 = \epsilon_A^1 C_A + \epsilon_B^1 C_B \quad \text{at } \lambda_1 \quad (7)$$

$$\text{Abs}_6 = \epsilon_A^2 C_A + \epsilon_B^2 C_B \quad \text{at } \lambda_2 \quad (8)$$

$\epsilon_A^1$ ,  $\epsilon_A^2$ ,  $\epsilon_B^1$  and  $\epsilon_B^2$ , the molar absorptivities are independently experimentally determinable quantities of compounds **A** and **B** using equation 2 from the concentration dependencies of absorbivity, the so called Beer-Lambert law.  $\text{Abs}_5$  and  $\text{Abs}_6$  are also experimentally determinable quantities from figure 3. Now equations 7 and 8 are two simultaneous equations with two unknowns in the mixture i.e.  $C_A$  and  $C_B$ . Using simple algebra one can eliminate one unknown to calculate the other. So first let  $C_B$  be eliminated. To do this let equation 7 be multiplied by  $\epsilon_B^2$  and equation 8 by  $\epsilon_B^1$ . Therefore we get

$$\epsilon_B^2 \cdot \text{Abs}_5 = \epsilon_B^2 \epsilon_A^1 C_A + \epsilon_B^2 \epsilon_B^1 C_B \quad (9)$$

$$\epsilon_B^1 \cdot \text{Abs}_6 = \epsilon_B^1 \epsilon_A^2 C_A + \epsilon_B^1 \epsilon_B^2 C_B \quad (10)$$

Subtraction of equation 10 from equation 9, the quantities  $\epsilon_B^2 \epsilon_B^1 C_B$  and  $\epsilon_B^1 \epsilon_B^2 C_B$  would be canceled and we get

$$\epsilon_B^2 \cdot \text{Abs}_5 - \epsilon_B^1 \cdot \text{Abs}_6 = C_A [\epsilon_B^2 \epsilon_A^1 - \epsilon_B^1 \epsilon_A^2] \quad (11)$$

$$\text{Therefore } C_A = \frac{\epsilon_B^2 \cdot \text{Abs}_5 - \epsilon_B^1 \cdot \text{Abs}_6}{[\epsilon_B^2 \epsilon_A^1 - \epsilon_B^1 \epsilon_A^2]} \quad (12)$$

Knowing all the quantities on right hand side of equation 12,  $C_A$ , the concentration of the compound **A** in the mixture could be obtained. Substituting  $C_A$  in either the equation 9 or 10 and knowing other quantities,  $C_B$ , the concentration of the compound **B** in the mixture could be calculated.

## 4. Practice

### 4.1. UV-Vis Spectra of $\text{KMnO}_4$ , $\text{K}_2\text{Cr}_2\text{O}_7$ and the Mixture of the Two

As a specific example for the practice in the laboratory we have performed the experiments using  $\text{KMnO}_4$ ,  $\text{K}_2\text{Cr}_2\text{O}_7$  and the mixture of the two. First we have recorded the spectra of  $\text{KMnO}_4$  ( $4.0 \times 10^{-4}$  M),  $\text{K}_2\text{Cr}_2\text{O}_7$  ( $4.8 \times 10^{-4}$  M) and the mixture (containing  $2.0 \times 10^{-4}$  M of  $\text{KMnO}_4$  and  $2.40 \times 10^{-4}$  M of  $\text{K}_2\text{Cr}_2\text{O}_7$ ) so that it is necessary to identify the wavelengths where the analysis is to be carried out. Figure 4 shows the spectra of these solutions in the range 250 nm to 650 nm. But unfortunately there was no absorbance at 525 nm for  $\text{K}_2\text{Cr}_2\text{O}_7$  which is the  $\lambda_{\text{max}}$  of  $\text{KMnO}_4$ . So it was thought worthwhile to choose another range of wavelength region for the analysis. Figure 5 is the reproduction of part of the figure 4 which shows that there are appreciable absorbivities for the two samples of solutions and the mixture.




Figure 4. UV-VIS spectra of  $\text{KMnO}_4$ ,  $\text{K}_2\text{Cr}_2\text{O}_7$  and the mixture




Figure 5. UV-Vis spectra of  $\text{KMnO}_4$  and  $\text{K}_2\text{Cr}_2\text{O}_7$  and the mixture

From figure 5 two wavelengths were identified  $\lambda_{\text{obs1}}$  (372 nm) and  $\lambda_{\text{obs2}}$  (290 nm) at which all the individual samples and the mixture have absorbencies [11]. First it is necessary to determine the molar absorbance of  $\text{KMnO}_4$  and  $\text{K}_2\text{Cr}_2\text{O}_7$ . These were determined at 372 nm and 290 nm for both the samples. In recording the spectra the 1 cm path length optical quartz cuvettes were used. Hence to determine the molar absorbencies ( $\epsilon$ ) again equation 2 is used. Figures 6 and 7 show the Beer-Lambert law plots of  $\text{KMnO}_4$  and  $\text{K}_2\text{Cr}_2\text{O}_7$ . From the slopes of these plots molar absorbencies ( $\epsilon$ ) of  $\text{KMnO}_4$  and  $\text{K}_2\text{Cr}_2\text{O}_7$  were obtained. They were  $805 \text{ mol}^{-1} \text{ cm}^{-1}$  ( $\epsilon_{\text{KMnO}_4}^1$ ) at 372 nm,  $1030 \text{ mol}^{-1} \text{ cm}^{-1}$  ( $\epsilon_{\text{KMnO}_4}^2$ ) at 290 nm for  $\text{KMnO}_4$ . And for  $\text{K}_2\text{Cr}_2\text{O}_7$  they were  $1652 \text{ mol}^{-1} \text{ cm}^{-1}$  ( $\epsilon_{\text{K}_2\text{Cr}_2\text{O}_7}^1$ ) at 372 nm and  $853 \text{ mol}^{-1} \text{ cm}^{-1}$  ( $\epsilon_{\text{K}_2\text{Cr}_2\text{O}_7}^2$ ) at 290 nm.

From figure 5, for the mixture  $\text{abs}_{\text{total}}$  at 372 nm is 0.732. Therefore

$$0.732 = \epsilon_{\text{KMnO}_4}^1 \cdot C_{\text{KMnO}_4} + \epsilon_{\text{K}_2\text{Cr}_2\text{O}_7}^1 \cdot C_{\text{K}_2\text{Cr}_2\text{O}_7} \quad (13)$$

$$0.732 = 805 \times C_{\text{KMnO}_4} + 1652 \times C_{\text{K}_2\text{Cr}_2\text{O}_7} \quad (14)$$

And at 290 nm

$$0.588 = \epsilon_{\text{KMnO}_4}^2 \cdot C_{\text{KMnO}_4} + \epsilon_{\text{K}_2\text{Cr}_2\text{O}_7}^2 \cdot C_{\text{K}_2\text{Cr}_2\text{O}_7} \quad (15)$$

$$0.588 = 1030 \times C_{\text{KMnO}_4} + 853 \times C_{\text{K}_2\text{Cr}_2\text{O}_7} \quad (16)$$

Therefore equations 14 and 16 are two simultaneous equations with two unknowns i.e.  $C_{\text{KMnO}_4}$  and  $C_{\text{K}_2\text{Cr}_2\text{O}_7}$ . First to eliminate  $C_{\text{K}_2\text{Cr}_2\text{O}_7}$  equation 14 is multiplied by 853 and equation 16 is multiplied by 1652. Then one would get

$$853 \times 0.732 = 853 \times 805 \times C_{\text{KMnO}_4} + 853 \times 1652 \times C_{\text{K}_2\text{Cr}_2\text{O}_7} \quad (17)$$

And

$$1652 \times 0.588 = 1652 \times 1030 \times C_{\text{KMnO}_4} + 1652 \times 853 \times C_{\text{K}_2\text{Cr}_2\text{O}_7} \quad (18)$$

Subtracting equation 17 from equation 18, the second terms on right hand side of the both equations would be cancelled out and rearranging for the concentration of permanganate, we get  $C_{\text{KMnO}_4} = 3.38 \times 10^{-4} \text{ M}$ . The concentration of permanganate actually taken in the mixture is only  $2.0 \times 10^{-4} \text{ M}$ . The deviation in this case most probably be due to the wavelength used is not the  $\lambda_{\text{max}}$  of  $\text{KMnO}_4$ . And substituting this value either in Equation 13 or 14 we get  $C_{\text{K}_2\text{Cr}_2\text{O}_7} = 2.8 \times 10^{-4} \text{ M}$ . And the concentration of dichromate used in the mixture is  $2.4 \times 10^{-4} \text{ M}$  which is in good agreement with analysis of the mixture.




Figure 6. Determination of molar absorptivity of  $\text{KMnO}_4$




Figure 7. Determination of molar absorptivity of  $\text{K}_2\text{Cr}_2\text{O}_7$

## REFERENCES

- [1] Spectrophotometric analysis of binary mixtures of urinary steroids, Joel Bitman, Jean-Pierre Rosselet, Alvira De M. Reddy, and Seymour Lieberman, *J. Biol. Chem.*, 1957, 225, 39-52
- [2] Spectrophotometric determination of binary mixtures of pseudoephedrine with some histamine H<sub>1</sub>-receptor antagonists using derivative ratio spectrum method, Hoda Mahgoub, Azza A Gazy, Fawzi A El-Yazbi, Mahmoud A El-Sayed, Rasha M Youssef, *Journal of Pharmaceutical and Biomedical Analysis*, 2003, 31, Pages 801–809
- [3] Analysis of binary mixtures of losartan potassium and hydrochlorothiazide by using high performance liquid chromatography, ratio derivative spectrophotometric and compensation technique, Nevin Erk, *Journal of Pharmaceutical and Biomedical Analysis*, 2001, 24, Pages 603–611
- [4] Spectrophotometric determination of sulphamethoxazole and trimethoprim (co-trimoxazole) in binary mixtures and in tablets, S Balyejusaa, RO Adomeb, and D Musokec, *African Health Science*, 2002, page 56-62
- [5] A Rapid Spectrophotometric Method to Resolve a Binary Mixture of Food Colorants Riboflavine and Sunset Yellow, Mahmure UST, *Turk J Chem*, (2004), 28, 325-333.
- [6] Simultaneous spectrophotometric determination of amlodipine besylate and atorvastatin calcium in binary mixture, R Sahu, and Vandana B Patel, *Indian J. Pharma. Chem.*, 2007, 69, Page 110-111
- [7] Spectrophotometric determination of binary mixtures of prednisolone with some antibiotics Abd El-Maboud I. Mohamed, Hesham Salem and Eman Maher, *Thai J. Pharm. Sci.* 2006, 30, 63-81
- [8] Simultaneous spectrophotometric determination of diclofenac potassium and methocarbamol in binary mixture using chemometric techniques and artificial neural networks, Ehab F. Elkady, *Drug Testing and Analysis*, 2011, 3, pages 228-233
- [9] [http://www.teacherschoice.com.au/maths\\_library/algebra/alg\\_10.htm](http://www.teacherschoice.com.au/maths_library/algebra/alg_10.htm)
- [10] J. D. J. Ingle and S. R. Crouch, *Spectrochemical Analysis*, Prentice Hall, New Jersey (1988)
- [11] It is not necessary to perform the analysis at the max of the samples. This is possible only when first sample has appreciable absorbance at the max of the second sample and vice-versa. If the analysis is carried out at other convenient wavelengths then one can call those wavelengths as obs.