

Simultaneous explanation of $R(D^{(*)})$ and $b \rightarrow s\mu^+\mu^-$: the last scalar leptoquarks standing

Andreas Crivellin,^a Dario Müller^{a,b} and Toshihiko Ota^c

^aPaul Scherrer Institut,
CH-5232 Villigen PSI, Switzerland

^bPhysik-Institut, Universität Zürich,
Winterthurerstrasse 190, CH-8057 Zürich, Switzerland

^cDepartment of Physics, Yachay Tech,
Hacienda San José s/n y Proyecto Yachay, 100119 San Miguel de Urcuquí, Ecuador

E-mail: andreas.crivellin@cern.ch, dario.mueller@psi.ch,
tota@yachaytech.edu.ec

ABSTRACT: Over the past years, experiments accumulated intriguing hints for new physics (NP) in flavor observables, namely in the anomalous magnetic moment of the muon (a_μ), in $R(D^{(*)}) = \text{Br}(B \rightarrow D^{(*)}\tau\nu)/\text{Br}(B \rightarrow D^{(*)}\ell\nu)$ and in $b \rightarrow s\mu^+\mu^-$ transitions, which are all at the $3 - 4\sigma$ level. In this article we point out that one can explain the $R(D^{(*)})$ anomaly using two scalar leptoquarks (LQs) with the same mass and coupling to fermions related via a discrete symmetry: an $SU(2)_L$ singlet and an $SU(2)_L$ triplet, both with hypercharge $Y = -2/3$. In this way, potentially dangerous contributions to $b \rightarrow s\nu\nu$ are avoided and non-CKM suppressed effects in $R(D^{(*)})$ can be generated. This allows for smaller overall couplings to fermions weakening the direct LHC bounds. In our model, $R(D^{(*)})$ is directly correlated to $b \rightarrow s\tau^+\tau^-$ transitions where an enhancement by orders of magnitude compared to the standard model (SM) is predicted, such that these decay modes are in the reach of LHCb and BELLE II. Furthermore, one can also naturally explain the $b \rightarrow s\mu^+\mu^-$ anomalies (including $R(K)$) by a $C_9 = -C_{10}$ like contribution without spoiling $\mu - e$ universality in charged current decays. In this case sizable effects in $b \rightarrow s\tau\mu$ transitions are predicted which are again well within the experimental reach. One can even address the longstanding anomaly in a_μ , generating a sizable decay rate for $\tau \rightarrow \mu\gamma$. However, we find that out of the three anomalies $R(D^{(*)})$, $b \rightarrow s\mu^+\mu^-$ and a_μ only two (but any two) can be explained simultaneously. We point out that a very similar phenomenology can be achieved using a vector leptoquark $SU(2)_L$ singlet with hypercharge $2/3$. In this case, no tuning between couplings is necessary, but the model is non-renormalizable.

KEYWORDS: Beyond Standard Model, Heavy Quark Physics

ARXIV EPRINT: [1703.09226](https://arxiv.org/abs/1703.09226)

Contents

1	Introduction	1
2	Model and observables	3
2.1	$R(D)$ and $R(D^*)$	4
2.2	$b \rightarrow s\ell^+\ell^-$ transitions	4
2.3	$B^0 - \bar{B}^0$ mixing	5
2.4	a_μ and $\tau \rightarrow \mu\gamma$	5
3	Phenomenological analysis	6
3.1	$R(D)$, $R(D^*)$ and $b \rightarrow s\tau^+\tau^-$	6
3.2	$b \rightarrow s\mu^+\mu^-$ and $b \rightarrow s\tau^\pm\mu^\mp$	7
3.3	a_μ and $\tau \rightarrow \mu\gamma$	8
4	Conclusions and outlook	8

1 Introduction

So far, the LHC did not directly observe any particles beyond the ones present in the SM of particle physics. However, we have intriguing hints for lepton flavor universality violating NP.¹ Most prominently, there exist deviations from the SM predictions in $b \rightarrow s\mu^+\mu^-$ at the $4 - 5\sigma$ level [2–4] and the combination of the ratios $R(D)$ and $R(D^*)$ differs by 3.9σ from its SM prediction [5]. Furthermore, the longstanding anomaly in a_μ (3.1σ [6]) also points towards NP.

$R(D)$ and $R(D^*)$ directly measure lepton flavor universality violation (LFUV), and in the fit to the $b \rightarrow s\mu^+\mu^-$ data also the LHCb measurement of $R(K)$ [7], which deviates by 2.6σ from the SM, points at LFUV. Therefore, it is well motivated to search for a simultaneous explanation of these two anomalies [8–20]. Furthermore, since a_e agrees with the SM prediction, also a_μ can be considered as a LFUV quantity and one can address it together with $R(D^{(*)})$ and/or $b \rightarrow s\mu^+\mu^-$ [13, 21–23].

Concerning $b \rightarrow s\mu^+\mu^-$, a solution is not particularly challenging, as one competes with a process which is in the SM loop and CKM suppressed, a rather small NP contribution, involving moderate couplings and not too light masses, is sufficient (like for example in Z' models [24–36], models with loop effects of heavy scalars and fermions [37, 38] and also by leptoquark models [9, 10, 12, 39–42]²). Also for the a_μ anomaly many possible solutions

¹See for example ref. [1] for a recent overview.

²In ref. [43] it was pointed out that NP effects in charm operators could account for the anomalies. However, this would lead to q^2 dependent effects and it could not explain signs for LFUV.

exist. Here, we would just like to stress that LQs provide a natural solution since they can give the desired large effect because of an m_t/m_μ enhancement [13, 44–47].

However, an explanation of $R(D)$ and $R(D^*)$ is getting more and more delicate. Since these processes are mediated in the SM already at tree-level, a rather large NP contribution is required to account for the $O(20\%)$ deviation. Therefore, new particles added to the SM for explaining $R(D)$ and $R(D^*)$ cannot be very heavy and must have sizable couplings. In the past, mainly three kinds of models with the following new particles have been proposed:

1. Charged Higgses [21, 48–52]
2. W' gauge bosons [11, 14, 53, 54]
3. Leptoquarks [9, 10, 12, 13, 15, 16, 19, 20, 42, 55–60]

Models with charged Higgses lead to (too) large effects in the total B_c lifetime [61] and, depending on the coupling structure, can also be disfavored by the q^2 distribution [62–64]. Interestingly, if the couplings of the charged Higgs are chosen in such a way that they are compatible with the measured q^2 distribution, these models are ruled out by direct searches [65].

Models with W' gauge bosons are also delicate because they necessarily involve Z' bosons due to $SU(2)_L$ gauge invariance. If the Z' width is not unnaturally large, these models are again ruled out by direct searches [11, 65].

In models with leptoquarks generating left-handed vector operators the coupling structure should be aligned to the bottom quark in order to avoid $b \rightarrow s\nu\nu$ bounds. However, in this case the effect in $R(D)$ and $R(D^*)$ is proportional to the small CKM element V_{cb} and large third generation couplings are required to account for the anomalies. These large third generation couplings lead again to stringent bounds from direct LHC searches [65] and electroweak precision observables [66]. In principle, these constraints can be avoided with right-handed couplings [59] (including possibly right-handed neutrinos [16]). However, in such solutions no interference with the SM appears and very large couplings, close to non-perturbativity, are required.

As stated above, LHC bounds from $\tau\tau$ searches can be avoided in case of non-CKM suppressed leptoquark contributions to $R(D)$ and $R(D^*)$. However, for single scalar leptoquark representations, this leads to unacceptably large effects in $b \rightarrow s$ transitions [59]. Therefore, we propose a novel solution to the $R(D^{(*)})$ problem in this article: we introduce two scalar leptoquarks with the same mass M and the same coupling strength to quarks and leptons; an $SU(2)_L$ singlet (Φ_1) and an $SU(2)_L$ triplet (Φ_3) both with hypercharge $Y = -2/3$. Here, the crucial observation is that Φ_1 and Φ_3 contribute with opposite relative sign to $R(D^{(*)})$ than to $b \rightarrow s\nu\nu$ processes such that the effect in $R(D^{(*)})$ is doubled while the contributions in $B \rightarrow K^{(*)}\nu\nu$ cancel at tree-level (see figure 1). Therefore, the couplings to the second quark generation can be larger, non-CKM suppressed effects $R(D^{(*)})$ are possible and the required overall coupling strength is much smaller such that the direct LHC bounds from $\tau\tau$ searches are significantly weakened and the remaining bounds from pair production of third generation LQs are still below the TeV scale [67, 68]. Furthermore, this solution results in a simple rescaling of the SM contributions, predicts




Figure 1. Feynman diagrams contributing to $b \rightarrow c\tau\nu$, $b \rightarrow s\nu\nu$ and $b \rightarrow s\ell\ell$ processes. Both LQs contribute to $b \rightarrow c\tau\nu$ and $b \rightarrow s\nu\nu$ but only Φ_3 to $b \rightarrow s\ell\ell$. Note that with our assumption on the couplings to fermions, the LQs interfere constructively (destructively) in $b \rightarrow c\tau\nu$ ($b \rightarrow s\nu\nu$).

naturally $R(D)/R(D)_{\text{SM}} = R(D^*)/R(D^*)_{\text{SM}}$ and leaves the q^2 distribution unchanged. Adding couplings to muons, we can also address the $b \rightarrow s\mu\mu$ anomalies with a $C_9 = -C_{10}$ like contribution. Finally, adding a (small) right-handed coupling of Φ_1 one can in principle explain a_μ .

This article is structured as follows: in the next section we will present the contributions of our model to all relevant observables. Afterwards, we perform a phenomenological analysis in section 3 before we conclude.

2 Model and observables

The scalar leptoquark singlet Φ_1 and the triplet Φ_3 couple to fermions in the following way:

$$L = \lambda_{fi}^{1L} \overline{Q}_f^c i\tau_2 L_i \Phi_1^\dagger + \lambda_{fi}^{3L} \overline{Q}_f^c i\tau_2 (\tau \cdot \Phi_3)^\dagger L_i + \text{h.c.} \quad (2.1)$$

As motivated in the introduction, we assume that both leptoquarks have the same mass M . In addition, to cancel their effect in $b \rightarrow s\nu\nu$ processes, we impose the discrete symmetry

$$\lambda_{jk}^L \equiv \lambda_{jk}^{1L}, \quad \lambda_{jk}^{3L} = e^{i\pi j} \lambda_{jk}^L, \quad (2.2)$$

on the couplings to fermions. Note that for Φ_1 there is in principle an additional coupling $\lambda_{fi}^R \overline{u}_f^c \ell_i \Phi_1^\dagger$ allowed. We will assume that this coupling is zero and neglect its effect till the discussion of a_μ where small values of λ_{fi}^R can be phenomenologically important due to m_t/m_μ enhanced effects. For our analysis we assume that the couplings λ_{fi}^L are given in the down-quark basis. I.e. after EW symmetry breaking the couplings to left-handed up-quarks involve CKM elements:

$$\lambda_{d_{fi}}^L \equiv \lambda_{fi}^L, \quad \lambda_{u_{fi}}^L = V_{fj}^* \lambda_{ji}^L. \quad (2.3)$$

We will now discuss the various relevant processes to which our model contributes. As already noted above, our model is constructed in such a way that we do not get tree-level contributions to $b \rightarrow s\nu\nu$ transitions, which we therefore omit in the following. We also neglect couplings to the first generation of quarks and leptons. For quarks, this is only possible in the interaction basis since CKM rotations induce either couplings to up or down quarks. However, charged current decays of Kaons or D mesons involve large CKM angles in the SM, making the relative effects of LQs small.

2.1 $R(D)$ and $R(D^*)$

We define the effective Hamiltonian for $b \rightarrow c\ell\nu$ transitions as

$$H_{\text{eff}}^{\ell_f\nu_i} = \frac{4G_F}{\sqrt{2}} V_{cb} C_L^{fi} [\bar{c}\gamma^\mu P_L b] [\bar{\ell}_f \gamma_\mu P_L \nu_i], \quad (2.4)$$

where in the SM $C_L^{fi} = \delta_{fi}$ and the contribution of our model is given by

$$C_L^{fi} = \frac{\sqrt{2}}{8G_F M^2} \frac{V_{cj}}{V_{cb}} \lambda_{3i}^L \lambda_{jf}^{L*} (1 + (-1)^j). \quad (2.5)$$

With these conventions we have

$$R(D^{(*)}) / R(D^{(*)})_{\text{SM}} \equiv X_{D^{(*)}} = \sum_{i=1}^3 (\delta_{3i} + C_L^{3i})^2, \quad (2.6)$$

assuming vanishing contributions to the muon and electron channels. This has to be compared to the experimental values of

$$R(D^*)_{\text{EXP}} = 0.316 \pm 0.016 \pm 0.010, \quad (2.7)$$

$$R(D)_{\text{EXP}} = 0.397 \pm 0.040 \pm 0.028, \quad (2.8)$$

and the corresponding SM predictions [69, 70]

$$R(D^*)_{\text{SM}} = 0.252 \pm 0.003, \quad (2.9)$$

$$R(D)_{\text{SM}} = 0.300 \pm 0.008. \quad (2.10)$$

2.2 $b \rightarrow s\ell^+\ell^-$ transitions

Using the effective Hamiltonian

$$H_{\text{eff}}^{\ell_f\ell_i} = -\frac{4G_F}{\sqrt{2}} V_{tb} V_{ts}^* \sum_{a=9,10} C_a^{fi} O_a^{fi},$$

$$O_{9(10)}^{fi} = \frac{\alpha}{4\pi} [\bar{s}\gamma^\mu P_L b] [\bar{\ell}_f \gamma_\mu (\gamma^5)\ell_i], \quad (2.11)$$

we have

$$C_9^{fi} = -C_{10}^{fi} = \frac{-\sqrt{2}}{2G_F V_{tb} V_{ts}^*} \frac{\pi}{\alpha} \frac{1}{M^2} \lambda_{3i}^L \lambda_{2f}^{L*}. \quad (2.12)$$

The allowed range at the 2σ level [2] (see also refs. [3, 4]) is given by

$$-0.18(-0.35) \geq C_9^{22} = -C_{10}^{22} \geq (-0.71) - 0.91, \quad (2.13)$$

at the $(1\sigma) 2\sigma$ level.

We will also need the process $B_s \rightarrow \tau^+\tau^-$. The current experimental limit is [71]

$$\text{Br}(B_s \rightarrow \tau^+\tau^-)_{\text{EXP}} \leq 6.8 \times 10^{-3}. \quad (2.14)$$

The SM prediction is given by [72, 73]

$$\text{Br}(B_s \rightarrow \tau^+\tau^-)_{\text{SM}} = (7.73 \pm 0.49) \times 10^{-7}, \quad (2.15)$$

and in our model we have

$$\text{Br}(B_s \rightarrow \tau^+ \tau^-) = \text{Br}(B_s \rightarrow \tau^+ \tau^-)_{\text{SM}} \left(1 + \frac{C_{10}^{33}}{C_{10}^{\text{SM}}} \right)^2, \quad (2.16)$$

with $C_{10}^{\text{SM}} \approx -4.3$ [74, 75]. For the analysis of $B \rightarrow K^{(*)} \tau \mu$ we will use the results of ref. [33].

2.3 $B^0 - \bar{B}^0$ mixing

Here we find

$$\begin{aligned} H_{\text{eff}} &= C_1 \bar{s} \gamma^\mu P_L b \bar{s} \gamma_\mu P_L b \\ C_1 &= \frac{-1}{128\pi^2} (\lambda_{23}^{L*} \lambda_{33}^L)^2 D_2(m_\tau^2, m_\tau^2, M^2, M^2) \end{aligned} \quad (2.17)$$

which corresponds to an effect of the order of 1%. This can be easily understood as follows: since we need an $\mathcal{O}(10\%)$ effect in $R(D^{(*)})$ at the amplitude level, this effects gets squared for $B^0 - \bar{B}^0$ and there are no enhancement factors, the final effect is around 1% and below the current sensitivity of approximately 10%.

2.4 a_μ and $\tau \rightarrow \mu \gamma$

In order to aim at an explanation of a_μ one needs a chirality enhanced effect. Therefore, let us add to eq. (2.1) the following term

$$\lambda_{fi}^R \bar{u}_f^c \ell_i \Phi_1^\dagger + h.c.. \quad (2.18)$$

In this case the numerically relevant m_t enhanced contribution to a_μ is given by

$$\delta a_\mu = \frac{m_\mu}{4\pi^2} \text{Re} [C_R^{22}], \quad (2.19)$$

with

$$C_L^{fi} = -\frac{N_c}{12M^2} m_t \lambda_{3f}^R \lambda_{3i}^{L*} \left(7 + 4 \log \left(\frac{m_t^2}{M^2} \right) \right), \quad (2.20)$$

and C_R^{23} is obtained from C_L^{23} by $L \leftrightarrow R$. We will assume that λ_{32}^R is small compared to λ_{32}^L .

The world average of the measurement of $a_\mu \equiv (g-2)_\mu/2$ is completely dominated by the Brookhaven experiment E821 [76] and is given by [77] $a_\mu^{\text{exp}} = (116\,592\,091 \pm 54 \pm 33) \times 10^{-11}$ where the first error is statistical and the second one is systematic. The current SM prediction is [6, 78–86] $a_\mu^{\text{SM}} = (116\,591\,811 \pm 62) \times 10^{-11}$ where almost the whole uncertainty is due to hadronic effects. This amounts to a discrepancy between the SM and the experimental value of

$$\delta a_\mu = a_\mu^{\text{exp}} - a_\mu^{\text{SM}} = (278 \pm 88) \times 10^{-11}, \quad (2.21)$$

i.e. a 3.1σ deviation.

For $\tau \rightarrow \mu \gamma$ the branching ratio reads

$$\text{Br}(\tau \rightarrow \mu \gamma) = \frac{\alpha m_\tau^3}{256\pi^4} \tau_\tau \left(|C_L^{23}|^2 + |C_R^{23}|^2 \right). \quad (2.22)$$

The current experimental bound is given by [87]

$$\text{Br}(\tau \rightarrow \mu \gamma) < 4.4 \times 10^{-8}. \quad (2.23)$$




Figure 2. Left: allowed regions by $R(D)$ and $R(D^*)$ in the $\lambda_{23}^L - \lambda_{33}^L$ plane for $M = 1$ TeV using the weighted average for $R(D^{(*)})_{\text{EXP}}/R(D^{(*)})_{\text{SM}}$. Note that already small couplings are sufficient to account for $R(D)$ and $R(D^*)$. Therefore, the bounds from LHC searches are weakened and the leptoquarks can also be easily heavier than 1 TeV and still explain the anomalies with couplings in the perturbative regime. Right: prediction for $B_s \rightarrow \tau\tau$ (red) as a function of $R(D^{(*)})/R(D^{(*)})_{\text{SM}}$. Here we neglected small CKM suppressed contributions.

3 Phenomenological analysis

3.1 $R(D)$, $R(D^*)$ and $b \rightarrow s\tau^+\tau^-$

Let us first consider the size of the couplings needed to explain $R(D)$ and $R(D^*)$. Here and in the following, we will assume them to be real. As we can see in the left plot of figure 2, we only need small couplings (of the order of 0.1 for 1 TeV leptoquarks) in order to explain $R(D)$, $R(D^*)$. This is possible because we avoid contributions to $b \rightarrow s\nu\nu$ and hence our effect in $b \rightarrow c\tau\nu$ does not need to be CKM suppressed. Therefore, the bounds from ref. [65] do not apply to our model and we are not in conflict with LHC bounds, especially because the LQs can be much heavier than 1 TeV while still possessing perturbative couplings and explaining $R(D^{(*)})$.

Next, note that neglecting small CKM factors, the contributions to $b \rightarrow c\tau\nu$ and $b \rightarrow s\tau\tau$ depend on the same product of couplings $\lambda_{23}^L\lambda_{33}^{L*}$ (modulus small CKM ratios). Therefore, we can express $B_s \rightarrow \tau^+\tau^-$ in terms of the effect in $R(D^{(*)})$:

$$\frac{\text{Br}(B_s \rightarrow \tau\tau)}{\text{Br}(B_s \rightarrow \tau\tau)_{\text{SM}}} = \left(1 + 2 \frac{\pi}{\alpha} \frac{V_{cb}}{V_{ts}^*} \frac{\sqrt{X_{D^{(*)}} - 1}}{C_{10}^{\text{SM}}} \right)^2. \quad (3.1)$$

The resulting numerical prediction for $B_s \rightarrow \tau^+\tau^-$ is shown in figure 2. We can see that the branching ratio can be enhanced by up to three orders of magnitude compared to the SM prediction. Therefore, even though it is experimentally challenging to search for, our model can be tested with $B_s \rightarrow \tau^+\tau^-$ measurements at LHCb. Also an enhancement of $B \rightarrow K^{(*)}\tau^+\tau^-$ in the same ballpark is predicted which could be tested at BELLE II.




Figure 3. Left: contours and excluded region for $B \rightarrow K\tau\mu = (B \rightarrow K\tau^+\mu^- + B \rightarrow K\tau^-\mu^+)/2$ for $C_9^{22} = -0.5$, i.e. assuming that C_9^{22} takes the central value obtained from the $b \rightarrow s\mu^+\mu^-$ fit. The colored regions are allowed by the various processes. For $R(D)$ and $R(D^*)$ we used again the weighted average for $R(D^{(*)})_{\text{EXP}}/R(D^{(*)})_{\text{SM}}$. Right: the contour lines show $\text{Br}[\tau \rightarrow \mu\gamma] \times 10^8$. The gray region is excluded by the current upper limit and (light) red region is allowed by a_μ at the (2σ) 1σ level. Note that both δa_μ and $\tau \rightarrow \mu\gamma$ are only a function of $\lambda_{33}^L/\lambda_{32}^L$ and therefore independent of $b \rightarrow s\mu^+\mu^-$ transitions.

3.2 $b \rightarrow s\mu^+\mu^-$ and $b \rightarrow s\tau^\pm\mu^\mp$

Let us now consider the effect of including $b \rightarrow s\mu^+\mu^-$ transitions in our analysis. In this case effects in $B \rightarrow D^{(*)}\mu\nu/B \rightarrow D^{(*)}e\nu$ are predicted if still addressing $R(D)$ and $R(D^*)$ simultaneously. We checked that the effect is at the per-mill level which is compatible with BELLE and BABAR measurements.³ However, interesting correlations with $b \rightarrow s\tau\mu$ processes appear. Here we find

$$C_9^{32} = -2\frac{\pi}{\alpha} \frac{V_{cb}}{V_{ts}^*} \frac{\lambda_{32}^L}{\lambda_{33}^L} \left(\sqrt{X_{D^{(*)}}} - 1 \right), \quad (3.2)$$

$$C_9^{23} = \frac{\lambda_{33}^L}{\lambda_{32}^L} C_9^{22}, \quad (3.3)$$

which depends only on the ratio $\lambda_{33}^L/\lambda_{32}^L$ as a free parameter. Note that the dependence on C_9^{22} is much weaker than on $X_{D^{(*)}}$. The resulting bounds and predictions are shown in the left plot of figure 3. We take the experimental limit [90]

$$\text{Br}[B \rightarrow K\tau\mu] < 4.8 \times 10^{-5}. \quad (3.4)$$

Note that $R(D^{(*)})$ can only be fully explained for $\lambda_{33}^L/\lambda_{32}^L > 1$.

³This is contrary to ref. [13] which cannot explain $R(D^{(*)})$ and $b \rightarrow s\mu^+\mu^-$ data simultaneously without violating the bounds from $B \rightarrow D^{(*)}\mu\nu/B \rightarrow D^{(*)}e\nu$ as pointed out in ref. [88]. However, this tension can be relieved with leptoquarks masses larger than 5 TeV [89].

3.3 a_μ and $\tau \rightarrow \mu\gamma$

Considering only the couplings λ^L the effect in $\tau \rightarrow \mu\gamma$ is negligibly small. Things get much more interesting if we aim at a simultaneous explanation of the anomalous magnetic moment of the muon. In this case chirally enhanced effects also appear in $\tau \rightarrow \mu\gamma$. We have

$$\text{Br}[\tau \rightarrow \mu\gamma] \geq \frac{\alpha m_\tau^3}{16\Gamma_\tau} \frac{a_\mu^2}{m_\mu^2} \left| \frac{\lambda_{33}^L}{\lambda_{32}^L} \right|^2. \quad (3.5)$$

Here we set $\lambda_{33}^R = 0$.

Note that $\text{Br}(\tau \rightarrow \mu\gamma)$ can only be enhanced by allowing λ_{33}^R to be different from zero, resulting in the \geq sign in eq. (3.5). The result is shown in the right plot of figure 3. Note that a_μ can only be explained for $\lambda_{33}^L/\lambda_{32}^L < 0.65$ (at the 2σ level). This is opposite to the case of $b \rightarrow s\mu^+\mu^-$ which can only be explained for $\lambda_{33}^L/\lambda_{32}^L > 1$. Therefore, we conclude that our model can explain out of the three anomalies $R(D^{(*)})$, $b \rightarrow s\mu^+\mu^-$ and a_μ only two simultaneously.

4 Conclusions and outlook

In this article we proposed a scalar leptoquark model which can give sizable effects on $R(D)$ and $R(D^*)$ without suffering from problems with $b \rightarrow s\nu\nu$, q^2 distributions in $R(D^{(*)})$, from large couplings in the non-perturbative regime or from tensions with direct LHC searches as it is the case for nearly all other models on the market. Our model predicts sizable branching fractions for $b \rightarrow s\tau^+\tau^-$ processes (of the order of 10^{-3}) being directly correlated to $R(D^{(*)})$.

Furthermore, the model can naturally explain $b \rightarrow s\mu^+\mu^-$ (including $R(K)$) via a $C_9 = -C_{10}$ contribution and therefore also predicts $R(K^*)$ to be significantly below the SM value. In case of a simultaneous explanation of $R(D^{(*)})$ with $b \rightarrow s\mu^+\mu^-$ we only get effects in $B \rightarrow D^{(*)}\mu\nu/B \rightarrow D^{(*)}e\nu$ at the per-mill level, but sizable ones in $b \rightarrow s\tau\mu$ processes (depending on only one free parameter), making them potentially observable at LHCb or BELLE II in the near future.

The tension in a_μ can be explained as well by adding a small right-handed coupling of the $SU(2)_L$ singlet LQ to tops and muons. As a consequence sizable rates for $\tau \rightarrow \mu\gamma$ are predicted. Here the dependence on the remaining free parameter is opposite to $b \rightarrow s\tau\mu$ excluding a simultaneous explanation of all three anomalies, i.e. out of $R(D^{(*)})$, $b \rightarrow s\mu^+\mu^-$ and a_μ our model can explain any two of them.

We stress that in general our approach of combining the $SU(2)_L$ singlet with the $SU(2)_L$ triplet is the only way of explaining $R(D)$ and $R(D^*)$, without violating $b \rightarrow s\nu\nu$ or direct LHC bounds, if the SM is extended by scalar LQs only. However, one can get the same phenomenology in B decays using a vector leptoquark $SU(2)_L$ singlet with hypercharge $2/3$ and couplings to left-handed fermions. In this case, no tuning between couplings is necessary and effects in $b \rightarrow s\nu\nu$ are automatically avoided. However, the model is non-renormalizable and while our model with scalar leptoquarks only predicts effects of the order of 1% in $B_s - \bar{B}_s$ mixing, this effect is much larger for the vector leptoquark [12].




Figure 4. Allowed regions for $R(D^{(*)})$ and $b \rightarrow s\nu\nu$ assuming independent couplings and masses for the leptquark singlet and triplet. Here $M_{1(3)}$ is the mass of Φ_1 (Φ_3).

Furthermore, adding right-handed couplings, the effect of the vector leptoquark in a_μ is only enhanced by the bottom mass but not by the top one.

In our model we assumed a discrete symmetry between the couplings of the two leptoquarks in order to cancel exactly the effect in $b \rightarrow s\nu\nu$. However, even if one disregards this assumption and allows for independent masses and couplings of Φ_1 and Φ_3 , a cancellation in $b \rightarrow s\nu\nu$ is still possible. In fact, as shown in figure 4 the constraints from $b \rightarrow s\nu\nu$ transitions still allow for an explanation of $R(D^{(*)})$ without severe fine-tuning. Therefore, our imposed symmetry does not to be exact in order to provide a valid explanation of the anomalies.

If one allows in addition for direct couplings (not only CKM induced couplings to up-quarks) to first generation quarks, a sizable effect in $K \rightarrow \pi\mu\mu/K \rightarrow \pi ee$ is possible which could be tested at NA62 or KOTO [91] and interesting correlations with $b \rightarrow d$ transitions occur. Therefore, a very detailed study of our model is important and promising in order to explore the many interesting effects which can be observed by ongoing and future experiments.

Acknowledgments

A.C. thanks Admir Greljo for useful discussions about LHC searches and Michael Spira for useful comments on the manuscript. The work of A.C. and D.M. is supported by an Ambizione Grant of the Swiss National Science Foundation (PZ00P2_154834).

Note added. After submission of this article, LHCb presented results for the ratio $R(K^*)$ [92] which are in agreement with the predictions of our model. Including the new measurement in the global fit to $b \rightarrow s\mu^+\mu^-$ data, the significance for NP in $C_9 = -C_{10}$ increased to 5.2σ [93].

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] A. Crivellin, *B-anomalies related to leptons and lepton flavour universality violation*, [PoS\(BEAUTY2016\)042](#) [[arXiv:1606.06861](#)] [[INSPIRE](#)].
- [2] W. Altmannshofer and D.M. Straub, *Implications of $b \rightarrow s$ measurements*, [arXiv:1503.06199](#) [[INSPIRE](#)].
- [3] S. Descotes-Genon, L. Hofer, J. Matias and J. Virto, *Global analysis of $b \rightarrow s\ell\ell$ anomalies*, [JHEP 06 \(2016\) 092](#) [[arXiv:1510.04239](#)] [[INSPIRE](#)].
- [4] T. Hurth, F. Mahmoudi and S. Neshatpour, *On the anomalies in the latest LHCb data*, [Nucl. Phys. B 909 \(2016\) 737](#) [[arXiv:1603.00865](#)] [[INSPIRE](#)].
- [5] Y. Amhis et al., *Averages of b -hadron, c -hadron and τ -lepton properties as of summer 2016*, [arXiv:1612.07233](#) [[INSPIRE](#)].
- [6] A. Nyffeler, *Precision of a data-driven estimate of hadronic light-by-light scattering in the muon $g - 2$: Pseudoscalar-pole contribution*, [Phys. Rev. D 94 \(2016\) 053006](#) [[arXiv:1602.03398](#)] [[INSPIRE](#)].
- [7] LHCb collaboration, *Test of lepton universality using $B^+ \rightarrow K^+\ell^+\ell^-$ decays*, [Phys. Rev. Lett. 113 \(2014\) 151601](#) [[arXiv:1406.6482](#)] [[INSPIRE](#)].
- [8] B. Bhattacharya, A. Datta, D. London and S. Shivashankara, *Simultaneous Explanation of the R_K and $R(D^{(*)})$ Puzzles*, [Phys. Lett. B 742 \(2015\) 370](#) [[arXiv:1412.7164](#)] [[INSPIRE](#)].
- [9] L. Calibbi, A. Crivellin and T. Ota, *Effective Field Theory Approach to $b \rightarrow s\ell\ell^{(\prime)}$, $B \rightarrow K^{(*)}\nu\bar{\nu}$ and $B \rightarrow D^{(*)}\tau\nu$ with Third Generation Couplings*, [Phys. Rev. Lett. 115 \(2015\) 181801](#) [[arXiv:1506.02661](#)] [[INSPIRE](#)].
- [10] S. Fajfer and N. Košnik, *Vector leptoquark resolution of R_K and $R_{D^{(*)}}$ puzzles*, [Phys. Lett. B 755 \(2016\) 270](#) [[arXiv:1511.06024](#)] [[INSPIRE](#)].
- [11] A. Greljo, G. Isidori and D. Marzocca, *On the breaking of Lepton Flavor Universality in B decays*, [JHEP 07 \(2015\) 142](#) [[arXiv:1506.01705](#)] [[INSPIRE](#)].
- [12] R. Barbieri, G. Isidori, A. Pattori and F. Senia, *Anomalies in B -decays and $U(2)$ flavour symmetry*, [Eur. Phys. J. C 76 \(2016\) 67](#) [[arXiv:1512.01560](#)] [[INSPIRE](#)].
- [13] M. Bauer and M. Neubert, *Minimal Leptoquark Explanation for the $R_{D^{(*)}}$, R_K and $(g - 2)_g$ Anomalies*, [Phys. Rev. Lett. 116 \(2016\) 141802](#) [[arXiv:1511.01900](#)] [[INSPIRE](#)].
- [14] S.M. Boucenna, A. Celis, J. Fuentes-Martin, A. Vicente and J. Virto, *Phenomenology of an $SU(2) \times SU(2) \times U(1)$ model with lepton-flavour non-universality*, [JHEP 12 \(2016\) 059](#) [[arXiv:1608.01349](#)] [[INSPIRE](#)].
- [15] D. Das, C. Hati, G. Kumar and N. Mahajan, *Towards a unified explanation of $R_{D^{(*)}}$, R_K and $(g - 2)_\mu$ anomalies in a left-right model with leptoquarks*, [Phys. Rev. D 94 \(2016\) 055034](#) [[arXiv:1605.06313](#)] [[INSPIRE](#)].
- [16] D. Bečirević, S. Fajfer, N. Košnik and O. Sumensari, *Leptoquark model to explain the B -physics anomalies, R_K and R_D* , [Phys. Rev. D 94 \(2016\) 115021](#) [[arXiv:1608.08501](#)] [[INSPIRE](#)].

- [17] S. Sahoo, R. Mohanta and A.K. Giri, *Explaining the R_K and $R_{D^{(*)}}$ anomalies with vector leptoquarks*, *Phys. Rev. D* **95** (2017) 035027 [[arXiv:1609.04367](#)] [[INSPIRE](#)].
- [18] B. Bhattacharya, A. Datta, J.-P. Guévin, D. London and R. Watanabe, *Simultaneous Explanation of the R_K and $R_{D^{(*)}}$ Puzzles: a Model Analysis*, *JHEP* **01** (2017) 015 [[arXiv:1609.09078](#)] [[INSPIRE](#)].
- [19] R. Barbieri, C.W. Murphy and F. Senia, *B-decay Anomalies in a Composite Leptoquark Model*, *Eur. Phys. J. C* **77** (2017) 8 [[arXiv:1611.04930](#)] [[INSPIRE](#)].
- [20] C.-H. Chen, T. Nomura and H. Okada, *Excesses of muon $g - 2$, $R_{D^{(*)}}$ and R_K in a leptoquark model*, [arXiv:1703.03251](#) [[INSPIRE](#)].
- [21] A. Crivellin, J. Heeck and P. Stoffer, *A perturbed lepton-specific two-Higgs-doublet model facing experimental hints for physics beyond the Standard Model*, *Phys. Rev. Lett.* **116** (2016) 081801 [[arXiv:1507.07567](#)] [[INSPIRE](#)].
- [22] G. Bélanger, C. Delaunay and S. Westhoff, *A Dark Matter Relic From Muon Anomalies*, *Phys. Rev. D* **92** (2015) 055021 [[arXiv:1507.06660](#)] [[INSPIRE](#)].
- [23] W. Altmannshofer, M. Carena and A. Crivellin, *$L_\mu - L_\tau$ theory of Higgs flavor violation and $(g - 2)_\mu$* , *Phys. Rev. D* **94** (2016) 095026 [[arXiv:1604.08221](#)] [[INSPIRE](#)].
- [24] S. Descotes-Genon, J. Matias and J. Virto, *Understanding the $B \rightarrow K^* \mu^+ \mu^-$ Anomaly*, *Phys. Rev. D* **88** (2013) 074002 [[arXiv:1307.5683](#)] [[INSPIRE](#)].
- [25] R. Gauld, F. Goertz and U. Haisch, *On minimal Z' explanations of the $B \rightarrow K^* \mu^+ \mu^-$ anomaly*, *Phys. Rev. D* **89** (2014) 015005 [[arXiv:1308.1959](#)] [[INSPIRE](#)].
- [26] A.J. Buras and J. Girrbach, *Left-handed Z' and Z FCNC quark couplings facing new $b \rightarrow s \mu^+ \mu^-$ data*, *JHEP* **12** (2013) 009 [[arXiv:1309.2466](#)] [[INSPIRE](#)].
- [27] A.J. Buras, F. De Fazio and J. Girrbach, *331 models facing new $b \rightarrow s \mu^+ \mu^-$ data*, *JHEP* **02** (2014) 112 [[arXiv:1311.6729](#)] [[INSPIRE](#)].
- [28] W. Altmannshofer, S. Gori, M. Pospelov and I. Yavin, *Quark flavor transitions in $L_\mu - L_\tau$ models*, *Phys. Rev. D* **89** (2014) 095033 [[arXiv:1403.1269](#)] [[INSPIRE](#)].
- [29] A. Crivellin, G. D'Ambrosio and J. Heeck, *Explaining $h \rightarrow \mu^\pm \tau^\mp$, $B \rightarrow K^* \mu^+ \mu^-$ and $B \rightarrow K \mu^+ \mu^- / B \rightarrow K e^+ e^-$ in a two-Higgs-doublet model with gauged $L_\mu - L_\tau$* , *Phys. Rev. Lett.* **114** (2015) 151801 [[arXiv:1501.00993](#)] [[INSPIRE](#)].
- [30] A. Crivellin, G. D'Ambrosio and J. Heeck, *Addressing the LHC flavor anomalies with horizontal gauge symmetries*, *Phys. Rev. D* **91** (2015) 075006 [[arXiv:1503.03477](#)] [[INSPIRE](#)].
- [31] C. Niehoff, P. Stangl and D.M. Straub, *Violation of lepton flavour universality in composite Higgs models*, *Phys. Lett. B* **747** (2015) 182 [[arXiv:1503.03865](#)] [[INSPIRE](#)].
- [32] D. Aristizabal Sierra, F. Staub and A. Vicente, *Shedding light on the $b \rightarrow s$ anomalies with a dark sector*, *Phys. Rev. D* **92** (2015) 015001 [[arXiv:1503.06077](#)] [[INSPIRE](#)].
- [33] A. Crivellin, L. Hofer, J. Matias, U. Nierste, S. Pokorski and J. Rosiek, *Lepton-flavour violating B decays in generic Z' models*, *Phys. Rev. D* **92** (2015) 054013 [[arXiv:1504.07928](#)] [[INSPIRE](#)].
- [34] A. Celis, J. Fuentes-Martin, M. Jung and H. Serodio, *Family nonuniversal Z' models with protected flavor-changing interactions*, *Phys. Rev. D* **92** (2015) 015007 [[arXiv:1505.03079](#)] [[INSPIRE](#)].

- [35] A. Crivellin, J. Fuentes-Martin, A. Greljo and G. Isidori, *Lepton Flavor Non-Universality in B decays from Dynamical Yukawas*, *Phys. Lett. B* **766** (2017) 77 [[arXiv:1611.02703](#)] [[INSPIRE](#)].
- [36] I. Garcia Garcia, *LHCb anomalies from a natural perspective*, *JHEP* **03** (2017) 040 [[arXiv:1611.03507](#)] [[INSPIRE](#)].
- [37] B. Gripaios, M. Nardecchia and S.A. Renner, *Linear flavour violation and anomalies in B physics*, *JHEP* **06** (2016) 083 [[arXiv:1509.05020](#)] [[INSPIRE](#)].
- [38] P. Arnan, L. Hofer, F. Mescia and A. Crivellin, *Loop effects of heavy new scalars and fermions in $b \rightarrow s\mu^+\mu^-$* , *JHEP* **04** (2017) 043 [[arXiv:1608.07832](#)] [[INSPIRE](#)].
- [39] B. Gripaios, M. Nardecchia and S.A. Renner, *Composite leptoquarks and anomalies in B-meson decays*, *JHEP* **05** (2015) 006 [[arXiv:1412.1791](#)] [[INSPIRE](#)].
- [40] D. Bećirević, S. Fajfer and N. Košnik, *Lepton flavor nonuniversality in $b \rightarrow s\ell^+\ell^-$ processes*, *Phys. Rev. D* **92** (2015) 014016 [[arXiv:1503.09024](#)] [[INSPIRE](#)].
- [41] I. de Medeiros Varzielas and G. Hiller, *Clues for flavor from rare lepton and quark decays*, *JHEP* **06** (2015) 072 [[arXiv:1503.01084](#)] [[INSPIRE](#)].
- [42] R. Alonso, B. Grinstein and J. Martin Camalich, *Lepton universality violation and lepton flavor conservation in B-meson decays*, *JHEP* **10** (2015) 184 [[arXiv:1505.05164](#)] [[INSPIRE](#)].
- [43] S. Jäger, K. Leslie, M. Kirk and A. Lenz, *Charming new physics in rare B-decays and mixing?*, [arXiv:1701.09183](#) [[INSPIRE](#)].
- [44] A. Djouadi, T. Kohler, M. Spira and J. Tutas, *(eb), (et) type leptoquarks at e p colliders*, *Z. Phys. C* **46** (1990) 679 [[INSPIRE](#)].
- [45] D. Chakraverty, D. Choudhury and A. Datta, *A Nonsupersymmetric resolution of the anomalous muon magnetic moment*, *Phys. Lett. B* **506** (2001) 103 [[hep-ph/0102180](#)] [[INSPIRE](#)].
- [46] K.-m. Cheung, *Muon anomalous magnetic moment and leptoquark solutions*, *Phys. Rev. D* **64** (2001) 033001 [[hep-ph/0102238](#)] [[INSPIRE](#)].
- [47] E. Coluccio Leskow, G. D'Ambrosio, A. Crivellin and D. Müller, *(g - 2) μ , lepton flavor violation and Z decays with leptoquarks: Correlations and future prospects*, *Phys. Rev. D* **95** (2017) 055018 [[arXiv:1612.06858](#)] [[INSPIRE](#)].
- [48] A. Crivellin, C. Greub and A. Kokulu, *Explaining $B \rightarrow D\tau\nu$, $B \rightarrow D^*\tau\nu$ and $B \rightarrow \tau\nu$ in a 2HDM of type-III*, *Phys. Rev. D* **86** (2012) 054014 [[arXiv:1206.2634](#)] [[INSPIRE](#)].
- [49] M. Tanaka and R. Watanabe, *New physics in the weak interaction of $\bar{B} \rightarrow D^{(*)}\tau\bar{\nu}$* , *Phys. Rev. D* **87** (2013) 034028 [[arXiv:1212.1878](#)] [[INSPIRE](#)].
- [50] A. Celis, M. Jung, X.-Q. Li and A. Pich, *Sensitivity to charged scalars in $B \rightarrow D^{(*)}\tau\nu_\tau$ and $B \rightarrow \tau\nu_\tau$ decays*, *JHEP* **01** (2013) 054 [[arXiv:1210.8443](#)] [[INSPIRE](#)].
- [51] A. Crivellin, A. Kokulu and C. Greub, *Flavor-phenomenology of two-Higgs-doublet models with generic Yukawa structure*, *Phys. Rev. D* **87** (2013) 094031 [[arXiv:1303.5877](#)] [[INSPIRE](#)].
- [52] C.-H. Chen and T. Nomura, *$\bar{B} \rightarrow D^{(*)}\tau\bar{\nu}$ and τ polarization in a generic two-Higgs-doublet model*, [arXiv:1703.03646](#) [[INSPIRE](#)].

- [53] S.M. Boucenna, A. Celis, J. Fuentes-Martin, A. Vicente and J. Virto, *Non-abelian gauge extensions for B-decay anomalies*, *Phys. Lett. B* **760** (2016) 214 [[arXiv:1604.03088](#)] [[INSPIRE](#)].
- [54] E. Megias, M. Quirós and L. Salas, *Lepton-flavor universality violation in R_K and $R_{D^{(*)}}$ from warped space*, *JHEP* **07** (2017) 102 [[arXiv:1703.06019](#)] [[INSPIRE](#)].
- [55] S. Fajfer, J.F. Kamenik, I. Nisandzic and J. Zupan, *Implications of Lepton Flavor Universality Violations in B Decays*, *Phys. Rev. Lett.* **109** (2012) 161801 [[arXiv:1206.1872](#)] [[INSPIRE](#)].
- [56] N.G. Deshpande and A. Menon, *Hints of R-parity violation in B decays into $\tau\nu$* , *JHEP* **01** (2013) 025 [[arXiv:1208.4134](#)] [[INSPIRE](#)].
- [57] Y. Sakaki, M. Tanaka, A. Tayduganov and R. Watanabe, *Testing leptoquark models in $\bar{B} \rightarrow D^{(*)}\tau\bar{\nu}$* , *Phys. Rev. D* **88** (2013) 094012 [[arXiv:1309.0301](#)] [[INSPIRE](#)].
- [58] N.G. Deshpande and X.-G. He, *Consequences of R-parity violating interactions for anomalies in $\bar{B} \rightarrow D^{(*)}\tau\bar{\nu}$ and $b \rightarrow s\mu^+\mu^-$* , *Eur. Phys. J. C* **77** (2017) 134 [[arXiv:1608.04817](#)] [[INSPIRE](#)].
- [59] X.-Q. Li, Y.-D. Yang and X. Zhang, *Revisiting the one leptoquark solution to the $R(D^{(*)})$ anomalies and its phenomenological implications*, *JHEP* **08** (2016) 054 [[arXiv:1605.09308](#)] [[INSPIRE](#)].
- [60] B. Dumont, K. Nishiwaki and R. Watanabe, *LHC constraints and prospects for S_1 scalar leptoquark explaining the $\bar{B} \rightarrow D^{(*)}\tau\bar{\nu}$ anomaly*, *Phys. Rev. D* **94** (2016) 034001 [[arXiv:1603.05248](#)] [[INSPIRE](#)].
- [61] R. Alonso, B. Grinstein and J. Martin Camalich, *Lifetime of B_c^- Constrains Explanations for Anomalies in $B \rightarrow D^{(*)}\tau\nu$* , *Phys. Rev. Lett.* **118** (2017) 081802 [[arXiv:1611.06676](#)] [[INSPIRE](#)].
- [62] M. Freytsis, Z. Ligeti and J.T. Ruderman, *Flavor models for $\bar{B} \rightarrow D^{(*)}\tau\bar{\nu}$* , *Phys. Rev. D* **92** (2015) 054018 [[arXiv:1506.08896](#)] [[INSPIRE](#)].
- [63] A. Celis, M. Jung, X.-Q. Li and A. Pich, *Scalar contributions to $b \rightarrow c(u)\tau\nu$ transitions*, *Phys. Lett. B* **771** (2017) 168 [[arXiv:1612.07757](#)] [[INSPIRE](#)].
- [64] M.A. Ivanov, J.G. Körner and C.-T. Tran, *Probing new physics in $\bar{B}^0 \rightarrow D^{(*)}\tau^-\bar{\nu}_\tau$ using the longitudinal, transverse and normal polarization components of the tau lepton*, *Phys. Rev. D* **95** (2017) 036021 [[arXiv:1701.02937](#)] [[INSPIRE](#)].
- [65] D.A. Faroughy, A. Greljo and J.F. Kamenik, *Confronting lepton flavor universality violation in B decays with high- p_T tau lepton searches at LHC*, *Phys. Lett. B* **764** (2017) 126 [[arXiv:1609.07138](#)] [[INSPIRE](#)].
- [66] F. Feruglio, P. Paradisi and A. Pattori, *Revisiting Lepton Flavor Universality in B Decays*, *Phys. Rev. Lett.* **118** (2017) 011801 [[arXiv:1606.00524](#)] [[INSPIRE](#)].
- [67] CMS collaboration, *Search for R-parity violating decays of a top squark in proton-proton collisions at $\sqrt{s} = 8$ TeV*, *Phys. Lett. B* **760** (2016) 178 [[arXiv:1602.04334](#)] [[INSPIRE](#)].
- [68] ATLAS collaboration, *A search for top squarks with R-parity-violating decays to all-hadronic final states with the ATLAS detector in $\sqrt{s} = 8$ TeV proton-proton collisions*, *JHEP* **06** (2016) 067 [[arXiv:1601.07453](#)] [[INSPIRE](#)].

- [69] S. Fajfer, J.F. Kamenik and I. Nisandzic, *On the $B \rightarrow D^* \tau \bar{\nu}_\tau$ Sensitivity to New Physics*, *Phys. Rev. D* **85** (2012) 094025 [[arXiv:1203.2654](#)] [[INSPIRE](#)].
- [70] HPQCD collaboration, H. Na, C.M. Bouchard, G.P. Lepage, C. Monahan and J. Shigemitsu, *$B \rightarrow D l \nu$ form factors at nonzero recoil and extraction of $|V_{cb}|$* , *Phys. Rev. D* **92** (2015) 054510 [[arXiv:1505.03925](#)] [[INSPIRE](#)].
- [71] LHCb collaboration, *Search for the decays $B_s^0 \rightarrow \tau^+ \tau^-$ and $B^0 \rightarrow \tau^+ \tau^-$* , *Phys. Rev. Lett.* **118** (2017) 251802 [[arXiv:1703.02508](#)] [[INSPIRE](#)].
- [72] C. Bobeth, M. Gorbahn, T. Hermann, M. Misiak, E. Stamou and M. Steinhauser, *$B_{s,d} \rightarrow l^+ l^-$ in the Standard Model with Reduced Theoretical Uncertainty*, *Phys. Rev. Lett.* **112** (2014) 101801 [[arXiv:1311.0903](#)] [[INSPIRE](#)].
- [73] C. Bobeth, *Updated $B_q \rightarrow \bar{\ell} \ell$ in the standard model at higher orders*, in *Proceedings of 49th Rencontres de Moriond on Electroweak Interactions and Unified Theories*, La Thuile Italy (2014), pg. 75 [[arXiv:1405.4907](#)] [[INSPIRE](#)].
- [74] C. Bobeth, M. Misiak and J. Urban, *Photonic penguins at two loops and m_t dependence of $BR[B \rightarrow X_s l^+ l^-]$* , *Nucl. Phys. B* **574** (2000) 291 [[hep-ph/9910220](#)] [[INSPIRE](#)].
- [75] T. Huber, E. Lunghi, M. Misiak and D. Wyler, *Electromagnetic logarithms in $\bar{B} \rightarrow X_s l^+ l^-$* , *Nucl. Phys. B* **740** (2006) 105 [[hep-ph/0512066](#)] [[INSPIRE](#)].
- [76] MUON G-2 collaboration, G.W. Bennett et al., *Final Report of the Muon E821 Anomalous Magnetic Moment Measurement at BNL*, *Phys. Rev. D* **73** (2006) 072003 [[hep-ex/0602035](#)] [[INSPIRE](#)].
- [77] PARTICLE DATA GROUP collaboration, C. Patrignani et al., *Review of Particle Physics*, *Chin. Phys. C* **40** (2016) 100001 [[INSPIRE](#)].
- [78] A. Czarnecki, B. Krause and W.J. Marciano, *Electroweak Fermion loop contributions to the muon anomalous magnetic moment*, *Phys. Rev. D* **52** (1995) R2619 [[hep-ph/9506256](#)] [[INSPIRE](#)].
- [79] A. Czarnecki, B. Krause and W.J. Marciano, *Electroweak corrections to the muon anomalous magnetic moment*, *Phys. Rev. Lett.* **76** (1996) 3267 [[hep-ph/9512369](#)] [[INSPIRE](#)].
- [80] C. Gnendiger, D. Stöckinger and H. Stöckinger-Kim, *The electroweak contributions to $(g-2)_\mu$ after the Higgs boson mass measurement*, *Phys. Rev. D* **88** (2013) 053005 [[arXiv:1306.5546](#)] [[INSPIRE](#)].
- [81] M. Davier, A. Hoecker, B. Malaescu and Z. Zhang, *Reevaluation of the Hadronic Contributions to the Muon $g-2$ and to $\alpha(M_Z)$* , *Eur. Phys. J. C* **71** (2011) 1515 [Erratum *ibid.* **72** (2012) 1874] [[arXiv:1010.4180](#)] [[INSPIRE](#)].
- [82] K. Hagiwara, R. Liao, A.D. Martin, D. Nomura and T. Teubner, *$(g-2)_\mu$ and $\alpha(M_Z^2)$ re-evaluated using new precise data*, *J. Phys. G* **38** (2011) 085003 [[arXiv:1105.3149](#)] [[INSPIRE](#)].
- [83] A. Kurz, T. Liu, P. Marquard and M. Steinhauser, *Hadronic contribution to the muon anomalous magnetic moment to next-to-next-to-leading order*, *Phys. Lett. B* **734** (2014) 144 [[arXiv:1403.6400](#)] [[INSPIRE](#)].
- [84] F. Jegerlehner and A. Nyffeler, *The Muon $g-2$* , *Phys. Rept.* **477** (2009) 1 [[arXiv:0902.3360](#)] [[INSPIRE](#)].

- [85] T. Aoyama, M. Hayakawa, T. Kinoshita and M. Nio, *Complete Tenth-Order QED Contribution to the Muon $g-2$* , *Phys. Rev. Lett.* **109** (2012) 111808 [[arXiv:1205.5370](#)] [[INSPIRE](#)].
- [86] G. Colangelo, M. Hoferichter, A. Nyffeler, M. Passera and P. Stoffer, *Remarks on higher-order hadronic corrections to the muon $g-2$* , *Phys. Lett. B* **735** (2014) 90 [[arXiv:1403.7512](#)] [[INSPIRE](#)].
- [87] BABAR collaboration, B. Aubert et al., *Searches for Lepton Flavor Violation in the Decays $\tau^\pm \rightarrow e^\pm \gamma$ and $\tau^\pm \rightarrow \mu^\pm \gamma$* , *Phys. Rev. Lett.* **104** (2010) 021802 [[arXiv:0908.2381](#)] [[INSPIRE](#)].
- [88] D. Bečirević, N. Košnik, O. Sumensari and R. Zukanovich Funchal, *Palatable Leptoquark Scenarios for Lepton Flavor Violation in Exclusive $b \rightarrow s \ell_1 \ell_2$ modes*, *JHEP* **11** (2016) 035 [[arXiv:1608.07583](#)] [[INSPIRE](#)].
- [89] Y. Cai, J. Gargalionis, M.A. Schmidt and R.R. Volkas, *Reconsidering the One Leptoquark solution: flavor anomalies and neutrino mass*, [arXiv:1704.05849](#) [[INSPIRE](#)].
- [90] BABAR collaboration, J.P. Lees et al., *A search for the decay modes $B^\pm \rightarrow h^\pm \tau^\pm l$* , *Phys. Rev. D* **86** (2012) 012004 [[arXiv:1204.2852](#)] [[INSPIRE](#)].
- [91] A. Crivellin, G. D'Ambrosio, M. Hoferichter and L.C. Tunstall, *Violation of lepton flavor and lepton flavor universality in rare kaon decays*, *Phys. Rev. D* **93** (2016) 074038 [[arXiv:1601.00970](#)] [[INSPIRE](#)].
- [92] R. Aaij et al., *Test of lepton universality with $B^0 \rightarrow K^{*0} \ell^+ \ell^-$ decays*, *JHEP* **08** (2017) 055 [[arXiv:1705.05802](#)] [[INSPIRE](#)].
- [93] B. Capdevila, A. Crivellin, S. Descotes-Genon, J. Matias and J. Virto, *Patterns of New Physics in $b \rightarrow s \ell^+ \ell^-$ transitions in the light of recent data*, [arXiv:1704.05340](#) [[INSPIRE](#)].