

General Disclaimer

One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

**SIMULTANEOUS STABILIZATION AND SIMULTANEOUS
POLE PLACEMENT BY NONSWITCHING DYNAMIC
COMPENSATION (Harvard Univ.) 24 p**
HC A02/BF A01

NS2-31031

CSCL 12A

**Unclas
G3/65 28774**

**SIMULTANEOUS STABILIZATION AND SIMULTANEOUS POLE-PLACEMENT
BY NONSWITCHING DYNAMIC COMPENSATION**

B. K. Ghosh*
Division of Applied Sciences
Harvard University
Cambridge, MA

C. I. Byrnes**
Department of Mathematics and
Division of Applied Sciences
Harvard University
Cambridge, MA



* This work is partially supported by NASA Grant No. NSG-2265.

** This work is partially supported by NASA Grant No. NSC-2265, the National Science Foundation Grant No. ECS-81-21428, and Air Force Grant No. AFOSR-81-0054.

1. INTRODUCTION AND STATEMENT OF THE MAIN RESULTS

The "simultaneous stabilization problem" - in either discrete or continuous time - consists in answering the following question:

Given an r -tuple $G_1(s), \dots, G_r(s)$ of $p \times m$ proper transfer functions, does there exist a compensator $K(s)$ such that the closed-loop systems $G_1(s)(I+K(s)G_1(s))^{-1}, \dots, G_r(s)(I+K(s)G_r(s))^{-1}$ are (internally) stable?

As pointed out in [13], this question arises in reliability theory, where $G_2(s), \dots, G_r(s)$ represents a plant $G_1(s)$ operating in various modes of failure and $K(s)$ is a nonswitching stabilizing compensator. Of course, for the same reason, it is important in the stability analysis and design of a plant which can be switched into various operating modes. The simultaneous stabilization problem can also apply to the stabilization of a nonlinear system which has been linearized at several equilibria. Finally, it has been shown [14], [20] that to solve the case $r=2$ is to solve the well-known problem considered by Youla et al in [21]: When can a single plant be stabilized by a stable compensator? This correspondence also serves to give some measure of the relative depth of this problem.

In order to describe the results obtained via this correspondence, we need some notation. First, set $n_i =$ McMillan degree of $G_i(s)$. In the scalar input-output setting ($m=p=1$), we regard each $G_i(s)$ as a point in \mathbb{R}^{2n_i+1} , viz. if

$$G_i(s) = p_i(s)/q_i(s), \text{ where}$$

$$p_i(s) = a_{0i} + \dots + a_{n_i i} s^{n_i}, \text{ and } q_i(s) = b_{1i} + \dots + b_{n_i i} s^{n_i-1} + s^{n_i}$$

then $G_1(s)$ corresponds to the vector $(a_{01}, \dots, a_{n_1 1}, b_{11}, \dots, b_{n_1 1}) \in \mathbb{R}^{2n_1+1}$.

Moreover, since p_1 and q_1 are relatively prime, this vector lies in the open dense set $\text{Rat}(n_1) \subset \mathbb{R}^{2n_1+1}$ (see [3] for the strictly proper case). In [14], Sacks and Murray used the techniques of fractional representations [8] and the correspondence mentioned above to give explicit inequalities defining the open set

$$U \subset \text{Rat}(n_1) \times \text{Rat}(n_2)$$

of pairs $(G_1(s), G_2(s))$ which are simultaneously stabilizable. In [20] Vidyasagar and Viswanadham showed, using similar techniques, that provided $\max(m, p) > 1$ the open set U of pairs $(G_1(s), G_2(s))$ which can be stabilized is in fact dense.

This can be made precise by topologizing a point $G_1(s)$ in the set

$$\sum_{m,p}^n = \{p \times m G_1(s) ; \text{degree } G_1(s) = n_1\}$$

as a vector in $\mathbb{R}^{(n_1+1)(mp)}$ via its Hankel parameters: If

$$G_1(s) = \sum_{j=0}^{\infty} H_{1j} s^{-j}$$

then $G_1(s)$ corresponds to the $n+1$ $p \times m$ block matrices $\{H_{10}, \dots, H_{1, n+1}\}$ which determines $G(s)$. It is known that $\sum_{m,p}^n$ is an $(n(m+p)+mp)$ -manifold (see [7], [12], [5]), although this is not important here. What is important is that $\sum_{m,p}^n$ is a topological space.

One of our main results concerns the generic stabilizability problem; that is,

Question 1.1. Fix m, p, r , and n_1 . Is the set U of r -tuples $G_1(s), \dots, G_r(s)$ which can be simultaneously stabilized open and dense in $\sum_{m,p}^{n_1} \times \dots \times \sum_{m,p}^{n_1}$?

It is also important to ask, for reasons of global robustness of algorithms finding such a compensator, for compensators with a fixed degree of complexity.

Question 1.2. Fix m, p, r , and n_i . What is the minimal value of q (if one exists) for which the set W_q of r -tuples which can be simultaneously stabilized, by a compensator of degree $\leq q$, is open and dense in $\sum_{m,p}^{n_1} \times \dots \times \sum_{m,p}^{n_r}$?

It should be noted that, in the case $r=1$, Question 1.2 is an outstanding, unsolved, classical problem. In this paper, we prove:

Theorem 1.1. In either discrete or continuous time, a sufficient condition for generic simultaneous stabilizability is

$$\max(m,p) \geq r \tag{1.1}$$

Indeed, if (1.1) holds, then the generic r -tuple can be stabilized by a compensator of degree less than or equal to q , where q satisfies:

$$q[\max(m,p)+1-r] \geq \sum_{i=1}^r n_i - \max(m,p) \tag{1.2}$$

In the case $r=1$, it is unknown whether generic stabilizability implies generic pole-assignability; that is, whether or not these properties of m, n , and p are really different (see [4]). Perhaps not surprisingly then, Theorem 1.1 follows from:

Theorem 1.2. A sufficient condition for generic simultaneous pole-assignability is (1.1), where the compensator $K(s)$ can be taken to be of degree q satisfying (1.2).

Here, simultaneous pole-assignability means the assignability of r sets of self-conjugate sets of numbers $\{s_{1i}, \dots, s_{n_1+q, i}\} \subset \mathbb{C}$. In fact sharper bounds on q can be obtained (see [18], [11]). Our proof relies on the recent pole-placement techniques derived for $r=1$ by P.K. Stevens in his thesis [18], which contains an improvement on existing results in the literature, see also [9], [17]. We shall prove Theorem 1.2 only in the strictly proper case; the proper case involves more technical arguments from algebraic geometry which can be found in [11]. We shall, however, give an independent proof of Theorem 1.1 in the nonstrictly proper case, based on the equivalence of generic stabilizability and existence of a solution to a generic "deadbeat control" problem, which we can solve if (1.1) is satisfied. This argument extends the argument given in [4] for the case $r=1$ and $q=0$.

Note that if $r=1$, then (1.1) is always satisfied in which case (1.2) implies

Corollary 1.3. (Brasch-Pearson [2]). The generic $p \times m$ plant $G(s)$ of degree n can be stabilized by a compensator of order q , where q satisfies

$$(q+1)\max(m,p) \geq n \quad (1.3)$$

If $r=2$ and $\max(m,p) > 1$, then (1.1) is again satisfied, so we obtain rather easily:

Corollary 1.4. (Vidyasagar-Viswanadham [20]). If $r=2$ and $\max(m,p) > 1$, then the generic pair $(G_1(s), G_2(s))$ is simultaneously stabilizable.

Moreover, in this case we know an upper bound on the order of the required compensator. For example, if $m=p=2$, $r=2$, then q can be taken to satisfy

$$q \geq n_1 + n_2 - 2$$

On the other hand, in [20] the explicit conditions defining the closed set

$$\sum_{m,p}^{n_1} \times \sum_{m,p}^{n_2} - U$$

of pairs not simultaneously stabilizable were derived. Such conditions can be derived from our proof, but instead we refer to [10], where Theorem 1.1 (excepting (1.2)) is proved by interpolation methods also yielding a set of explicit conditions in the range $r \leq \max(m,p)$.

Finally, we prove that the condition (1.1) is sharp in the following sense.

Theorem 1.5. If $\min(m,p) = 1$, then for fixed m,p,r and n_i the following statements are equivalent for proper plants:

- (i) $q \in \mathbb{N}$ satisfies $q(\max(m,p) + 1 - r) + \sum_{i=1}^r n_i$;
- (ii) the generic r -tuple $(G_1(s), \dots, G_r(s))$ is simultaneously stabilizable in discrete or continuous time by a compensator of degree $\leq q$;
- (iii) the generic r -tuple $(G_1(s), \dots, G_r(s))$ is simultaneously stabilizable in discrete or continuous time.

In the strictly proper case it follows that (i)-(iii) is also equivalent to generic simultaneous pole assignability. This holds in the proper case as well, but requires a separate argument [11].

Corollary 1.6. If $\min(m,p) = 1$ and $r \leq \max(m,p)$ then the generic r -tuple is simultaneously stabilizable by a compensator of order precisely given by the least integer q satisfying (1.2).

As a further corollary, we obtain one of the results obtained by Saks and Murray in [], see also [15]:

Corollary 1.7. (Saks-Murray). Suppose $m=p=1$ and $r=2$. Simultaneous stabilizability is not a generic property.

We remark that these results hold also over the field \mathbb{C} of complex numbers - in particular, the complex analogue of Corollary 1.7 dispels a folklore conjecture concerning simultaneous stabilization using compensators with complex coefficients.

Finally, over any field, the method of proof of Theorem 1.2 gives linear equations for a compensator simultaneously placing $r(n+q)$ poles when the generic hypothesis is satisfied.

2. POLE PLACEMENT AND THE GENERALIZED SYLVESTER MATRIX: A PROOF OF THEOREM 1.2

In this section we proceed to prove Theorem 1.2. Note that Theorem 1.1 and Corollaries 1.3 and 1.4 follow immediately in the strictly proper case from this theorem. Without any loss of generality we can assume that $m \geq p$, for, if $K(s)$ stabilizes $G_1^t(s)$ then $K^t(s)$

ORIGINAL PAGE IS
OF POOR QUALITY

stabilizes $G_1(s)$.

Suppose, first of all, that $p=1$, so that we are given a set of r, m input 1 output plants of McMillan degree $\leq n$ represented as

$$\left[\begin{array}{c} \frac{\sum_{i=0}^n p_{1i}^k s^i}{\sum_{i=0}^n p_{m+pi}^k s^i} \\ \frac{\sum_{i=0}^n p_{2i}^k s^i}{\sum_{i=0}^n p_{m+pi}^k s^i} \\ \dots \\ \frac{\sum_{i=0}^n p_{m+p-1,i}^k s^i}{\sum_{i=0}^n p_{m+p,i}^k s^i} \end{array} \right] \quad (2.1)$$

for $k=1,2,\dots,r$. A 1 input, m output, compensator of McMillan degree $\leq q$ is represented as

$$\left[\begin{array}{c} \frac{\sum_{i=0}^q a_{1i} s^i}{\sum_{i=0}^q a_{m+pi} s^i} \\ \frac{\sum_{i=0}^q a_{2i} s^i}{\sum_{i=0}^q a_{m+pi} s^i} \\ \dots \\ \frac{\sum_{i=0}^q a_{m+p-1,i} s^i}{\sum_{i=0}^q a_{m+p,i} s^i} \end{array} \right] \quad (2.2)$$

Note that in (2.1) and (2.2) the coefficients $p_{ji}^k \forall k$ and a_{ji} has been defined up to a nonzero scale factor. Moreover, for a strictly proper plant or compensator, $p_{jn_i}^k = 0, a_{jq} = 0 \forall j = 1, \dots, m+p-1; k=1, \dots, r$.

The associated return difference equation, $\det(I+K(s)G_k(s)) = 0$ is given by

$$\Pi_k(s) = \sum_{j=1}^{m+p} \left[\sum_{i=0}^n p_{ji}^k s^i \right] \left[\sum_{i=0}^q a_{ji} s^i \right] \quad (2.3)$$

$$\forall k = 1, 2, \dots, r$$

A generic r -tuple of plants define a mapping χ , via equation (2.3), between the plant parameters and the coefficient of the return difference polynomials given by

$$\chi : \mathbb{R}^{(q+1)(m+p)} \rightarrow \mathbb{R}^{r(n+q+1)} \quad (2.4)$$

where

ORIGINAL PAGE IS
OF POOR QUALITY

$$\chi(A_0, \dots, A_q) = (A_0, \dots, A_q) \begin{bmatrix} P_0 & \dots & \dots & \dots & P_n \\ & P_0 & & & P_n \\ & & \cdot & & \\ & & & P_0 & \\ & & & & P_n \end{bmatrix} \quad (2.5)$$

where

$$A_i = (a_{1i}, a_{2i}, \dots, a_{m+pi}) \quad (2.6)$$

$$P_i = \begin{bmatrix} 1 & 1 & \dots & \dots & P_{1i}^r \\ P_{1i}^1 & P_{1i}^2 & \dots & \dots & P_{1i}^r \\ 1 & 2 & \dots & \dots & P_{2i}^r \\ P_{2i}^1 & P_{2i}^2 & \dots & \dots & P_{2i}^r \\ \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots \\ 1 & 2 & \dots & \dots & P_{m+pi}^r \\ P_{m+pi}^1 & P_{m+pi}^2 & \dots & \dots & P_{m+pi}^r \end{bmatrix} \quad (2.7)$$

The matrix in the right hand side of (2.5) is classically known as the generalized Sylvester matrix and is of order $(q+1)(m+p) \times r(n+q+1)$. For $r=1$ its rank has been analyzed by Bitmead, Kailath, Kung in [1]. In particular, for a generic plant, it is known to have full rank. For $r \geq 1$, we have the following:

Lemma 2.1. The generalized Sylvester matrix is of full rank for a generic r -tuple.

Proof: See Appendix I.

ORIGINAL PAGE IS
OF POOR QUALITY

Lemma 2.2. Assume $\min(m,p) = 1$. A sufficient condition for generic pole assignment, for an r tuple of strictly proper plants by a proper compensator is given by

$$(q+1)(m+p-r) \geq \sum_{i=1}^r n_i - r + 1 \quad (2.8)$$

Proof: We prove this Lemma assuming for notational convenience that $n_i = n \forall i=1, \dots, r$ and analyze the mapping χ as defined by (2.4), (2.5). Assume

$$a_{m+p,q} = -1, \quad p_{m+p,n}^k = 1 \forall k=1, \dots, r$$

and that the coefficient of s^{n+q} in all the r return difference polynomials (2.3) has been normalized to 1.

Thus a sufficient condition for generic pole assignment is that χ' is onto. Here the mapping

$$\chi' : \mathbb{R}^{(q+1)(m+p)-1} \rightarrow \mathbb{R}^{(n+q)} \quad (2.9)$$

is given by

$$\chi(A_0, \dots, A_{q-1}, A'_q) = (A_0, \dots, A_{q-1}, A'_q) \begin{bmatrix} p_0 & \dots & \dots & \dots & p_n \\ & p_0 & \dots & \dots & p_n \\ & & \ddots & & \\ & & & p_0 & \dots & \dots & p_n \\ & & & & p'_0 & \dots & \dots & p'_n \end{bmatrix} \quad (2.10)$$

where

$$A'_q = (a_{1q}, a_{2q}, \dots, a_{m+p-1q})$$

and p'_i is obtained from p_i by deleting its $(m+p)^{\text{th}}$ row.

By Lemma 2.1 the matrix in the right-hand side of (2.10) is of full rank for a generic r -tuple of plants, and has the order $(q+1)(m+p-1) \times r(n+q)$. Therefore, a sufficient condition for generic pole placement is given by

$$(q+1)(m+p) - 1 \geq r(n+q) \quad (2.11)$$

which is same as (2.8) for $n_i = n \forall i=1, \dots, r$.

The proof of Theorem 1.2 now proceeds by a reduction to the case $\min(m,p) = 1$, which has been treated in Lemmas 2.1-2.2. This procedure, which is called "vectoring down", is adopted from the case $r=1$, studied in P.K. Stevens thesis [18].

Lemma 2.3. Given an r -tuple of $p \times m$ plants $G_i(s)$ of degrees n_i , each with distinct simple poles, there is an open dense set of $1 \times p$ vectors $v \in \mathbb{R}^p$ such that $vG_i(s)$ has degree n_i .

Proof: If $r=1$, then we may expand $G(s)$

$$G(s) = \sum_{i=1}^n \frac{R_i}{s-\lambda_i}$$

in a partial fraction expansion, where $\lambda_i \in \mathbb{C}$ and each R_i has rank 1. Now, the set U_1 of real vectors v such that $vR_1 \neq 0$ is clearly open and dense in \mathbb{R}^p . Defining U_2, \dots, U_n similarly, set

$V = \bigcap_{i=1}^n U_i$. Thus, V is an open dense set of vectors with the required property.

If $r > 1$, one obtains, as above, sets V_1, \dots, V_r in \mathbb{R}^p having an open dense intersection $\bigcap_{i=1}^r V_i$. Q.E.D.

Lemma 2.4. Given an r -tuple of $p \times m$ plants $G_i(s)$ there exists a constant gain output feedback k such that the closed loop systems $G_i(s)(I + kG_i(s))^{-1}$ have distinct simple poles.

Proof: For $r=1$, the set W_1 of K such that the closed loop system has simple poles is the complement in \mathbb{R}^{mp} of an algebraic set. It is well known [2] that this set is nonempty; therefore, W_1 is open and dense. Taking any K in the open dense set $\bigcap_{i=1}^r W_i$ gives the desired conclusion. Q.E.D.

Thus, choosing any $(v, K) \in \mathbb{R}^p \times \mathbb{R}^{mp}$ we have a mapping from an open dense set

$$\phi(v, k) : \sum_{m,p}^n \times \dots \times \sum_{m,p}^n \rightarrow \sum_{m,1}^n \times \dots \times \sum_{m,1}^n$$

$$\phi(v, k) (G_i(s))_{i=1}^r = (vG_i(s)(\phi + KG_i(s))^{-1})_{i=1}^r$$

which is rational in the Hankel parameters (H_{ij}) of (G_i) . Applying Lemmas 2.1-2.2 to the case $\min(m,p)=1$, i.e. $\sum_{m,1}^n \times \dots \times \sum_{m,1}^n$, gives - via composition with ϕ - an open dense set of $\sum_{m,p}^n \times \dots \times \sum_{m,p}^n$ which can be simultaneously pole-assigned. Q.E.D.

3. GENERIC STABILIZABILITY CONDITION OF AN r -TUPLE OF PROPER PLANTS

In this section we proceed to prove Theorem 1.1 independent of Theorem 1.2. We first show that the following three statements are equivalent.

I. A generic r -tuple of proper plants is stabilizable with respect to the open left half plane by a proper compensator of degree $\leq q$.

II. A generic r -tuple of proper plants is stabilizable with respect to the interior of the unit disc, by a proper compensator of degree $\leq q$.

III. A generic r -tuple of proper plants is pole assignable at the origin by a proper compensator of degree $\leq q$.

Lemma 3.1. I \Leftrightarrow II

Proof: Consider the conformal transformation

$$\phi(s) = (s+1)/(s-1) \tag{3.1}$$

which maps the r -tuple of proper plants g_1, g_2, \dots, g_r onto the r -tuple of proper plants g'_1, \dots, g'_r where $g'_i(s) = g_i(\phi(s))$ except for the algebraic set of plants satisfying - " $g_i(s)$ has a pole at $s=1$ for some $i=1, \dots, r$ ". The proof now follows from the two facts.

1. $\phi(s)$ maps the open left half plane onto the interior of the unit disc.

2. The mapping

$$(g_1, \dots, g_r) \mapsto (g'_1, \dots, g'_r)$$

and its inverse, map the generic r -tuple of proper plants to the generic r -tuple of proper plants.

Lemma 3.2. II \Leftrightarrow III

Proof: Sufficiency is clear and follows by an analagous argument of Lemma 3.1 with $\phi(s) = s + a$, $a > 0$, $a \in \mathbb{R}$.

To prove necessity, we have the following: For each $r = 1, 2, \dots$ (shown easily by assuming statement II and considering $\phi(s) = as$, $a > 0$, $a \in \mathbb{R}$). \exists an open dense set of U_r of r -tuple of plants for which there exist a compensator of degree $\leq q$ which places the poles in the interior of the disc D_r of radius $1/r$ centered at the origin. Consider the set

$$U = \bigcap_{r=1}^{\infty} U_r$$

Clearly, U is a dense set by the Baire Category Theorem [13]. Since the mapping χ given by (2.4) is linear, it has a closed image. Moreover, every r -tuple of plants in U admits a sequence of compensators which places the poles arbitrary close to the origin. Since image of χ is closed, U is contained in a set V of all r -tuple of plants for which there exists a compensator which places the poles at the origin. By the Tarski [19]-Seidenberg [16] theory of elimination over \mathbb{R} , V is indeed defined by union and/or intersection of sets given by polynomial equations or inequations $f_{\alpha} > 0$, $f_{\beta} = 0$. Finally, since U is dense in V , $f_{\beta}(U) = 0 \Rightarrow f_{\beta} \equiv 0$ so that V is defined by strict polynomial inequalities. Hence V is open. Moreover, since U is dense, V is also dense.

Lemma 3.3. For a generic r -tuple ($r \leq m+p$) of $\min(m,p) = 1$ plants

$$\text{III} \Leftrightarrow (q+1)(m+p-r) \geq r(n-1)+1$$

Proof: The only nontrivial part is to prove sufficiency for the case

$$r(n+q) < (q+1)(m+p) < r(n+q+1)$$

(The other cases follow easily from the fact that the associated Sylvester's matrix is of full rank for a generic r-tuple.)

To prove sufficiency, for the above case we want to show that the vector

$$(0, 0, \dots, 0, s_1, s_2, \dots, s_r)$$

$$\longleftarrow r(n+q) \qquad \longleftarrow r \qquad \longrightarrow$$

indeed belongs to the image of χ (defined by (2.5)) for some $s_i \neq 0, i=1, \dots, r$.

Partition the Sylvester's matrix in (2.5) as $[S_1 S_2]$ where S_1 is of order $(q+1)(m+p) \times r(n+q)$. Clearly we are solving the pair of equations

$$[A_0, \dots, A_q] S_1 = [0, \dots, 0] \tag{3.2}$$

$$[A_0, \dots, A_q] S_2 = [s_1, \dots, s_r] \tag{3.3}$$

We claim that for a generic r-tuple of plants (3.2) has a solution for a nonzero vector A_q for otherwise if $A_q = 0$ we have

$$[A_0, \dots, A_{q-1}] S'_1 = (0, \dots, 0) \tag{3.4}$$

where S'_1 is of order $q(m+p) \times r(n+q)$ obtained by deleting the last $m+p$ rows of S_1 . From (3.4) $(A_0, \dots, A_{q-1}) = \underline{0}$ since S'_1 is of full rank generically. Thus the only solution of (3.2) is the zero vector which is a contradiction since the kernel of S_1 is at least of dimension 1. On the other hand, for $A_q \neq 0$, for a generic r-tuple of plants the right-hand side of (3.3) is a vector none of whose entries are zero.

Theorem 1.1 then follows from Lemma 3.1, 3.2, 3.3 and the vectoring down technique used in the proof of Theorem 1.2 in Section 2.

4. PROOF OF THEOREM 1.5

To say there exists $q \in \mathbb{N}$ satisfying (1.2) is to $\max(m, p) >$
Thus, (ii) follows from (i) by Theorem 1.1.

(ii) \Rightarrow (iii) since (iii) is weaker than (ii).

By Lemma 3.1, in order to prove (iii) \Rightarrow (i) it suffices to assume that $G_1(s), \dots, G_r(s)$ are simultaneously stabilizable in continuous-time.

Proposition 4.1. The generic $(m+1)$ -tuple of $1 \times m$ proper continuous time plants of degree n is not simultaneously stabilizable by a proper compensator of finite (but not a priori bounded) degree.

Proof: Consider the domain of (simultaneous) stability

$$\mathcal{D} = \{(c_{11}, \dots, c_{1n}, \dots, c_{r, n_r}) : \sum_{j=0}^{n_i+q} c_{i,j} s^j \text{ has all roots in } D_1\}$$

and its convex hull $\Omega(\mathcal{D}) \subset \mathbb{R}^{n_1+q} \times \dots \times \mathbb{R}^{n_r+q}$. Clearly, a necessary condition for generic simultaneous stabilizability is

$$\text{image}(\chi_\eta) \cap \Omega(\mathcal{D}) \neq \emptyset,$$

for an open dense set of η . Since

ORIGINAL PAGE IS
OF POOR QUALITY

$$\Omega(\mathcal{D}) = \{(c_{ij}) : c_{ij} > 0\}$$

it will suffice to prove:

Lemma 4.2. If $r = m + p$, then there exists an open set of r -tuples η such that image (χ_η) contains no vector with only positive entries.

We fix the value of q and construct the associated Sylvester matrix S . We claim that the open set E of plants defined by

$$E \triangleq \{(P_0, P_1, \dots, P_n) \mid P_0^{-1} P_j \forall j = 1, \dots, n \text{ has all the entries negative}\}$$

cannot be stabilized by a proper compensator of degree $\leq q$.

Suppose the above is not true, then there exist $\eta \in E$, such that

$$\text{image}(\chi_\eta) \cap \Omega(\mathcal{D}) \neq \emptyset$$

or in other words $\exists \underline{\alpha} \mid \alpha_i > 0 \forall i = 1, r(n+q+1)$ and

$$\underline{a} S = \underline{\alpha} \tag{4.1}$$

has a solution. Writing S as

$$S \triangleq [S' \quad S'']$$

where

$$S' = \begin{bmatrix} P_0 & P_1 & \dots & P_q \\ 0 & P_0 & \dots & P_{q-1} \\ \dots & \dots & \dots & \dots \\ 0 & 0 & & P_0 \end{bmatrix} \tag{4.2}$$

and $P_j = 0$ for all $j > n$

ORIGINAL PAGE IS
OF POOR QUALITY

Equation (4.1) can be written as

$$\underline{a}' [I \mid S'^{-1} S''] = \underline{\alpha} \quad (4.3)$$

where S'^{-1} is given as follows

$$S'^{-1} = \begin{bmatrix} X_0 & X_1 & \dots & X_q \\ & X_0 & \dots & X_{q-1} \\ \dots & \dots & \dots & \dots \\ & & & X_0 \end{bmatrix}$$

where $X_0 = P_0^{-1}$

$$-(P_1, P_2, \dots, P_{r+1}) \begin{bmatrix} X_r \\ X_{r-1} \\ \vdots \\ X_1 \end{bmatrix} = X_{r+1} \quad \forall r = 0, \dots, q-1$$

$$P_j = P_0^{-1} P_j \quad ; \quad j = 1, \dots, q$$

The identity matrix of order $(q+1)(m+p)$ in (4.3) forces \underline{a}' to have all the entries positive. Moreover, since $\eta \in E, S'^{-1} S''$ has all its entries negative so that $\underline{a}'(S'^{-1} S'')$ has all the entries negative which is a contradiction since $\underline{\alpha}$ is a positive vector.

Finally it is shown that E is not an empty set. For a fixed $P_0 = P_0^*$ choose the vector δ to be so that $P_0^{*-1} \delta$ has all its entries negative. Let

$$P_j^* = (\delta, \delta, \dots, \delta) \quad j = 1, \dots, n$$

← m+p →

ORIGINAL PAGE IS
OF POOR QUALITY

so that

$$(P_0^*, P_1^*, \dots, P_n^*) \in E$$

Q.E.D.

Remark: If $\text{image}(\chi_\eta)$ is affine hyperplane, then the necessary condition

$$\text{image}(\chi_\eta) \cap \Omega(\mathcal{D}) \neq \emptyset$$

of course is sufficient, i.e. implies

$$\text{image}(\chi_\eta) \cap \mathcal{D} \neq \emptyset$$

This fact was used by Chen, together with

Lemma 4.3. (Chen [6]) If $r=1$, $\Omega(\mathcal{D}) = \{(c_1, \dots, c_n) : c_i > 0\}$

to give precise conditions for stabilizability in the case $r=1$, $q=0$, $\min(m,p)=1$, and $\max(m,p)=n-1$. This technique can be adapted in the cases $r \geq 1$ to give explicit conditions - in certain cases - defining the open set of simultaneously stabilizable plants when $r > \max(m,p)$, see [11].

Note that Corollary 1.6 now follows from our previous results on the generic rank of the generalized Sylvester matrix, while Corollary 1.7 follows either from Theorem 1.5 or Proposition 4.1.

ORIGINAL QUALITY
OF POOR QUALITY

APPENDIX I: PROOF OF LEMMA 2.1

The generalized Sylvester matrix is co-ordinatized by $r(n+1)(m+p)$ parameters, and it is sufficient to show the existence of one principal minor with nonvanishing determinant.

By reordering the rows and columns, the generalized Sylvester matrix can be written as

$$\bar{S} = [Q_1, Q_2, \dots, Q_{m+p}]^T \quad (1)$$

where

$$Q_i = [P_{i1}, P_{i2}, \dots, P_{ir}] \quad (2)$$

$$P_{jk} = \begin{bmatrix} P_{j0}^k & P_{j1}^k & \dots & P_{jn}^k & & 0 \\ & \ddots & \ddots & \ddots & \ddots & \\ & & 0 & P_{j0}^k & P_{j1}^k & \dots & P_{jn}^k \end{bmatrix} \begin{matrix} \updownarrow \\ (q+1) \\ \updownarrow \end{matrix} \quad (3)$$

← (n+q+1) →

in the notation of (2.7). Moreover, each p_{jk} is referred to as a 'block' of S .

Define a set M of matrices as follows: "m belongs to M provided m is obtainable from one of the matrices p_{jk} in (3) either by deleting the first α_1 columns or the last α_2 rows $\alpha_1, \alpha_2 > 0$."

Proposition A.1. Every element m of M has the property that there exists a principal minor $m_p \in M$ of m , a coordinate p_m^* and an integer j_m^* such that $p_m^{*j_m^*}$ is a summand in $\det m_p$ where j_m^* is the order of the minor.

Proof: Clear from the structure of p_{jk} .

The following is an algorithm to construct a principal minor with nonidentically-vanishing determinant.

Algorithm:

Set $S = \bar{S}$, Initialize $\xi = 0$

1. Set $\xi = \xi + 1$.
2. Look at P_{11} . Obtain the principal minor m of P_{11} , satisfying Proposition 1. If there is more than one possible choice, choose the one containing the first column. Define $\alpha_{\xi} = p_m^*$ and $j_{\xi} = j_m^*$.
3. Delete the rows and columns corresponding to the coordinate p_m^* from S . Renumber the blocks of the resulting matrix and call it S . (Every block of S is to be identified as a minor of the corresponding block in \bar{S} obtained by row or column deletion.)
4. Do the same "delete" operation as in step 3 in \bar{S} .
5. If S is empty, terminate. Otherwise go to 6.
6. Set $k = \xi$.
Construct the principal minor \bar{m}_p of \bar{S} by choosing those elements of \bar{S} whose corresponding row and column has been deleted in Step 6.

Proposition A.2. During the execution of the above algorithm, S can always be decomposed into blocks belonging to M .

Proof: Clearly \bar{S} satisfies the above proposition, since each block p_{jk} belongs to M . Each iteration of the algorithm deletes either the

first α_1 columns of the first block column of S or the last α_2 rows of the first block row of S . The proposition thus follows from the definition of M .

Proposition A.3. \bar{m}_p constructed in Step 6 of the algorithm has a nonidentically-vanishing determinant.

Proof: We prove the proposition by showing that $\det \bar{m}_p$ has a summand given by $\prod_{i=1}^k \alpha_i^{j_i}$, in the notation of the algorithm. This is clear, however, by observing that in the ξ^{th} iteration the matrix S has a principal minor, the determinant of which has the summand $\prod_{i=\xi}^k \alpha_i^{j_i}$, where k is defined in Step 6 of the algorithm.

BIBLIOGRAPHY

- [1] R.R. Bitmead, S.Y. Kung, B.D.O. Anderson and T. Kailath, "Greatest Common Divisors via Generalized Sylvester and Bezout Matrices," IEEE Trans. Aut. Contr. AC-23 (1978) 1043-1047.
- [2] F.M. Brasch and J.B. Pearson, "Pole Placement Using Dynamic Compensator," IEEE Trans. Aut. Contr. AC-15 (1970) 34-43.
- [3] R.W. Brockett, "Some Geometric Questions in the Theory of Linear Systems," IEEE Trans. on Aut. Contr. AC-21 (1976) 449-455.
- [4] C.I. Byrnes and B.D.O. Anderson, "Output Feedback and Generic Stabilizability," submitted to SIAM J. of Control.
- [5] C.I. Byrnes and N.E. Hurt, "On the Moduli of Linear Dynamical Systems," Adv. in Math, Suppl. Series Vol. 4 (1978) 83-122; also in Modern Mathematical System Theory, MIR Press, Moscow, 1978 (in Russian).
- [6] R. Chen, Ph.D. Dissertation, University of Florida, 1979.
- [7] J.M.C. Clark, "The Consistent Selection of Local Coordinates in Linear System Identification," Proc. JACC, Purdue, 1976, 576-580.
- [8] C.A. Desoer, R.W. Liu, J.J. Murray and R. Saeks, "Feedback System Design: The Fractional Representation Approach to Analysis and Synthesis," IEEE Trans. Aut. Contr. AC-25 (1980) 401-412.
- [9] T.E. Djaferis, "Generic Pole Assignment Using Dynamic Output Feedback," preprint.
- [10] B.K. Ghosh, "Simultaneous Stabilization and Its Connection with the Problem of Interpolation by Rational Functions," submitted for publication.
- [11] B.K. Ghosh, Ph.D. Dissertation (under preparation).
- [12] M. Hazewinkel and R.E. Kalman, "On Invariants, Canonical Forms and Moduli for Linear Constant Finite-Dimensional Dynamical Systems," in Lecture Notes in Econo. - Math. System Theory Vol. 131, Springer-Verlag, Berlin, 1976, 48-60.
- [13] H.L. Royden, Real Analysis, MacMillan Company, N.Y., 1968.
- [14] R. Saeks and J.J. Murray, "Fractional Representation, Algebraic Geometry, and the Simultaneous Stabilization Problem," IEEE Trans. Aut. Contr. AC-27 (1982) 895-903.

- [15] R. Saeks, J.J. Murray, O Chua and C. Karmokolias, "Feedback System Design: The Single Variate Case," unpublished notes, Texas Tech Univ. 1980.
- [16] A. Seidenberg, "A New Decision Method for Elementary Algebra," Ann. Math 60 (1954) 365-374.
- [17] H. Seraji, "Design of Pole Placement Compensators for Multivariable Systems," Automatica 16 (1980) 335-338.
- [18] P.K. Stevens, Algebro-Geometric Methods for Linear Multivariable Feedback Systems, Ph.D. Dissertation, Harvard University, 1982.
- [19] A. Tarski, "A Decision Method for Elementary Algebra and Geometry," Berkeley note, 1951.
- [20] M. Vidyasagar and N. Viswanadham, "Algebraic Design Techniques for Reliable Stabilization," to appear.
- [21] D. Youla, J. Bongiorno and C. Lu, "Single Loop Feedback Stabilization of Linear Multivariable Dynamic Plant," Automatica 10 (1974) 155-173.