

Simultaneous Topography-guided PRK Followed by Corneal Collagen Cross-linking for Keratoconus

George D. Kymionis, MD, PhD; Georgios A. Kontadakis, MD, MSc; George A. Kounis, PhD; Dimitra M. Portaliou, MD; Alexandra E. Karavitaki, MD; Michael Magarakis, MD; Sonia Yoo, MD; Ioannis G. Pallikaris, MD, PhD

ABSTRACT

PURPOSE: To present the results after simultaneous photorefractive keratectomy (PRK) followed by corneal collagen cross-linking (CXL) for progressive keratoconus.

METHODS: Ten patients (14 eyes) with progressive keratoconus were prospectively treated with customized topography-guided PRK with the Pulzar Z1 (wavelength 213 nm, CustomVis) immediately followed by corneal collagen CXL with the use of riboflavin and ultraviolet A irradiation.

RESULTS: Mean follow-up was 10.69 ± 5.95 months (range: 3 to 16 months). Mean preoperative spherical equivalent refraction (SE) was -3.03 ± 3.23 diopters (D) and defocus was 4.67 ± 3.29 D; at last follow-up SE and defocus were statistically significantly reduced to -1.29 ± 2.05 D and 3.04 ± 2.53 D, respectively ($P < .01$). Preoperative mean (logMAR) uncorrected visual acuity was 0.99 ± 0.81 and best spectacle-corrected visual acuity was 0.21 ± 0.19 , which improved postoperatively to 0.16 ± 0.15 and 0.11 ± 0.15 , respectively. The mean steepest keratometry was reduced from 48.20 ± 3.40 D preoperatively to 45.13 ± 1.80 D at last follow-up.

CONCLUSIONS: Simultaneous PRK followed by CXL seems to be a promising treatment capable of offering functional vision in patients with keratoconus. [*J Refract Surg.* 2009;25:xxx-xxx.]

doi:10.3928/1081597X-20009

Keratoconus is a progressive corneal disorder that can lead to severe vision deterioration through the development of irregular astigmatism and corneal scarring. Thus far, techniques that have been employed to improve vision in patients with keratoconus include the use of spectacles, rigid gas permeable contact lenses, intracorneal ring segment implantation,^{1,2} and, in the most severe cases, lamellar or penetrating keratoplasty.

It has been shown that the progression of the disorder can be impeded by riboflavin-ultraviolet A (UVA) corneal collagen cross-linking (CXL).^{3,4} The application of CXL is capable of stabilizing the cornea in keratoconus and in postoperative refractive ectasia for up to 5 years after the procedure, but topography and visual outcomes of treated patients show only minimal improvement.⁵

The irregular astigmatism in keratoconus can be confronted with the use of customized laser ablations that restore the refractive properties of the anterior corneal surface. The application of customized topography-guided surface ablation has been reported in patients with stable or subclinical keratoconus with promising visual outcomes.⁶⁻⁸ The disadvantage of ablative procedures is that tissue removal might lead to further destabilization of corneal biomechanics and progression of the ectatic disorder. Until recently, the combination of CXL with laser ablation has been reported only as a two-step procedure of CXL with consequent photorefractive keratectomy (PRK).⁹

In this case series, we present a new technique and the results of simultaneous customized topography-guided surface ablation followed by CXL in patients with keratoconus.

From the Institute of Vision and Optics University of Crete, Greece (Kymionis, Kontadakis, Kounis, Portaliou, Karavitaki, Magarakis, Pallikaris); and Bascom Palmer Eye Institute, Cornea and External Diseases, Miami, Fla (Yoo).

Dr Kymionis was supported by a grant from the Hellenic Society of Intraocular Implant and Refractive Surgery.

The authors have no financial or proprietary interest in the materials presented herein.

Correspondence: George D. Kymionis, MD, PhD, Institute of Vision and Optics, University of Crete, Medical School, 71003 Heraklion, Crete, Greece. Tel: 302810394656; Fax: 30281039465; E-mail: kymionis@med.uoc.gr

TABLE 1
Attempted Correction and Maximum Ablation Depth* in 14 Eyes That Underwent Topography-guided PRK and Cross-linking

Patient No.	Attempted Correction (D)	Ablation Depth (μm)
1	-1.00	34
2	-1.75	22
3 (OD)	-1.00	31
3 (OS)	0.50	15
4 (OD)	-2.00	46
4 (OS)	-1.00	30
5	-0.50	20
6	-0.50	25
7	-3.25	46
8	-1.00	36
9	-1.50	30
10	-0.25	15
11	-2.75	50
12	-1.50	16

*Ablation zone 4.5 to 6.0 mm.

PATIENTS AND METHODS

In this pilot study, 10 patients (14 eyes) with progressive keratoconus were included. Mean patient age was 28 years (range: 20 to 39 years). Inclusion criteria were progressive keratoconus, hard contact lens and full spectacle correction intolerance, expected central corneal thickness (CCT) after PRK $>400 \mu\text{m}$, and no other corneal pathological signs. Keratoconus was described as progressive when there was an increase in the cone apex keratometry of -0.75 diopters (D) or alteration of -0.75 D in the spherical equivalent refraction, in a period of at least 6 months. Patients were informed about the experimental nature of the intervention, the possible outcomes, and the current clinical experience, and provided written consent according to the Declaration of Helsinki and institutional guidelines.

CLINICAL EVALUATION

Preoperative evaluation consisted of general and ocular health history assessment; autorefractometry and autokeratometry (Canon autorefractor; Canon USA Inc, Lake Success, NY); corneal topography (iTrace; Tracey Technologies, Houston Tex); assessment of uncorrected visual acuity (UCVA), best-spectacle corrected

visual acuity (BSCVA), manifest and cycloplegic refraction; scotopic pupillometry; central ultrasound pachymetry (Corneo Gage Plus; Sonogage, Cleveland, Ohio); and slit-lamp examination of the anterior and posterior segments of the eye.

SURGICAL PROCEDURE

All procedures were performed by the same surgeon (G.D.K.). After topical anesthesia with tetracaine 1% and oxybuprocaine 0.4% eye drops, the epithelium was mechanically removed within an 8.5-mm diameter. A solid-state laser with a wavelength of 213 nm (Pulzar Z1; CustomVis, Perth, Wash) was used for the PRK procedure. The wavelength is generated using a major Nd:YAG laser system of 1064 nm and through special cultivated crystals the 213 nm is finally used. The customization was performed based on the corneal higher order aberrations. System software allows the use of a percentage of customization from 0% to 100%. Using 0% would be equivalent to a conventional laser treatment and 100% would be equivalent to a full customized treatment. Adjusting this percentage, attempted correction and ablation zone could lower the maximum depth of tissue removed. Maximum ablation depth was 50 μm . Attempted correction and maximum ablation depth are shown in Table 1.

Riboflavin (0.1% solution 10 mg riboflavin-5-phosphate in 10 mL dextran-T-500 20% solution) was applied every 3 minutes for approximately 20 minutes until the stroma was completely penetrated and the aqueous was stained yellow (riboflavin shielding). A commercially available UVA system (UV-X illumination system version 1000; IROC AG, Zurich, Switzerland) was used for UVA irradiation. Before treatment, the intended 3 mW/cm^2 surface irradiance (5.4 J/cm^2 surface dose after 30 minutes) was calibrated using the UVA meter YK-34UV (Lutron Electronics Co Inc, Coopersburg, Pa), which is supplied with the UV-X device. During treatment, riboflavin solution was applied every 5 minutes to ensure saturation.

After treatment, a bandage contact lens was applied until the epithelium healed completely, followed by application of fluorometholone 0.1% eye drops (FML Liquifilm; Allergan, Irvine, Calif) twice daily for 2 weeks.

RESULTS

All values are expressed as mean \pm standard deviation. Visual acuity is expressed as logarithm of minimum angle of resolution (logMAR).

Mean follow up was 10.69 ± 5.95 months (range: 3 to 16 months). Mean preoperative spherical equivalent refraction was -3.03 ± 3.23 diopters (D) and defocus was 4.67 ± 3.29 D, and at last follow-up both were statisti-




Figure 1. Preoperative best spectacle-corrected visual acuity versus postoperative uncorrected visual acuity (efficacy) following topography-guided photorefractive keratectomy (PRK) and corneal cross-linking.




Figure 2. Change in best spectacle-corrected visual acuity (safety) following topography-guided photorefractive keratectomy and corneal cross-linking.

cally significantly ($P < .01$) reduced to -1.29 ± 2.05 D and 3.04 ± 2.53 D, respectively. Preoperative mean UCVA and BSCVA were 0.99 ± 0.81 and 0.21 ± 0.19 , respectively, which improved at last follow-up to 0.16 ± 0.15 and 0.11 ± 0.15 , respectively. Postoperatively, all eyes had BSCVA of 20/100 (Snellen) or better (Fig 1). No eye lost >2 lines of BSCVA whereas 10% gained >3 lines postoperatively (Fig 2). Preoperatively, 93% of patients had BSCVA of 0.3 or less (logMAR) whereas all eyes presented with UCVA of 0.3 or better at last follow-up.

Figure 3 represents a scattergram of attempted versus achieved refraction.

keratoconus with the main drawback being that it is a tissue removal technique. In stable or forme fruste keratoconus, surface ablation has been reported with promising results. Several studies report the application of excimer laser ablation to correct astigmatism in patients with stable keratoconus. Alpíns and Stamatelatos⁶ report a series of 32 eyes treated with photostigmatic refractive keratectomy and followed up for up to 10 years with no evidence of keratoconic progression. Koller et al⁷ treated 11 eyes with customized PRK and reported improvement in refractive astigmatism, topography, and quality of vision in all patients.

The mean steepest keratometry was reduced from 48.20 ± 3.40 D preoperatively to 45.13 ± 1.80 D at last follow-up (Table 2, Fig 4).

DISCUSSION

Treatment of keratoconus is an issue consisting of two parameters—corneal biomechanical stability and optical inefficiency of the irregular cornea. The goal of our study was to develop a technique to treat keratoconic patients and offer them both stability and functional vision. Reaching functional vision consists of improving UCVA and BSCVA and normalization of corneal topography, which indicates that these patients are less dependent on contact lenses to achieve higher postoperative vision quality.

Stabilization of corneas with ectatic disorders can be achieved with the application of CXL. The method is based on the absorption of UVA radiation by the cornea after the photosensitizer riboflavin is infused in the stroma.^{10,11} Despite stabilizing the cornea, the topography and visual performance of patients treated with CXL show minimal improvement.

The application of laser ablation has been introduced formerly for the correction of




Figure 3. Scattergram of attempted versus achieved refraction following topography-guided photorefractive keratectomy and corneal cross-linking.




Figure 4. Preoperative axial map (bottom right), postoperative axial map (top right), and difference axial map (left), showing significant topographic improvement following topography-guided photorefractive keratectomy and corneal cross-linking.

Twenty-five eyes treated with topography-guided PRK reported by Cennamo et al⁸ also had improved topography and visual outcomes.

Given that the CXL procedure is capable of stiffening

up to 50 μm were performed, whereas the ablation depth was mapped out by modifying the target correction (expected central corneal thickness after PRK and prior CXL $>400 \mu\text{m}$).

the cornea and halting keratoconus progression, the combination of laser ablation and CXL provides a potential treatment for keratoconus. Kanellopoulos and Binder⁹ proposed a two-step procedure with corneal CXL and PRK after a 1-year interval. There are three limitations with this approach. First, the stiffened cross-linked corneal tissue is removed in a second step by the PRK (potentially decreasing the possible benefits of CXL). Second, the efficacy of this approach is limited because the corneal ablation rate could be different in cross-linked corneas than in the virgin cornea (this could lead to unpredictable refractive results). Third, there is an increased possibility of postoperative PRK haze formation (following CXL, the anterior stroma is repopulated by new keratocytes after 6 months).

Due to these aspects, we considered that the best option for treating keratoconus is topography-guided PRK immediately followed by CXL. The main advantage of this technique is that ablation does not interfere with the already cross-linked part of the cornea. On the contrary, cross-linking of the ablated stroma offers the advantage of depopulating keratocytes in the anterior stroma, which could reduce the possibility of haze formation.¹²

The major consideration in the planning of this procedure was the postoperative corneal thickness. Photorefractive keratectomy was planned based on the patient's corneal thickness. Restricted treatments of

TABLE 2

Pre- and Postoperative Data of 14 Eyes That Underwent Topography-guided PRK and Cross-linking

Parameter	Mean±SD (Range)	
	Preoperative	Last Follow-up
Manifest refraction (D)	-3.03±3.23 (1.38 to -12.88)	-1.29±2.05 (1.25 to -4.00)
Defocus (D)	4.67±3.29 (14.50 to 1.00)	3.04±2.53 (6.75 to 0.75)
UCVA (logMAR)	0.99±0.81 (2 to 0.16)	0.16±0.15 (0.32 to 0)
BSCVA (logMAR)	0.21±0.19 (0.66 to -0.04)	0.11±0.15 (0.3 to -0.08)
Keratometry (flat/steep) (D)	43.62±2.40/48.20±3.40	41.00±1.20/45.13±1.80

UCVA = uncorrected visual acuity, BSCVA = best spectacle-corrected visual acuity

Patients treated with simultaneous topography-guided PRK with CXL showed a rapid and significant improvement in UCVA and BSCVA. Topographic evaluation of these patients showed marked improvement of irregularity. Even with relatively small correction due to the thickness limitations, patients may benefit.

Simultaneous PRK followed by CXL seems to be a promising treatment. Performing this technique with careful observance of safety aspects may offer patients with keratoconus the opportunity to gain functional vision, avoid complications of long-term contact lens use, and also reduce the need for corneal transplantation.

AUTHOR CONTRIBUTIONS

Study concept and design (G.D.K., G.A.K., S.Y., I.G.P.); data collection (D.M.P., A.E.K., M.M.); interpretation and analysis of data (G. Kounis); drafting of the manuscript (G.D.K., G. Kontadakis, G. Kounis, D.M.P., A.E.K., M.M.); critical revision of the manuscript (S.Y., I.G.P.); statistical expertise (G. Kontadakis, G. Kounis); supervision (I.G.P.)

REFERENCES

- Kymionis GD, Siganos CS, Tsiklis NS, Anastasakis A, Yoo SH, Pallikaris AI, Astyrakakis N, Pallikaris IG. Long-term follow-up of Intacs in keratoconus. *Am J Ophthalmol.* 2007;143:236-244.
- Coskunseven E, Kymionis GD, Tsiklis NS, Atun S, Arslan E, Jankov MR, Pallikaris IG. One-year results of intrastromal corneal ring segment implantation (KeraRing) using femto-second laser in patients with keratoconus. *Am J Ophthalmol.* 2008;145:775-779.
- Wollensak G, Spoerl E, Seiler T. Riboflavin/ultraviolet-a-induced collagen crosslinking for the treatment of keratoconus. *Am J Ophthalmol.* 2003;135:620-627.
- Wollensak G, Spoerl E, Seiler T. Stress-strain measurements of human and porcine corneas after riboflavin-ultraviolet-A-induced cross-linking. *J Cataract Refract Surg.* 2003;29:1780-1785.
- Raiskup-Wolf F, Hoyer A, Spoerl E, Pillunat LE. Collagen cross-linking with riboflavin and ultraviolet-A light in keratoconus: long-term results. *J Cataract Refract Surg.* 2008;34:796-801.
- Alpins N, Stamatelatos G. Customized photoastigmatic refractive keratectomy using combined topographic and refractive data for myopia and astigmatism in eyes with forme fruste and mild keratoconus. *J Cataract Refract Surg.* 2007;33:591-602.
- Koller T, Iseli HP, Donitzky C, Ing D, Papadopoulos N, Seiler T. Topography-guided surface ablation for forme fruste keratoconus. *Ophthalmology.* 2006;113:2198-2202.
- Cennamo G, Intravaja A, Boccuzzi D, Marotta G, Cennamo G. Treatment of keratoconus by topography-guided customized photorefractive keratectomy: two-year follow-up study. *J Refract Surg.* 2008;24:145-149.
- Kanellopoulos AJ, Binder PS. Collagen cross-linking (CCL) with sequential topography-guided PRK: a temporizing alternative for keratoconus to penetrating keratoplasty. *Cornea.* 2007;26:891-895.
- Wollensak G. Crosslinking treatment of progressive keratoconus: new hope. *Curr Opin Ophthalmol.* 2006;17:356-360.
- Spoerl E, Huhle M, Seiler T. Induction of cross-links in corneal tissue. *Exp Eye Res.* 1998;66:97-103.
- Kymionis GD, Diakonis VF, Kalyvianaki M, Portaliou D, Siganos C, Kozobolis VP, Pallikaris AI. One-year follow-up of corneal confocal microscopy after corneal cross-linking in patients with post laser in situ keratomileusis ectasia and keratoconus. *Am J Ophthalmol.* 2009;147:774-778.e1.

AUTHOR QUERIES

Was this paper presented at the Fourth International Congress of Corneal Cross Linking?