

Abstract Submitted
for the MAR16 Meeting of
The American Physical Society

Singlet–Triplet Energy Gaps for Diradicals from Particle–Particle Random Phase Approximation YANG YANG, DEGAO PENG, Duke Univ, ERNEST DAVIDSON, University of Washington, WEITAO YANG, Duke Univ — The particle–particle random phase approximation (pp-RPA) has been applied to the calculation of vertical and adiabatic singlet–triplet energy gaps for a variety of categories of diradicals, including diatomic diradicals, carbene-like diradicals, disjoint diradicals, four- π -electron diradicals, and benzynes are calculated. Except for some excitations in four- π -electron diradicals, where four-electron correlation may play an important role, the singlet–triplet gaps are generally well predicted by pp-RPA. With a relatively low $O(r^4)$ scaling, the pp-RPA with DFT references outperforms spin-flip configuration interaction singles. It is similar to or better than the (variational) fractional-spin method. For small diradicals such as diatomic and carbene-like ones, the error of pp-RPA is slightly larger than noncollinear spin-flip time-dependent density functional theory (NC-SF-TDDFT) with LDA or PBE functional. However, for disjoint diradicals and benzynes, the pp-RPA performs much better and is comparable to NC-SF-TDDFT with long-range corrected ω PBEh functional and spin-flip configuration interaction singles with perturbative doubles (SF-CIS(D)). In particular, with a correct asymptotic behavior and being almost free from static correlation error, the pp-RPA with DFT references can well describe the challenging ground state and charge transfer excitations of disjoint diradicals in which almost all other DFT-based methods fail. Therefore, the pp-RPA could be a promising theoretical method for general diradical problems.

Yang Yang
Duke Univ

Date submitted: 04 Nov 2015

Electronic form version 1.4