

SINGULAR POLYNOMIALS FOR FINITE REFLECTION GROUPS

C. F. DUNKL, M. F. E. DE JEU, AND E. M. OPDAM

ABSTRACT. The Dunkl operators involve a multiplicity function as parameter. For generic values of this function the simultaneous kernel of these operators, acting on polynomials, is equal to the constants. For special values, however, this kernel is larger. We determine these singular values completely and give partial results on the representations of G that occur in this kernel.

1. INTRODUCTION AND NOTATIONS

Let \mathfrak{a} be a real vector space of finite dimension N , equipped with an inner product (\cdot, \cdot) . Let $G \subset O(\mathfrak{a}(\cdot, \cdot))$ be a finite (real) reflection group. We may and will assume that (\cdot, \cdot) is G -invariant. Let R be the corresponding root system, where we will assume that $(\alpha, \alpha) = 2$ for all $\alpha \in R$. Choose and fix a positive system R_+ in R .

Let $\mathcal{P} = \mathbf{C}[\mathfrak{a}]$ denote the polynomial functions on \mathfrak{a} . \mathcal{P} has a natural grading $\mathcal{P} = \bigoplus_{n \geq 0} \mathcal{P}_n$; we let $\mathcal{P}_+ = \bigoplus_{n \geq 1} \mathcal{P}_n$. There is a natural representation of G in $\text{Aut}_{\mathbf{C}}(\mathcal{P})$, given by

$$(g \cdot f)(x) := f(g^{-1}x) \quad (g \in G, f \in \mathcal{P}, x \in \mathfrak{a}).$$

The G -action preserves the degree.

Any $\xi \in \mathfrak{a}$ defines an element $\xi^* \in \text{Hom}_{\mathbf{C}}(\mathfrak{a}, \mathbf{C}) = \mathfrak{a}^* = \mathcal{P}_1$ by $\xi^*(\eta) = (\xi, \eta)$ ($\eta \in \mathfrak{a}$). Let K be the complex vectorspace of \mathbf{C} -valued G -invariant functions on R . An element of K is called a multiplicity function (the reason for this terminology is a connection of the theory of Dunkl operators with the harmonic analysis for the Cartan motion group; the values k_α of $k \in K$ are then determined by the multiplicities of the restricted roots). To each $\xi \in \mathfrak{a}$ and $k \in K$ we assign an operator $T_\xi(k) \in \text{End}_{\mathbf{C}}(\mathcal{P})$ as follows:

$$T_\xi(k)f = \partial_\xi f + \sum_{\alpha \in R_+} k_\alpha(\alpha, \xi) \frac{f - r_\alpha f}{\alpha^*} \quad (f \in \mathcal{P}),$$

where ∂_ξ is the directional derivative corresponding to ξ and r_α is the reflection in the hyperplane orthogonal to α .

These operators were introduced in [D2]. In that paper it was shown that $\{T_\xi(k) \mid \xi \in \mathfrak{a}\}$ is a commuting family of operators, for all fixed $k \in K$. They are homogeneous of degree -1 and G -equivariant:

$$(1.1) \quad g \circ T_\xi(k) \circ g^{-1} = T_{g\xi}(k) \quad (\xi \in \mathfrak{a}, g \in G).$$

Received by the editors February 4, 1994.

1991 *Mathematics Subject Classification.* Primary 20F55, 33C80, 33C35, 20C30.

During the preparation of this paper Dunkl was partially supported by NSF grant DMS-9103214.

For $k = 0$, Dunkl’s operators reduce to the ordinary directional derivatives, so the simultaneous kernel of the $T_\xi(k)$ ($\xi \in \mathfrak{a}$) consists of the constants alone. For generic $k \in K$ this is still true, but there exist exceptional k for which this simultaneous kernel is strictly larger than the constants. This paper is concerned with the study of such singular values. Let us introduce some terminology. For $k \in K$, put $S(k) = \{p \in \mathcal{P}_+ \mid T_\xi(k)p = 0 \ \forall \xi \in \mathfrak{a}\}$. If $S(k) \neq \{0\}$, then k is called *singular*, and *regular* otherwise. We use the notation K^{sing} (resp. K^{reg}) for the set of all singular (resp. regular) $k \in K$. Similarly, a polynomial $p \in \mathcal{P}_+$ is called *singular* if $p \in S(k)$ for some $k \in K$. Put $S = \sum_{k \in K} S(k)$. As a consequence of the homogeneity and the G -equivariance of the $T_\xi(k)$ one may split S according to homogeneous degree and G -type: $S = \bigoplus_{n \geq 1, \tau \in \widehat{G}} S_{n, \tau}$, where \widehat{G} denotes the dual of G . There is an easy first observation on K^{sing} . Let $\{e_1, \dots, e_N\}$ be an orthonormal basis for \mathfrak{a} . Then

$$(1.2) \quad \sum_{i=1}^N e_i^* T_{e_i}(k) = \sum_{i=1}^N e_i^* \partial_{e_i} + \sum_{\alpha \in R_+} k_\alpha (1 - r_\alpha).$$

Due to the G -invariance of k , $\sum_{\alpha \in R_+} k_\alpha (1 - r_\alpha)$ is a central element in the group algebra of G . Hence it acts as a scalar $\lambda_\tau(k)$ in any irreducible representation $\tau \in \widehat{G}$. If $k_\alpha = k \in \mathbb{C}$ ($\forall \alpha \in R$), then it is easy to see that $\lambda_\tau(k \cdot 1_R) = k \sum_{\alpha \in R_+} (1 - \chi_\tau(r_\alpha) / \chi_\tau(1))$, where χ_τ denotes the character of τ . Now if $p \in S(k)$ is of isotype τ and homogeneous degree m , then p is annihilated by the operator in (1.2); this gives the inclusion $S_{m, \tau} \subset S(-m / \lambda_\tau(1_R))$ in the case of equal k_α . In other words, we can determine K^{sing} completely if we know the embedding degrees of all $\tau \in \widehat{G}$ in S . One might hope that $K^{\text{sing}} = \{-m / \lambda_\tau(1_R) \mid m > 0, \tau \in \widehat{G}, \tau \text{ occurs in } \mathcal{P}_m\}$, but this is false (the easiest counterexample is furnished by $G = S_4$). Let us note that $\lambda_\tau(1_R)$ is an integer; this follows from general representation theory of finite groups. It is easy to see that this integer is nonpositive and equal to zero iff $\tau = \text{triv}$. So K^{sing} will consist of strictly negative rationals in the case of equal k_α .

One of the reasons for our interest in S is the fact that $S(k)$ turns out to be the only obstruction for the existence of a linear isomorphism $V(k): \mathcal{P} \rightarrow \mathcal{P}$, homogeneous of degree 0, that intertwines the action of $T_\xi(k)$ and the ordinary derivative ∂_ξ on \mathcal{P} (see Corollary 3.5). There are many other different descriptions for S or K^{sing} and their role with regard to the Dunkl operators. For example, K^{sing} is also the set of poles of the simultaneous $T_\xi(k)$ -eigenfunction $\text{Exp}_G(\lambda, k, x)$, considered as a function of $k \in K$ (see [O]). Using monodromy arguments this set of poles was described in terms of the zeros of the Poincaré polynomial of the coinvariant algebra of G [O, Theorem 9.7]. This leads to an explicit description of K^{sing} . If $k_\alpha = k \ \forall \alpha \in R$, then the result is

$$K^{\text{sing}} = \bigcup_{m \geq 1} \bigcup_{i=1}^N \bigcup_{j=1}^{d_i-1} \left\{ \frac{j}{d_i} - m \right\},$$

where the d_i are the primitive degrees of G . Yet another meaning for this set K^{sing} is the following. As a consequence of the commutativity of the $T_\xi(k)$ for fixed k , there exists an algebra homomorphism from \mathcal{P} to $\text{End}_{\mathbb{C}}(\mathcal{P})$ which sends ξ^* to $T_\xi(k)$ and 1 to the identity. If we define the bilinear form $(\cdot, \cdot)_k$

on $\mathcal{P} \times \mathcal{P}$ by $(p, q)_k = (p(T(k))(q))(0)$, then K^{sing} is the set of all values for which this form is degenerate.

The purpose of this paper is twofold. First of all, we explain these different interpretations of the space S and the set K^{sing} . Secondly, we try to get a better understanding of the structure of K^{sing} with regard to the representation theory of G . The optimal result in this direction would be a list of irreducibles $\tau \in \widehat{G}$ that can occur in S , and for each of these a description of the function $m \mapsto \dim_{\mathbb{C}}(S_{m, \tau})$. As yet this seems out of reach, but we will be able to give estimates for $\dim_{\mathbb{C}}(S_{m, \tau})$ entirely in terms of decomposition numbers for Hecke algebras.

2. THE EXISTENCE OF INTERTWINING OPERATORS

The present section is concerned with the existence of operators that intertwine the action of two sets of operators in a graded vectorspace. The presence of a grading for which the various actions are homogeneous enables one to use induction. This fact and an additional assumption (formulated in the language of differential forms) are surprisingly powerful tools for the construction of certain intertwining operators. The underlying ideas were used more or less parenthetically in [D3]; this section can be regarded as an abstraction and extension of the proof of Theorem 3.11 in [loc. cit.]. The results show that, in the context of this section, there is an equivalence between the commutativity of sets of operators, the triviality of the first cohomology-group of a certain complex and the existence of intertwining operators. We will use this equivalence to give a new proof of the commutativity of Dunkl's operators and the existence of the intertwining operator (see [loc. cit.]).

Let L be a vectorspace with a grading $L = \bigoplus_{i=0}^{\infty} L_i$. Put $L_{-1} = 0$. Let k be a positive integer and suppose $\{\partial_j\}_{j=1}^k$ are linear operators in L such that

(i) $\partial_j L_i \subset L_{i-1}$ ($j = 1, \dots, k; i = 0, 1, \dots$).

(ii) $\bigcap_{j=1}^k \text{Ker}(\partial_j) = L_0$.

A sequence $\{l_i\}_{i=1}^k$ in L will be called a ∂ -closed 1-form if $\partial_j l_i = \partial_i l_j$ ($i, j = 1, \dots, k$); it is a ∂ -exact 1-form if there exists $l \in L$ such that $\partial_i l = l_i$ ($i = 1, \dots, k$). Suppose in addition that

(iii) every ∂ -closed 1-form is ∂ -exact.

Theorem 2.1. *Let $\{\delta_j\}_{j=1}^k$ be linear operators in L such that*

(i') $\delta_j(L_i) \subset L_{i-1}$ ($j = 1, \dots, k; i = 0, 1, \dots$);

(ii') the δ_j commute.

Let $\{\partial_j\}_{j=1}^k$ be linear operators in L satisfying (i), (ii), and (iii) above and let $A: L_0 \mapsto L_0$ be linear. Then there exists a unique linear map $B: L \mapsto L$ such that:

(a) B extends A ;

(b) $\partial_j B = B \delta_j$ ($j = 1, \dots, k$);

(c) $B L_i \subset L_i$ ($i = 0, 1, \dots$).

If A is injective and $\bigcap_{j=1}^k \text{Ker}(\delta_j) = L_0$, then B is injective.

Proof. Existence. We will construct a family of operators $B_i: L_i \mapsto L_i$ ($i = 0, 1, \dots$) such that:

(α) $B_0 = A$;

(β) $\partial_j B_i = B_{i-1} \delta_j$ ($j = 1, \dots, k; i = 1, 2, \dots$);

(γ) $B_i L_i \subset L_i$ ($i = 0, 1, \dots$).

Assuming for the moment that this has been done, we can take $B|_{L_i} = B_i$ and extend by linearity to obtain an endomorphism of L , which satisfies (a), (b), and (c).

The construction is by induction. Put $B_0 = A$. Next, let $l \in L_1$. Consider the 1-form $\{A\delta_i l\}_{i=1}^k$. Since $A\delta_i l \in L_0$, this 1-form is trivially ∂ -closed by (i), so there exists $l' \in L$ such that $\partial_i l' = A\delta_i l$ ($i = 1, \dots, k$). Properties (i) and (ii) imply that any such l' is in $L_0 \oplus L_1$. By (i), we may take l' in L_1 and it is then uniquely determined as a consequence of (ii). We can thus define $B_1 l$ to be the unique element in L_1 satisfying $\partial_i B_1 l = A\delta_i l$ ($i = 1, \dots, k$). The uniqueness makes it clear that $B_1: L_1 \mapsto L_1$ is linear. For the induction step, assume that B_0, \dots, B_d ($d \geq 1$) have been constructed and satisfy (α), (β), and (γ) (for $i = 0, \dots, d$). Let $l \in L_{d+1}$ and consider the 1-form $\{B_d \delta_i l\}_{i=1}^k$. Since the δ_i commute, we have, using the induction hypothesis

$$\partial_j B_d \delta_i l = B_{d-1} \delta_j \delta_i l = B_{d-1} \delta_i \delta_j l = \partial_i B_d \delta_j l.$$

Hence the form is ∂ -closed and there exists $l' \in L$ such that $\partial_i l' = B_d \delta_i l$ ($i = 1, \dots, k$). The same argument as above shows that we can determine l' uniquely by requiring it to be in L_{d+1} . Again, the map $B_{d+1}: L_{d+1} \mapsto L_{d+1}$, defined by $B_{d+1} l = l'$ is linear and by construction we have $\partial_i B_{d+1} l = \partial_i l' = B_d \delta_i l$ ($i = 1, \dots, k$). This completes the induction step.

Uniqueness. Let B' be any other operator in L satisfying (a), (b), and (c) in the Theorem. Put $B'_d = B'|_{L_d}$. We prove by induction that $B'_d = B_d$ ($d = 0, 1, \dots$). For $d = 0$, this is just (a). Assuming $B'_i = B_i$ ($i = 0, \dots, d$), we have for $l \in L_{d+1}$ arbitrary

$$\partial_j (B_{d+1} - B'_{d+1})l = (B_d - B'_d)\delta_j l = 0 \quad (j = 1, \dots, k),$$

so $(B_{d+1} - B'_{d+1})l \in L_0$ by (ii). But then $(B_{d+1} - B'_{d+1})l \in L_0 \cap L_{d+1}$ by (c), which is the zero subspace since $d + 1 \geq 1$.

We now prove the last statement of the theorem. Put $L^d = \bigoplus_{i=0}^d L_i$ ($d = 0, 1, \dots$). We prove, again by induction, that B is injective on L^d . For $d = 0$, this is the injectivity of A . Assuming the statement for $i = 0, \dots, d$, let $l \in L^{d+1}$ and suppose $Bl = 0$. Then, by (b), $B\delta_j l = 0$ ($j = 1, \dots, k$). Since $\delta_j l \in L^d$, $\delta_j l = 0$ ($j = 1, \dots, k$) by the induction hypothesis. So, by the assumption on the simultaneous kernel of the δ_j , l is in L_0 . But then $Al = Bl = 0$, so $l = 0$.

Corollary 2.2. *Let L and the L_i be as above and suppose in addition that $\dim(L_i) < \infty$ ($i = 1, 2, \dots$). Suppose that $\{\partial_j\}_{j=1}^k$ are linear operators such that:*

- (i) $\partial_j L_i \subset L_{i-1}$ ($j = 1, \dots, k; i = 0, 1, \dots$);
- (ii) $\bigcap_{j=1}^k \text{Ker}(\partial_j) = L_0$;
- (iii) every ∂ -closed 1-form is ∂ -exact;
- (iv) the ∂_j commute.

Suppose $\{\delta_j\}_{j=1}^k$ are linear operators in L such that

- (i') $\delta_j L_i \subset L_{i-1}$ ($j = 1, \dots, k; i = 0, 1, \dots$);
- (ii') $\bigcap_{j=1}^k \text{Ker}(\delta_j) = L_0$.

Then the following are equivalent:

(a) The δ_j commute.

(b) Every δ -closed 1-form is δ -exact.

(c) There exists an automorphism A of L_0 that extends to an automorphism B of L such that $\partial_j B = B\delta_j$ ($j = 1, \dots, k$).

(d) Every automorphism A of L_0 uniquely extends to an automorphism B of L such that $\partial_j B = B\delta_j$ ($j = 1, \dots, k$) and $BL_i = L_i$ ($i = 0, 1, \dots$).

Proof. (a) \Rightarrow (d): Theorem 2.1 applies. Since B is injective on the L_i , it is an automorphism by the finite dimensions of the L_i . The implications (d) \Rightarrow (c) and (c) \Rightarrow (a) are obvious.

(b) \Rightarrow (d): Apply Theorem 2.1 with the roles of the ∂_j and δ_j interchanged. Again the finite dimensions of the L_i ensure that B is an automorphism.

(c) \Rightarrow (b): This follows from (iii). \square

3. THE PERTURBED DE RHAM COMPLEX

We consider the differential forms on \mathfrak{a} with polynomial coefficients, on which we introduce a tentative boundary operator $d(k)$ (based on the Dunkl operators). A calculation and subsequent appeal to the results of the previous section will enable us to conclude that the Dunkl operators commute; it is then immediate that $d(k)$ is a boundary operator. We conclude with an alternative characterization of singular polynomials.

For $l = 0, 1, \dots$, let $K^l = \mathcal{P} \otimes_{\mathbb{C}} \bigwedge_l \mathfrak{a}^*$ (this K^l has no relation to the space K of multiplicity functions) and put $K^\bullet = \bigoplus_{l=0}^N K^l$. Let $\{e_1, \dots, e_N\}$ be an orthonormal basis for \mathfrak{a} . We will denote a typical generator of K^l by $p \otimes de_{i_1}^* \wedge \dots \wedge de_{i_l}^*$, where $p \in \mathcal{P}$ is a polynomial. An element $g \in G$ acts on K^l as follows:

$$(g \otimes g) \cdot (p \otimes de_{i_1}^* \wedge \dots \wedge de_{i_l}^*) = (g \cdot p) \otimes d(g \cdot e_{i_1}^*) \wedge \dots \wedge d(g \cdot e_{i_l}^*).$$

We introduce operators $d(k): K^l \mapsto K^{l+1}$ by the action on generators:

$$d(k)(p \otimes de_{i_1}^* \wedge \dots \wedge de_{i_l}^*) = \sum_{j=1}^N (T_{e_j}(k)p) \otimes de_j^* \wedge de_{i_1}^* \wedge \dots \wedge de_{i_l}^*.$$

Lemma 3.1. $d(k)$ commutes with the G -action.

Proof. The well-known fact that $d(0)$ commutes with the action of $GL(\mathfrak{a})$ follows from an easy computation which is based on the linearity of the map $\xi \mapsto \partial_\xi$ and on the $GL(\mathfrak{a})$ -equivariance of the ordinary directional derivatives. Hence, the same computation applies for $d(k)$ and G , using the G -equivariance of the $T_\xi(k)$. \square

We recall the definition of the Koszul-operator $\partial: K^l \mapsto K^{l-1}$:

$$\partial(p \otimes de_{i_1}^* \wedge \dots \wedge de_{i_l}^*) = \sum_{r=1}^l (-1)^{r+1} e_{i_r}^* p \otimes de_{i_1}^* \wedge \dots \wedge de_{i_{r-1}}^* \wedge de_{i_{r+1}}^* \wedge \dots \wedge de_{i_l}^*.$$

Proposition 3.2. On K^l , we have

$$(\partial d(k) + d(k)\partial) = l(1 \otimes 1) + \left(\sum_{j=1}^N e_j^* \partial e_j \right) \otimes 1 + \sum_{\alpha \in R_+} k_\alpha (1 \otimes 1 - r_\alpha \otimes r_\alpha).$$

Proof. Let $\omega = p \otimes de_{i_1}^* \wedge \cdots \wedge de_{i_l}^* \in K^l$. We calculate as follows:

$$\begin{aligned}
 &(\partial d(k) + d(k)\partial)\omega \\
 &= \partial \left(\sum_{j=1}^N T_{e_j}(k)p \otimes de_j^* \wedge de_{i_1}^* \wedge \cdots \wedge de_{i_l}^* \right) \\
 &\quad + d(k) \left(\sum_{r=1}^l (-1)^{r+1} e_r^* p \otimes de_{i_1}^* \wedge \cdots \wedge de_{i_{r-1}}^* \wedge de_{i_{r+1}}^* \wedge \cdots \wedge de_{i_l}^* \right) \\
 &= \sum_{j=1}^N e_j^* T_{e_j}(k)p \otimes de_{i_1}^* \wedge \cdots \wedge de_{i_l}^* \\
 &\quad + \sum_{j=1}^N \sum_{r=1}^l (-1)^r e_r^* T_{e_j}(k)p \otimes de_j^* \wedge de_{i_1}^* \wedge \cdots \wedge de_{i_{r-1}}^* \wedge de_{i_{r+1}}^* \wedge \cdots \wedge de_{i_l}^* \\
 &\quad + \sum_{j=1}^N \sum_{r=1}^l (-1)^{r+1} T_{e_j}(k)(e_r^* p) \otimes de_j^* \wedge de_{i_1}^* \wedge \cdots \wedge de_{i_{r-1}}^* \wedge de_{i_{r+1}}^* \wedge \cdots \wedge de_{i_l}^*.
 \end{aligned}$$

An easy calculation shows that

$$(3.1) \quad [T_{e_j}(k), e_i^*] = \delta_{ij} + \sum_{\alpha \in R_+} k_\alpha(\alpha, e_i)(\alpha, e_j)r_\alpha,$$

so

$$\begin{aligned}
 (\partial d(k) + d(k)\partial)\omega &= \sum_{j=1}^N e_j^* T_{e_j}(k)p \otimes de_{i_1}^* \wedge \cdots \wedge de_{i_l}^* \\
 &\quad + \sum_{j=1}^N \sum_{r=1}^l (-1)^{r+1} \left(\delta_{ji_r} + \sum_{\alpha \in R_+} k_\alpha(\alpha, e_{i_r})(\alpha, e_j)r_\alpha \right) p \\
 &\quad \quad \quad \otimes de_j^* \wedge de_{i_1}^* \wedge \cdots \wedge de_{i_{r-1}}^* \wedge de_{i_{r+1}}^* \wedge \cdots \wedge de_{i_l}^* \\
 &= \sum_{j=1}^N e_j^* T_{e_j}(k)p \otimes de_{i_1}^* \wedge \cdots \wedge de_{i_l}^* \\
 &\quad + \sum_{r=1}^l (-1)^{r+1} p \otimes de_{i_r}^* \wedge de_{i_1}^* \wedge \cdots \wedge de_{i_{r-1}}^* \wedge de_{i_{r+1}}^* \wedge \cdots \wedge de_{i_l}^* \\
 &\quad + \sum_{\alpha \in R_+} k_\alpha \sum_{j=1}^N \sum_{r=1}^l (-1)^{r+1} (\alpha, e_{i_r})(\alpha, e_j)(r_\alpha p) \\
 &\quad \quad \quad \otimes de_j^* \wedge de_{i_1}^* \wedge \cdots \wedge de_{i_{r-1}}^* \wedge de_{i_{r+1}}^* \wedge \cdots \wedge de_{i_l}^* \\
 &= \sum_{j=1}^N e_j^* T_{e_j}(k)p \otimes de_{i_1}^* \wedge \cdots \wedge de_{i_l}^* + l\omega \\
 &\quad + \sum_{\alpha \in R_+} k_\alpha \sum_{r=1}^l (-1)^{r+1} (\alpha, e_{i_r})(r_\alpha p) \\
 &\quad \quad \quad \otimes de_{i_r}^* \wedge de_{i_1}^* \wedge \cdots \wedge de_{i_{r-1}}^* \wedge de_{i_{r+1}}^* \wedge \cdots \wedge de_{i_l}^*.
 \end{aligned}$$

Now note that

$$\begin{aligned} &(r_\alpha \otimes r_\alpha)(p \otimes de_{i_1}^* \wedge \cdots \wedge de_{i_l}^*) \\ &= (r_\alpha p) \otimes de_{i_1}^* \wedge \cdots \wedge de_{i_l}^* + \sum_{r=1}^l (-1)^r (\alpha, e_{i_r}) (r_\alpha p) \\ &\quad \otimes d\alpha^* \wedge de_{i_1}^* \wedge \cdots \wedge de_{i_{r-1}}^* \wedge de_{i_{r+1}}^* \wedge \cdots \wedge de_{i_l}^*. \end{aligned}$$

Thus, if we invoke (1.2), we see that

$$(\partial d(k) + d(k)\partial)\omega = \sum_{j=1}^N (e_j^* \partial_{e_j} p) \otimes de_{i_1}^* \wedge \cdots \wedge de_{i_l}^* + l\omega + \sum_{\alpha \in R_+} k_\alpha (1 \otimes 1 - r_\alpha \otimes r_\alpha)\omega.$$

□

For $m = 0, 1, \dots$, let \mathcal{P}_m denote the homogeneous polynomials of degree m , and put $K_m^l = \mathcal{P}_m \otimes_{\mathbb{C}} \bigwedge_l \mathfrak{a}^*$. The K_m^l are invariant under the G -action, so there is a further splitting $K_m^l = \bigoplus_{\tau \in \widehat{G}} K_{m,\tau}^l$, where $K_{m,\tau}^l$ is the τ -isotypical component in K_m^l . If $\omega \in L_{m,\tau}^l$, then $(\partial d(k) + d(k)\partial)\omega = (l + m + \lambda_\tau(k))\omega$. It follows from Corollary 2.2 in [D3] that $\text{Re}(\lambda_\tau(k)) \geq 0$ if $\text{Re}(k) \geq 0$. Note that K^\bullet is a direct sum of finite-dimensional subcomplexes $K^\bullet(n) = \bigoplus_{l+m=n} K_m^l$.

Corollary 3.3. *Suppose that $\text{Re}(k) \geq 0$, $\omega \in K^l (l \geq 1)$ and $d(k)\omega = 0$. Then $\omega \in d(k)K^{l-1}$.*

Proof. By Lemma 3.1 and homogeneity, the statement reduces to the case that $\omega \in K_{m,\tau}^l$. But for such ω the Corollary is obvious from the above remarks. □

Corollary 3.4. *We have*

$$T_\xi(k)T_\eta(k) = T_\eta(k)T_\xi(k) \quad (\xi, \eta \in \mathfrak{a}, k \in K)$$

in $\text{End}_{\mathbb{C}}(\mathcal{P})$.

Proof. By linearity, it is sufficient to prove this for $\xi = e_i$ and $\eta = e_j$. Furthermore, as a consequence of the polynomial dependence on k , we may assume that $\text{Re}(k) \geq 0$. In the notation of Corollary 2.2, take $L_i = \mathcal{P}_i$, $\partial_j = T_{e_j}(0)$ and $\delta_j = T_{e_j}(k)$; the conditions (i) to (iv), (i') and (ii') are then satisfied. We verify (b) in Corollary 2.2. To this end, let $\{p_j\}_{j=1}^N$ be a closed 1-form (in the terminology of Corollary 2.2). Consider $\omega = \sum_{j=1}^N p_j \otimes de_j^* \in K^1$; then $d(k)\omega = 0$. By Corollary 3.3, ω is exact and we conclude (a) in Corollary 2.2. □

As a consequence of Corollary 3.4 we see that $d(k)$ is a boundary operator on K^\bullet (this statement is in fact equivalent to Corollary 3.4); hence we may form the cohomology groups $H^i(K^\bullet, d(k))$ of the complex $(K^\bullet, d(k))$.

Corollary 3.5. *The following are equivalent:*

1. $k \in K^{\text{reg}}$.
2. *There exists a linear isomorphism $V(k)$ of \mathcal{P} , homogeneous of degree 0, such that $V(k)|_{\mathcal{P}_0} = \text{id}$ and $T_\xi(k)V(k) = V(k)\partial_\xi \quad \forall \xi \in \mathfrak{a}$.*
3. *There exists an endomorphism $V(k)$ of \mathcal{P} , homogeneous of degree 0, such that $V(k)|_{\mathcal{P}_0} = \text{id}$ and $T_\xi(k)V(k) = V(k)\partial_\xi \quad \forall \xi \in \mathfrak{a}$.*

If $V(k)$ exists, it is uniquely determined by the four properties in 2.

Proof. $1 \Rightarrow 2$. Combine Corollary 3.4 and Corollary 2.2.

$2 \Rightarrow 3$. This is trivial.

$3 \Rightarrow 2$. Using the intertwining property, it is easily seen by induction with respect to the homogeneous degree that $V(k)$ is necessarily injective.

$2 \Rightarrow 1$. This is trivial.

The uniqueness statement is part of Corollary 2.2(d). \square

Corollary 3.6. *The following are equivalent:*

1. $H^i(K^\bullet, d(k)) = 0 \quad \forall i > 0$.
2. $\dim_{\mathbb{C}} H^0(K^\bullet, d(k)) = 1$.
3. $k \in K^{\text{reg}}$.

Proof. This is a consequence of Corollary 3.5, except for the implication $1 \Rightarrow 2$. But this follows from the fact that the Euler characteristic of $K^\bullet(n)$ is zero if $n > 0$. \square

In the case where $k_\alpha = k \in \mathbb{C} \ (\forall \alpha \in R)$ we can give an alternative description for the singular polynomials:

Theorem 3.7. *Let $p \in \mathcal{P}_{n,\tau}$ (the τ -isotypical component of \mathcal{P}_n). Then the following are equivalent:*

1. $(\exists k \in \mathbb{C} \text{ such that } d(k)p = 0) \vee (\tau = 1)$.
2. $\exists k_1, k_2 \in \mathbb{C}, k_1 \neq k_2 \text{ such that } d(k_1)d(k_2)p = 0$.
3. $\forall k_1, k_2 \in \mathbb{C} \text{ we have } d(k_1)d(k_2)p = 0$.

Proof. $1 \Rightarrow 3$. By continuity, it is sufficient to prove this for $k_1 \neq k$. Then there exists $\alpha \in \mathbb{C}$ such that $k_2 = \alpha k_1 + (1 - \alpha)k$. Now the result follows from the observation that $d(k_2) = \alpha d(k_1) + (1 - \alpha)d(k)$ and the fact that $d^2(k_1) = 0$ (see Corollary 3.4). If $\tau = 1$, then $d(k_2)p = d(k_1)p$.

$3 \Rightarrow 2$. This is trivial.

$2 \Rightarrow 1$. If $d(k_1)d(k_2)p = 0$, then $\partial d(k_1)d(k_2)p = 0$. Since $\partial p = 0$ we have, using Proposition 3.1 and 3.2,

$$\begin{aligned} (\partial d(k_1) + d(k_1)\partial)d(k_2)p &= d(k_1)(\partial d(k_2) + d(k_2)\partial)p \\ &\Rightarrow ((m + k_1\lambda_\tau)d(k_2) - (m + k_2\lambda_\tau)d(k_1))p = 0 \\ &\Rightarrow \tau = 1 \vee \left(\tau \neq 1 \wedge d\left(\frac{k_2(m + k_1\lambda_\tau) - k_1(m + k_2\lambda_\tau)}{\lambda_\tau(k_1 - k_2)}\right)p = 0 \right) \\ &\Rightarrow \tau = 1 \vee \left(\tau \neq 1 \wedge d\left(-\frac{m}{\lambda_\tau}\right)p = 0 \right). \quad \square \end{aligned}$$

Corollary 3.8. *Let $\{e_1, \dots, e_N\}$ be a basis for \mathfrak{a} . Put*

$$\Delta_i = \sum_{\alpha \in R_+} (\alpha, e_i) \frac{1 - r_\alpha}{\alpha^*} \quad (i = 1, \dots, N).$$

Then

$$\begin{aligned} \{p \in \mathcal{P}_+ | (\partial_i \Delta_j - \partial_j \Delta_i)p &= 0 \ \forall i, j\} \\ &= \{p \in \mathcal{P}_+ | (\Delta_i \partial_j - \Delta_j \partial_i)p = 0 \ \forall i, j\} \\ &= \text{Span}(\{p \in \mathcal{P}_+ | \exists k \in \mathbb{C} \text{ s.t. } T_\xi(k)p = 0 \ \forall \xi \in \mathfrak{a}\} \cup I_+) \end{aligned}$$

where I_+ denotes the invariant polynomials with zero constant term.

Proof. This follows from Theorem 3.7 if one notes that $\Delta_i = T_{e_i}(1) - T_{e_i}(0)$. \square

4. THE SINGULAR SET

In this section we determine the singular set. A bilinear form on \mathcal{P} (introduced in [D4]), together with the work of Heckman [H] on shift operators, will enable us to reduce the question to the determination of the zero locus of a certain polynomial. This zero locus already occurred in the work of Opdam [O] as the poles of the generalized Bessel function. The results in [loc. cit], combined with the explicit form of a certain Poincaré polynomial (introduced and determined by Macdonald), turn out to be sufficient to determine the aforementioned polynomial explicitly in all cases.

As a consequence of the commutativity of the $T_\xi(k)$ for fixed k , there exists an algebra homomorphism from \mathcal{P} to $\text{End}_{\mathbb{C}}(\mathcal{P})$ which sends ξ^* to $T_\xi(k)$ and 1 to the identity. If $p \in \mathcal{P}$, we will denote the corresponding operator by $p(T(k))$. We can thus make the following definition:

$$(p, q)_k = (p(T(k))q)(0) \quad (p, q \in \mathcal{P}).$$

Let us collect some basic properties in the following lemma.

Lemma 4.1. 1. $(\cdot, \cdot)_k$ is symmetric.

2. $(\xi^* p, q)_k = (p, T_\xi(k)q)_k \quad (\xi \in \alpha, p, q \in \mathcal{P}).$

3. $(T_\xi(k)p, q)_k = (p, \xi^* q)_k \quad (\xi \in \alpha, p, q \in \mathcal{P}).$

4. $(g \cdot p, g \cdot q)_k = (p, q)_k \quad (g \in G, p, q \in \mathcal{P}).$

5. $(\mathcal{P}_{r,\tau}, \mathcal{P}_{s,\sigma})_k = 0$ if $r \neq s$ or $\tau \neq \sigma$.

Proof. 1. This was proved in [D4].

2. Obvious from the definition.

3. This follows from part 1 and 2.

4. This is a consequence of (1.1).

5. It is obvious that homogeneous polynomials of different degree are orthogonal. The statement concerning the G -type follows from 4 and the fact that the characters of the representations of G take the same value on g and g^{-1} for all $g \in G$. This is a consequence of a result of Springer [S], stating that any $g \in G$ can be written as $g = \theta_1 \theta_2$ with $\theta_1^2 = \theta_2^2 = 1$, which shows that g and g^{-1} are in the same conjugacy class. \square

The form $(\cdot, \cdot)_k$ can be degenerate. We therefore define

$$\text{Rad}(k) = \{p \in \mathcal{P} \mid (p, q)_k = 0 \quad \forall q \in \mathcal{P}\}.$$

Remark 4.2. It follows from Lemma 4.1 that $\text{Rad}(k)$ is a graded ideal in \mathcal{P} which is invariant under the $T_\xi(k)$ and the G -action. If $p \in \text{Rad}(k)$, then $\prod_{g \in G} g \cdot p \in \text{Rad}(k)$; thus, if $\text{Rad}(k) \neq 0$, $\text{Rad}(k)$ contains a nonzero G -invariant and hence all G -types. Note that the only constant in $\text{Rad}(k)$ is 0.

Proposition 4.3. *The following are equivalent:*

1. $k \in K^{\text{reg}}$.

2. $\text{Rad}(k) = 0$.

Proof. $1 \Rightarrow 2$. If $\text{Rad}(k) \neq 0$, take any nonzero homogeneous $p \in \text{Rad}(k)$ of minimal degree. By the properties of $\text{Rad}(k)$ and the choice of p , p must be annihilated by the $T_{\xi}(k)$, which is a contradiction.

$2 \Rightarrow 1$. This is obvious. \square

As it turns out, K^{sing} is invariant under translations over negative integers. An important ingredient for our proof of this fact is the following theorem of Heckman [H]. Let \mathcal{P}^G denote the G -invariant polynomials, and let $\pi = \prod_{\alpha \in R_+} \alpha^*$ be the fundamental alternating polynomial. The multiplicity function that is constant with value 1 will be denoted by 1_R .

Theorem 4.4. *For all $p, q \in \mathcal{P}^G$ we have*

$$p(T(k))(\pi q) = \pi p(T(k + 1_R))q.$$

Corollary 4.5. *Let $p, q \in \mathcal{P}^G$. Then $(\pi p, \pi q)_k = (\pi, \pi)_k(p, q)_{k+1_R}$.*

Proof. We may assume that p and q are homogeneous and of equal degree. Then

$$\begin{aligned} (\pi p, \pi q)_k &= (\pi, p(T(k))\pi q)_k = (\pi, \pi p(T(k + 1_R))q)_k \\ &= (\pi(T(k))\pi, p(T(k + 1_R))q)_k. \end{aligned}$$

Now both arguments in the right-hand side are constants, so the pairing is just the product of these constants. \square

Corollary 4.6. *If $k + 1_R \in K^{\text{sing}}$, then $k \in K^{\text{sing}}$.*

Proof. If $k + 1_R \in K^{\text{sing}}$, then $\text{Rad}(k + 1_R) \neq 0$ by Proposition 4.3. By Remark 4.2, we see that $\text{Rad}(k + 1_R)$ contains an invariant nonzero q , homogeneous of some degree n . From Corollary 4.5, we infer that $(\pi p, \pi q)_k = 0$ for all $p \in \mathcal{P}^G$, homogeneous of degree n . But then πq is orthogonal to the isotypical component (in degree $n + |R_+|$) corresponding to the determinant representation of G . Thus, as a consequence of Lemma 4.1.5, $\pi q \in \text{Rad}(k)$ and $k \in K^{\text{sing}}$. \square

Lemma 4.7. *Let $k \in K^{\text{sing}}$ and put $n_0 = \max_{n \in \mathbb{N}} \{n | k + n \cdot 1_R \in K^{\text{sing}}\}$. Then $(\pi, \pi)_{k+n_0 \cdot 1_R} = 0$.*

Proof. Recall that $k \in K^{\text{reg}}$ as soon as $\text{Re}(k) \geq 0$, so n_0 is finite. Since $k + n_0 \cdot 1_R \in K^{\text{sing}}$, we see by Remark 4.2 that there exists a nonzero $q \in \mathcal{P}^G$ such that $(\pi p, \pi q)_{k+n_0 \cdot 1_R} = 0$ for all $p \in \mathcal{P}^G$. Thus, by Corollary 4.5 we have $(\pi, \pi)_{k+n_0 \cdot 1_R}(p, q)_{k+(n_0+1) \cdot 1_R} = 0$ for all $p \in \mathcal{P}^G$. Now $(p, q)_{k+(n_0+1) \cdot 1_R}$ cannot be zero for all $p \in \mathcal{P}^G$. For, if this were the case, then

$$q \in \text{Rad}(k + (n_0 + 1) \cdot 1_R)$$

by Lemma 4.1.5 and consequently $k + (n_0 + 1) \cdot 1_R \in K^{\text{sing}}$ by Proposition 4.3, contradicting the definition of n_0 . Thus $(\pi, \pi)_{k+n_0 \cdot 1_R} = 0$. \square

Theorem 4.8. *Let $K^0 = \{k \in K | (\pi, \pi)_k = 0\}$. Then $K^{\text{sing}} = \bigcup_{n=0}^{\infty} \{K^0 - n \cdot 1_R\}$.*

Proof. From Lemma 4.7 we have “ \subset ”. For the reverse inclusion, note that if $(\pi, \pi)_k = 0$, then $\pi \in \text{Rad}(k)$ (for reasons of G -type) so $k \in K^{\text{sing}}$. Thus “ \supset ” is a consequence of Corollary 4.6. \square

The sets K^{sing} and K^0 already occurred in [O], where they were denoted by Z and Z_R . In fact, the description of K^{sing} in Theorem 4.8 and the results in

[loc. cit, §9] show “by inspection” that K^{sing} is the pole set of the generalized Bessel function. There is also a more direct way to see that this must be the case. To this end, fix $k \in K$ and consider the problem of finding a family of G -invariant formal power series $\{J_G(\lambda, k, \xi) | \lambda \in \mathfrak{a}_\mathbb{C}\}$ in ξ such that

$$(4.1) \quad \begin{cases} p(T(k))J_G(\lambda, k, \cdot) = p(\lambda)J_G(\lambda, k, \cdot) & (\forall \lambda \in \mathfrak{a}_\mathbb{C}, \forall p \in \mathcal{P}^G), \\ J_G(\lambda, k, 0) = 1 & (\forall \lambda \in \mathfrak{a}_\mathbb{C}). \end{cases}$$

Evaluating at zero, a solution of (4.1) apparently has the following reproducing property:

$$(J_G(\lambda, k, \cdot), p)_k = p(\lambda) \quad (\forall \lambda \in \mathfrak{a}_\mathbb{C}, \forall p \in \mathcal{P}^G).$$

From this one sees that one must have $k \in K^{\text{reg}}$ and that $J_G(\lambda, k, \cdot)$ is then uniquely determined for all λ , hence equal to $V(k) \frac{1}{|G|} \sum_{g \in G} \exp(g\lambda, \cdot)$. On the other hand, if $k \in K^{\text{reg}}$ then $V(k) \frac{1}{|G|} \sum_{g \in G} \exp(g\lambda, \cdot)$ satisfies (4.1). Hence (4.1) has a formal solution if and only if $k \in K^{\text{reg}}$ and this solution is then equal to $V(k) \frac{1}{|G|} \sum_{g \in G} \exp(g\lambda, \cdot)$. This family of formal power series is in fact meromorphic on $\mathfrak{a}_\mathbb{C} \times K \times \mathfrak{a}_\mathbb{C}$; the poles occur exactly when the restriction of $V(k)$ to the G -invariants has a pole, i.e., in K^{sing} . But (4.1) is just the defining equation for the generalized Bessel function, so K^{sing} and the pole set of the generalized Bessel function coincide as claimed.

In view of Theorem 4.8, all that remains to be done is the determination of K^0 . If R is reducible, then one sees easily that $(\pi, \pi)_k$ factors as the product of the corresponding polynomials for the irreducible components of R . Hence we may assume that R is irreducible, implying that K is at most two-dimensional.

We recall some definitions and results from [O].

Let $B \in \mathbb{C}[K]$ be defined by $B(k) = (\pi, \pi)_k$ and introduce the polynomial b in one variable by $b(s) = B(s1_R)$. Then (see [loc. cit.]) b has degree $|R_+|$. Let $R = \bigcup_{i=1}^m C_i$ be the decomposition of R into minimal G -orbits. For $g \in G$, let $l_i(g)$ be the number of simple reflections r_α with $\alpha \in C_i$ that occur in a minimal expression for g (this is well defined). For $k = \sum_{i=1}^m k_i 1_{C_i} \in K$, put $k^{l(g)} = \prod_{i=1}^m k_i^{l_i(g)}$, and let $P_G(k) = \sum_{g \in G} k^{l(g)}$ be the Poincaré polynomial of G . These polynomials were introduced and determined by Macdonald [M]. If one takes the k_i all equal, they reduce to the Poincaré polynomial of the coinvariant algebra of G . From the results in [O] we extract the following.

Theorem 4.9. 1. $K^{\text{sing}} \subset K^P$, where K^P is the zero-locus of $k \mapsto P_G(\exp(2\pi ik))$ and $\exp(2\pi ik)$ is to be calculated componentwise.

2. $b(k) = |G| \prod_{i=1}^m \prod_{j=1}^{d_i-1} (d_i k + j)$, where the d_i are the primitive degrees of G .

If $m = 1$, Theorem 4.9.2 answers our question. So let us assume that $m = 2$. It follows from Macdonald’s list that K^P consists of a countable union of lines. Since in particular $K^0 \subset K^P$, the following lemma allows us to conclude that B must be a product of these lines.

Lemma 4.10. *Let L be an algebraically closed uncountable field. Let $F \in L[X, Y]$ be nonconstant and suppose that $V(F) \subset \bigcup_{i=1}^\infty V(F_i)$ where the F_i are irreducible (here $V(P)$ denotes the algebraic set determined by P). Then F is the product of a finite number of the F_i up to a constant.*

Proof. We use results of §1.6 in [Fu]. First of all, we may assume that F is irreducible. Since $V(F) = \bigcup_{i=1}^{\infty} (V(F) \cap V(F_i))$ and $V(F)$ is uncountable, there is F_{i_0} such that $V(F) \cap V(F_{i_0})$ is infinite. Hence at least one of the irreducible components of $V(F) \cap V(F_{i_0})$ is an irreducible curve $V(C)$. But then it is a well-known consequence of the Nullstellensatz that F and C must be multiples of each other. Since the same argument applies to F_{i_0} and C , the proof is finished. \square

Thus B splits as a product of linear factors, to be taken from the lines constituting K^P . Now inspection of Macdonald’s list shows that each of these lines is of the form $pk_1 + qk_2 + r$ where p and q are both real (in fact integer) and nonnegative. This implies that B factors as a product of exactly R_+ (the degree of b) linear factors, each of which (by Theorem 4.9.2) intersects $C.1_R$ in $(-1, 0)$.

Let us show how we can now determine B for the dihedral groups $I_2(2m)$ of order $4m$. There are $2m$ positive roots. From Macdonald’s list we have $P_G(t_1, t_2) = (1 + t_1)(1 + t_2)(1 - u^m)/(1 - u)$ where $u = t_1t_2$. It is then easy to see that K^P consists of three sets of lines:

1. $k_1 = \frac{1}{2} + l$ ($l \in \mathbf{Z}$).
2. $k_2 = \frac{1}{2} + l$ ($l \in \mathbf{Z}$).
3. $k_1 + k_2 = \frac{l}{m}$ ($l \in \mathbf{Z}, m \nmid l$).

By inspection, there are only $2m$ lines that intersect $C.1_R$ in $(-1, 0)$, namely $k_1 + \frac{1}{2}$, $k_2 + \frac{1}{2}$, and $k_1 + k_2 + \frac{l}{m}$ ($j = 1, \dots, m-1, m+1, \dots, 2m-1$). So B is known up to a constant, which then follows from Theorem 4.9.2. This method also works for B_n and F_4 . We summarize the results in the following theorem.

Theorem 4.11. *Let R be irreducible. If there is only one G -orbit in R , then*

$$(\pi, \pi)_k = |G| \prod_{i=1}^n \prod_{j=1}^{d_i-1} (d_i k + j),$$

where the d_i are the primitive degrees of G . In the remaining cases $(\pi, \pi)_k$ is equal to

B_n :

$$2^{\frac{1}{2}n(n+1)} n! \left\{ \prod_{j=0}^{n-1} \prod_{l=1}^{j+1} (2jk_1 + 2k_2 + 2l - 1) \right\} \cdot \left\{ \prod_{j=1}^{n-1} \prod_{l=1}^j ((j+1)k_1 + l) \right\},$$

where k_1 corresponds to the long root in the crystallographic root system.

F_4 :

$$\begin{aligned} & 2^{20} 3^4 \cdot \prod_{i=1,2} \{ (2k_i + 1)(3k_i + 1)(3k_i + 2)(2k_i + 2u + 1) \\ & \qquad \qquad \qquad \cdot (2k_i + 2u + 3)(2k_i + 2u + 5) \} \\ & \cdot (2u + 1)^2(2u + 3)^2(4u + 1)(4u + 3)(4u + 5)(4u + 7) \\ & \cdot (6u + 1)(6u + 5)(6u + 7)(6u + 11) \end{aligned}$$

where $u = k_1 + k_2$. License: Copyrighted material. Redistribution and reuse are permitted by the publisher, provided the original work is properly cited. For more information, see <https://www.ams.org/journal-terms-of-use>

$I_2(2m)$:

$$4m^2(2k_1 + 1)(2k_2 + 1) \prod_{\substack{j=1 \\ j \neq m}}^{2m-1} (m(k_1 + k_2) + j).$$

5. SINGULAR POLYNOMIALS: ONE-DIMENSIONAL REPRESENTATIONS AND THE DIHEDRAL GROUPS

The singular set was determined in the previous section for arbitrary G . In general, the set of singular polynomials (as in the introduction) is still unknown. We can, however, determine the singular polynomials completely if G is a dihedral group. For general G , we are able to give a description of the singular polynomials that transform according to a one-dimensional representation of G .

We recall the definition of the k -Laplacian:

$$\Delta(k) = \sum_{i=1}^N T_{\xi_i}^2(k),$$

where $\{\xi_1, \dots, \xi_N\}$ is any orthonormal basis for \mathfrak{a} . If $f \in \mathcal{P}_n$ and $k \in K$, we say that f has a k -Laplace decomposition if f can be written as

$$f = \sum_{j \geq 0} |x|^{2j} f_{n-2j},$$

with $f_{n-2j} \in \mathcal{P}_{n-2j}$ and $\Delta(k)f_{n-2j} = 0$ for all j . We will investigate the values of k sufficient for existence and uniqueness of decompositions of this kind. The result will be helpful in the dihedral case. Put $\gamma = \sum_{\alpha \in R_+} k_\alpha$. Suppose that $\tau \in \widehat{G}$ and that τ is realized on $V_n \subset \mathcal{P}_n$. Let $n_0 = \min_{j \in \mathbb{N}} \{n - 2j \mid \tau \text{ occurs in } \mathcal{P}_{n-2j}\}$. If $n = n_0$, then $\Delta(k)V_n = (0)$, and the decomposition is trivial (but unique).

Lemma 5.1. *Each $f \in V_n$ has a unique k -Laplace decomposition if $-(\frac{N}{2} + \gamma) \neq n_0, n_0 + 1, n_0 + 2, \dots, n - 2$.*

Proof. By Theorem 1.11 in [D1], we have the decomposition

$$f = \sum_{j \leq n/2} |x|^{2j} f_{n-2j},$$

valid for generic values of k , with

$$f_{n-2j} = \left(4^j j! \binom{\frac{N}{2} + \gamma + n - 2j}{j} \right)^{-1} \cdot \sum_{i \leq \lfloor n/2 \rfloor - j} \left(4^i i! \binom{-\frac{N}{2} - \gamma - n + 2j + 2}{i} \right)^{-1} |x|^{2i} \Delta(k)^{i+j} f$$

and $\Delta(k)f_{n-2j} = 0$. The verification is based on the identity

$$\sum_{j=0}^l \frac{1}{j!(l-j)!(A-2j)_i(-A+2j+2)_{l-j}} = 0$$

for $l = 1, 2, \dots$. There are poles at $A = 2, 3, \dots, 2l$. The sum appears as the coefficient of $\Delta(k)^l f$, hence it must be valid for $n - 2l \geq n_0$, that is $l \leq \frac{(n-n_0)}{2}$. This occurs if $\frac{N}{2} + \gamma + n$ avoids $2, 3, 4, \dots, n - n_0$, as claimed. (The effect of the forbidden values is to allow the situation $\Delta(k)(|x|^{2j} f_{n-2j}) = 0$ and $\Delta(k)f_{n-2j} = 0$, resulting in "too many" harmonics of degree n .) \square

The one-dimensional representations. Suppose G has two orbits C_1 and C_2 in R and k has the values k_1 and k_2 , respectively. There is an alternating polynomial for each class, namely $p_i = \prod_{\alpha \in C_i, \alpha \in R_+} \alpha^*$. For $\alpha \in R$ we have $r_\alpha p_i = -p_i$ if $\alpha \in C_i$ and $r_\alpha p_i = p_i$ otherwise. The present discussion specializes easily to the case that G has only one orbit in R .

Recall the definition of the k -gradient:

$$\nabla(k)f = \nabla f + \sum_{\alpha \in R_+} k_\alpha \left(\frac{1 - r_\alpha}{\alpha^*} f \right) \alpha.$$

Obviously, f is singular if and only if $\nabla(k)f = 0$ for some k .

Proposition 5.2. *The singular polynomials of the isotypes of the nontrivial one dimensional representations of G are $p_1^{2n_1+1}$ (if $k_1 = -n_1 - \frac{1}{2}$), $p_2^{2n_2+1}$ (if $k_2 = -n_2 - \frac{1}{2}$) and $p_1^{2n_1+1} p_2^{2n_2+1}$ (if $k_1 = -n_1 - \frac{1}{2}$ and $k_2 = -n_2 - \frac{1}{2}$) for arbitrary $n_1, n_2 = 0, 1, 2, \dots$.*

Proof. The general function of the same isotype as $p_1 p_2$ is $p_1 p_2 g$ where g is invariant. Then

$$\nabla(k)(p_1 p_2 g) = p_1 p_2 \nabla g + p_1 p_2 g (1 + 2k) \sum_{\substack{\alpha \in C_1 \\ \alpha \in R_+}} \frac{\alpha}{\alpha^*} + p_1 p_2 g (1 + 2k_2) \sum_{\substack{\alpha \in C_2 \\ \alpha \in R_+}} \frac{\alpha}{\alpha^*}.$$

For $p_1 p_2 g$ to be singular, g must satisfy

$$\frac{\nabla g}{g} = -(1 + 2k_1) \sum_{\substack{\alpha \in C_1 \\ \alpha \in R_+}} \frac{\alpha}{\alpha^*} - (1 + 2k_2) \sum_{\substack{\alpha \in C_2 \\ \alpha \in R_+}} \frac{\alpha}{\alpha^*}.$$

Thus, g is a scalar multiple of $|p_1|^{-(1+2k_1)} |p_2|^{-(1+2k_2)}$. For this to be an invariant polynomial we must have $-1 - 2k_1 = 2n_1$, $-1 - 2k_2 = 2n_2$ for nonnegative integers n_1, n_2 . The other cases $p_1^{2n_1+1}$, $p_2^{2n_2+1}$ are shown similarly.

The dihedral groups. The dihedral groups $I_2(m)$ act on $\mathbf{R}^2 \simeq \mathbf{C}$. Let $\omega = \exp(2\pi i/m)$, then the group G consists of the reflections $z \mapsto \bar{z}\omega^j$ and the rotations $z \mapsto z\omega^j$ ($0 \leq j \leq m - 1, z \in \mathbf{C}$). G has one conjugacy class of reflections when m is odd, and two classes when m is even. When m is odd, we use k to denote the value of the parameter and write $T(k), \bar{T}(k)$ for the operators of Definition 3.1 in [D2]. The singular polynomials f are those for which $T(k)f = \bar{T}(k)f = 0$. In [D2] we already determined the solutions of $\bar{T}(k)f = 0$, for any k . Essentially we only have to apply the condition $T(k)f = 0$.

The dihedral groups, odd case. Let m be odd ($m \geq 3$). $\text{Span}(\{z^l, \bar{z}^l\})$ and $\text{Span}(\{z^{m-l}, \bar{z}^{m-l}\})$ are of the same isotype for $1 \leq l \leq m - 1$ (and this includes all two dimensional representations of G). For each such l and

each $n = 0, 1, 2, \dots$, there is a unique (up to scalar multiplication) polynomial of isotype l , degree $nm + l$, in the kernel of $\overline{T}(k)$, namely $f_{nm+l}(z) = z^l C_n^{(k, k+1)}(z^m)$ (see [D2, p. 181]). Here $C_n^{(\delta, \varepsilon)}(z)$ is the Heisenberg polynomial defined by

$$C_n^{(\delta, \varepsilon)}(z) = \sum_{j=0}^n \frac{(\delta)_j (\varepsilon)_{n-j}}{j!(n-j)!} \overline{z}^j z^{n-j} \quad (z \in \mathbb{C}),$$

generated by

$$(1 - t\overline{z})^{-\delta} (1 - tz)^{-\varepsilon} = \sum_{n=0}^{\infty} C_n^{(\delta, \varepsilon)}(z) t^n.$$

The polynomial $C_n^{(\delta, \varepsilon)}$ is nonzero, unless $\delta = -n_1, \varepsilon = -n_2$ with $n_1, n_2 = 0, 1, 2, \dots$ and $n > n_1 + n_2$. Now $T(k)f_{nm+l}(z) = (l + mn + mk)f_{nm+l-1}(z)$. Setting $k = -n - l/m$ gives the singular polynomial ($k \notin \mathbb{Z}$, so $C_n^{(k, k+1)}(z^m)$ is not zero). The complex conjugate of f_{nm+l} is the other basis element for the singular polynomials of degree $nm + l$. There are no other singular polynomials of this isotype and degree. They would be harmonic, since $\Delta(k) = 4T(k)\overline{T}(k)$, but by Lemma 5.1 there are only two linearly independent harmonic polynomials provided $-(1 + mk)$ avoids $n_0, n_0 + 1, \dots, nm + l - 2$, and of course $-1 - mk = -1 + mn + l$.

By Lemma 5.1 the alternating polynomial $p(z) = z^m - \overline{z}^m$, gives rise to the singular polynomials $(z^m - \overline{z}^m)^{2n+1}$ with $k = -\frac{1}{2} - n$.

This is the entire list of singular polynomials because the only other representation of G is the trivial one.

The dihedral groups, even case. Let $m \geq 2$ be even and let $G = I_2(2m)$. Put $\omega = \exp(\pi i/m)$. Associate the parameter values k_1 to the reflections $z \mapsto \overline{z}\omega^{2j}$ and k_2 to $z \mapsto \overline{z}\omega^{2j+1}$ ($0 \leq j \leq m - 1$). We will write $\Delta(k_1, k_2)$, $T(k_1, k_2)$ and $\overline{T}(k_1, k_2)$ for the operators. For each $l \in \{1, 2, \dots, m - 1\}$ there is an isotype realized on $\text{span}\{z^l, \overline{z}^l\}$ and on $\text{span}\{z^{2m-l}, \overline{z}^{2m-l}\}$.

The polynomials of degree $sm + l$ ($1 \leq l \leq m - 1, s = 0, 1, 2, \dots$) annihilated by $\overline{T}(k_1, k_2)$ are (see 3.19 in [D2]) $z^l f_s(z^m)$ where

$$\begin{aligned} f_{2n}(re^{i\theta}) &= r^{2n} P^{(k_1 - \frac{1}{2}, k_2 - \frac{1}{2})}(\cos 2\theta) + \left(\frac{i}{2}\right) r^{2n} (\sin 2\theta) P_{n-1}^{(k_1 + \frac{1}{2}, k_2 + \frac{1}{2})}(\cos 2\theta), \\ f_{2n+1}(re^{i\theta}) &= r^{2n} \left((n + k_1 + \frac{1}{2})(r \cos \theta) P_n^{(k_1 - \frac{1}{2}, k_2 + \frac{1}{2})}(\cos 2\theta) \right. \\ &\quad \left. + i(n + k_2 + \frac{1}{2})(r \sin \theta) P_n^{(k_1 + \frac{1}{2}, k_2 - \frac{1}{2})}(\cos 2\theta) \right) \\ &\quad (r \geq 0, 0 \leq \theta < 2\pi, n = 0, 1, 2, \dots). \end{aligned}$$

The Jacobi polynomial is

$$P_n^{(\delta, \varepsilon)}(t) = \frac{(\delta + 1)_n}{n!} {}_2F_1 \left(\begin{matrix} -n, n + \delta + \varepsilon + 1 \\ \delta + 1 \end{matrix}; \frac{1-t}{2} \right);$$

there are no poles (parameter values) and the polynomial is of degree n , provided $(n + \delta + \varepsilon + 1)_n \neq 0$. This condition applies in the present case. By the formula in [D2, p. 183] we have

$$T(k_1, k_2) z^l f_s(z^m) = (l + ms + m(k_1 + k_2)) z^{l-1} f_s(z^m).$$

Thus for $k_1 + k_2 = -s - l/m$, $z^l f_s(z^m)$ (and its complex conjugate) is a singular polynomial. The parameters (δ, ε) in the Jacobi polynomials in f_{2n} , f_{2n+1} satisfy $n + \gamma + \delta + 1 = n + k_1 + k_2 + s$ with $s = 0, 1$ or 2 , and $k_1 + k_2 \notin \mathbf{Z}$.

There are no other singular (or harmonic) polynomials for these parameter values and this degree. By Lemma 5.1, $-(1 + m(k_1 + k_2))$ must avoid $n_0, n_0 + 1, \dots, ms + l - 2$ and $-(1 + m(k_1 + k_2)) = ms + l - 1$. The singular polynomials of alternating isotype are $(z^m - \bar{z}^m)^{2n_1+1}$, $(z^m + \bar{z}^m)^{2n_2+1}$ and the product of these two, with $k_1 = -n_1 - \frac{1}{2}$ for the first, $k_2 = -n_2 - \frac{1}{2}$ for the second and both conditions for the product. Lemma 5.1 does not apply in the product case. Here $-1 - m(k_1 + k_2)$ must avoid $2m, 2m + 1, \dots, 2m(n_1 + n_2 + 1) - 2$ but here $k_1 + k_2 = -n_1 - n_2 - 1$ and $m(n_1 + n_2 + 1)$ appears in the list if $n_1 + n_2 \geq 2$. Again we have described all the singular polynomials.

6. ESTIMATES FOR $\dim_{\mathbf{C}}(S_{m,\tau})$ IF $k_\alpha = k \ \forall \alpha \in R$

Recall the following constructions from [D5]. Let $\mathfrak{h} = \mathfrak{a} \otimes_{\mathbf{R}} \mathbf{C}$. The form (\cdot, \cdot) extends to a bilinear form on \mathfrak{h} , which will again be denoted by (\cdot, \cdot) . Any $\xi \in \mathfrak{h}$ gives rise, via this form, to $\xi^* \in \mathfrak{h}^*$ as in the Introduction. Put $\mathfrak{h}^{\text{reg}} = \{\xi \in \mathfrak{h} \mid \alpha^*(\xi) \neq 0 \ \forall \alpha \in R\}$. Let $\mathcal{O}(\mathfrak{h}^{\text{reg}})$ denote the holomorphic functions on $\mathfrak{h}^{\text{reg}}$. Let $\tau \in \widehat{G}$, and let V_τ be the corresponding representation space of dimension d_τ . Define the following connection $\nabla^\tau(k)$ on $\mathcal{O}(\mathfrak{h}^{\text{reg}}) \otimes V_\tau$ (the holomorphic sections in the trivial bundle $\mathfrak{h}^{\text{reg}} \times V_\tau$):

$$\nabla_\xi^\tau(k)(f \otimes v) = (\partial_\xi f) \otimes v + k \sum_{\alpha \in R_+} (\alpha, \xi) \left(\frac{f}{\alpha^*} \right) \otimes (1 - \tau(r_\alpha))v.$$

This connection is integrable and has regular singularities.

Let ρ denote the natural representation of G on $\mathcal{O}(\mathfrak{h}^{\text{reg}})$:

$$(g \cdot f)(z) = f(g^{-1}z) \quad (g \in G, f \in \mathcal{O}(\mathfrak{h}^{\text{reg}}), z \in \mathfrak{h}^{\text{reg}}).$$

Then $\nabla^\tau(k)$ is equivariant for $\rho \otimes \tau$ in the sense that

$$\nabla_\xi^\tau(k)(\rho(g)f \otimes \tau(g)v) = (\rho(g) \otimes \tau(g))(\nabla_{g^{-1}\xi}^\tau(k)(f \otimes v)).$$

Hence we can view $\nabla^\tau(k)$ as the inverse image of a connection $\overline{\nabla^\tau(k)}$ on $\overline{\mathcal{V}}_\tau = \mathcal{O}(G \backslash \mathfrak{h}^{\text{reg}}) \otimes_{\mathbf{C}} \overline{V}_\tau$, where \overline{V}_τ is the local system on $X = G \backslash \mathfrak{h}^{\text{reg}}$ defined by $\overline{V}_\tau = \mathfrak{h}^{\text{reg}} \times_G V_\tau$. The fundamental group of X is the braid group of G (see [B]). The monodromy representation of $\overline{\nabla^\tau(k)}$ factorizes through the Hecke algebra $H_G(q(k))$ where $q(k) = \exp(-2\pi i k)$, and where we use the convention that the generators of $H_G(q)$ satisfy $(T_i - 1)(T_i + q) = 0$. The resulting representation of $H_G(q)$ is called $Y_\tau(k)$. From the construction it is clear that Y_τ is defined over the ring \mathcal{E} of entire functions on $K \cong \mathbf{C}$. By deformation theoretical arguments it was shown in [D5] that $Y_\tau(k)$ is irreducible and principal when $\Psi_l(q^{-1}) = \frac{q^{lR_+} P_G(q^{-1})}{d_\tau(q^{-1})} \neq 0$ (here d_τ is the generic degree of τ and P_G is the Poincaré polynomial of the coinvariant algebra of G). Hence loss of irreducibility of some $Y_\tau(k)$ can occur only if $P_G((q(k))^{-1}) = 0$. Consequently, the zeros of $P_G(q^{-1})$ correspond to nonsemisimple specializations of H_G (see [GU, Y]). Also observe that the trivial representation is *never* principal if $H_G(q)$ is not semisimple.

We will now apply results of the theory of algebras defined over local domains that are complete. Let $k_0 \in K \cong \mathbb{C}$, and write $x = k - k_0$. Let $\mathcal{R} = \mathbb{C}\{x\}$. (It is important that \mathcal{R} is a complete local principal ideal domain for these statements and this is the reason that we assume $k_\alpha = k \forall \alpha$). Let $\mathcal{K} = \mathbb{C}\{\{x\}\} = \{\sum_{k \geq N} c_k x^k \mid N \in \mathbb{Z}, c_k \in \mathbb{C}\}$ be its quotient field. Define $X_\tau = \mathbf{R} \otimes_\varepsilon Y_\tau$ and $A = \mathcal{R} \otimes_{\mathbb{C}[q, q^{-1}]} H_G$. Finally put $\bar{X}_\tau = X_\tau / (x)\mathcal{R}$ (a module for $A/(x)\mathcal{R} = \bar{A} = H_G(q(k_0))$), and $(X_\tau)_{\mathcal{K}} = X_\tau \otimes_{\mathcal{R}} \mathcal{K}$ (a module for $A_{\mathcal{K}} = \mathcal{K} \otimes_{\mathcal{R}} A = \mathcal{K} \otimes_{\mathbb{C}[q, q^{-1}]} H_G$). Clearly $\{(X_\tau)_{\mathcal{K}}\}_{\tau \in \widehat{G}}$ is a complete set of representatives for the irreducibles of $A_{\mathcal{K}}$. We are now in the situation as described on p. 67 in [Fe].

Definition 6.1. Let *triv* be the one-dimensional representation of $H_G(q)$ that is defined by sending the T_i to 1. The multiplicity of *triv* $\in (\bar{A})_{\text{irr}}$ in a decomposition series of \bar{X}_τ will be denoted by $\delta_\tau(q_0)$. The multiplicity of *triv* in \bar{X}_τ as a submodule (i.e., the multiplicity of *triv* in the socle of \bar{X}_τ) will be denoted by $\Delta_\tau(k_0)$.

This definition makes sense because by [Fe, Theorem 17.7] the number $\delta_\tau(q_0)$ depends on τ and q_0 only. In fact, these decomposition numbers [Fe, p. 67] have a different interpretation in terms of representation theory of A by virtue of a theorem of Brauer. In our situation we have a one-to-one correspondence between principal indecomposable modules of A and $\bar{A} = H_G(q_0)$ (since [Fe, Theorem 13.7] applies). Let U denote the principal indecomposable module of A that corresponds to *triv* $\in (A)_{\text{irr}}$ (hence \bar{U} is principal indecomposable for $\bar{A} = H_G(q_0)$ and $U/\text{rad}(U) = \bar{U}/\text{rad}(\bar{U}) = \text{triv}$). We have

Theorem 6.2 (Brauer, see [Fe, Theorem 17.8]).

$$U_{\mathcal{K}} \cong \bigoplus_{\tau \in \widehat{G}} \delta_\tau(q_0) (X_\tau)_{\mathcal{K}}.$$

Corollary 6.3. If k_0 is such that $H_G(q_0)$ is not semisimple, then there exists at least one $\tau \in \widehat{G}$, $\tau \neq \text{triv}$ such that $\delta_\tau(q_0) \neq 0$.

Proof. Suppose that $\delta_\tau(q_0) = 0$ if $\tau \neq \text{triv}$. Clearly $\delta_{\text{triv}}(q_0) = 1$ and by Theorem 6.2 we obtain that $U_{\mathcal{K}} \cong (\text{triv})_{\mathcal{K}}$. This implies that $U = \text{triv}$, and we thus conclude that *triv* is principal. This contradicts the results of Yamane in [Y]. \square

Remarks 6.4. 1. $\delta_\tau(q_0) = 0$ if $\Psi_\tau(q_0^{-1}) \neq 0$, since Yamane proved that $\Psi_\tau(q_0^{-1}) \neq 0$ implies that \bar{X}_τ is principal and irreducible.

2. Of course $\Delta_\tau(k_0) \leq \delta_\tau(q_0)$ and $\Delta_\tau(k_0) = 0$ if $k_0 > 0$.

Proposition 6.5. Assume $\tau \neq \text{triv}$ and $k_0 \notin \frac{1}{2}\mathbb{N}$. Then $\dim_{\mathbb{C}} S_\tau(k_0) = d_\tau \Delta_\tau(k_0)$.

Proof. Choose an explicit unitary matrix realization M_τ of τ on \mathbb{C}^{d_τ} (where d_τ is the dimension of the representation space of τ), and let $(\mathcal{Z}^\tau, \bar{\nabla}^\tau(k_0))$ be the associated connection as above. Let $F \subset S(k_0)$ be a copy of τ embedded in $S(k_0)$. There exists a 1-dimensional space of column vectors $(f_1, \dots, f_{d_\tau})^t (f_i \in F)$ such that

$$(6.1) \quad (f_i^g)_{i=1, \dots, d_\tau}^t = M_\tau(g^{-1})(f_1, \dots, f_{d_\tau})^t$$

and $T_\xi(k_0)f = 0 \forall i$ implies $\nabla_\xi^t(k_0)(f_1, \dots, f_{d_\tau})^t = 0$. Hence $\sigma = (f_1, \dots, f_{d_\tau})^t$ is in $(\mathcal{Z}^\tau)^{\nabla^t(k_0)}$ and because of (6.1) we see that σ is fixed for the monodromy action. This sets up a one-to-one correspondence between $(Y_\tau(k_0))^{\text{triv}} (= (\bar{X}_\tau)^{\text{triv}})$ and vectors $(f_1, \dots, f_{d_\tau})^t$ of elements $f_i \in S(k_0)$ satisfying (6.1). Namely, suppose on the other hand that we are given $\sigma = (f_1, \dots, f_{d_\tau})^t \in (\mathcal{Z}^\tau)^{\nabla^t(k_0)}$ that is trivial for monodromy. The exponents along the walls for the equation $\nabla_\xi^t(k_0)\sigma = 0$ are 0 and $-2k$. Since we assume that $k \notin \frac{1}{2}\mathbb{N}$ we may conclude that f_i extends holomorphically along the walls, and that (6.1) holds. Finally we consider the equation $(\sum_{i=1}^N x_i \nabla_i^t(k_0))\sigma = 0$. It follows that f_i is homogeneous of degree $-k_0 \lambda_\tau$. Hence $-k_0 \lambda_\tau \in \mathbb{N}$, $f_i \in \mathcal{P}_{-k_0 \lambda_\tau}$, and f_i satisfies $T_\xi(k_0)f_i = 0$ ($\forall i$). The result follows easily from this one-to-one correspondence. \square

Corollary 6.6. *If $\tau \neq \text{triv}$ then $\dim_{\mathbb{C}}(S_{m, \tau}) \leq d_\tau \delta_\tau(\exp(\frac{2\pi i m}{\lambda_\tau}))$. Hence $m \mapsto \dim_{\mathbb{C}}(S_{m, \tau})$ is bounded by a function that is periodic in m with period λ_τ .*

Proof. $S_{m, \tau} = S_\tau(-\frac{m}{\lambda_\tau})$. \square

Remark 6.7. We hope that in fact equality holds. This would be very satisfying, since this would mean that the equations of Corollary 3.8 completely describe the decomposition of $(U)_{\mathcal{R}}$.

Examples 6.8.

1. $\tau = \text{sign representation}$. Then $\delta_\tau(q) = 1$ if $q = -1$, and zero otherwise. Moreover $\lambda_\tau = |R|$, so $q = \exp(\frac{2\pi i m}{\lambda_\tau}) = -1$ iff $m = (p - \frac{1}{2})|R|$ ($p \in \mathbb{N}$). On the other hand, if we let $\pi = \prod_{\alpha \in R_+} \alpha^*$ denote the fundamental alternating polynomial, then

$$T_\xi \left(\frac{1}{2} - p \right) \pi^{2p-1} = \left(\partial_\xi - (1 - 2p) \sum_{\alpha \in R_+} (\alpha, \xi) \frac{1}{\alpha^*} \right) \pi^{2p-1} = (\pi^{2p-1} \partial_\xi) \pi = 0.$$

Hence $\dim_{\mathbb{C}}(S_{m, \tau}) = 1$ if $m = (p - \frac{1}{2})|R|$ and zero otherwise.

2. *The dihedral cases.* According to [CIK], the two-dimensional representations for $H_{I_2(m)}(q)$ can be defined explicitly as follows. Let σ_1, σ_2 be the standard generators. Then put $\pi_j(\sigma_1) = \begin{pmatrix} 1 & -q \\ 0 & -q \end{pmatrix}$ and $\pi_j(\sigma_2) = \begin{pmatrix} -q & 0 \\ -(\zeta^j + \zeta^{-j}) & 1 \end{pmatrix}$ where $\zeta = \exp(\frac{2\pi i}{m})$ and $j = 1, \dots, \frac{m-1}{2}$ (if m is odd) or $j = 1, \dots, \frac{m}{2} - 1$ (if m is even). From this we find that $\delta_{\pi_j(q)} = 1$ if $q = \zeta^{\pm j}$ and zero otherwise (if $q = \zeta^{\pm j}$ then $\pi_j(\sigma_i)(q, q + 1)^t = -q(q, q + 1)^t$ ($i = 1, 2$)). Furthermore, it is easy to see that $\lambda_{\pi_j} = m$ ($\forall j$). Hence we infer from $q_j^\pm = \zeta^{\pm j} = \exp(-2\pi i (\frac{m_j^\pm}{m}))$ that $m_j^\pm = mp \pm j$ ($p \in \mathbb{N}$). By Corollary 6.6 we see that π_j can occur in S_n only if $n = mp \pm j$ for some $p \in \mathbb{N}$ and that the multiplicity is at most 1. According to the results of §5 this multiplicity equals 1 for all n of the form $n = mp \pm j$ ($p \in \mathbb{N}$). Hence, equality holds in Corollary 6.6 in this example.

3. *The symmetric groups.* We discuss some conjectures for the irreducible representations of $G = S_n$ (the Weyl group of type A_{n-1}) which appear in the space of singular polynomials. Recall that such representations correspond to partitions $\mu = (\mu_1, \mu_2, \dots, \mu_j)$ with $\mu_i \in \mathbb{Z}_+, \mu_1 \geq \mu_2 \geq \dots \geq \mu_j > 0$ and $n = |\mu| = \sum_i \mu_i$. We will use μ as the label for the representation so that $\chi_{(\mu)} = 1$ and $\chi_{(\mu)}$ is the sign character.

The conjecture is that the partitions μ for which $S_\mu(k_0) \neq (0)$ for some k_0 are those with three or more (nonzero) parts of the form $\mu = (sd - 1, d - 1, \dots, d - 1, r)$ with $s = 1, 2, 3, \dots, 0 < r \leq d - 1$ and $k_0 = -j/d$ with $\gcd(j, d) = 1$ ($j \geq 1$), and all of the two-part partitions (μ_1, μ_2) with $k_0 = -j/(\mu_1 + 1)$ for $\gcd(j, \mu_1 + 1) < (\mu_1 + 1)/\mu_2$. Here \gcd denotes greatest common divisor.

The evidence for this conjecture comes from the Specht modules for the Hecke algebra $H_{S_n}(q)$ constructed by Dipper and James [DJ1, DJ2]. Because of our different defining relations (we use generators T_1, T_2, \dots, T_{n-1} satisfying $T_i^2 = q1 + (1 - q)T_i$, $T_i T_{i+1} T_i = T_{i+1} T_i T_{i+1}$ ($1 \leq i \leq n - 2$), $T_i T_j = T_j T_i$ for $1 \leq i < i + 2 \leq j \leq n - 1$, so that the eigenvalues of the generators are 1 and $-q$, rather than q and -1), our partition labels are the transposes of those in [DJ1, DJ2].

For each partition μ , Dipper and James construct an $H_{S_n}(q)$ -module M^μ whose basis vectors correspond to certain tableaux with shape μ , and then an irreducible submodule S^μ , the Specht module. The action of the generators T_i on this module is defined by the so-called Garnir relations. We are concerned with the existence of an $H_{S_n}(q)$ -invariant element of the dual of S^μ , that is, one which annihilates $v - T_i v$ for each basis element v of S^μ and $1 \leq i \leq n - 1$. From the Garnir relations one can show that the desired property holds exactly when the Gaussian polynomial (in the parameter q) $[\mu_j + 1]_q = 0$ for each (nonzero) part μ_j and $1 \leq i \leq \mu_{j+1}$ (provided $\mu_{j+1} > 0$). The coefficient of q^l in the Gaussian polynomial $[\mu_j + 1]_q = \prod_{l=1}^{\mu_j} (q^{\mu_j + 1 - l} - 1) / (q^l - 1)$ is the number of subsets E of size i of the set $\{1, 2, \dots, m\}$ for which $|\{(n_1, n_2) | n_1 \in E, n_2 \in E^c, n_1 > n_2\}| = l$. This is related to the action of T_i on S^μ .

For a partition μ , let $Z(\mu)$ be the common zero set of $\{[\mu_j + 1]_q | 1 \leq i \leq \mu_{j+1}\}$. Then $Z(\mu) = \emptyset$ except for the partitions described above. Let $P(j) = \{q \in \mathbb{C} | q^j = 1 \text{ and } q^s \neq 1 \text{ for any } s = 1, 2, \dots, j - 1\}$ (that is, the primitive j th roots of unity). For $\mu = (sd - 1, d - 1, \dots, d - 1, r)$ (with $s \geq 1, 0 < r \leq d - 1$), $Z(\mu) = P(d)$. For $\mu = (\mu_1, \mu_2)$, $Z(\mu)$ is the union of all $P(d)$ such that d divides $(\mu_1 + 1)$ and $d > \mu_2 > 0$.

It is not difficult to see that for each n and $\mu_1 \leq n - 1$ and each divisor d of $\mu_1 + 1$ there is exactly one partition μ ($|\mu| = n$) with $P(d) \subset Z(\mu)$, and so each zero of $(q^{\mu_1 + 1} - 1)/(q - 1)$ is associated to a unique μ . Recall the equation $q = -e^{2\pi i k}$.

Several tasks remain to be done to prove our conjecture. One must establish a correspondence between the Specht modules and the monodromy representation Y_μ (see the beginning of §6), and further show that there is a function with trivial monodromy (a singular polynomial) exactly when the dual of the Specht module has an invariant vector, as described above.

Another problem is to prove the existence of singular polynomials in $S_\mu(k)$ for the appropriate degrees of homogeneity, that is, $-k\lambda_\mu$ (see (1.2) for the definition of λ_μ). The zero-set of each factor of the Poincaré series $\prod_{d=2}^n ((q^d - 1)/(q - 1))$ of S_n corresponds to the union of zero-sets $Z(\mu)$ with $\mu_1 = d - 1$. By use of Young's formula we see that

$$\lambda_\mu = (n - \mu_1)((\mu_1 + 1) + l(\mu_2 + 1)) - \mu_2(\mu_2 + 1)l(l + 1)/2$$

for $\mu = (\mu_1, \mu_2, \dots, \mu_2)$ with $(l + 1)$ parts or $\mu = (\mu_1, \mu_2, \dots, \mu_2, r)$ with

$(l+2)$ parts, and this supports the conjectured link to the singular values $-j/d$, $\gcd(j, d) = 1$ (when $\mu_1 = sd - 1$, $\mu_2 = d - 1$).

REFERENCES

- [B] E. Brieskorn, *Die Fundamentalgruppe des Raumes der regulären Orbits einer komplexen Spiegelungsgruppe*, Invent. Math. **12** (1971), 57–61.
- [CIK] C. W. Curtis, N. Iwahori, and R. W. Kilmoyer, *Hecke algebras and the characters of parabolic type of finite groups with BN-pairs*, Publ. Math. Inst. Hautes Etudes Sci. **40** (1971), 81–116.
- [D1] C. F. Dunkl, *Reflection groups and orthogonal polynomials on the sphere*, Math. Z. **197** (1988), 33–60.
- [D2] ———, *Differential-difference operators associated to reflection groups*, Trans. Amer. Math. Soc. **311** (1989), 167–183.
- [D3] ———, *Operators commuting with Coxeter group actions on polynomials*, Invariant Theory and Tableaux (D. Stanton, ed.), Springer-Verlag, 1990, pp. 107–117.
- [D4] ———, *Integral kernels with reflection group invariance*, Canad. J. Math. **43** (1991), 1213–1227.
- [D5] ———, *Differential-difference operators and monodromy representations of Hecke algebras*, Pacific J. Math. **159** (1993), 271–298.
- [DJ1] R. Dipper and G. James, *Representations of Hecke algebras of general linear groups*, Proc. London Math. Soc. (3) **52** (1986), 20–52.
- [DJ2] ———, *Blocks and idempotents of Hecke algebras of general linear groups*, Proc. London Math. Soc. (3) **54** (1987), 57–82.
- [Fe] W. Feit, *The representation theory of finite groups*, North-Holland, 1982.
- [Fu] W. Fulton, *Algebraic curves*, Addison-Wesley, 1989.
- [GU] A. Gyoja and K. Uno, *On the semisimplicity of Hecke algebras*, J. Math. Soc. Japan **41** (1989), 75–79.
- [H] G. J. Heckman, *A remark on the Dunkl differential-difference operators*, Harmonic Analysis on Reductive Groups (W. Barker and P. Sally, eds.), Birkhäuser, Basel, 1991, pp. 181–191.
- [M] I. G. Macdonald, *The Poincaré series of a Coxeter group*, Math. Ann. **199** (1972), 161–174.
- [O] E. M. Opdam, *Dunkl operators, Bessel functions and the discriminant of a finite Coxeter group*, Comput. Math. **85** (1993), 333–373.
- [S] T. A. Springer, *Regular elements of finite reflection groups*, Invent. Math. **25** (1974), 159–198.
- [Y] H. Yamane, *Irreducible projective modules of the Hecke algebras of finite Coxeter groups*, J. Algebra **127** (1989), 373–384.

(C. F. Dunkl) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF VIRGINIA, CHARLOTTESVILLE, VIRGINIA 22903

E-mail address: cfd5z@virginia.edu

(M. F. E. de Jeu and E. M. Opdam) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF LEIDEN, P. O. BOX 9512, 2300 RA LEIDEN, THE NETHERLANDS

E-mail address: jeu@rulwinw.leidenuniv.nl

E-mail address: opdam@rulwinw.leidenuniv.nl