

1 **Title Page**

2 **The title:**

3 Sleep and physical activity in relation to all-cause, cardiovascular disease, and cancer mortality
4 risk.

5 **Full name, postal address and e-mail of the corresponding author:**

6 Emmanuel Stamatakis.

7 Postal address: Hub D17, Charles Perkins Centre L6 West, the University of Sydney, NSW
8 Australia 2006

9 E-mail: emmanuel.stamatakis@sydney.edu.au

10 **Full name, department, institution, city and country of all co-authors:**

11 Bo-Huei Huang, MSc^a, Mitch J Duncan, PhD^{b, c}, Peter A Cistulli, PhD^d, Natasha Nassar, PhD
12 ^d, Mark Hamer, PhD^e, Emmanuel Stamatakis, PhD^a

13 ^a Charles Perkins Centre, School of Health Sciences, Faculty of Medicine and Health, the
14 University of Sydney, Camperdown, Australia

15 ^b Priority Research Centre for Physical Activity and Nutrition, the University of Newcastle,
16 Callaghan, Australia

17 ^c School of Medicine & Public Health; Faculty of Health and Medicine, the University of
18 Newcastle, Callaghan, Australia

19 ^d Charles Perkins Centre, Sydney Medical School, Faculty of Medicine and Health, the
20 University of Sydney, Camperdown, Australia

21 ^e Institute Sport Exercise Health, Division of Surgery and Interventional Science, University
22 College London, London, UK

23 **Word count, excluding title page, abstract, references, figures and tables: 3,341**

1 **ABSTRACT**

2 **Objectives** Although both physical inactivity and poor sleep are deleteriously associated with
3 mortality, the joint effects of these two behaviours remain unknown. This study aimed to
4 investigate the joint association of physical activity (PA) and sleep with all-cause/cause-
5 specific mortality risks.

6 **Methods** 380,055 participants (55.9±8.1 years, 55% female) from the UK Biobank were
7 included. Baseline PA levels were categorized as high, medium, low, and no moderate-to-
8 vigorous PA (MVPA) based on public health guidelines. We categorized sleep into healthy,
9 intermediate, and poor with an established composited sleep score of chronotype, sleep,
10 insomnia, snoring, and daytime sleepiness. We derived twelve PA–sleep combinations,
11 accordingly. Mortality risks were ascertained to May 2020 for all-cause, total cardiovascular
12 disease (CVD), CVD sub-types (coronary heart disease, hemorrhagic stroke, ischemic stroke),
13 as well as total cancer and lung cancer.

14 **Results** After an average follow-up of 11.1 years, sleep scores showed dose-response
15 associations with all-cause, total CVD, and ischemic stroke mortality. Compared to high PA-
16 healthy sleep group (reference), the no MVPA-poor sleep group had the highest mortality risks
17 for all-cause (hazard ratio [95% confidence intervals], (1.39 [1.20 to 1.61]), total CVD (1.44
18 [1.09 to 1.88]), total cancer (1.30 [1.06 to 1.59]), and lung cancer (1.59 [1.08 to 2.34]). The
19 deleterious associations of poor sleep with all outcomes, except for stroke, was amplified with
20 lower PA.

21 **Conclusion** The detrimental associations of poor sleep with both all-cause and cause-specific
22 mortality risks are exacerbated by low PA, suggesting likely synergistic effects. Our study
23 supports the need to target both behaviours in research and clinical practice.

24 **Keywords:**

25 sleep patterns; cohort study; risk factor; synergistic effect.

1 **KEY MESSAGES**

2 **what are the new findings?**

3 Among middle-aged UK adults without obstructive sleep apnea history or class III obesity,
4 PA at levels above the WHO guideline (600 MET-mins/wk) threshold eliminated most of the
5 deleterious associations of poor sleep with mortality.

6 **how might it impact on clinical practice in the near future?**

7 As emerging evidence supports a synergistic effect of sleep and PA on health outcomes,
8 future trials concurrently targeting both behaviours are warranted.

1 INTRODUCTION

2 Both physical inactivity and poor sleep are deleteriously associated with all-cause,
3 cardiovascular diseases (CVD), and cancer mortality.[1-5] Both behaviours are complex, and
4 their health effects are dependent upon various characteristics. For example, health-related
5 sleep characteristics include duration, quality, and timing. Sleep duration has shown a
6 curvilinear relationship (U-shaped) with all-cause, CVD, and cancer mortality, with the lowest
7 risk observed among those obtaining between 7-8 hours per night.[1-3, 6] Based on a Swedish
8 cohort (n = 70,973) with a 15-year follow-up, Bellavia et al. (2014) suggested that a low
9 physical activity (PA) level could exaggerate the detrimental association between inadequate
10 sleep duration and all-cause and CVD mortality risk.[6] Although sleep quality and timing
11 (e.g., snoring and chronotype) are also associated with all-cause and cause-specific
12 mortality,[7] to our knowledge, very few studies have examined the synergistic effect of PA
13 and sleep quality on mortality while the results were inconsistent.[8, 9] The lack of a
14 standardized sleep measurement might hinder the investigation of sleep concerning health and
15 the joint effects with other behaviours.[3]

16 Besides the independent health effects of sleep duration, quality, and timing, these sleep
17 characteristics have shown a potential joint effect on health outcomes. Poor sleep quality, in
18 combination with either inadequate or excess sleep,[7, 10] is associated with increased
19 mortality risk. Based on the Sleep Heart Health Study with an 11.4-year follow-up (n = 4,994),
20 Bertisch et al. (2018) found that only the combination of inadequate sleep duration and poor
21 sleep quality was associated with incident CVD risk.[11] This interaction between sleep
22 characteristics, as well as the heterogeneity of the sleep characteristics chosen between studies,
23 might produce less consistent results.[3, 12] Recently, Fan et al. (2020) proposed a novel score

1 integrating both duration, quality, and timing into a single sleep measurement.[12] They
2 demonstrated its capacity to identify high-risk populations for incident CVD based on the UK
3 Biobank cohort with an 8.5-year follow-up (n = 385,292).

4 PA and sleep could be co-dependent and influence health conditions through related
5 pathways.[6, 8, 9, 13-16] For example, PA might improve health outcomes through improving
6 metabolic fitness (e.g., reducing insulin resistance), maintaining a stable circadian rhythm and
7 a healthy sleep pattern, and enhancing energy expenditure.[8, 14, 15] Prolonged sleep duration
8 may decrease the time available for PA due to the time-dependency between the two
9 behaviours.[6] Beyond their independent health effects, the potential joint effects of these two
10 key behaviours remain largely unknown.[17] We investigated the joint association of PA and
11 a novel comprehensive sleep score with all-cause and subtype-specific CVD and cancer
12 mortality risks.

13

14 **METHODS**

15 The UK Biobank is a prospective cohort of 502,616 participants of adults aged 37-73
16 years, recruited from 22 centres across the UK between 2006-2010, with ethical approval by
17 the National Research Ethics Service (Ref 11/NW/0382). Participants completed
18 questionnaires, interviews, physical measurements at baseline assessments, and consented to
19 the use of their de-identified data and access to their national health-related hospital and death
20 records. Detailed methods are described elsewhere.[18]

21 **Outcomes**

1 Date of death was obtained from death certificates through data linkage with national
2 datasets from the National Health Service (NHS) Information Centre (England and Wales) and
3 the NHS Central Register Scotland (Scotland). The cause of death (including both primary and
4 contributory causes of death) was defined based on the international classification of diseases,
5 10th revision (ICD10)(**Supplementary File, Table S1**). In short, total CVD was defined as
6 diseases of the circulatory system, excluding hypertension, diseases of arteries and lymph.
7 Total cancer was defined as neoplasms, excluding *in situ*, benign, uncertain, or non-well-
8 defined cancers. Three CVD sub-types and one cancer site which have been previously linked
9 to sleep disorders in published systematic reviews were also defined: coronary heart disease,
10 hemorrhagic stroke, ischemic stroke, and lung cancer.[5, 16] The censoring date was May
11 2020. Participants were followed up from the date of attendance at the recruitment centre to
12 the date of death or censorship, whichever came first. The data linkage was updated two or
13 three times per year.

14 **Exposures**

15 PA was quantified using the modified short-form International Physical Activity
16 Questionnaire (IPAQ), which assessed the duration and frequency of PA in leisure time.[19]
17 Weekly PA was summarized using weekly total Metabolic Equivalent Task (MET), calculated
18 by multiplying the MET value of activity by the number of PA hours per week. Based on the
19 lower and upper limits of the World Health Organization (WHO) PA guideline,[19] PA was
20 categorized as low (0 to < 600 MET-mins/wk), medium (600 to < 1200 MET-mins/wk), and
21 high (\geq 1200 MET-mins/wk). Another no moderate-to-vigorous PA (MVPA) category was
22 further defined to reflect the potential health benefits of insufficient PA compared to no
23 MVPA.[4, 21]

1 A novel healthy sleep score, comprising five sleep characteristics, was applied.[12]
2 Morning chronotype, adequate sleep duration (7-8 hr/d), not usually insomnia, no snoring, and
3 no frequent daytime sleepiness, represented healthy sleep characteristics. **Supplementary File,**
4 **Table S2** provides a detailed questionnaire and definition of each item. Participants were
5 scored from 0 to 5, according to their number of the healthy characteristics and were
6 categorized into three groups: "healthy sleep" (≥ 4 sleep score); "intermediate sleep" (2-3 sleep
7 score); and "poor sleep" (≤ 1 sleep score). This score has been used in the UK Biobank to
8 distinguish high-risk middle-aged populations for incident CVD.[12] We generated twelve
9 combined joint categories of PA and sleep accordingly (**Supplementary File, Table S3**).

10 **Covariates**

11 To reduce the effect of potential confounding, demographics and contextual covariates were
12 selected based on previous literature,[3, 12, 15] and included age, sex, socioeconomic status,
13 employment status, body mass index (BMI), cigarette smoking, vegetable and fruit intake,
14 alcohol consumption, sedentary behaviour, mental health issues, and PA or sleep scores when
15 applicable. **Supplementary File, Table S4** describes the definitions in detail.

16 **Statistics and data analysis**

17 All tests were performed using SAS 9.4 or R 3.6.3 software and were two-sided.
18 Descriptive statistics are presented stratified by the sleep score, while chi-square tests, analysis
19 of variance (ANOVA), and Kruskal-Wallis tests were conducted as applicable. Kaplan-Meier
20 estimation, the scaled Schoenfeld residuals, and supremum tests of functional forms were used
21 to examine the proportional hazards assumption, although no noticeable violations were
22 observed. We also performed models excluding 15 potential outliers with extreme values of
23 confounding detected by the $dff\beta$ measure-based influence diagnostics. Since the effect sizes of

1 all the effects were unaffected after exclusion (data not shown), the present analyses reserved
2 them in all models.

3 Both the independent and joint association between PA and sleep scores with mortality
4 were examined using Cox-proportional hazard models, with a high PA level, healthy sleep, or
5 the combination of both as the reference as applicable. Each of the described analyses started
6 with a minimally adjusted Cox model including only age and sex as covariates (Model 1), and
7 a second model additionally adjusted for socioeconomic status, employment status, body mass
8 index (BMI), cigarette smoking, vegetable and fruit intake, alcohol consumption, sedentary
9 behaviour, mental health issues, and PA or sleep scores when applicable (Model 2).[3, 12, 15]
10 In the sensitivity analysis, we further examined the potential synergistic effect between the two
11 main exposures by performing additional analysis of both the additive and multiplicative
12 interactions between the two primary exposures (PA and sleep).[22]

13

14 **RESULTS**

15 For this study, participants were excluded if they had history of total CVD (n = 30,696),
16 total cancer (n = 21,257), or both (n = 2,585) at baseline (**Supplementary File, Figure S1**).
17 Participants were also excluded if they had obstructive sleep apnea history (n = 1,825), class
18 III obesity (body mass index (BMI) ≥ 40 kg/m²) (n = 7,937), missing or unusable data on sleep
19 characteristics (n = 10,633), PA (n = 35,754), or any confounders (n = 10,737). We also
20 excluded those who died from coronavirus disease (COVID-19) (either a primary or
21 contributory cause) (n = 235). To reduce the potential reverse causality,[23] we further
22 excluded those who died in the first two years of follow-up (n = 902). The comparison between
23 included and excluded samples was also provided in **Supplementary File, Table S5**. A total

1 of 380,055 participants (55.9 ± 8.1 years old, 55% female) were included in the present study.
2 Among those, the average follow-up time was 11.1 ± 1.2 years, with 15,503 all-cause deaths
3 (4,095 events from total CVD and 9,064 from total cancer, including 1,932 from coronary heart
4 diseases, 359 from hemorrhagic stroke, 450 from ischemic stroke, and 1,595 from lung cancer).
5 **Table 1** presents the characteristics of participants stratified by sleep scores. The majority (56%)
6 of the participants had a healthy sleep, followed by having intermediate (42%) or poor sleep
7 (3%), demonstrating the same distribution as the work by Fan et al. (2020).[12] Individuals
8 with different sleep categorizations had significant differences in all the covariates. Participants
9 who were younger, female, faced less socioeconomic deprivation, were employed in non-shift
10 work, never smoked, had higher vegetable and fruit intakes, drank less alcohol, sat less, had no
11 mental health issues, and were more PA, tended to have healthier sleep scores.

12 **Independent association of exposures with mortality**

13 **Table 2** (and **Supplementary File, Figure S2**) shows the independent (and mutually
14 adjusted) association of sleep scores and PA with mortality risks. There was a dose-response
15 increase toward poorer sleep in mortality risks for all-cause, total CVD, and ischemic stroke.
16 Relative to healthy sleep, with full adjustment for selected confounders and PA levels (Model
17 2), poor and intermediate sleep was associated with higher mortality risks for all-cause (hazard
18 ratio, HR, [95% confidence intervals]: 1.05 [1.02 to 1.09], for intermediate sleep; 1.23 [1.13 to
19 1.34], for poor sleep) and total CVD (1.09 [1.03 to 1.17], for intermediate sleep; 1.39 [1.19 to
20 1.62], for poor sleep). Poor sleep was further associated with ischemic stroke mortality (1.94
21 [1.29 to 2.94]), while intermediate sleep was further associated with coronary heart diseases
22 (1.16 [1.06 to 1.27]).

1 Compared to participants with a high PA level, those with all the other PA levels had
2 an incrementally higher risk for all-cause mortality after full adjustment (1.05 [1.01 to 1.10],
3 for medium level; 1.08 [1.02 to 1.14], for low level; 1.25 [1.20 to 1.31], for no MVPA,
4 respectively). Those with no MVPA also had higher risks for all the other conditions except
5 for hemorrhagic stroke (for all-cause, 1.25 [1.20 to 1.31]; for total CVD, 1.31 [1.21 to 1.42];
6 for total cancer, 1.16 [1.10 to 1.23]; for coronary heart diseases, 1.35 [1.21 to 1.52]; for
7 ischemic stroke, 1.38 [1.07 to 1.77]; and for lung cancer, 1.34 [1.19 to 1.53]).

8 **Joint association of exposures with mortality**

9 **Figure 1 and 2, and Supplementary File, Table S6 and S7** illustrate the HR for each
10 condition of exposure combinations compared to the referent high PA-healthy sleep group.
11 After full adjustment, participants with no MVPA had higher mortality risks for all outcomes
12 except for hemorrhagic stroke, while those mutually with poor sleep had the highest risks (for
13 all-cause, 1.57 [1.35 to 1.82]; for total CVD, 1.67 [1.27 to 2.19]; for total cancer, 1.45 [1.18 to
14 1.77]; for coronary heart diseases, 1.59 [1.07 to 2.37]; for ischemic stroke, 2.96 [1.43 to 6.11];
15 and for lung cancer, 1.91 [1.30 to 2.81], respectively). Participants with low PA-poor sleep
16 combination had higher mortality risks for all-cause, total CVD, coronary heart disease, and
17 lung cancer. In the sensitivity analysis, we found statistically significant additive interaction in
18 total CVD, and multiplicative interactions in all the outcomes except for stroke
19 **(Supplementary File, Table S8 and S9).**

20

21 **DISCUSSION**

22 To our knowledge, this is the first prospective study investigating the association of a
23 composite score reflecting both sleep duration, quality, and timing with mortality outcomes, as

1 well as the largest investigation examining the joint effect of sleep and PA with all-cause and
2 cause-specific mortality. The PA measurement in the UK Biobank captured frequency,
3 intensity, and duration (time) in the leisure-time domain. Our results suggest that, among
4 middle-aged adults, poorer sleep, after adjustment for PA, remained associated with higher
5 mortality risks for all-cause, total CVD, and ischemic stroke. Besides, there was a potential
6 synergistic effect of sleep and PA with mortality risks for all-cause, total CVD, total cancer,
7 and lung cancer. Compared to no MVPA, levels of PA above the lower threshold recommended
8 by WHO (600 MET-min/wk),[20] appeared to eliminate most of the detrimental associations
9 of poor sleep and mortality.

10 **Comparison with other studies**

11 Sleep duration, quality, and timing are interrelated dimensions of healthy sleep
12 characteristics ^{7 9 11 16}. [7-,12, 16] Previous meta-analyses and systematic reviews based on
13 single sleep characteristics provided mixed results in relation to poor sleep and all-cause
14 mortality risk.[3, 24] Yin et al. (2017) suggested that inadequate sleep duration (shorter or
15 longer than 7 hr/d) was associated with all-cause mortality risk (pooled relative risk, RR: 1.06
16 [1.04 to 1.07] per hour reduction for short sleepers, and 1.13 [1.11 to 1.15] per hour increment
17 for long sleepers).[24] Alternatively, Kwok et al. (2018) proposed that only long sleepers (≥ 8
18 hr/d) but not short sleepers (< 7 hr/d) had a higher risk for all-cause mortality (*e.g.*, pooled RR:
19 1.14 [1.05 to 1.25] for 9-hr sleeper) compared to healthy sleepers (7 hr/d), while sleep quality
20 showed no significant association with mortality risk (pooled RR: 1.03 [0.93 to 1.14]).[3] With
21 a multicomponent sleep characteristic score, the present study shows that people with poor or
22 intermediate sleep, with adjustment for PA, still had a 23% or 5% higher risk for all-cause
23 mortality and a 39% or 9% higher risk for total CVD mortality (**Table 2**). As poor sleep was

1 further associated with 94% higher mortality risk for ischemic stroke but not for hemorrhagic
2 stroke, our finding extends the work by Fan et al. (2020), which did not distinguish the subtypes
3 of stroke.[12] The adverse health effects of poor sleep remain even within participants with a
4 high PA level (**Figure 1**). Besides, our results indicate that not participating in any MVPA
5 could result in 16% to 38% higher mortality risks for all-cause, total CVD, total cancer, and
6 the sub-types (**Table 2**). In agreement with a previous prospective study and public health
7 guidelines,[4, 21] our finding suggested that a below-guideline MVPA level could be
8 beneficial.

9 Recent studies have suggested the synergistic health effect of physical inactivity and
10 poor sleep.[6, 8, 9, 15, 17] Cassidy et al. (2016) reported that people with CVD or type 2
11 diabetes tended to simultaneously report physical inactivity and poor sleep, based on a cross-
12 sectional analysis of the same cohort as the present study.[15] In a 15-year follow-up of a
13 Taiwanese cohort (n = 341,248), short sleep duration (defined as < 6 hr) increased all-cause
14 and CVD mortality risk only among physically inactive participants (< 450 MET-mins/wk).[9]
15 Wennman et al. (2017) found that the combination of insufficient sleep duration (≤ 6 hr) and
16 low leisure-time PA (< 450 MET-mins/wk) was associated with higher mortality risks for all-
17 cause (HR: 1.49 [1.05 to 2.11]) and CVD (HR: 1.98 [1.25 to 3.12]), compared to the referent
18 combination of adequate sleep (6.5 to 8.5 hr) and high leisure-time PA (≥ 450 MET-mins/wk),
19 based on a male cohort comprising 60% of former elite athletes with a 26-year follow-up (n =
20 1,638).[8] However, the same study also suggested no synergistic effect of sleep quality and
21 PA on mortality. Likewise, Xiao et al. (2014) found no significant relative excess risk of
22 insufficient sleep (< 7 hr) and low MVPA (<1 hr/wk) for all-cause mortality, based on a large
23 US cohort with a 14-year follow-up (n = 239,896).[25] The present study extended the findings
24 from the above studies by capturing different perspectives of sleep and death causes. Compared

1 to those with the high PA-healthy sleep combination, participants with the no MVPA-poor
2 sleep combination had higher mortality risks (*e.g.*, 57% for all-cause; 67% for total CVD; 45%
3 for total cancer). Moreover, the detrimental effect of poorer sleep with mortality risks could
4 potentially be exaggerated among participants with lower PA levels (**Figure 1 and 2;**
5 **Supplementary File, Table S6 to S9**).

6 Several explanations of the deleterious associations of poor sleep with health have been
7 put forward,[6, 8, 9, 15, 16] although none of them has been widely established. In particular,
8 the mechanisms surrounding the deleterious associations of prolonged sleep duration with
9 mortality outcomes are controversial.[5] One possibility is that these associations are not causal
10 and are due to residual confounding, *e.g.*, fatigue, sleep fragmentation, substance use, or
11 undiagnosed mental disorders.[26] Another hypothetical mechanism is that prolonged sleep
12 duration might compromise the time available for health-enhancing behaviours such as PA.[15]
13 In the present sample, however, the average total PA volume was not very different in
14 prolonged sleepers compared to those in the adequate sleep range (adequate vs prolonged sleep:
15 43.0 vs 42.2 MET-hrs/wk), despite the large difference in average sleep times between the two
16 groups (adequate vs prolonged sleepers: 7.4 vs 9.3 hr). We used a composite sleep score that
17 relied on self-reported data and did not allow us to incorporate time sequence or patterns of the
18 sleep-related behaviours in this study.[27] Future studies featuring wearable devices that
19 measure sleep and physical activity will shed further light on the time-dependent aspects of
20 these behaviours and their impact on health. Although an emerging body of evidence suggests
21 that excessive PA might worsen sleep and potentially increase cardiovascular health risks,[28-
22 30] the proportion of high PA (≥ 1200 MET-mins/wk) in the present study was higher in those
23 with the healthiest sleep score (**Table 1**). Our results also showed that, among participants with
24 PA higher than the upper limit of the WHO guideline (1200 MET-mins /week),[20] PA may

1 attenuate the deleterious effects of poor sleep. However, in populations at the very top end of
2 the activity spectrum, *e.g.*, athletes, excessive PA might compromise sleep.[28] The
3 combination of very high PA and poor sleep could, in the long term, even increase the risks for
4 injuries and infections.[30] As the point past which PA might be harmful is unknown,[29] we
5 acknowledge that our results may not be applicable to extremely active populations.

6 **Study strengths and limitations**

7 The strengths of the present study include the prospective study design with large
8 sample size and long follow up and extensive measurement of covariates, which allowed us to
9 perform several sensitivity analyses to strengthen our interpretation. The novel sleep score
10 integrating various aspects of the sleep characteristics provides a convenient yet thorough way
11 for investigating the joint effects of sleep with other time-dependent behaviours like PA. Our
12 study had some potential limitations. Firstly, the observational nature of our data precludes a
13 definitive causal interpretation of our findings, although we took several measures to minimize
14 risks for reverse causalities and confounding, such as exclusion criteria and a wide range of
15 covariates included. Second, our exposures were measured by self-reported, random
16 measurement error may have biased results toward the null, and thereby underestimating the
17 true magnitude of the associations. Third, all the exposures and confounders were collected at
18 recruitment and were assumed to be relatively constant over the course of the follow-up period.
19 In a UK Biobank sub-sample with at least one repeated measurement of both exposures over a
20 seven-year follow-up ($n = 35,466$), 59% and 68% of the participants maintained their PA and
21 sleep, respectively. Future studies investigating changes of the two behaviours over time will
22 further elucidate the synergistic effects of the two behaviours. Fourth, the PA measurement of
23 the UK Biobank did not include other domains like occupation, transportation, and

1 household.[15, 31] The relatively low sample size in some joint groups (*e.g.*, the low PA-poor
2 sleep combination) may have compromised our statistical power to detect the differences in
3 risk. Lastly, the UK Biobank had a response rate of 5.5%, and it is not representative of the UK
4 population.[32] However, a recent UK Biobank study showed that poor representativeness does
5 not materially affect the associations of physical activity and other lifestyle behaviours with
6 mortality.[33]

7

8 **CONCLUSIONS**

9 Poor sleep was associated with a higher risk for all-cause and cause-specific mortality,
10 and these risks were markedly exacerbated among participants with insufficient PA. Meeting
11 the lower threshold of the current PA guidelines (600 MET-mins/wk) eliminated most of the
12 deleterious associations of poor sleep with mortality. Our results support the value of
13 interventions to concurrently target PA and sleep to improve health. Future prospective studies
14 with device-based sleep and PA assessments and trials concurrently targeting both behaviours
15 are warranted.

1 **Acknowledgement:** The authors gratefully thank all the participants and professionals
2 contributing to the UK Biobank.

3 **Funding:** ES is funded by a National Health and Medical Research Council
4 (NHMRC)Leadership 2 Fellowship (APP1194510). MJD is supported by a Career
5 Development Fellowship (APP1141606). The funding sources had no involvement in study
6 design; in the collection, analysis and interpretation of data; in the writing of the report; or in
7 the decision to submit the article for publication.

8 **Competing interests:** None declared.

9 **References**

- 10 1 Ma QQ, Yao Q, Lin L, et al. Sleep duration and total cancer mortality: a meta-analysis of
11 prospective studies. *Sleep Med* 2016;27-28:39-44.
- 12 2 Dohrn M, Sjöström M, Kwak L, et al. Accelerometer-measured sedentary time and physical
13 activity—a 15 year follow-up of mortality in a Swedish population-based cohort. *J Sci*
14 *Med Sport* 2018;21(7):702-07.
- 15 3 Kwok CS, Kontopantelis E, Kuligowski G, et al. Self-Reported Sleep Duration and Quality
16 and Cardiovascular Disease and Mortality: A Dose-Response Meta-Analysis. *J Am*
17 *Heart Assoc* 2018;7(15):e008552.
- 18 4 US Department of Health and Human Services. 2018 Physical Activity Guidelines
19 Advisory Committee Scientific Report U.S. Department of Health and Human
20 Services. Available: [https://health.gov/our-work/physical-activity/current-](https://health.gov/our-work/physical-activity/current-guidelines/scientific-report/)
21 [guidelines/scientific-report/](https://health.gov/our-work/physical-activity/current-guidelines/scientific-report/) [Accessed Aug 2020].
- 22 5 Jike M, Itani O, Watanabe N, et al. Long sleep duration and health outcomes: A systematic
23 review, meta-analysis and meta-regression. *Sleep Med Rev* 2018;39:25-36.
- 24 6 Bellavia A, Akerstedt T, Bottai M, et al. Sleep duration and survival percentiles across
25 categories of physical activity. *Am J Epidemiol* 2014;179(4):484-91.
- 26 7 Chien KL, Chen PC, Hsu HC, et al. Habitual sleep duration and insomnia and the risk of
27 cardiovascular events and all-cause death: report from a community-based cohort.
28 *Sleep* 2010;33(2):177-84.

- 1 8 Wennman H, Kronholm E, Heinonen OJ, et al. Leisure Time Physical Activity and Sleep
2 Predict Mortality in Men Irrespective of Background in Competitive Sports. *Prog*
3 *Prev Med (N Y)* 2017;2(6):e0009.
- 4 9 Li-Jung C, Mark H, Yun-Ju L, et al. Can physical activity eliminate the mortality risk
5 associated with poor sleep? A 15-year follow-up of 341,248 MJ Cohort participants. *J*
6 *Sport Health Sci* 2021. Available: <https://doi.org/10.1016/j.jshs.2021.03.001>
7 [Accessed Apr 2021].
- 8 10 Rod NH, Kumari M, Lange T, et al. The Joint Effect of Sleep Duration and Disturbed
9 Sleep on Cause-Specific Mortality: Results from the Whitehall II Cohort Study. *PLOS*
10 *ONE* 2014;9(4):e91965.
- 11 11 Bertisch SM, Pollock BD, Mittleman MA, et al. Insomnia with objective short sleep
12 duration and risk of incident cardiovascular disease and all-cause mortality: Sleep
13 Heart Health Study. *Sleep* 2018;41(6)
- 14 12 Fan M, Sun D, Zhou T, et al. Sleep patterns, genetic susceptibility, and incident
15 cardiovascular disease: a prospective study of 385 292 UK biobank participants. *Eur*
16 *Heart J* 2020;41(11):1182-89.
- 17 13 Kline CE. The bidirectional relationship between exercise and sleep: implications for
18 exercise adherence and sleep improvement. *Am J Lifestyle Med* 2014;8(6):375-79.
- 19 14 Huang BH, Hamer M, Duncan MJ, et al. The bidirectional association between sleep and
20 physical activity: A 6.9 years longitudinal analysis of 38,601 UK Biobank
21 participants. *Prev Med* 2021;143:106315.
- 22 15 Cassidy S, Chau JY, Catt M, et al. Cross-sectional study of diet, physical activity,
23 television viewing and sleep duration in 233,110 adults from the UK Biobank; the
24 behavioural phenotype of cardiovascular disease and type 2 diabetes. *BMJ Open*
25 2016;6(3):e010038.
- 26 16 Stone CR, Haig TR, Fiest KM, et al. The association between sleep duration and cancer-
27 specific mortality: A systematic review and meta-analysis. *Cancer Causes Control*
28 2019;30(5):501-25.
- 29 17 Rayward AT, Duncan MJ, Brown WJ, et al. A cross-sectional cluster analysis of the
30 combined association of physical activity and sleep with sociodemographic and health
31 characteristics in mid-aged and older adults. *Maturitas* 2017;102:56-61.

- 1 18 Sudlow C, Gallacher J, Allen N, et al. UK biobank: an open access resource for
2 identifying the causes of a wide range of complex diseases of middle and old age.
3 *PLoS Med* 2015;12(3):e1001779.
- 4 19 Craig CL, Marshall AL, Sjostrom M, et al. International physical activity questionnaire:
5 12-country reliability and validity. *Med Sci Sports Exerc* 2003;35(8):1381-95.
- 6 20 World Health Organization. Physical Activity and Adults: World Health Organization.
7 Available: https://www.who.int/dietphysicalactivity/factsheet_adults/en/ [Accessed
8 Aug 2020].
- 9 21 O'Donovan G, Lee IM, Hamer M, et al. Association of "Weekend Warrior" and Other
10 Leisure Time Physical Activity Patterns With Risks for All-Cause, Cardiovascular
11 Disease, and Cancer Mortality. *JAMA Intern Med* 2017;177(3):335-42.
- 12 22 Knol MJ, VanderWeele TJ. Recommendations for presenting analyses of effect
13 modification and interaction. *Int J Epidemiol* 2012;41(2):514-20.
- 14 23 Strain T, Wijndaele K, Sharp SJ, et al. Impact of follow-up time and analytical approaches
15 to account for reverse causality on the association between physical activity and
16 health outcomes in UK Biobank. *Int J Epidemiol* 2020;49(1):162-72.
- 17 24 Yin J, Jin X, Shan Z, et al. Relationship of Sleep Duration With All-Cause Mortality and
18 Cardiovascular Events: A Systematic Review and Dose-Response Meta-Analysis of
19 Prospective Cohort Studies. *J Am Heart Assoc* 2017;6(9):e005847.
- 20 25 Xiao Q, Keadle SK, Hollenbeck AR, et al. Sleep duration and total and cause-specific
21 mortality in a large US cohort: interrelationships with physical activity, sedentary
22 behavior, and body mass index. *Am J Epidemiol* 2014;180(10):997-1006.
- 23 26 Ding D, Rogers K, Macniven R, et al. Revisiting lifestyle risk index assessment in a large
24 Australian sample: should sedentary behavior and sleep be included as additional risk
25 factors? *Prev Med* 2014;60:102-6.
- 26 27 Huang B-H, Inan-Eroglu E, Hamer M, et al. Joint associations of device-measured
27 physical activity and sleep duration with cardiometabolic health in the 1970 British
28 Cohort Study. *J Sci Med Sport* 2020;23(12):1191-6.
- 29 28 Falck RS, Stamatakis E, Liu-Ambrose T. The athlete's sleep paradox prompts us to
30 reconsider the dose-response relationship of physical activity and sleep. *Br J Sports*
31 *Med* 2021. Available: <https://doi.org/10.1136/bjsports-2020-103835> [Accessed Mar
32 2021].

- 1 29 Eijsvogels TMH, Thompson PD, Franklin BA. The "Extreme Exercise Hypothesis":
2 Recent Findings and Cardiovascular Health Implications. *Curr Treat Options*
3 *Cardiovasc Med* 2018;20(10):84.
- 4 30 Walsh NP, Halson SL, Sargent C, et al. Sleep and the athlete: narrative review and 2021
5 expert consensus recommendations. *Br J Sports Med* 2020. Available:
6 <https://doi.org/10.1136/bjsports-2020-102025> [Accessed Mar 2021].
- 7 31 DiPietro L, Al-Ansari SS, Biddle SJH, et al. Advancing the global physical activity
8 agenda: recommendations for future research by the 2020 WHO physical activity and
9 sedentary behavior guidelines development group. *Int J Behav Nutr Phys Act*
10 2020;17(1):143-32 Fry A, Littlejohns TJ, Sudlow C, et al. Comparison of
11 sociodemographic and health-related characteristics of UK Biobank participants with
12 those of the general population. *Am J Epidemiol* 2017;186(9):1026-34.
- 13 33 Stamatakis E, Owen KB, Shepherd L, Drayton B, Hamer M, Bauman AE. Is Cohort
14 Representativeness Passé? Poststratified Associations of Lifestyle Risk Factors with
15 Mortality in the UK Biobank. *Epidemiology* 2021;32(2):179.

1 **Figure legends**

2 **Figure 1.** The Joint Association of Physical Activity and Sleep Scores with Mortality for All-
3 Cause, Total Cardiovascular Disease, and Total Cancer (n = 380,055)

4
5 a, all-cause; b, total cardiovascular disease; c, total cancer. Physical activity levels were
6 categorized based on public health guidelines: no MVPA (those reported 0 MET-mins/wk from
7 MVPA, regardless of total MET-mins/wk); low (< 600 MET-mins/wk); medium (600 to <
8 1200 MET-mins/wk); and high (\geq 1200 MET-mins/wk). Sleep scores were categorized into:
9 poor, 0~1; intermediate, 2~3; healthy, 4~5.

10 Models were adjusted for age, sex, socioeconomic status, employment status, BMI, cigarette
11 smoking, vegetable and fruit intake, alcohol consumption, sedentary behaviour, and mental
12 health.

13

14 **Figure 2.** The Joint Association of Physical Activity and Sleep Scores with Mortality for
15 Sub-Type Conditions (n = 380,055)

16

17 a, coronary heart disease; b, hemorrhagic stroke; c, ischemic stroke; d, lung cancer. Physical
18 activity levels were categorized based on public health guidelines: no MVPA (those reported
19 0 MET-mins/wk from MVPA, regardless of total MET-mins/wk); low (< 600 MET-mins/wk);
20 medium (600 to < 1200 MET-mins/wk); and high (\geq 1200 MET-mins/wk). Sleep scores were
21 categorized into: poor, 0~1; intermediate, 2~3; healthy, 4~5.

22 Models were adjusted for age, sex, socioeconomic status, employment status, BMI, cigarette
23 smoking, vegetable and fruit intake, alcohol consumption, sedentary behaviour, and mental
24 health.

Text tables

Table 1. The Descriptive Statistics by Sleep Scores (n = 380,055)

Characteristics	All	Sleep Scores ^a		
		Poor	Intermediate	Healthy
n (%)	380,055 (100)	10,166 (3)	158,103 (42)	211,786 (56)
Follow-up (yr) (mean (SD))	11.1 (1.2)	11.0 (1.4)	11.1 (1.3)	11.1 (1.2)
Age (mean (SD))	55.9 (8.1)	56.1 (7.7)	56.1 (7.9)	55.6 (8.2)
Male (n (%))	171,315 (45)	890 (48)	76,819 (49)	89,606 (42)
Body mass index (kg/m ²) (mean (SD))	26.9 (4.1)	28.7 (4.5)	27.5 (4.2)	26.4 (4.0)
Socioeconomic Status (mean (SD)) ^b	-1.5 (3.0)	-0.8 (3.3)	-1.4 (3.1)	-1.6 (2.9)
Vegetable and Fruit Intake (serves/d) (mean (SD))	4.4 (2.9)	4.2 (3.1)	4.3 (3.0)	4.5 (2.9)
Sedentary Behaviour (hr/d) (mean (SD))	4.7 (2.4)	5.5 (2.8)	5.0 (2.5)	4.5 (2.3)
Mental Health Issue (n (%)) ^c	127,943 (34)	5,108 (50)	59,207 (37)	63,628 (30)

Cigarette Smoking (n (%))				
Never	213,618 (56)	4,758 (47)	82,786 (52)	126,074 (60)
Previous Smoker	128,597 (34)	3773 (37)	56,135 (36)	68,689 (32)
Current Smoker	37,840 (10)	1,635 (16)	19,182 (12)	17,023 (8)
Employment Status (n (%))				
Retired/not in the workforce	144,090 (38)	4,626 (46)	61,893 (39)	77,571 (37)
Employed not in shift work	197,803 (52)	4,225 (42)	78,439 (50)	115,139 (54)
Employed in night shift work	19,365 (5)	752 (7)	9,584 (6)	9,029 (4)
Employed in day shift work	18,797 (5)	563 (6)	8,187 (5)	10,047 (5)
Physical Active (n (%)) ^d				
No MVPA	59,541 (16)	2,510 (25)	28,289 (18)	28,742 (14)
Low	9,298 (10)	1,291 (13)	17,757 (11)	20,250 (10)
Medium	57,771 (15)	1,384 (14)	24,110 (15)	32,277 (15)
High	223,445 (59)	4,981 (49)	87,947 (56)	130,517 (62)
Alcohol Consumption (n (%)) ^e				

Never	13,843 (4)	389 (4)	5,567 (4)	7,887 (4)
Previous Drinker	11,512 (3)	525 (5)	4,980 (3)	6,007 (3)
Occasional Drinker	81,191 (21)	2,555 (25)	33,930 (22)	44,706 (21)
Within Guideline	130,901 (34)	2,712 (27)	50,203 (32)	77,986 (37)
Above Guideline	84,936 (22)	1,967 (19)	35,409 (22)	47,560 (23)
Above Double Guideline	57,672 (15)	2,018 (20)	28,014 (18)	27,640 (13)

a. Sleep scores were categorized into: poor, 0~1; intermediate, 2~3; healthy, 4~5. There were significant differences ($p < 0.05$) across sleep score groups in all the characteristics shown in the table.

b. Townsend area deprivation index was used.

c. Had ever seen a doctor or psychiatrist for nerves, anxiety or depression.

d. Categorization based on public health guidelines: no MVPA (those reported 0 MET-mins/wk from MVPA, regardless of total MET-mins/wk); low (< 600 MET-mins/wk); medium (600 to < 1200 MET-mins/wk); and high (≥ 1200 MET-mins/wk).

e. The UK guideline (14 UK units/wk) was used.

MET = metabolic equivalent task; MVPA = moderate-to-vigorous physical activity

Table 2. The Independent (and Mutually Adjusted) Associations of Physical Activity and Sleep Scores with Mortality (n = 380,055)

Exposure ^a	Hazard Ratio for Mortality Risks (95% confidence interval)							
	All-cause		Total cardiovascular disease		Total cancer			
	Model 1 ^b	Model 2 ^c	Model 1 ^b	Model 2 ^c	Model 1 ^b	Model 2 ^c	Model 1 ^b	Model 2 ^c
Sleep Pattern								
Poor	1.56	1.23	1.92	1.39	1.35	1.13		
(n = 10,166)	(1.44, 1.70)	(1.13, 1.34)	(1.65, 2.23)	(1.19, 1.62)	(1.20, 1.52)	(1.00, 1.27)		
Intermediate	1.16	1.05	1.24	1.09	1.12	1.03		
(n = 158,103)	(1.13, 1.20)	(1.02, 1.09)	(1.17, 1.33)	(1.03, 1.17)	(1.08, 1.17)	(0.99, 1.08)		
Healthy (n = 211,786)				1.00				
	Coronary heart diseases		Hemorrhagic stroke		Ischemic stroke		Lung cancer	
Poor	1.83	1.26	1.23	1.05	2.46	1.94	1.93	1.25
(n = 10,166)	(1.46, 2.29)	(1.00, 1.59)	(0.67, 2.26)	(0.57, 1.95)	(1.64, 3.70)	(1.29, 2.94)	(1.51, 2.46)	(0.98, 1.60)

Intermediate	1.34	1.16	1.12	1.04	1.18	1.06	1.29	1.05
(n = 158,103)	(1.22, 1.46)	(1.06, 1.27)	(0.90, 1.38)	(0.84, 1.29)	(0.97, 1.43)	(0.87, 1.29)	(1.17, 1.43)	(0.95, 1.16)
Healthy (n = 211,786)					1.00			
Physical Activity	All-cause		Total cardiovascular disease		Total cancer			
No MVPA	1.39	1.25	1.52	1.31	1.28	1.16		
(n = 59,541)	(1.33, 1.44)	(1.20, 1.31)	(1.41, 1.65)	(1.21, 1.42)	(1.21, 1.35)	(1.10, 1.23)		
Low	1.1	1.08	1.14	1.09	1.07	1.03		
(n = 9,298)	(1.04, 1.16)	(1.02, 1.14)	(1.02, 1.26)	(0.98, 1.21)	(0.99, 1.14)	(0.96, 1.11)		
Medium	1.05	1.05	1.07	1.06	1.03	1.02		
(n = 57,771)	(1.01, 1.10)	(1.01, 1.10)	(0.98, 1.17)	(0.97, 1.16)	(0.97, 1.09)	(0.96, 1.08)		
High (n = 223,445)					1.00			
	Coronary heart diseases		Hemorrhagic stroke		Ischemic stroke		Lung cancer	
No MVPA	1.61	1.35	1.32	1.26	1.51	1.38	1.68	1.35

(n = 59,541)	(1.44, 1.81)	(1.21, 1.52)	(1.01, 1.73)	(0.96, 1.66)	(1.18, 1.92)	(1.07, 1.77)	(1.48, 1.89)	(1.19, 1.53)
Low	1.13	1.08	0.95	0.97	1.16	1.16	1.13	1.09
(n = 9,298)	(0.97, 1.32)	(0.93, 1.26)	(0.65, 1.39)	(0.67, 1.43)	(0.83, 1.61)	(0.83, 1.61)	(0.96, 1.35)	(0.92, 1.30)
Medium	0.99	0.98	1.02	1.04	1.46	1.47	1.03	1.04
(n = 57,771)	(0.86, 1.13)	(0.86, 1.13)	(0.75, 1.38)	(0.77, 1.41)	(1.14, 1.88)	(1.15, 1.89)	(0.89, 1.20)	(0.89, 1.21)
High (n = 223,445)					1.00			

a. Physical activity levels were categorized based on public health guidelines: no MVPA (those reported 0 MET-mins/wk from MVPA, regardless of total MET-mins/wk); low (< 600 MET-mins/wk); medium (600 to < 1200 MET-mins/wk); and high (\geq 1200 MET-mins/wk).


Sleep scores were categorized into: poor, 0~1; intermediate, 2~3; healthy, 4~5.

b. Adjusted for age and sex.


c. Further adjusted for socioeconomic status, employment status, BMI, cigarette smoking, vegetable and fruit intake, alcohol consumption, sedentary behaviour, mental health, and mutually adjusted for sleep scores or physical activity levels as appropriate.

BMI = body mass index; MET = metabolic equivalent task; MVPA = moderate-to-vigorous physical activity.


1a




1b




1c




2a




2b




2c




2d





Supplementary Figure S1. Participants recruitment flowchart


(a) All-cause mortality



Supplementary Figure S2. The Independent (and Mutually Adjusted) Associations of All Selected Variables with Mortality (n = 380,055) of (a) All-Cause, (b) Total Cardiovascular Disease, and (c) Total Cancer.

* 'Townsend area deprivation index' served as an indicator of socioeconomic status, with higher scores indicating greater socioeconomic deprivation.


(b) Total cardiovascular disease mortality



Supplementary Figure S2 (cont.). The Independent (and Mutually Adjusted) Associations of All Selected Variables with Mortality (n = 380,055) of (a) All-Cause, (b) Total Cardiovascular Disease, and (c) Total Cancer.

* 'Townsend area deprivation index' served as an indicator of socioeconomic status, with higher scores indicating greater socioeconomic deprivation.

(c) Total cancer mortality



Supplementary Figure S2 (cont.). The Independent (and Mutually Adjusted) Associations of All Selected Variables with Mortality (n = 380,055) of (a) All-Cause, (b) Total Cardiovascular Disease, and (c) Total Cancer.

* 'Townsend area deprivation index' served as an indicator of socioeconomic status, with higher scores indicating greater socioeconomic deprivation..

Supplementary Table S1. The Definition of Diseases

Variable	Definition
Total cardiovascular disease (CVD)	The total CVD definition is aligned with Zao et al. (2020) (https://doi.org/10.1136/bmj.m2031). In brief, the ICD-10 code described in the Chapter IX Diseases of the circulatory system (I0, I11, I13, I20-I51, I60-I69) or the ICD-9 code within (390.0-398.9/ 402.0- 402.9/ 404.0-404.9/ 411.1-436.9) were included.
Sub-type CVD	
Coronary heart diseases	The ICD-10 code described in Chapter IX Diseases of the circulatory system – Ischemic heart disease (I20-I25) was included.
Hemorrhagic stroke	The definition of hemorrhagic stroke (both intracerebral and subarachnoid) is aligned with UK Biobank. The ICD-10 code beginning with ('I60' 'I61') was included.
Ischemic stroke	The definition of ischemic stroke is aligned with UK Biobank. The ICD-10 code beginning with ('I63' 'I64') was included.
Total cancers	The definition of total cancer excludes in situ, benign, uncertain, or non-well-defined cancers. In brief, the ICD-10 code beginning with ('C0' 'C1' 'C2' 'C3' 'C4' 'C5' 'C6' 'C70' 'C71' 'C72' 'C73' 'C74' 'C75' 'C7A' 'C8' 'C9') or the ICD-9 code within (140.0-194.9/ 199.0-209.3) were included.
Sub-type malignant neoplasm	
Lung cancer	The ICD-10 code beginning with ('C33' 'C34') was included.
Others	
Obstructive sleep apnea	The ICD-10 code 'G47.3' and ICD-9 code '327.2' and '780.5' were applied.
Coronavirus disease (COVID-19)	The ICD-10 code 'U07.1' and 'U07.2' were applied.

Primary/secondary inpatient diseases diagnosis is based on both ICD-9 and ICD-10 code. All death cause was based on both primary and secondary cause of death with ICD-10 code only.

Supplementary Table S2. The Scoring System of Sleep

Characteristics	UK BioBank Code	UK BioBank Questionnaire	Healthy Answer (%)	Unhealthy Answer (%)
Chronotype	1180	Do you consider yourself to be?	Definitely a "morning" person; More a "morning" than "evening" person. (57%)	More an "evening" than a "morning person"; Definitely an "evening" person. (43%)
Sleep Duration	1160	About how many hours sleep do you get in every 24 hours? (please include naps)	7-8 hr/d. (69%)	<7 or >=9 hr/d. (31%)
Insomnia	1200	Do you have trouble falling asleep at night or do you wake up in the middle of the night?	Never/rarely; Sometimes (73%)	Usually (27%)
Snoring	1210	Does your partner or a close relative or friend complain about your snoring?	No (60%)	Yes (40%)
Daytime Sleepiness	1220	How likely are you to doze off or fall asleep during the daytime when you don't mean to? (e.g. when working, reading or driving)	Never/rarely; Sometimes (98%)	Often; All the Time (2%)

Supplementary Table S3. Number of Mortality Events by Joint Category of Physical Activity and Sleep Scores (n = 380,055)

Joint Category ^a	N With/Without Events						
	All-cause	Total cardiovascular disease	Total cancer	Coronary heart diseases	Hemorrhagic stroke	Ischemic stroke	Lung cancer
No MVPA, Poor (n = 2,510)	187/2,323	56/2,454	99/2,411	26/2,484	4/2,506	8/2,502	28/2,482
No MVPA, Intermediate (n = 28,289)	1,570/26,719	469/27,820	851/27,438	237/28,052	36/28,253	43/28,246	201/28,088
No MVPA, Healthy (n = 28,742)	1,293/27,449	325/28,417	738/28,004	155/28,587	31/28,711	37/28,705	138/28,604
Low, Poor (n = 1,291)	79/1,212	28/1,263	43/1,248	15/1,276	1/1,290	1/1,290	11/1,280
Low, Intermediate (n = 17,757)	765/16,992	197/17,560	442/17,315	95/17,662	13/17,744	17/17,740	72/17,685
Low, Healthy (n = 20,250)	689/19,561	182/20,068	406/19,844	83/20,167	17/20,233	24/20,226	72/2,0178
Medium, Poor (n = 1,384)	67/1,317	21/1,363	35/1,349	7/1,377	1/1,383	6/1,378	4/1,380
Medium, Intermediate (n = 24,110)	1,050/23,060	279/23,831	605/23,505	125/23,985	24/24,086	34/24,076	103/24,007
Medium, Healthy (n = 32,277)	1,128/31,149	283/31,994	675/31,602	121/32,156	27/32,250	43/32,234	110/32,167
High, Poor (n = 4,981)	262/4,719	80/4,901	130/4,851	33/4,948	5/4,976	11/4,970	28/4,953
High, Intermediate (n = 87,947)	3,704/84,243	996/86,951	2,176/85,771	506/87,441	86/87,861	109/87,838	384/87,563
High, Healthy (n = 130,517)	4,709/125,808	1,179/129,338	2,864/127,653	529/129,988	114/130,403	117/130,400	444/130,073
Overall (n = 380,055)	15,503/364,552	4,095/375,960	9,064/370,991	1,932/378,123	359/379,696	450/379,605	1,595/378,460

a. Physical activity levels were categorized based on public health guidelines: No MVPA (those reported 0 MET-mins/wk from MVPA, regardless of total MET-mins/wk); low (< 600 MET-mins/wk); medium (600 to < 1200 MET-mins/wk); and high (\geq 1200 MET-mins/wk). Sleep scores were categorized into: poor, 0~1; intermediate, 2~3; healthy, 4~5.

Supplementary Table S4. The Resource and Definition of the Selected Covariates (except for Age, Sex, and BMI)

Covariates	UK BioBank Code	Description	Type	Category
Socioeconomic status	189	The existing variable 'Townsend area deprivation index' (189) served as an indicator of socioeconomic status, with higher scores indicating greater socioeconomic deprivation.	Continuous	-
Vegetable and fruit intake	1299, 1309	The sum of the daily servings of 'salad /raw vegetable intake' (1299) and 'fresh fruit intake' (1309) served as a proxy of a healthy diet.	Continuous	-
Sedentary behaviour	1070, 1080, 1090	The total daily hours of television viewing (1070), computer use (1080), and driving (1090) as a marker of sedentary behaviour.	Continuous	-
Mental health issue	2090, 2100	Participants having ever seen a doctor (GP) (2090) or psychiatrist (2100) for nerves, anxiety, or depression were classified as having mental health issues.	Categorical	Yes; No
Cigarette smoking	20116	The existing variable 'smoking status' (20116) was applied.	Categorical	never; previous smoker; current smoker
Employment	6142 3426 826	We derived four groups based on 'Current employment status' (6142), 'Job involves night shift work' (3426), and 'Job involves shift work' (826).	Categorical	retired/not in the workforce; employed not in shift work; employed in night shift work; employed in day shift work
Alcohol consumption	1558, 1568, 1578, 1588, 1598, 1608, 5364. 20117	The level of overall alcohol consumption as the number of UK units of alcohol (10 mL/unit) consumed per week was calculated; participants were categorized based on the consumption according to the UK guideline (14 UK units/wk).	Categorical	never; previous drinker; occasional drinker; within guidelines (14 units per week); above guidelines (14-28 units/wk); double guidelines (>28 units/wk).

Supplementary Table S5. Comparison of the Characteristics between Included and Excluded Samples.

Characteristics	Included Sample (n = 380,055)	Excluded Sample (n = 122,561)	Missing Count in Excluded Sample
Age (mean (SD))	55.9 (8.1)	58.60 (7.7)	-
Male (n (%))	171,315 (45)	57848 (47)	-
BMI (kg/m ²) (mean (SD))	26.9 (4.1)	29.13 (6.17)	3,194
Socioeconomic Status (mean (SD)) ^a	-1.5 (3.0)	-0.65 (3.4)	627
Vegetable and Fruit Intake (serves/d) (mean (SD))	4.4 (2.9)	4.43 (3.2)	8,801
Sedentary Behaviour (hr/d) (mean (SD))	4.7 (2.4)	5.10 (2.6)	7,763
Mental Health Issue (n (%)) ^b	127,943 (34)	45257 (38)	2,179
Cigarette Smoking (n (%))			2,952
Never	213,618 (56)	59,970 (50)	
Previous Smoker	128,597 (34)	44,494 (37)	
Current Smoker	37,840 (10)	15,145 (13)	
Employment Status (n (%))			5,759
Retired/not in the workforce	144,090 (38)	65,557 (56)	
Employed not in shift work	197,803 (52)	39,725 (34)	
Employed in night shift work	19,365 (5)	6,006 (5)	
Employed in day shift work	18,797 (5)	5,514 (5)	
Alcohol Consumption (n (%)) ^c			1,658
Never	13,843 (4)	8,551 (7)	
Previous Drinker	11,512 (3)	6,599 (5)	
Occasional Drinker	81,191 (21)	32,704 (27)	
Within Guideline	130,901 (34)	36,254 (30)	
Above Guideline	84,936 (22)	21,627 (18)	
Above Double Guideline	57,672 (15)	15,168 (13)	
Sleep Score (n (%)) ^d			12,458
Poor	10,166 (3)	5,401 (5)	
Intermediate	158,103 (42)	53,394 (48)	
Healthy	211,786 (56)	51,308 (47)	
Physical Active (n (%)) ^e			47,153
No MVPA	59,541 (16)	9871 (13)	
Low	9,298 (10)	14344 (19)	
Medium	57,771 (15)	12290 (16)	
High	223,445 (59)	38903 (52)	

a. Townsend area deprivation index was used.

b. Had ever seen a doctor or psychiatrist for nerves, anxiety or depression.

c. The UK guideline (14 UK units/wk) was used.

d. Sleep scores were categorized into: poor, 0~1; intermediate, 2~3; healthy, 4~5.

e. Categorization based on public health guidelines: no MVPA (those reported 0 MET-mins/wk from MVPA, regardless of total MET-mins/wk); low (< 600 MET-mins/wk); medium (600 to < 1200 MET-mins/wk); and high (≥ 1200 MET-mins/wk).

Supplementary Table S6. The Joint Association of Physical Activity and Sleep Scores with Mortality for All-Cause, Total Cardiovascular Disease, and Total Cancer (n = 380,055)

Joint Category ^a	Hazard Ratio for Mortality Risks (95% CI)					
	All-cause		Total cardiovascular disease		Total cancer	
	Model 1 ^b	Model 2 ^c	Model 1 ^b	Model 2 ^c	Model 1 ^b	Model 2 ^c
No MVPA, Poor (n = 2,510)	2.18 (1.88, 2.52)	1.57 (1.35, 1.82)	2.64 (2.02, 3.45)	1.67 (1.27, 2.19)	1.89 (1.55, 2.31)	1.45 (1.18, 1.77)
No MVPA, Intermediate (n = 28,289)	1.56 (1.48, 1.65)	1.33 (1.25, 1.40)	1.88 (1.69, 2.09)	1.50 (1.34, 1.67)	1.39 (1.29, 1.5)	1.19 (1.10, 1.29)
No MVPA, Healthy (n = 28,742)	1.3 (1.22, 1.38)	1.22 (1.15, 1.30)	1.34 (1.19, 1.52)	1.22 (1.07, 1.38)	1.21 (1.12, 1.31)	1.13 (1.04, 1.22)
Low, Poor (n = 1,291)	1.8 (1.44, 2.24)	1.46 (1.17, 1.83)	2.58 (1.78, 3.76)	1.91 (1.31, 2.78)	1.6 (1.18, 2.16)	1.35 (1.00, 1.82)
Low, Intermediate (n = 17,757)	1.24 (1.15, 1.34)	1.13 (1.05, 1.22)	1.28 (1.10, 1.49)	1.11 (0.96, 1.30)	1.19 (1.07, 1.31)	1.07 (0.97, 1.19)
Low, Healthy (n = 20,250)	1.04 (0.96, 1.12)	1.05 (0.97, 1.13)	1.11 (0.95, 1.30)	1.11 (0.95, 1.30)	1 (0.90, 1.11)	0.99 (0.89, 1.1)
Medium, Poor (n = 1,384)	1.33 (1.05, 1.69)	1.15 (0.90, 1.46)	1.66 (1.08, 2.56)	1.33 (0.86, 2.05)	1.15 (0.82, 1.6)	1.01 (0.72, 1.41)
Medium, Intermediate (n = 24,110)	1.19 (1.12, 1.28)	1.11 (1.04, 1.19)	1.26 (1.11, 1.44)	1.14 (1.00, 1.30)	1.13 (1.04, 1.24)	1.06 (0.97, 1.15)
Medium, Healthy (n = 32,277)	1.04 (0.97, 1.11)	1.04 (0.98, 1.11)	1.06 (0.93, 1.21)	1.06 (0.93, 1.21)	1.01 (0.93, 1.1)	1.01 (0.93, 1.09)
High, Poor (n = 4,981)	1.44 (1.27, 1.63)	1.21 (1.06, 1.37)	1.74 (1.39, 2.18)	1.37 (1.09, 1.72)	1.18 (0.99, 1.40)	1.02 (0.86, 1.22)
High, Intermediate (n = 87,947)	1.12 (1.07, 1.17)	1.03 (0.99, 1.08)	1.18 (1.09, 1.29)	1.07 (0.98, 1.16)	1.09 (1.03, 1.15)	1.02 (0.96, 1.07)
High, Healthy (n = 130,517)	1.00					

^a Physical activity levels were categorized based on public health guidelines: No MVPA (those reported 0 MET-min/wk from MVPA, regardless of total MET-min/wk); low (< 600 MET-min/wk); medium (600 to < 1200 MET-min/wk); and high (≥ 1200 MET-min/wk). Sleep scores were categorized into: poor, 0~1; intermediate, 2~3; healthy, 4~5.

^b Adjusted for age and sex.

^c Further adjusted for socioeconomic status, employment status, BMI, cigarette smoking, vegetable and fruit intake, alcohol consumption, sedentary behaviour, and mental health.

Supplementary Table S7. The Joint Association of Physical Activity and Sleep Scores with Mortality for Sub-Type Conditions (n = 380,055)

Joint Category ^a	Hazard Ratio for Mortality Risks (95% CI)							
	Coronary heart disease		Hemorrhagic stroke		Ischemic stroke		Lung cancer	
	Model 1 ^b	Model 2 ^c	Model 1 ^b	Model 2 ^c	Model 1 ^b	Model 2 ^c	Model 1 ^b	Model 2 ^c
No MVPA, Poor (n = 2,510)	2.70 (1.82, 4.00)	1.59 (1.07, 2.37)	1.95 (0.72, 5.29)	1.63 (0.60, 4.44)	3.98 (1.95, 8.15)	2.96 (1.43, 6.11)	3.48 (2.38, 5.10)	1.91 (1.30, 2.81)
No MVPA, Intermediate (n = 28,289)	2.11 (1.81, 2.46)	1.63 (1.40, 1.91)	1.48 (1.02, 2.16)	1.35 (0.92, 1.98)	1.78 (1.25, 2.52)	1.55 (1.08, 2.21)	2.13 (1.80, 2.52)	1.47 (1.24, 1.75)
No MVPA, Healthy (n = 28,742)	1.47 (1.23, 1.76)	1.31 (1.09, 1.56)	1.24 (0.83, 1.84)	1.20 (0.81, 1.80)	1.53 (1.06, 2.22)	1.47 (1.01, 2.14)	1.46 (1.20, 1.76)	1.25 (1.03, 1.51)
Low, Poor (n = 1,291)	3.09 (1.85, 5.15)	2.20 (1.31, 3.69)	0.94 (0.13, 6.75)	0.86 (0.12, 6.15)	0.96 (0.13, 6.88)	0.81 (0.11, 5.78)	2.66 (1.46, 4.84)	1.87 (1.02, 3.40)
Low, Intermediate (n = 17,757)	1.34 (1.07, 1.66)	1.15 (0.92, 1.44)	0.91 (0.52, 1.62)	0.89 (0.50, 1.58)	1.17 (0.70, 1.94)	1.09 (0.65, 1.82)	1.27 (0.99, 1.63)	1.02 (0.79, 1.31)
Low, Healthy (n = 20,250)	1.12 (0.89, 1.42)	1.13 (0.90, 1.43)	1.07 (0.64, 1.79)	1.12 (0.67, 1.86)	1.54 (0.99, 2.39)	1.60 (1.03, 2.49)	1.16 (0.91, 1.49)	1.18 (0.92, 1.52)
Medium, Poor (n = 1,384)	1.22 (0.58, 2.57)	0.96 (0.46, 2.03)	0.84 (0.12, 5.99)	0.79 (0.11, 5.67)	4.86 (2.14, 11.03)	4.24 (1.86, 9.66)	0.85 (0.32, 2.27)	0.67 (0.25, 1.79)
Medium, Intermediate (n = 24,110)	1.24 (1.02, 1.51)	1.12 (0.92, 1.36)	1.15 (0.74, 1.79)	1.12 (0.72, 1.75)	1.58 (1.08, 2.32)	1.49 (1.01, 2.18)	1.25 (1.01, 1.55)	1.08 (0.87, 1.33)
Medium, Healthy (n = 32,277)	1.02 (0.84, 1.25)	1.03 (0.84, 1.25)	1.01 (0.66, 1.54)	1.03 (0.68, 1.57)	1.65 (1.16, 2.33)	1.67 (1.18, 2.37)	1.07 (0.87, 1.32)	1.08 (0.88, 1.33)
High, Poor (n = 4,981)	1.57 (1.10, 2.23)	1.21 (0.85, 1.73)	1.16 (0.48, 2.85)	1.03 (0.42, 2.52)	2.47 (1.33, 4.58)	2.07 (1.11, 3.86)	1.64 (1.12, 2.41)	1.14 (0.78, 1.68)
High, Intermediate (n = 87,947)	1.31 (1.16, 1.48)	1.17 (1.04, 1.32)	1.11 (0.84, 1.47)	1.05 (0.79, 1.39)	1.32 (1.02, 1.72)	1.22 (0.94, 1.59)	1.24 (1.08, 1.42)	1.05 (0.91, 1.20)
High, Healthy (n = 130,517)					1.00			

^a Physical activity levels were categorized based on public health guidelines: No MVPA (those reported 0 MET-mins/wk from MVPA, regardless of total MET-mins/wk); low (< 600 MET-mins/wk); medium (600 to < 1200 MET-mins/wk); and high (≥ 1200 MET-mins/wk). Sleep scores were categorized into: poor, 0~1; intermediate, 2~3; healthy, 4~5.

^b Adjusted for age and sex.

^c Further adjusted for socioeconomic status, employment status, BMI, cigarette smoking, vegetable and fruit intake, alcohol consumption, sedentary behaviour, and mental health.

Supplementary Table S8. Analyses on Interaction of Sleep Scores ^a and Physical Activity ^b with Mortality Risks for All-Cause, Total Cardiovascular Disease, and Total Cancer (n = 380,055)

	Healthy Sleep		Intermediate and Poor Sleep		Hazard Ratio ^c (95% CI) for Sleep Scores within Strata of Physical Activity Level
	N With/Without Events	Hazard Ratio ^c (95% CI)	N With/Without Events	Hazard Ratio ^c (95% CI)	
All-cause death					
Any MVPA	6,526/183,044	1.00	5,927/137,470	1.06 (1.02, 1.10) <i>P</i> = 0.0021	1.06 (1.01, 1.11) <i>P</i> = 0.0021
No MVPA	1,293/28,742	1.20 (1.13, 1.28) <i>P</i> < .0001	1,757/30,799	1.33 (1.26, 1.40) <i>P</i> < .0001	1.10 (1.03, 1.19) <i>P</i> = 0.0073
Hazard Ratio ^c (95% CI) for No MVPA within Strata of Sleep		1.20 (1.13, 1.28) <i>P</i> < .0001		1.33 (1.26, 1.40) <i>P</i> < .0001	
Total cardiovascular disease death					
Any MVPA	1,644/183,044	1.00	1,601/137,470	1.08 (1.01, 1.16) <i>P</i> = 0.0236	1.08 (1.01, 1.16) <i>P</i> = 0.0236
No MVPA	325/28,742	1.19 (1.05, 1.34) <i>P</i> = 0.0050	525/30,799	1.48 (1.34, 1.63) <i>P</i> < .0001	1.24 (1.08, 1.43) <i>P</i> = 0.0021
Hazard Ratio ^c (95% CI) for No MVPA y within Strata of Sleep		1.19 (1.05, 1.34) <i>P</i> = 0.0050		1.48 (1.34, 1.63) <i>P</i> < .0001	
Total cancer death					
Any MVPA	3,945/183,044	1.00	3,431/137,470	1.03 (0.99, 1.08) <i>P</i> = 0.1763	1.03 (0.99, 1.08) <i>P</i> = 0.1763
No MVPA	738/28,742	1.13 (1.04, 1.22) <i>P</i> = .0033	9,50/30,799	1.21 (1.13, 1.30) <i>P</i> < .0001	1.08 (0.98, 1.19) <i>P</i> = 0.1324
Hazard Ratio ^c (95% CI) for No MVPA y within Strata of Sleep		1.13 (1.04, 1.22) <i>P</i> = .0033		1.21 (1.13, 1.30) <i>P</i> < .0001	

Measure of interaction on additive scale: for all-cause death, RERI (95% CI) = 0.07 (-0.03, 0.16), AP (95% CI) = 0.05 (-0.02, 0.12), and S = 1.26 (0.89, 1.79); for cardiovascular diseases death, RERI (95% CI) = 0.21 (0.01, 0.40), AP (95% CI) = 0.14 (0.02, 0.26), and S = 1.76 (0.93, 3.31); for total cancers death, RERI (95% CI) = 0.05 (-0.07, 0.18), AP (95% CI) = 0.05 (-0.05, 0.14), and S = 1.34 (0.66, 2.73).

Measure of interaction on multiplicative scale: for all-cause death, hazard ratios (HR) (95% CI) = 1.25 (1.19, 1.32), *P* < .0001; for cardiovascular diseases death, HR (95% CI) = 1.36 (1.23, 1.51), *P* < .0001; for total cancers death, HR (95% CI) = 1.17 (1.09, 1.26), *P* < .0001.

^a Sleep scores were categorized into: poor, 0~1; intermediate, 2~3; healthy, 4~5.

^b Physical activity levels were categorized based on MET-mins/wk from MVPA, regardless of total MET-mins/wk.

^c Hazard ratios were adjusted for age, sex, socioeconomic status, employment status, BMI, cigarette smoking, vegetable and fruit intake, alcohol consumption, sedentary behaviour, and mental health.

Supplementary Table S9. Analyses on Interaction of Sleep Scores ^a and Physical Activity ^b with Sub-Type Conditions Mortality (n = 380,055)

	Healthy Sleep		Intermediate and Poor Sleep		Hazard Ratio ^c (95% CI) for Sleep Scores within Strata of Physical Activity Level
	N With/Without Events	Hazard Ratio ^c (95% CI)	N With/Without Events	Hazard Ratio ^c (95% CI)	
Coronary heart diseases death					
Any MVPA	733/183,044	1.00	781/137,470	1.15 (1.04, 1.27) <i>P</i> = 0.0081	1.15 (1.04, 1.27) <i>P</i> = 0.0081
No MVPA	155/28,742	1.28 (1.08, 1.52) <i>P</i> = 0.0055	263/30,799	1.60 (1.38, 1.84) <i>P</i> < .0001	1.24 (1.02, 1.52) <i>P</i> = 0.0300
Hazard Ratio ^c (95% CI) for No MVPA within Strata of Sleep		1.28 (1.08, 1.52) <i>P</i> = 0.0055		1.60 (1.38, 1.84) <i>P</i> < .0001	
Hemorrhagic stroke death					
Any MVPA	158/183,044	1.00	130/137,470	1.02 (0.81, 1.29) <i>P</i> = 0.8688	1.02 (0.81, 1.29) <i>P</i> = 0.8688
No MVPA	31/28,742	1.18 (0.80, 1.75) <i>P</i> = 0.3908	40/30,799	1.35 (0.95, 1.93) <i>P</i> = 0.0944	1.14 (0.71, 1.83) <i>P</i> = 0.5804
Hazard Ratio ^c (95% CI) for No MVPA within Strata of Sleep		1.18 (0.80, 1.75) <i>P</i> = 0.3908		1.35 (0.95, 1.93) <i>P</i> = 0.0944	
Ischemic stroke death					
Any MVPA	184/183,044	1.00	178/137,470	1.12 (0.91, 1.38) <i>P</i> = 0.2868	1.12 (0.91, 1.38) <i>P</i> = 0.2868
No MVPA	37/28,742	1.25 (0.88, 1.79) <i>P</i> = 0.2143	51/30,799	1.42 (1.03, 1.94) <i>P</i> = 0.0315	1.13 (0.74, 1.73) <i>P</i> = 0.5724
Hazard Ratio ^c (95% CI) for No MVPA within Strata of Sleep		1.25 (0.88, 1.79) <i>P</i> = 0.2143		1.42 (1.03, 1.94) <i>P</i> = 0.0315	
Lung cancer death					
Any MVPA	626/183,044	1.00	602/137,470	1.02 (0.91, 1.15) <i>P</i> = 0.6868	1.02 (0.91, 1.15) <i>P</i> = 0.6868
No MVPA	138/28,742	1.21 (1.00, 1.45) <i>P</i> = 0.0476	2,29/30,799	1.46 (1.25, 1.71) <i>P</i> < .0001	1.21 (0.98, 1.50) <i>P</i> = 0.0765
Hazard Ratio ^c (95% CI) for No MVPA within Strata of Sleep		1.21 (1.00, 1.45) <i>P</i> = 0.0476		1.46 (1.25, 1.71) <i>P</i> < .0001	

Measure of interaction on additive scale: for coronary heart diseases death, RERI (95% CI) = 0.17 (-0.13, 0.46), AP (95% CI) = 0.11 (-0.07, 0.29), and S = 1.39 (0.74, 2.64); for hemorrhagic stroke death, RERI (95% CI) = 0.15 (-0.48, 0.78), AP (95% CI) = 0.11 (-0.34, 0.56), and S = 1.72 (0.11, 26.53); for ischemic stroke death, RERI (95% CI) = 0.04 (-0.56, 0.64), AP (95% CI) = -0.03 (-0.39, 0.45), and S = 1.12 (0.24, 5.31); for lung cancer death, RERI (95% CI) = 0.23 (-0.06, 0.53), AP (95% CI) = 0.16 (-0.03, 0.35), and S = 2.01 (0.63, 6.35).

Measure of interaction on multiplicative scale: for coronary heart diseases death, HR (95% CI) = 1.39 (1.21, 1.60), *P* < .0001; for hemorrhagic stroke death, HR (95% CI) = 1.33 (0.93, 1.90), *P* = 0.1211; for ischemic stroke death, hazard ratios (HR) (95% CI) = 1.26 (0.92, 1.73), *P* = 0.1438; for lung cancer death, HR (95% CI) = 1.43 (1.22, 1.67), *P* < .0001.

^a Sleep scores were categorized into: poor, 0~1; intermediate, 2~3; healthy, 4~5.

^b Physical activity levels were categorized based on MET-mins/wk from MVPA, regardless of total MET-mins/wk.

^c Hazard ratios were adjusted for age, sex, socioeconomic status, employment status, BMI, cigarette smoking, vegetable and fruit intake, alcohol consumption, sedentary behaviour, and mental health.