

1 **Title:** Sleep and recovery in team sport:  
2 current sleep related issues facing  
3 professional team-sport athletes  
4  
5 **Submission Type:** Brief Review  
6  
7 **Authors:** Hugh H.K. Fullagar<sup>1</sup>,  
8 Rob Duffield<sup>2</sup>,  
9 Sabrina Skorski<sup>1</sup>,  
10 Aaron J Coutts<sup>2</sup>,  
11 Ross Julian<sup>1</sup>  
12 and Tim Meyer<sup>1</sup>  
13  
14 **Institutions:** <sup>1</sup>. Institute of Sport and  
15 Preventive Medicine, Saarland  
16 University, Germany  
17 <sup>2</sup>. Sport & Exercise Discipline  
18 Group, UTS: Health, University  
19 of Technology, Australia  
20  
21 **Corresponding author** <sup>1</sup>. Hugh Fullagar  
22 Institute of Sport and Preventive  
23 Medicine, Saarland University,  
24 GEB. B82  
25 66123 Saarbrücken, Germany  
26 Email:  
27 [hughfullagar@uni-saarland.de](mailto:hughfullagar@uni-saarland.de)  
28 Phone: 0681-302 70400  
29 Fax: 0681-302 4296  
30  
31 **Running head:** Sleep and recovery in team sport  
32  
33 **Abstract word count:** 242  
34  
35 **Text only word count:** 4497  
36  
37 **References:** 50  
38  
39 **Number of Figures:** 2  
40  
41 **Number of Tables:** 1  
42  
43  
44  
45  
46  
47  
48  
49  
50

51 **ABSTRACT**

52 Whilst the effects of sleep loss on performance have previously  
53 been reviewed, the effects of disturbed sleep on recovery  
54 following exercise are far less reported. Specifically, the  
55 interaction between sleep and physiological and psychological  
56 recovery in team-sport athletes is not well understood.  
57 Accordingly, the aim of the present review is to examine the  
58 current evidence of sleep and the potential role it may play in  
59 post-exercise recovery, with a tailored focus on professional  
60 team-sport athletes. Recent studies show that team-sport  
61 athletes are at high risk of poor sleep during and following  
62 competition. Although limited published data is available, these  
63 athletes also appear particularly susceptible to reductions in  
64 both sleep quality and duration following night competition and  
65 periods of heavy training. However, studies examining the  
66 relationship between sleep and recovery in such situations are  
67 lacking. Indeed, further observational sleep studies in team-  
68 sport athletes are required to confirm these concerns. Naps,  
69 sleep extension and sleep hygiene practices appear  
70 advantageous to performance; however, future proof of concept  
71 studies are now required to determine the efficacy of these  
72 interventions on the post-exercise recovery. Moreover, more  
73 research is required to understand how sleep interacts with  
74 numerous recovery responses within team-sport environments.  
75 This is pertinent given the regularity at which these teams  
76 encounter challenging scenarios during the course of a season.  
77 Taken collectively, this review will examine the factors that  
78 compromise sleep during a season and following competition,  
79 and discuss strategies which may help improve sleep in team-  
80 sport athletes.

81  
82 **KEYWORDS:**

83 Regeneration, exercise, stress, soccer, circadian rhythms,  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100

## 101 **1. Introduction**

102 High performance team-sport athletes endure numerous  
103 physiological, psychological and neuromuscular stressors  
104 during training and competition.<sup>1</sup> It is logical that these athletes  
105 balance these stressors with appropriate recovery to maximise  
106 performance and adaptation, whilst also minimising the injury  
107 risk.<sup>2</sup> A crucial part of this stress-recovery balance is the  
108 management of an athlete's sleep, especially during intense  
109 training and competition.<sup>3</sup> However, whilst the interest  
110 afforded to the relationship between sleep and athletic  
111 performance is well documented,<sup>4</sup> the evidence underpinning  
112 the role of sleep in recovery is less understood. This is  
113 surprising from both a scientific and applied perspective given  
114 athletes often rate sleep as their most important recovery  
115 strategy.<sup>5</sup>

116 There are three key factors which determine the  
117 recuperative outcome of sleep; the duration (total sleep time),  
118 quality and phase (circadian timing) of sleep.<sup>6</sup> A 'healthy' night  
119 of sleep has been suggested to be 7-9 h.<sup>7</sup> In addition to  
120 duration, sleep quality is also critical for optimal health and  
121 restorative functioning.<sup>7</sup> Although a clear definition is not  
122 readily available, sleep quality can best be outlined as the  
123 personal satisfaction of the sleep experience.<sup>7</sup> Further, the  
124 timing of sleep will also influence the effectiveness of the sleep  
125 bout. The timing of an individual's preferred bedtime in turn  
126 affects their circadian rhythms (i.e. body temperature, hormone  
127 regulation), which can impact both sleep duration and quality.<sup>6</sup>  
128 From an athletic perspective, disturbances to one or all of these  
129 collective aspects of sleep are suggested to affect the post-  
130 exercise recovery process.<sup>6</sup> For instance, it has been shown that  
131 a reduction in the quantity and quality of sleep hinders the  
132 capacity of rugby-league footballers to recover for the demands  
133 of ensuing training and competitive bouts.<sup>8</sup> Thus, it may be  
134 paramount for team-sport athletes to be aware of situations  
135 where disturbed sleep duration, quality or phase may affect  
136 ensuing recovery.

137 A reduction in sleep duration and/or quality in  
138 individual athletes prior to,<sup>9-11</sup> and during competition<sup>12</sup> has  
139 been recently documented. Whilst there is less information  
140 available on team-sport athletes, Lastella et al<sup>13</sup> reported a  
141 mean sleep duration of 7.0 h per night in 58 elite Australian  
142 team-sport athletes during a typical training phase, ~ one hour  
143 less than the recommended 8 h per night. Further to these  
144 findings, sleep disruption or deprivation can occur for team-  
145 sport athletes, particularly during short- or long-haul travel,<sup>14-16</sup>  
146 congested competition schedules,<sup>1</sup> and training or playing at  
147 night,<sup>17</sup> presenting the potential for compromised recovery.<sup>3,8</sup>  
148 Indeed, sleep loss in team-sport athletes is often affected by  
149 these situational factors,<sup>18</sup> with many professional teams  
150 currently facing the challenge of coping with these specific, but

151 recurring stressors. For example, Major League baseballers  
152 play every two days combined with repeated travel across the  
153 United States, which provide conditions that are not conducive  
154 to optimal sleep.<sup>19</sup> Similarly, the majority of European soccer  
155 tournaments are commonly played at night, resulting in late  
156 night finishes and players subjectively reporting sleep loss.<sup>20</sup>  
157 These observations of altered sleep in team-sport athletes are  
158 also supported by objective evidence of post-competitive sleep  
159 disturbance in elite rugby union players<sup>17</sup> and professional  
160 Australian soccer players,<sup>16</sup> and a recent report that 52.3% of  
161 elite (individual and team sport) athletes experience sleep  
162 disturbances following late matches or training sessions.<sup>18</sup>  
163 Collectively, these data suggest that although ‘normal’ sleep  
164 patterns may be sufficient, under specific, recurring  
165 circumstances there are cases for reduced sleep durations and  
166 quality in team-sport athletes.

167 At present, the importance of sleep as a recovery  
168 method in team-sport athletes (i.e. return to baseline of psycho-  
169 physiological and performance parameters following exercise  
170 and disrupted sleep) is unclear. In particular, there is little  
171 analysis of the role sleep plays in the post-exercise recovery  
172 process during various situations where sleep is compromised.  
173 Whilst the literature examining the interaction between sleep  
174 and recovery in athletes is increasing (Figure 1), there have  
175 been no critical reviews of these factors in the context of  
176 training and competition demands of team-sport athletes.  
177 Accordingly, the aim of the current study was to examine the  
178 evidence of the potential role sleep may play in post-exercise  
179 recovery, with a specific focus on professional team-sport  
180 athletes. As such, an analysis of situations which may  
181 continually compromise sleep throughout a season and/or one-  
182 off post-competition sleep disturbance is provided. Strategies to  
183 alleviate such issues facing team-sport athletes are also  
184 addressed. For this review, it is important here to discern the  
185 difference between *recovery* and *performance*. From an athletic  
186 perspective, *performance* in absolute terms refers to the  
187 magnitude to which the athlete completes certain tasks within  
188 their sporting domain. These can include but are not limited to  
189 competition performance (e.g. goals scored by a footballer),  
190 predictors of performance (e.g. sprinting speed) and surrogate  
191 measures of performance (e.g. counter movement jump score).  
192 The effects of sleep loss on performance trials involve baseline  
193 performance measures followed by a sleep loss  
194 intervention/sleep control condition and then final performance  
195 measures the next morning/days. Comparatively, *recovery*  
196 refers to the *return* to baseline of performance parameters  
197 following a distinct exercise bout and disrupted sleep (e.g.  
198 return of creatine kinase to baseline values following a rugby  
199 match). Thus, the main discernible difference between  
200 performance and recovery is that recovery experiments follow a

201 distinct time-course analysis from a prior stressor (i.e. match  
202 play). This makes making them suitable for the assessment of  
203 the health, wellbeing and readiness to perform of team-sport  
204 athletes.

205

206

207

208

INSERT FIGURE ONE

209

210

211

## 212 **2. Sleep and recovery for team-sport athletes**

213 A typical night of sleep is comprised of approximately  
214 90-min cycles divided into periods of rapid-eye-movement  
215 (REM), and non-REM (NREM) sleep. Whilst REM sleep has a  
216 role in periodic brain activation, localized recuperative  
217 processes and emotional regulation, the role for NREM sleep is  
218 proposed to assist with energy conservation and nervous system  
219 recuperation.<sup>21</sup> Taken collectively, there is considerable  
220 evidence supporting the recuperative nature of sleep in  
221 restoring molecular homeostasis, cellular maintenance and  
222 synaptic plasticity.<sup>6,21,22</sup> From an athletic perspective, this  
223 implicates that disturbances to either the timing of sleep phases,  
224 or the quality and duration of sleep within these phases, can  
225 result in the hindrance of psychological and physical recovery  
226 following an exercise bout.<sup>6</sup> This would seem especially  
227 pertinent for field-based team sports that are typically exposed  
228 to prolonged bouts of intermittent-sprint activity during both  
229 high-intensity training and competition. Logically, exposure to  
230 such activity will increase the need for recovery and  
231 subsequently increase the overall requirement for sleep.<sup>13</sup>

232 From this perspective, it seems rational to first  
233 investigate the sleep-wake behaviour of team-sport athletes  
234 during and following training, and competition periods. Mah et  
235 al<sup>23</sup> reported mean average sleep durations of  $6.7 \pm 1.0$  h in  
236 collegiate basketballers during a competitive season. Similarly,  
237 Lastella et al<sup>13</sup> found a sample of 58 elite Australian team-sport  
238 athletes slept for a mean duration of  $7.0 \pm 1.2$  h during a regular  
239 training phase. With regard to sleep following competition,  
240 Eagles et al<sup>17</sup> found a significant reduction in sleep duration on  
241 game nights compared to non-game nights.<sup>17</sup> Juliff et al<sup>18</sup>  
242 reported that more than half of a sample of 283 elite individual  
243 and team-sport elite athletes (of which 210 were from team  
244 sports) endured sleep disturbances following a late training  
245 session or match.<sup>18</sup> In support of this, sleep duration and quality  
246 were significantly reduced on the night of away matches  
247 compared to the night prior in elite Australian soccer players.<sup>16</sup>  
248 Whilst caution needs to be taken in comparing these studies  
249 (i.e. due to differences in sleep-assessment methodologies), it  
250 seems reasonable to assume sleep in team-sport athletes is

251 dependent on many factors. These could include the type of  
252 sport, training demands, age, time of season and team culture.<sup>13</sup>  
253 Taken collectively, high performance team-sport athletes are  
254 considered susceptible to sleep loss during training periods and  
255 following match play (especially at night). Whilst such insight  
256 is important, further descriptive research of sleep with high  
257 performance team-sport athletes is required to confirm this,  
258 most importantly for the nights following competition.

259         Recent studies have also shown that sleep restriction  
260 following team-sport competition affects the time course of  
261 recovery for both performance and psychophysiological  
262 measures. For instance, Skein and colleagues<sup>8</sup> investigated the  
263 effect of sleep deprivation (0 h sleep) compared with normal  
264 sleep (~8 h) on the physiological and perceptual recovery of  
265 eleven rugby-league footballers following competitive matches  
266 in a randomised cross-over design. Overall, sleep deprivation  
267 negatively affected recovery with significant impairments  
268 observed in mean and peak countermovement jump height and  
269 cognitive reaction time. Although sleep deprivation was  
270 excessive, this study highlights the increased physiological load  
271 during wakefulness following sleep loss in team sports, and in  
272 turn, suppression of cognitive function and lower body power.  
273 Similarly, Fowler et al<sup>16</sup> reported significant reductions in sleep  
274 duration and quality, along with an impaired stress-recovery  
275 balance, on the night of a match compared to the night prior for  
276 away matches. Whilst additional literature is lacking in team-  
277 sport athletes, there is further evidence of this relationship in  
278 individual athletes. For instance, significant reductions in sleep  
279 quantity and efficiency were associated with increased fatigue  
280 and impaired exercise capacity in a group of ten functionally-  
281 overreached elite synchronized swimmers.<sup>24</sup> Furthermore,  
282 McMurray and Brown<sup>25</sup> investigated the cardiovascular and  
283 metabolic responses of five participants during submaximal  
284 exercise following 24 h of sleep deprivation. They reported  
285 increased minute ventilation and oxygen uptake during the  
286 recovery period, suggesting negative effects of sleep loss on  
287 physiological recovery.<sup>25</sup> Nonetheless, the evidence as to how  
288 sleep interacts with multi-factorial recovery responses within  
289 high performance team-sport environments is currently lacking.  
290 In particular, there is little longitudinal objective sleep data  
291 available in the scientific literature. This is surprising given this  
292 would appear the first step in understanding the relationship  
293 between sleep and recovery.

294         Finally, since a variety of other recovery strategies are  
295 utilised in sport, some studies have also examined the  
296 interaction between sleep and these protocols. For instance,  
297 Robey et al<sup>26</sup> reported that cold water immersion post-training  
298 does not affect subsequent sleep duration, onset or efficiency.  
299 However, the mechanisms between the interaction of sleep and  
300 other recovery protocols are difficult to determine, due to an

301 abundance of confounding factors (e.g. protocol type, timing,  
302 facilities). Further research and practical investigation within  
303 professional environments which address whether it is more  
304 advantageous to use a recovery protocol which enhances sleep  
305 and/or whether a combination of these protocols enhances the  
306 recovery process is warranted. This is especially pertinent  
307 given the wide prevalence of these methods in team sports.

308

### 309 **3. Sleep-related issues facing team-sport athletes**

310 As summarised in Figure 2, the following section  
311 outlines particular situations where sleep is at risk of  
312 compromise in team sport athletes. Whilst acknowledging the  
313 previous work done in this area but also recognising the  
314 absence of published data over prolonged periods, this gives  
315 particular relevance to situations during a season and/or one-off  
316 post-competition sleep disturbance.

317

318

319

320

INSERT FIGURE TWO

321

322

323

324

325

#### 326 **3.1 Team-sport matches played at night**

327 As often determined by television scheduling, numerous  
328 team-sports schedule the completion of matches at night.  
329 Indeed, the pure timing of matches (i.e. some matches in the  
330 Spanish La Liga commence at 22:00) will force players into  
331 later bedtimes.<sup>27</sup> Furthermore, since physical activity promotes  
332 arousal, it has long been assumed exercising during the evening  
333 hours produces a greater number of sleep disturbances than  
334 exercising during daylight.<sup>28</sup> Team-sport athletes also have  
335 extensive post-game commitments such as press conferences,  
336 recovery practises and social functions, which could lead to  
337 later bedtimes and disrupt sleep duration and quality.<sup>1</sup> As  
338 alluded to previously, Juliff et al<sup>18</sup> found 52.3% of a sample of  
339 283 elite individual (n=73) and team-sport (n=210) athletes  
340 reported sleep disturbances following a night training  
341 session/match. Moreover, 59.1% of team-sport athletes  
342 reported that that did not use a strategy to overcome these sleep  
343 disturbances.<sup>18</sup> Furthermore, a recent review on regenerative  
344 interventions used in professional soccer explains that many  
345 medical doctors report players lose sleep following night  
346 matches, which include findings on elite Bundesliga soccer  
347 players subjectively reporting reduced sleep duration and  
348 quality.<sup>20</sup> Notwithstanding these findings, the anecdotal  
349 evidence of athletes reporting sleep disturbances following  
350 night competition outweighs that documented in the literature;

351 thus, further research in elite athletic populations is required to  
352 confirm this.

353         Recent data shows that performing maximal aerobic  
354 exercise in the evening results in elevated sleep onset latency,  
355 awakenings, and REM sleep latency - suggesting poorer overall  
356 sleep quality in judo competitors.<sup>29</sup> Whilst several  
357 physiological variables are elevated prior to sleep onset  
358 following late-night vigorous exercise (suggesting possible  
359 effects on cardiac autonomic control and metabolic function<sup>30</sup>),  
360 delayed sleep onset can also be caused by mental stimulation or  
361 cognitive fatigue.<sup>22</sup> Moreover, given pain is a significant  
362 predictor of a poor night's sleep,<sup>31</sup> it is likely prolonged late-  
363 night, high-intensity exercise (equivalent to match situations)  
364 will incur sleep disturbances throughout the night as a result of  
365 pain and soreness. This is of particular relevance for heavy  
366 contact sports such as American football, ice hockey, and rugby  
367 union. It should be noted that there is opposing evidence on the  
368 effect of competing at night on sleep. For instance, Roach et  
369 al<sup>32</sup> reported no effect of two night (19:00-21:00) matches on  
370 sleep in elite junior soccer players. Similarly, Robey et al<sup>33</sup>  
371 found no effect of early evening high-intensity training on the  
372 subsequent sleep quality or duration in elite youth soccer  
373 players. In light of this, it should be recognised that the  
374 mechanisms behind the effect of exercise (and timing) on sleep  
375 are complex due to the main confounding variable (amongst  
376 others) of the stress induced by the exercise itself. From an  
377 applied perspective, future research must first focus on  
378 providing objective evidence (e.g. acute and chronic  
379 measurements of actigraphy) on whether disturbances  
380 following match play at night occur. Researchers might also  
381 focus on the effects of disrupted sleep following match play in  
382 team-sport athletes and attempt to delineate the mechanisms  
383 responsible. At present, practitioners should also be aware of  
384 the intra-individual variability in sleep requirement and  
385 chronotype (those who arise early in the morning vs. those who  
386 prefer later bedtimes). Accommodating these differences within  
387 a team environment is difficult as it may require more  
388 individualised approaches. Indeed, this would be even more  
389 pertinent for team scheduling training the day after a game. For  
390 instance, it is common after a loss for coaches to train some of  
391 their athletes hard the day after match as 'punishment'. This  
392 may create recovery concerns given players will sleep  
393 differently after these night matches.

394

### 395 **3.2 Sleep and travel fatigue**

396         Cumulative sleep loss occurs as a consequence of travel  
397 during busy periods, which tends to lead to accumulative  
398 fatigue over a season.<sup>34</sup> Travel fatigue is dependent on the  
399 distance and frequency of travel, and the length of the season. It  
400 should be noted that travel-induced fatigue is separate to jet-lag



401 fatigue, with the main difference being jet-lag comprises an  
402 effect of time-zone change.<sup>34</sup> The influences of jet-lag arising  
403 from long-haul international travel in elite athletes have been  
404 discussed previously<sup>34,35</sup> and thus will not be further addressed  
405 here. Sleep disturbances during or following travel can result in  
406 reductions in mood, acute fatigue and difficulty in initiating  
407 sleep at the arrival destination.<sup>34</sup> For team-sports, the method,  
408 mode, distance and timing of travel varies greatly and is largely  
409 dependent on scheduling, team budget and the coach's  
410 preference.<sup>36</sup> Many teams, particularly in America and  
411 Australia, endure one-way short haul domestic or international  
412 travel up to 6 h prior to or following competition.<sup>19,37,38</sup> In  
413 addition to sleep disturbances, travelling can result in  
414 detrimental health, impaired mood, dehydration and loss of  
415 motivation all of which can affect recovery.<sup>34</sup> Of further  
416 concern, it has been shown that baseball teams whose circadian  
417 rhythms are more synchronised to optimal performance times  
418 are more likely to be successful, indicating either a negative  
419 effect of travel and/or desynchronised body-clock  
420 functioning.<sup>19</sup> However, it should be noted that these data do  
421 not actually outline any physical or perceptual response to the  
422 travel, limiting its implication in athlete recovery.

423 Empirical data describing the effect of short-haul air  
424 travel on sleep, performance and the ensuing recovery in these  
425 situations is largely unknown. For instance, the sleep quantity  
426 and quality of players following away competition performance  
427 remains unclear, with short-haul air travel (1-3 h) affecting  
428 perceived sleep quality,<sup>37</sup> whereas some soccer players report  
429 earlier mean bed times after short-haul air travel (~5 h) and an  
430 away match.<sup>16</sup> Competition performance, along with reduced  
431 physical demands, appears to be greater at home compared to  
432 away (in American football<sup>38</sup>, baseball<sup>19</sup>, rugby league<sup>14</sup> and  
433 soccer<sup>16</sup>) suggesting either a negative effect of travel or a  
434 circadian advantage.<sup>35</sup> However, extrapolating these effects to  
435 determinations of match performance is difficult due to other  
436 external factors and the inter-match variability in opposition  
437 and match intensity. Whilst there have been few empirical  
438 studies, the available data suggests that short-haul travel has  
439 minimal effect on physiological and perceptual recovery (e.g.  
440 no significant effect on YoYo Intermittent Recovery level 1 test  
441 performance), with more regular or longer periods of travel  
442 (e.g. 24-h international transfers) more likely to result in  
443 negative responses.<sup>15</sup> Whilst short-haul air travel appears to  
444 have negligible effects on post-match physiological recovery,  
445 the effect on perceptual markers of fatigue and sleep patterns  
446 following competition performance is equivocal. If these  
447 parameters decline, they can negatively influence training  
448 intensity or volume during ensuing sessions due to decreased  
449 motivation.<sup>39</sup> Given the myriad of conflicting demands whilst  
450 experiencing travel and sleep loss (e.g. treatment, timing of

451 training, recovery practices), it can be difficult for coaches to  
452 manage the most appropriate schedule for their team the day  
453 after a match. Indeed more research is required to clarify the  
454 acute and chronic effects of cumulative travel (e.g. over a  
455 season) on sleep and psychological and physiological recovery  
456 parameters of professional team-sport athletes.

457

### 458 **3.3 Sleep and congested competition schedules**

459 Excessive exercise loads can disturb the stress-recovery  
460 balance and result in performance decrements and injury  
461 occurrence.<sup>2</sup> For example, during periods of heavy match  
462 congestion in soccer, there is an increased injury risk for  
463 players when they play two matches per week rather than one.<sup>40</sup>  
464 In this regard, English Premier League may compete in up to  
465 five competitions at once – which likely impacts on players’  
466 sleep behaviour. Congested schedules are also present  
467 throughout American sports such as baseball, hockey and  
468 basketball. During these periods of high physical workloads,  
469 there is a potential for a reduction in sleep duration and quality.  
470 For example, it has been shown that as the effects of increased  
471 baseball match exposure accumulate towards the end of the  
472 season strike zone judgement is impaired, which suggests a  
473 fatigue-induced decline in performance; with sleep believed to  
474 be one of the main symptoms responsible.<sup>41</sup>

475 Sleep has also been suggested to be sensitive to exercise  
476 overload - with high training volumes associated with greater  
477 sleep disruptions.<sup>42</sup> Although no published data is yet apparent  
478 in team-sport cases, Netzer et al<sup>43</sup> found significant increases in  
479 the REM sleep onset latency and decreases in REM sleep of  
480 well trained cyclists following training and a competitive 120-  
481 150 km race, compared to no training or competition.  
482 Following this, it is logical that when team-sport athletes  
483 compete in a greater number of matches within a short period,  
484 exercise-induced muscle damage will accumulate (dependant  
485 also on exercise intensity), characterised by decreased  
486 neuromuscular function, increased perceptual fatigue and  
487 increases in perceived soreness which can disrupt sleep.<sup>1</sup>  
488 Moreover, if there are several events in short succession, the  
489 continual anticipation of competition can also negate sleep.<sup>18</sup>  
490 However, at present, there is little research that describes or  
491 quantifies the effect of these changes on the subsequent  
492 recovery, particularly in team-sports undertaking congested  
493 fixture scheduling. Future investigations into the time course of  
494 recovery following sleep loss would be particularly pertinent to  
495 team sports such as baseball and cricket, since these athletes  
496 can play on consecutive days and could be at a high risk of  
497 cognitive impairments (e.g. reduction in reaction time).

498

499

500

### 501 **3.4 Sleep and disturbances to training adaptation**

502 Since sleep loss impedes muscle protein accumulation,  
503 the ability of skeletal muscle to adapt and repair can be  
504 hindered – which likely limits training adaptations.<sup>3,6,44</sup> This  
505 may be concerning during the pre-season for team-sport  
506 athletes given sleep disturbances are present during higher  
507 training volumes.<sup>42</sup> Since sleep loss can also affect vigour,  
508 mood and perceptual awareness,<sup>39</sup> early training sessions could  
509 cause reductions in motivation and consequently reduce  
510 optimal training performance and subsequent adaptations.<sup>45</sup>  
511 Furthermore, if the stress-recovery balance of team-sport  
512 athletes is disrupted by either an increase in training load/stress  
513 or inadequate recovery, it may lead to an overreached, or even  
514 overtrained state.<sup>2</sup> Interestingly, disturbed sleep is believed to  
515 be one of many symptoms of either overreaching or the  
516 overtraining syndrome.<sup>2</sup> In a recent study by Hauswirth et al<sup>46</sup>,  
517 it was found that objective measures of sleep duration,  
518 efficiency and immobile time were all negatively altered in a  
519 group of functionally overreached tri-athletes. There was also a  
520 higher prevalence of upper respiratory tract infections within  
521 this group, implying an association between the two; however  
522 whether impaired sleep and illness occurrence are  
523 consequences, or simply symptoms or coincidental  
524 associations, of overreaching remains unknown.<sup>46</sup> In light of  
525 this, practitioners are encouraged to monitor the sleeping  
526 patterns of their athletes in high periods of stress either through  
527 subjective sleep diaries and/or wristwatch actigraphy.<sup>5</sup>

528 Since sleep loss can hinder the learning of new skills,  
529 affect emotional regulation and disrupt cognitive function,<sup>6</sup> it is  
530 likely that sleep is also important for optimising cognitive  
531 training adaptations in team-sport athletes. For instance, sleep  
532 is critical for memory retention, neural plasticity, and has been  
533 shown to improve visual discrimination and motor adaptation.<sup>22</sup>  
534 Therefore, it is likely that disturbing sleep during intense  
535 training or skill acquisition periods (e.g. pre-season) will  
536 encumber adaption in skill-based tasks with high  
537 neurocognitive reliance.<sup>4</sup> However, objective evidence to  
538 support this suggestion is not currently present. Therefore,  
539 future research (with well controlled randomised-control trials)  
540 into the effects of sleep disruption on acute or chronic  
541 cognitive-based training adaptations in athletic populations is  
542 required.

543

## 544 **4. Sleep strategies for team-sport athletes**

### 545 **4.1. Napping**

546 In an attempt to recover from sleep debt, a commonly  
547 utilised sleep strategy amongst team-sport athletes is the  
548 restorative nap. Naps have been shown to improve alertness,  
549 sleepiness, short-term memory and accuracy during reaction  
550 time tests.<sup>47</sup> Furthermore, Waterhouse et al<sup>47</sup> found

551 improvements in mean sprint performance following a 30 min  
552 post-lunch nap after 4-5 h of sleep restriction. On the basis of  
553 this, it has been proposed athletes take a post-lunch nap to  
554 ameliorate the performance deficits caused by ultradian  
555 biological rhythms that occur within the circadian cycle.<sup>39,47</sup> As  
556 such, it appears napping behaviours have many benefits and  
557 should be undertaken where necessary in team-sport  
558 environments. An example would be for soccer players to have  
559 a nap after lunch if they are playing a match at night. However,  
560 it is critical that if naps are implemented within a team-sport  
561 environment they balance the need to enhance performance  
562 whilst not disturbing subsequent sleep patterns, as this could  
563 hinder the recovery process following training or competition.  
564 Indeed, whilst napping appears advantageous for performance  
565 (e.g. napping prior to competition), more research is required to  
566 evaluate its possible effectiveness in recovery.

567

#### 568 **4.2 Sleep extension**

569 Extending sleep during normal sleep times is another  
570 strategy to alleviate the decrements in physiological and  
571 cognitive performance caused by sleep loss. Mah et al<sup>23</sup> found  
572 faster sprint and reaction times and improved shooting  
573 accuracy, energy and mood following approximately three  
574 weeks of sleep extension (mean + 110 min) in eleven  
575 basketball players, indicating its use as a viable option for  
576 enhancing team-sport performance. Moreover, extending sleep  
577 improves psychological wellbeing thus optimising athletes'  
578 mental preparedness for competition.<sup>23</sup> However, obtaining  
579 extra sleep can be difficult, because increased sleep onset  
580 latency and mood effects can be nullified due to earlier  
581 bedtimes. Thus, if an athlete is not sleep deprived it is possible  
582 that extending sleep will reap no benefit. The timing of this  
583 sleep intervention could also influence the effects of sleep  
584 extension depending on the sleep chronotype of the athlete.  
585 Additionally, more research assessing whether sleep extension  
586 during periods of high-training load is a useful tool to ensure  
587 appropriate recovery is required. Such research would be  
588 pertinent in assisting players achieve higher sustained  
589 intensities in subsequent exercise bouts (i.e. during pre-season).

590

#### 591 **4.3 Sleep hygiene protocols**

592 Identifying and modifying the factors that contributes to  
593 improve sleep quality (improving sleep hygiene) in team-sport  
594 athletes can also assist in ameliorating the detrimental effect of  
595 sleep loss and potentially enhance recovery. Sleep hygiene  
596 strategies have been shown to improve sleep quality and onset  
597 latency in university students and reduced sleep irregularity in  
598 adolescents, although the effect of numerous components of  
599 sleep hygiene in normal sleepers is mixed.<sup>48</sup> From an athletic  
600 perspective, little is known about the interaction between these

601 sleep hygiene strategies and the recovery of exercise and  
602 psychological parameters. Preliminary evidence indicates  
603 adhering to some of the previous sleep hygiene  
604 recommendations improves sleep quantity, resulting in a  
605 reduction in perceived soreness and fatigue in elite tennis  
606 players.<sup>49</sup> Furthermore, regulating sleep-wake times helps  
607 synchronise the circadian timing system, improving sleep  
608 quality and quantity.<sup>50</sup> As pre-competition worry and anxiety  
609 are evident in athletes,<sup>10,18</sup> it may be of benefit to utilise self-  
610 confidence tools (i.e. meditation) to manage anxiety and stress,  
611 as these correlate with improved sleep.<sup>50</sup> Identifying each  
612 individuals best sleep habits (e.g. bed comfort) are also  
613 pertinent, as unfamiliar environments may reduce sleep  
614 quality.<sup>50</sup> Such recommendations are similar to those designed  
615 for team-sport athletes who endure constant travel.<sup>34</sup> It is well  
616 known sleep onset is prolonged by noise, light and extreme  
617 temperatures, with athletes reporting noise and light as the two  
618 most important factors to their sleep quality.<sup>10</sup> Since the use of  
619 technology just prior to sleeping promotes afferent signals from  
620 the retina to the pineal gland, inhibiting the secretion of  
621 melatonin and delaying sleep onset, the avoidance of bedtime  
622 technology (and thus reducing arousal and physiological  
623 excitement) has been recommended to improve sleep onset.<sup>50</sup>  
624 As part of a healthy sleep protocol, several nutritional  
625 recommendations have also been proposed to assist with sleep  
626 onset. For instance, a recent review by Halson<sup>5</sup> proposed diets  
627 high in carbohydrates and protein may result in shorter sleep  
628 latencies and improved sleep quality, respectively.<sup>5</sup> Whilst  
629 there is a clear need for nutrition during the post-exercise  
630 recovery period, the interaction between foods consumed post-  
631 exercise and the ensuing sleep and recovery timeline is unclear.  
632 Indeed, the effects of nutrition are intricately complex and  
633 beyond the scope of this review (see Halson<sup>5</sup> for further detail).

634

## 635 **5. Future research**

636 Currently, there is insufficient evidence to conclusively  
637 describe the role of sleep for post-exercise recovery and  
638 resultant performance outcomes. As such, the first step in  
639 understanding this contribution is for the utilisation of long-  
640 term observational field studies through the use of subjective  
641 sleep diaries and/or actimetry in various situations. This will  
642 help to identify areas where sleep may be an issue in team-sport  
643 athletes. Once this specific context is known, it is important to  
644 understand the interaction sleep has with variables within the  
645 high performance athletic environment during situations where  
646 sleep is an issue. This requires both randomised-cross over  
647 trials which investigate the measurement of sleep and the post-  
648 exercise recovery timeline (both physiological and  
649 psychological), and also case studies in high performance team-  
650 sport athletes. Future work within this field could also focus on

651 understanding the mechanisms involved and providing  
652 appropriate interventions to improve sleep and the ensuing  
653 recovery process.

654

## 655 **6. Practical recommendations for team-sport athletes**

656 The following recommendations (Table 1) are based on  
657 the literature within this review. However, the authors  
658 recognise that there is a lack of research examining the  
659 interactions between sleep and recovery in athletes.  
660 Nonetheless, there seems little risk but much (potential) benefit  
661 in following these recommendations. It is perhaps most  
662 important to tailor interventions toward individual athletes.

663

664

665

666

667

INSERT TABLE ONE

668

669

670

671

## 672 **7. Conclusion**

673 While sleep is commonly reported to be critical for  
674 recovery from intense exercise and/or competition by athletes,  
675 coaches and scientists, the current understanding of the effect  
676 of sleep on the recovery profile, especially in athletic  
677 populations, remains unclear. There is evidence to suggest elite  
678 athletes lose sleep prior to and during competition periods.  
679 Further, although limited published data is available, team-  
680 sport athletes appear to be susceptible to reductions in sleep  
681 quality and duration during and following competition  
682 (especially at night), during periods of congested fixture  
683 scheduling and following longer forms of travel. Given the  
684 regularity at which numerous professional teams might  
685 encounter these situations throughout a season, they may  
686 encumber the players sleep and recovery. The efficacy of  
687 interventions to improve sleep, such as sleep hygiene protocols  
688 and sleep extension appear advantageous - but require further  
689 investigation in situations relevant to professional team sports.  
690 These interventions may be suited to specific situations when  
691 the risk of compromised sleep is higher (i.e. playing at home or  
692 away, at night and/or inclusive of travel). This is especially  
693 pertinent with regards to the recovery of exercise parameters.  
694 Indeed, since research in this area is lacking, further research  
695 into the role of sleep and recovery in team sports is warranted.

696

## 697 **Acknowledgments:**

698 No funding was provided which contributed to the  
699 development of this manuscript. The authors declare that there  
700 are no conflicts of interest. Hugh Fullagar is supported by a

701 “Science and Health in Soccer” scholarship funded by the  
702 DAAD (German Academic Exchange Service).

703

704 **REFERENCES:**

705 1. Nédélec M, McCall A, Carling C, et al. Recovery in  
706 Soccer. *Sports Med.* 2012;43(1):9-22.

707 2. Kellmann M. Preventing overtraining in athletes in  
708 high-intensity sports and stress/recovery monitoring. *Scand J*  
709 *Med Sci Sports.* 2010;20:95-102.

710 3. Halson SL. Nutrition, sleep and recovery. *Eur J Sport*  
711 *Sci.* 2008;8(2):119-26.

712 4. Fullagar HHK, Skorski S, Duffield R, et al. Sleep and  
713 athletic performance: The effects of sleep loss on exercise  
714 performance, and physiological and cognitive responses to  
715 exercise. *Sports Med.* 2014:[Epub ahead of print].

716 5. Halson SL. Sleep in elite athletes and nutritional  
717 interventions to enhance sleep. *Sports Med.* 2014;44(Suppl  
718 1):S13–S23.

719 6. Samuels C. Sleep, recovery, and performance: The new  
720 frontier in high-performance athletics. *Neurol Clin.*  
721 2008;26(1):169-80.

722 7. Krystal AD, Edinger JD. Measuring sleep quality. *Sleep*  
723 *Med.* 2008;9(S1):S10–S7.

724 8. Skein M, Duffield R, Minett G, et al. The effect of  
725 overnight sleep deprivation after competitive rugby league  
726 matches on postmatch physiological and perceptual recovery.  
727 *Int J Sport Physiol Perform.* 2013;8:556-64.

728 9. Erlacher D, Ehrlenspiel F, Adegbesan OA, et al. Sleep  
729 habits in German athletes before important competitions or  
730 games. *J Sports Sci.* 2011;29(8):859-66.

731 10. Venter RE. Perceptions of team athletes on the  
732 importance of recovery modalities. *Eur J Sport Sci.*  
733 2014;14:S69-S76.

734 11. Lastella M, Lovell GP, Sargent C. Athletes'  
735 precompetitive sleep behaviour and its relationship with  
736 subsequent precompetitive mood and performance. *Eur J Sport*  
737 *Sci.* 2014:1-8.

738 12. Lastella M, Roach G, Halson SL, et al. Sleep/wake  
739 behaviour of endurance cyclists before and during competition.  
740 *J Sports Sci.* 2014;8:1-7.

741 13. Lastella M, Roach G, Halson SL, et al. Sleep/wake  
742 behaviours of elite athletes from individual and team sports.  
743 *Eur J Sport Sci.* 2014:[Epub ahead of print].

744 14. McGuckin TA, Sinclair WH, Sealey RM, et al. The  
745 effects of air travel on performance measures of elite Australian  
746 rugby league players. *Eur J Sport Sci.* 2014;14:S116-S22.

747 15. Fowler P, Duffield R, Vaile J. Effects of simulated  
748 domestic and international air travel on sleep, performance and  
749 recovery. *Scand J Med Sci Sports.* 2014:[Epub ahead of print].

- 750 16. Fowler P, Duffield R, Vaile J. Effects of domestic air  
751 travel on technical and tactical performance and recovery in  
752 soccer. *Int J Sport Physiol Perform.* 2014;9(3):378-86.
- 753 17. Eagles A, McLellan C, Hing W, et al. Changes in sleep  
754 quantity and efficiency in professional rugby union players  
755 during home based training and match-play. *J Sports Med Phys*  
756 *Fitness.* 2014;[Epub ahead of print].
- 757 18. Juliff LE, Halson SL, Peiffer JJ. Understanding sleep  
758 disturbance in athletes prior to important competitions. *J Sci*  
759 *Med Sport.* 2014;[Epub ahead of print].
- 760 19. Winter WC, Hammond WR, Green NH, et al.  
761 Measuring circadian advantage in Major League Baseball: a  
762 10-year retrospective study. *Int J Sports Physiol Perform.*  
763 2009;4(3):394-401.
- 764 20. Meyer T, Wegmann M, Poppendieck W, et al.  
765 Regenerative interventions in professional football. *Sport*  
766 *Orthop Traumatol.* 2014;30:112-8.
- 767 21. Vyazovskiy VV, Delogu A. NREM and REM Sleep:  
768 Complementary roles in recovery after wakefulness. *The*  
769 *Neuroscientist.* 2014;[Epub ahead of print].
- 770 22. Stickgold R. Sleep-dependent memory consolidation.  
771 *Nature.* 2005;437:1272-8.
- 772 23. Mah CD, Mah KE, Kezirian EJ, et al. The effects of  
773 sleep extension on the athletic performance of collegiate  
774 basketball players. *Sleep.* 2011;34(7):943-50.
- 775 24. Schaal K, Le Meur Y, Louis J, et al. Whole-body  
776 cryostimulation limits overreaching in elite synchronized  
777 swimmers. *Med Sci Sports Exerc.* 2014;[Epub ahead of print].
- 778 25. McMurray RG, Brown CF. The effect of sleep loss on  
779 high intensity exercise and recovery. *Aviat Space Environ Med.*  
780 1984;55(11):1031-5.
- 781 26. Robey E, Dawson B, Halson S, et al. Effect of evening  
782 postexercise cold water immersion on subsequent sleep. *Med*  
783 *Sci Sport Exerc.* 2013;45(7):1394-402.
- 784 27. Nédélec M, McCall A, Carling C, et al. Recovery in  
785 soccer: Part II-recovery strategies. *Sports Med.* 2013;43(1):9-  
786 22.
- 787 28. Porter JM, Horne JA. Exercise and sleep behaviour - a  
788 questionnaire approach. *Ergonomics.* 1981;24:511-21.
- 789 29. Souissi N, Chtourou H, Aloui A, et al. Effects of time-  
790 of-day and-partial sleep deprivation on short term maximal  
791 performances of judo competitors. *J Strength Cond Res.*  
792 2013;27(9):2473-80.
- 793 30. Myllymaki T, Kyröläinen H, Savolainen K, et al.  
794 Effects of vigorous late-night exercise on sleep quality and  
795 cardiac autonomic activity. *J Sleep Res.* 2011;20(1):146-53.
- 796 31. Raymond I, Nielsen TA, Lavigne G. Quality of sleep  
797 and its daily relationship to pain intensity in hospitalized adult  
798 burn patients. *Pain.* 2001;92:381-8.



- 799 32. Roach GD, Schmidt WF, Aughey RJ, et al. The sleep of  
800 elite athletes at sea level and high altitude: a comparison of sea-  
801 level natives and high-altitude natives (ISA3600). *Br J Sports*  
802 *Med.* 2013;47:i114-i20.
- 803 33. Robey E, Dawson B, Halson S, et al. Sleep quantity and  
804 quality in elite youth soccer players: A pilot study. *Eur J Sport*  
805 *Sci.* 2013:1-8.
- 806 34. Samuels C. Jet lag and travel fatigue: a comprehensive  
807 management plan for sport medicine physicians and high-  
808 performance support teams. *Clin J Sport Med.* 2012;22:268-73.
- 809 35. Reilly T. How can travelling athletes deal with jet-lag?  
810 *Kinesiology.* 2009;41:128-35.
- 811 36. Bishop D. The effects of travel on team performance in  
812 the Australian national netball competition. *J Sci Med Sport.*  
813 2004;7(1):118-22.
- 814 37. Richmond LK, Dawson B, Stewart G, et al. The effect  
815 of interstate travel on the sleep patterns and performance of  
816 elite Australian Rules footballers. *J Sci Med Sport.*  
817 2007;10(4):252-8.
- 818 38. Smith RS, Guilleminault C, Efron B. Circadian rhythms  
819 and enhanced athletic performance in the National Football  
820 League. *Sleep.* 1997;20(5):362-5.
- 821 39. Reilly T, Edwards B. Altered sleep-wake cycles and  
822 physical performance in athletes. *Physiol Behav.* 2007;90(2-  
823 3):274-84.
- 824 40. Dupont G, Nedelec M, McCall A, et al. Effect of 2  
825 soccer matches in a week on physical performance and injury  
826 rate. *Am J Sport Med.* 2010;38(9):1752-8.
- 827 41. Kutscher S, Song Y, Wang L, et al. Validation of a  
828 statistical model predicting possible fatigue effect in Major  
829 League Baseball. *Sleep.* 2013;36:A408.
- 830 42. Taylor SR, Rogers GG, Driver HS. Effects of training  
831 volume on sleep, psychological, and selected physiological  
832 profiles of elite female swimmers. *Med Sci Sport Exer.*  
833 1997;29(5):688-93.
- 834 43. Netzer N, Kristo D, Steinle H, et al. REM sleep and  
835 catecholamine excretion: a study in elite athletes. *Eur J Appl*  
836 *Physiol.* 2001;84:521-6.
- 837 44. Dattilo M, Antunes HKM, Medeiros A, et al. Sleep and  
838 muscle recovery: Endocrinological and molecular basis for a  
839 new and promising hypothesis. *Med Hypotheses.*  
840 2011;77(2):220-2.
- 841 45. Sargent C, Halson S, Roach GD. Sleep or swim? Early-  
842 morning training severely restricts the amount of sleep obtained  
843 by elite swimmers. *Eur J Sport Sci.* 2014;14:S310-S5.
- 844 46. Hausswirth C, Louis J, Aubry A, et al. Evidence of  
845 disturbed sleep and increased illness in overreached endurance  
846 athletes. *Med Sci Sport Exer.* 2014;46(5):1036-45.
- 847 47. Waterhouse J, Atkinson G, Edwards B, et al. The role of  
848 a short post-lunch nap in improving cognitive, motor, and

- 849 sprint performance in participants with partial sleep  
850 deprivation. *J Sports Sci.* 2007;25(14):1557-66.
- 851 48. Stepanski E, Wyatt J. Use of sleep hygiene in the  
852 treatment of insomnia. *Sleep Med Rev.* 2003;7(3):215-25.
- 853 49. Duffield R, Murphy A, Kellett A, et al. Recovery from  
854 repeated on-court tennis sessions; combining cold water  
855 immersion, compression and sleep recovery interventions. *Int J*  
856 *Sport Physiol Perform.* 2013:[Epub ahead of print].
- 857 50. Malone SK. Early to bed, early to rise?: An exploration  
858 of adolescent sleep hygiene practices. *J School Nursing.*  
859 2011;27(5):348-54.

860

861

862 **FIGURE CAPTIONS**

863

864 **Figure 1:** The increase in the number of sleep, athlete and  
865 recovery publications over the past eight years. The  
866 solid fill lines illustrate the amount of literature which  
867 appears following a Pub Med database search using the  
868 terms “sleep”, “recovery” and “athlete” in all fields for  
869 each calendar year.

870

871 **Figure 2:** A schematic representation of the commonly  
872 encountered situations in team sports which may  
873 compromise sleep patterns and potentially recovery.  
874 Theoretical effects of these situations are also  
875 described; however it should be noted more research is  
876 required to confirm the majority of these effects.

877