

Sleep Disturbances in Chronic Pain: Neurobiology, Assessment, and Treatment in Physical Therapist Practice

Jo Nijs, Olivier Mairesse, Daniel Neu, Laurence Leysen, Lieven Danneels, Barbara Cagnie, Mira Meeus, Maarten Moens, Kelly Ickmans, Dorien Goubert

Among people with chronic pain, insomnia is highly prevalent, closely related to the mechanism of central sensitization, characterized by low-grade neuroinflammation, and commonly associated with stress or anxiety; in addition, it often does not respond effectively to drug treatments. This review article applies the current understanding of insomnia to clinical practice, including assessment and conservative treatment of insomnia in people with chronic pain. Cognitive-behavioral therapy for insomnia can be efficacious for improvements in sleep initiation, sleep maintenance, perceived sleep quality, and pain interference with daily functioning in people with chronic pain. A recent systematic review concluded that with additional training, physical therapist–led cognitive-behavioral interventions are efficacious for low back pain, allowing their implementation within the field. Cognitive-behavioral therapy for insomnia, as provided to people with chronic pain, typically includes education, sleep restriction measures, stimulus control instructions, sleep hygiene, and cognitive therapy.

J. Nijs, PT, MT, PhD, Pain in Motion International Research Group, Department of Physiotherapy, Human Physiology and Anatomy, Faculty of Physical Education & Physiotherapy, the Department of Physical Medicine and Physiotherapy, University Hospital Brussels, Belgium, and Vrije Universiteit Brussel, Building F-Kima, Laarbeeklaan 103, BE-1090 Brussels, Belgium. Address all correspondence to Dr. Nijs at: Jo.Nijs@vub.be.

O. Mairesse, PhD, Brugmann University Hospital, Sleep Laboratory and Unit for Chronobiology U78, Université Libre de Bruxelles - Vrije Universiteit Brussel, Brussels, Belgium, and the Department of Experimental and Applied Psychology (EXTO), Vrije Universiteit Brussel.

D. Neu, PhD, Brugmann University Hospital, Sleep Laboratory and Unit for Chronobiology U78, Université Libre de Bruxelles - Vrije Universiteit Brussel, and UNI Neuroscience Institute, ULB312 Faculty of Medicine, and ULB388 Faculty of Motor Sciences, Université Libre de Bruxelles.

L. Leysen, PT, MT, MSc, Pain in Motion International Research Group, and the Department of Physiotherapy, Human Physiology and Anatomy, Faculty of Physical Education & Physiotherapy, Vrije Universiteit Brussel.

L. Danneels, PT, PhD, Department of Rehabilitation Sciences and Physiotherapy, Faculty of Medicine, Ghent University, Ghent, Belgium.

B. Cagnie, PT, PhD, Department of Rehabilitation Sciences and Physiotherapy, Faculty of Medicine, Ghent University.

M. Meeus, PT, PhD, Pain in Motion International Research Group, Department of Rehabilitation Sciences and Physiotherapy, Faculty of Medicine, Ghent University.

Author information continues on next page.



Post a comment for this article at:
<https://academic.oup.com/ptj>

Insomnia Intervention for Chronic Pain

M. Moens, MD, PhD, Department of Neurosurgery and Radiology, University Hospital, and the Department of Manual Therapy, Faculty of Medicine and Pharmacy, Vrije Universiteit Brussel.

K. Ickmans, PT, PhD, Pain in Motion International Research Group, Department of Physiotherapy, Human Physiology and Anatomy, Faculty of Physical Education & Physiotherapy, Vrije Universiteit Brussel, and the Department of Physical Medicine and Physiotherapy, University Hospital Brussels.

D. Goubert, PT, PhD, Pain in Motion International Research Group, Department of Physiotherapy, Human Physiology and Anatomy, Faculty of Physical Education & Physiotherapy, Vrije Universiteit Brussel, and Department of Rehabilitation Sciences and Physiotherapy, Faculty of Medicine, Ghent University.

[Nijs J, Mairesse O, Neu D, et al. Sleep disturbances in chronic pain: neurobiology, assessment, and treatment in physical therapist practice. *Phys Ther.* 2018;98:325–335.]

© 2018 American Physical Therapy Association

Accepted: February 5, 2018

Submitted: June 11, 2017

Conservative and pharmacological strategies for chronic pain management offer, at best, modest effect sizes in reducing pain and related disabilities^{1–3}; thus, there is a need for improved care. A comprehensive approach to chronic pain management should also address pain-associated conditions, such as comorbid insomnia, a topic seldom addressed in the physical therapy literature. Indeed, sleep is increasingly recognized as a plausible therapeutic target for a range of chronic conditions, including chronic pain.^{4,5} If left untreated, insomnia can represent a barrier for effective chronic pain management.⁶ Chronic pain is described as debilitating pain with a duration of at least 3 months.

In the absence of other intrinsic sleep disorders and inadequate opportunity or circumstances for sleep (eg, shift work), insomnia in adults is defined as more than 30 minutes of sleep latency and/or minutes awake after sleep onset for more than 3 days per week for more than 3 months.^{5–7} Insomnia is highly prevalent among people with chronic pain, with 53% to 90% having a clinically significant degree of insomnia.^{8–11} People with chronic back pain are 18 times more likely to experience clinically defined insomnia than those without chronic back pain.¹⁰

Insomnia is closely related to pain severity in people with chronic (low back) pain.¹⁰ People with whiplash-associated disorders have insomnia problems such as difficulties initiating sleep and difficulties maintaining sleep, which are related to cognitive disturbances, such as reduced information processing speed.¹² Insomnia is usually considered a consequence of chronic pain, but research findings show that sleep disturbances may have a bidirectional relationship with chronic pain.^{6,9,13}

Insomnia may act as both a precipitating factor and a perpetuating factor.⁵ Population-based longitudinal studies showed that sleep impairments predicted the onset and exacerbation of chronic pain.¹³ On the basis of the available literature, it was concluded that sleep impairments predict chronic pain more

strongly than pain predicts sleep impairments.¹³ Insomnia is independently associated with the perceived impact of pain on daily functioning and life satisfaction in people with chronic spinal pain.⁵ Lastly, people with chronic pain spontaneously engage in more physical activity following a better night of sleep¹⁴ (Fig. 1). The latter suggests that improving nighttime sleep may be a novel avenue for promoting greater daytime physical activity in people with chronic pain.¹⁴

Despite the growing body of scientific literature suggesting a high prevalence of insomnia in individuals with chronic pain, many pain treatment programs propose little other than sedative pain and/or hypnotic neuropharmacological drugs for the treatment of comorbid insomnia.¹⁰ These drugs include anticonvulsants, nonbenzodiazepines, benzodiazepines, and tricyclic antidepressants. These drugs are problematic for 2 reasons. First, the efficacy and safety of drug treatment for insomnia in people with chronic pain has not been established. Furthermore, the usage of hypnotic medications is associated with a number of side effects, including sedation, daytime drowsiness, dizziness, lightheadedness, and cognitive and psychomotor impairments.^{15,16} Pharmacological strategies directed at nociceptive mechanisms do not yield improvements in sleep in people with chronic pain.^{17–20} Moreover, many trials in people with headaches or musculoskeletal pain found that medication may disturb sleep and even increase pain.²¹ Finally, reliance on pharmacological treatments for insomnia often opposes patients' preferences for non-pharmacological alternatives.²²

Consequently, there is a need to inform physical therapists about the close interaction between insomnia and the chronic pain experience. Here, we aim to update the reader with the current understanding of the neurobiology behind the interaction between insomnia and chronic pain and suggest a conservative approach to the management of insomnia in people with chronic pain within a physical therapy setting. Henceforth, clinical guidelines for the




Figure 1. Clinical importance of insomnia in people with chronic pain.^{5,6,8-11,14}

assessment and treatment of insomnia in people with chronic pain within a physical therapy setting will be provided. These recommendations for assessment and treatment of insomnia are based on the available scientific evidence and the authors' clinical expertise.

How Sleep Can Influence Chronic Pain: Neuroinflammation, Central Sensitization, and Beyond Sleep and Central Sensitization: A Bidirectional Interaction

Central sensitization implies increased neuronal response to stimuli in the central nervous system (ie, central hyperexcitability).²³⁻²⁵ Although the concept of central sensitization originates primarily from laboratory, the awareness is growing that it should be part of our clinical reasoning in daily practice.²⁶⁻²⁸ There is a close interaction between central sensitization and sleep disturbances in people with chronic pain. A single night of total sleep deprivation has been shown to induce generalized hyperalgesia and increase state anxiety in healthy people.^{29,30} Likewise, sleep curtailment impairs endogenous nociceptive-inhibitory function and increases

spontaneous pain in healthy people.³¹ These findings suggest that sleep disturbances might not only perpetuate central nervous system hyperexcitability in people with chronic pain, but may also serve as an aetiological factor. This may in turn ensue a vicious cycle: poor sleep lowers pain thresholds,^{6,31} which then contributes to hyperalgesia³¹ and subsequent increased incidence and/or severity of insomnia.⁶

The link between central sensitization and insomnia is underscored by various research findings. People with knee osteoarthritis have more central sensitization when they have comorbid insomnia.³² The same study revealed that people with osteoarthritis who have reduced sleep efficiency and higher catastrophizing show more signs of central sensitization.³² Likewise, in a large sample of 961 people with chronic pain, pain-related sleep interference was significantly associated with pain hypersensitivity as measured by quantitative sensory testing, but not with spinal nociceptive hypersensitivity as typically assessed by means of the electromyography response to an electrical stimulus (nociceptive withdrawal reflex).³³ Pain hypersensitivity was assessed using pain threshold to electrical

stimulation. Moreover, the magnitude of pain-related interference with sleep was associated with increased risk of pain hypersensitivity in people with chronic pain.³³ The authors concluded that improving sleep and sleep quality may be important for the management of pain hypersensitivity, potentially leading to pain reduction.³³ In people with temporomandibular joint disorder, insomnia is associated with hyperalgesia at a nonorofacial site, suggesting that insomnia may be linked with central sensitization.³⁴

Roles of Dopaminergic, Serotonergic, and Opioidergic Systems in Explaining Insomnia in People With Chronic Pain

The link between insomnia and central sensitization might imply overlapping mechanisms in the central nervous system, for instance mesolimbic dopaminergic pathways playing a cardinal role in both sleep regulation and endogenous analgesia.³⁵ Hence, decreased dopamine availability may explain both sleep disturbances and dysfunctional endogenous analgesia in people with chronic pain. Serotonergic pathways are also implicated in shared modulatory mechanisms of pain and sleep regulations. Serotonin is critical for top-down orchestration of endogenous analgesia³⁶ as well as of circadian rhythm control.³⁷ Serotonergic dysfunction can result in altered patterns of circadian behavior or even contribute to a disruption of sleep-wake homeostasis³⁷ along with dysfunctional endogenous analgesia.³⁸ While it is unlikely that either of the 2 mentioned neurotransmitters is the sole factor underlying the links between pain and insomnia, both seem to play major roles in explaining sleep disturbances in people with chronic pain. Essentially, a possible dysfunction in either dopaminergic or serotonergic systems fits within the view that central sensitization is exacerbated or even initiated by sleep disturbances. Preclinical studies support this view by revealing that selective melatonin MT2 receptor partial agonists hold analgesic properties through modulation of "on/off" cells of the brainstem descending antinociceptive systems in neuropathic pain models.^{39,40}

Opioidergic signaling is also implicated in shared modulatory mechanisms of pain and sleep regulations. Opioids are known to influence sleep-wake regulation at least in part mediated by central opioid input to the ventrolateral preoptic nucleus, a key cell group for producing behavioral sleep.⁴¹ Morphine decreases the total amount of deep sleep and rapid eye movement sleep in humans through inhibiting the firing rate of sleep-promoting neurons in the ventrolateral preoptic area.⁴²

Poor Sleep Results in Low-Grade Central Nervous System Inflammation

In addition, our current understanding of sleep-related neuro-immunology provides potential links between sleep impairments and pain. While healthy sleep facilitates immune functions, impaired sleep quality or quantity can result in low-grade inflammatory responses.⁴³⁻⁴⁵ This low-grade inflammatory response as a consequence of sleep deprivation includes increased levels of interleukin 6, prostaglandin E2^{44,45} and nitric oxide⁴⁶ possibly mediated by cerebral microglia.⁴⁶ Allegedly, even low levels of inflammatory cytokines are known to potentially affect brain function.⁴⁷ These cytokines interfere with central nervous system-mediated fatigue⁴⁸ and correlate with observations of increased sensitivity to painful stimuli following sleep restriction.^{30,44,49} Taken together, sleep deprivation conveys a glia-mediated low-grade inflammatory response leading toward increased sensitivity to pain as observable in people with chronic pain.⁵⁰ The nature of neuro-immunologic links between sleep and pain may be important for the inclusion of pain neuroscience education in people with chronic pain (ie, empower patients to invest time in cognitive-behavioral therapy [CBT] for insomnia [CBT-I]).

Role of Stress in Explaining Insomnia in People With Chronic Pain

Stress and sleep are consistently interconnected, as evidenced in numerous studies reporting strong associations between anxiety levels and insomnia severity,^{51,52} such as in people with chronic low back pain.¹⁰ Daily life stress

(eg, worries about the next morning's workload) can negatively impact sleep.⁵³ Similarly, major stressful life events and/or traumatic events such as natural disasters, combat, or a traffic accident can result in sleep architecture alterations reflecting poor sleep.⁵³ Increased nighttime arousal and decreased sleep efficiency are among the most sensitive sleep variables in response to stressors.⁵³

In addition, health anxiety has also been described as a significant predictor of insomnia severity.¹⁰ Health anxiety refers to a specific type of anxiety characterized by excessive concerns about one's health (eg, a preoccupation with fears of having a serious disease, despite medical reassurance). It may contribute to aggravate or even trigger insomnia by inducing hypervigilance and arousal by activation of a cascade of cognitive and behavioral processes such as selective attention to threats and negative appraisal.¹⁰ Likewise, depression accounts in part for the variance in sleep quality among people with chronic pain.^{54,55}

The preceding paragraphs have documented how stress can impair sleep, but sleep itself can be a stressor. Individuals with insomnia often show high levels of apprehension about bedtime and performance anxiety in an attempt to control the process of sleep onset; and even engage in catastrophic thinking about potential consequences of poor sleep.⁵⁶ The latter may in turn result in a marked decrease of coping abilities with every day life's stressors. For these reasons, it comes as no surprise that sleep management is mostly included as a part of stress management programs and vice versa. In a successful trial of CBT-I in people with chronic pain, improvers not only showed a significant increase in perceived sleep efficiency but also a decrease in self-reported levels of distress,⁵⁷ suggesting that effective coping with stress is a key factor for sleep improvements.

The preceding text has explained how insomnia is of significance to people with chronic pain, with special emphasis on the neurobiology of these interactions. The following text explains

how the current understanding of insomnia in people with chronic pain is applied to clinical practice (including assessment and CBT-I).

Applying Science to Practice I: Assessment of Insomnia in People With Chronic Pain in Clinical Practice

Depending on the education received, physical therapists may not be trained to screen for insomnia, and patients are not referred to physical therapists for insomnia either. Yet physical therapists are advised to briefly question people with chronic pain about sleep latency and minutes awake after sleep. This can allow them to estimate whether the patient complies with the definition of insomnia (ie, more than 30 minutes of sleep latency and/or minutes awake after sleep onset for more than 3 days per week for more than 3 months⁵⁻⁷). It is advised to determine from the patient if the insomnia began before or after the pain condition and consider how the pain condition is influencing their insomnia symptoms. In addition, the Insomnia Severity Index generates reliable and valid data to quantify the subjective harshness of insomnia.⁵⁸ A cutoff level of 14 (ie, range of 0-14 vs 15-28) has been proposed for the clinical significance of chronic insomnia in patients presenting with chronic back pain,¹⁰ and greater than 10 was recommended as the cutoff in a community sample.⁵⁹ In case of suspected insomnia the physical therapists should refer the patient to a sleep physician who can use polysomnography to ascertain whether underlying primary sleep disorders are present.^{6,60} Such primary sleep disorders include sleep-related breathing or movement disorders, circadian sleep-wake disorders, etc. The polysomnography is not used to diagnose insomnia, but to rule out other sleep disorders.

In case of diagnosed or suspected insomnia disorder, physical therapists should then explore whether insomnia coexists with the chronic pain experience. This is achieved by ascertaining that insomnia is reported

by the patient to originate after and/or to be aggravated by the current pain condition.⁶ If this is the case and the physical therapist is planning to provide CBT-I, a thorough questioning of the patient's sleep habits and difficulties is required. Such an interview may include the items listed in Table 1. Table 1 comprises not only items for questioning the patient about sleep habits and difficulties, but a guide on how to interpret the obtained information and treatment implications.

Whether the patient has been or is currently a substance abuser is not only important because of its potentially negative impact on biological functions of sleep, but also because past or current substance abusers' dopaminergic circuits may be altered as a consequence of repeated activations associated with consumption.³⁵ In addition to sleep history taking, therapists may consider using self-reported tools for the assessment of sleep parameters. Such self-reported tools include sleep diaries, questionnaires like the Insomnia Severity Index, the Pittsburgh Sleep Quality Index and/or the Epworth sleepiness scale.^{61,62}

Applying Science to Practice II: Conservative Treatment of Insomnia in People With Chronic Pain

CBT is effective for the treatment of chronic pain.^{2,63-65} CBT is a psychosocial intervention that aims at changing unhelpful thoughts, beliefs, and attitudes in order to improve their coping skills, self-regulation, and healthy behavior. CBT is widely used for the treatment of chronic pain and depression. A systematic review concluded with high-quality evidence that with additional training, physical therapist-led cognitive-behavioral interventions are effective for low back pain.⁶⁶ CBT-I is a specific form of CBT for a specific condition, one that is a candidate to be considered for psychologically informed practice delivered by physical therapists.⁶⁷

Available Evidence for Conservative Sleep Treatment in People With Chronic Pain

A meta-analysis of available studies examining the effects of nonpharmacological treatments of insomnia for cancer and non-cancer-related chronic pain conditions showed that addressing sleep resulted in large immediate improvements in sleep quality, relatively small reductions in pain and fatigue and moderate decrease of depression symptoms.⁴ The improvements in sleep quality and fatigue were maintained after a 1-year follow-up.⁴

CBT-I is the recommended evidence-based treatment for chronic primary insomnia.⁶⁰ The advantage over sedative drugs or classical hypnotics mainly lies in the sustainability of treatment effects and a lower risk for side effects.⁶⁰ Based on the available treatment studies in people with chronic pain, a similar picture arises. The first report of a treatment trials targeted at sleep for people with chronic pain dates back to 2010.^{5,68} CBT-I results in improvements in various sleep parameters (eg, self-reported sleep onset latency, time spent awake after sleep onset, sleep efficiency, and sleep quality), sleep and pain interference with daily functioning in people with chronic pain.^{5,68,69} Improvements were maintained at the 3-month follow-up.⁶⁸ The clinical relevance of the treatment effects was emphasized by the large difference between group responders (78% in the CBT-I-group vs 22% in the control group) and non-responders (42% vs 11%).⁵ The effect sizes varied from moderate to large for sleep parameters and pain interference, but were small for pain intensity.⁵

Should CBT only target sleep, or should it target both pain and sleep? The results from 2 smaller pilot studies are in support of the latter: combining CBT for pain with CBT-I was feasible and produced significant improvements in sleep, disability from pain, pain interference, depression, and fatigue.^{6,70} Importantly, the combined intervention appeared to have a strong advantage over CBT for pain alone and modest advantage over CBT-I alone in reducing

insomnia severity in people with chronic pain.⁶ The gains in insomnia severity and pain interference were maintained at 1- and 6-months follow-up.⁷⁰

Taken together, recently accumulated knowledge in the field has progressed and contributed to increase our understanding about the interplay between sleep and pain. It has been shown that CBT-I is efficacious for the improvement of sleep duration, maintenance of sleep, perceived sleep quality,⁶⁰ and pain interference with daily functioning⁵ in people with chronic pain. Additionally, there is relatively little available evidence about the optimal care-path for sleep related problems in the context of chronic pain.¹¹ There is a need for an improved reporting of procedural information allowing clinicians to replicate study findings in daily routine.⁶⁶ To these extents, the subsequent question is: what should be the exact content of such a CBT-I in people with chronic pain?

Introduction to CBT-I for People With Chronic Pain and Insomnia

CBT-I includes sleep and bedtime restrictions, stimulus control instructions, education for sleep and promoting good sleep habits (or sleep hygiene), teaching relaxation skills, and cognitive therapy.^{5,68,71} Each of these main tenets of CBT-I are introduced and described in Table 2 (to be used together with Tab. 1 to allow individually tailored CBT-I). For more details and a comprehensive guide for clinicians, readers are referred to an available manual.⁷² Clinicians who wish to provide CBT-I can benefit from additional training and are advised to attend an educational course.

Addressing poor sleep routine by scheduling a new sleep pattern¹¹ and poor sleep efficiency (eg, 8 hours in bed and only 4 hours of sleep)¹¹ can be done by providing classical bedtime restriction approaches.^{5,11} In case of impaired sleep efficiency and inappropriate sleep routines, this includes sticking to the same getting up time (or advancing the latter) as usual, but getting into bed at a later time when sufficiently sleepy, in order to reduce the amount of time spent in bed and increase homeostatic

Insomnia Intervention for Chronic Pain

Table 1.

Determination of a Patient's Sleep Habits and Difficulties, a Guide for Interpreting the Obtained Information, and Treatment Implications^a

Item	Explanation	Mode of Assessment	Sample Questions	Interpretation	Treatment Implications
(Non) restorative sleep	Recuperative sleep or not; whether a patient feels refreshed at final awakening.	Questioning/PSQI	Do you feel refreshed when you wake up?	Ideally, good sleep should be refreshing. Nonrestorative sleep can be a symptom of disordered sleep.	If a patient has recurrent sensations of nonrestorative sleep or unrefreshing morning arousals, then further inquiry about sleep is indicated.
Time in bed ⁸⁰	Amount of time spent in bed.	Questioning/sleep diary/PSQI/actigraphy	How much time do you spend in bed over the course of 24 h?	Interpretation should take into account time spent in and out of bed as well as time spent awake in bed (ie, sleep efficiency).	Treatment implications rely on the interpretation of various sleep items taken together; the amount of time spent in bed at night alone has no treatment implication per se.
Time out of bed ⁸⁰	Amount of time spent (awake or asleep) out of bed at night.	Questioning/sleep diary/actigraphy	How much time do you spend out of bed during the night?	If a patient spends more than half an hour out of bed at night, then further questioning about what he or she does while out of bed is required. In general, this problem may be related to poor sleep quality.	Unless the time spent out of bed is part of stimulus control instructions (Tab. 2), efforts should be made to limit time out of bed at night. Activities during time out of bed at night should be relaxing and not physically or emotionally stimulating and should aim to not further interfere with sleep.
Sleep latency ⁸⁰	Time from "lights off" to sleep onset.	Questioning/sleep diary/actigraphy	How long does it take for you to fall asleep?	Ideally, sleep latency should not be longer than 30 min.	If sleep latency appears to be recurrently longer than 30 min, then treatment should be aimed at decreasing sleep latency. This goal can be achieved by, eg, delaying bedtime (ie, increasing sleep pressure), improving preparation, reducing the level of stimulating activities or food (energy) intake, or using relaxation techniques in the hour before going to bed.
Awake after sleep onset ⁸⁰	Time spent awake in bed after sleep onset.	Questioning/sleep diary/actigraphy	How much time do you spend awake in bed after falling asleep?	If a patient spends more than half an hour awake in bed after sleep onset, then this problem should be addressed.	Time spent awake in bed after sleep onset can be reduced by applying bedtime restriction/sleep compression. A patient who lies awake for more than half an hour should be advised to leave the bedroom and do something relaxing until feeling sleepy again. Motivational interviewing should be applied to promote behavioral change.
Early-morning awakening ⁸⁰	Time spent awake prior to planned wake-up time.	Questioning/sleep diary/actigraphy	Do you wake up prior to your planned wake-up time?	Time spent awake in bed prior to getting out of bed should be as short as possible. Ideally, a patient should get out of bed as soon as possible after final arousal.	A patient who stays in bed awake after final arousal should be advised to get out of bed as soon as possible to prevent wake inertia and daytime fatigue. Motivational techniques should be applied to promote behavioral change.
Number of awakenings ⁸⁰	Number of conscious arousals experienced per night.	Questioning/sleep diary/actigraphy	Do you wake up during the night and, if so, how many times?	On average, a patient should not consciously wake up more than 2 times during the allotted sleep period. Waking up once per night (eg, for a bathroom visit), however, may not be a clinically relevant issue.	Various parts of CBT-I (Tab. 2) can decrease the number of arousals. If a patient has recurrent awakenings (>2) and/or awakenings lasting more than half an hour in total, then the strategies used for time spent awake in bed after sleep onset should be used.
Sleep efficiency ⁸¹	(Total sleep time/total time in bed) × 100.	Calculation		Ideally, sleep efficiency should be >90%; >85% is acceptable given adequate sleep duration.	If a patient has reduced sleep efficiency, then bedtime restriction (Tab. 2) may be required. The aim is to find a compromise with respect to a realistic estimation of quantitative sleep needs.
Sleep hygiene	Recommended behavioral practices intended to promote improved sleep quality.	Questioning	Do you monitor time while being awake in bed? Do you consume stimulant beverages prior to bedtime? Do you intensively exercise close to bedtime? What do you eat before going to bed?	Stimulants (eg, caffeine or nicotine) close to bedtime may impair sleep onset and/or maintenance. A small amount of aerobic exercise could improve sleep quality; however, intense workouts close to bedtime should be avoided. Food that has a high caloric load and/or that affects absorption processes in the digestive tract can cause sleep disruption.	Sleep hygiene (Tab. 2) instructions are needed to explain that clock monitoring in bed, food and beverage consumption, and intense exercise before bedtime may disrupt sleep initiation and maintenance. Motivational interviewing should be applied to promote behavioral change.

(Continued)

Table 1.
Continued

Item	Explanation	Mode of Assessment	Sample Questions	Interpretation	Treatment Implications
Napping	Whether a patient sleeps or lies down with the intention to sleep, at what time, how long, and how often.	Questioning/ sleep diary/ actigraphy	Do you sleep during the day or evening and, if so, how long and how often?	In a patient with insomnia, napping generally is not recommended. It may be acceptable if the nap is ≤20 min (ie, “power nap”) and has appropriate timing with respect to circadian rhythms (ie, not later than 15.30 h). Otherwise, a nap can lower sleep pressure or disrupt nighttime sleep. A nap includes falling asleep before bedtime (eg, while watching TV).	Sleep hygiene (Tab. 2) instructions are needed to explain that daytime sleeping may disrupt circadian rhythms or lower sleep pressure at bedtime. A patient should avoid daytime sleeping beyond the duration of a power nap.
Sleep environment	Physical bedroom characteristics, such as darkness, temperature, noise/quietness, and humidity.	Questioning	Is your bedroom sufficiently dark (ie, even after sunrise)? What is the temperature in your bedroom? Is it sufficiently quiet in your bedroom?	Ambient light may interfere with sleep quality and/or continuity. Ideally, the bedroom temperature should be 16°C–19°C, and the bedroom should be isolated from noise disturbances.	Sleep hygiene (Tab. 2) instructions are needed to explain and improve the sleep environment.
Perceived sleep quality ¹¹	Subjective sleep quality.	Questioning/ sleep diary/ PSQI/rating	How well do you sleep?	If a patient perceives sleep quality as satisfying, then further inquiry may not be compulsory. If not, then sleep quality can be used to monitor a therapeutic response.	A lack of improvement in sleep quality over time may indicate that the treatment response to CBT-I is not evolving as intended.
Beliefs and attitudes with respect to the nature/ etiology of the sleep problem	Patient’s erroneous causal attributions with respect to insomnia.	Questioning/ DBAS	What do you think causes your sleep problem?	Assess whether sleep beliefs are adaptive or prevent behavioral change.	If a patient has maladaptive sleep beliefs, then general sleep education (Tab. 2) is indicated.
Beliefs and attitudes regarding the evolution of the sleep problem	Whether a patient expects improvement or not.	Questioning/ DBAS	Do you think that your sleep can improve?	Assess whether beliefs and attitudes regarding the evolution of insomnia are adaptive or prevent behavioral change.	Maladaptive beliefs and attitudes regarding the evolution of the sleep problem should be addressed with general sleep education (Tab. 2).
Identification of (life) stressors ⁸²	Whether a patient has experienced life stressors and is currently struggling to cope with certain (daily) stressors.	Questioning	Are you struggling to deal with daily stressors and, if so, which ones?	Interpret the obtained information to assess whether stress is an issue for a patient. If stress is an issue, then assess whether stress influences sleep.	If stress has a negative impact on a patient’s sleep, then relaxation training/ stress management should be included in the CBT-I treatment (Tab. 2).
Sleep medication	Whether a patient uses any hypnotic or sedative drug (including any synthetic or herbal treatment).	Questioning/ reviewing medical record	Are you using any drug treatments for your sleep? If so, do you consider them useful? Would you like to pursue or discontinue them?	Sole administration of sleep medication is not recommended for treating chronic insomnia but may provide some temporary benefit or an improved therapeutic response (bimodal therapy).	A patient’s perception of sleep medication may interfere with counseling and CBT-I. If a patient wishes to discontinue drug treatment, then systematic withdrawal may be initiated. If not, then treatment may be supplemented by CBT-I and withdrawal may be postponed (ie, when the patient is improving, showing increased self-efficacy in regaining sleep efficiency and sleep quality). In any case, therapists may discuss the treatment plan with the prescriber to prevent contradictory messages.
Substance abuse ⁸²	Past or present substance abuse.	Questioning/ reviewing medical record	Are you currently using or have you been using any substances or drugs (including nicotine)?	Substances may have a negative impact on biological functions of sleep. Past or current substance abuse may interfere with adrenergic, cholinergic, and/or dopaminergic pathways.	Motivational interviewing should be applied to promote behavioral change.

^aCBT-I = cognitive-behavioral therapy for insomnia, DBAS = Dysfunctional Beliefs and Attitudes About Sleep, PSQI = Pittsburgh Sleep Quality Index.

Table 2.

Components of Cognitive-Behavioral Therapy for Insomnia in People With Chronic Pain and Comorbid Insomnia

Component	Description/Content
General sleep education ⁸³	Explaining the importance of sleep and the behavioral neuroscience behind sleep, including the role of melatonin in initiating/sustaining deep sleep, the role of daylight, and the influence of regular sleeping hours.
Sleep (or bedtime) restriction therapy ⁸⁴	Manipulation of homeostatic sleep drive to consolidate sleep via sleep restriction ⁶⁰ : initially limiting the amount of time spent in bed to an amount equal to the average sleep time for 1 wk. ⁸⁵ Once sleep becomes more efficient, total sleep time is incrementally increased on a week-to-week basis, and the sleep window is adjusted in time. ⁸⁵
	In a broader sense, sleep restriction therapy also includes the alteration of circadian regularity and alignment, ⁶⁰ ie, imposing a regular wake-up time in an attempt to synchronize “strong” (fairly unresponsive to light) and “weak” (depending on exposure to bright light) endogenous clocks by modulating behavior.
Stimulus control instructions ⁸⁴	Application of operant and classical conditioning principles via stimulus control instructions, ⁶⁰ including restricting bedroom behaviors to sleep and sex, ⁸⁵ limiting the amount of time spent awake in bed or in the bedroom, ⁸⁵ and promoting counterconditioning by ensuring that the bed and the bedroom environment are tightly coupled with sleepiness and sleep. ⁸⁵
Sleep hygiene instructions ⁸⁶	Replacement of sleep-interfering behaviors with sleep-promoting behaviors through sleep hygiene education and behavior change counseling. ⁶⁰
(Sleep-specific) cognitive therapy ⁸⁴	Modification of maladaptive sleep-related cognition. ⁶⁰
Relaxation training/stress management ⁸³	Teaching patients to cope better with stress, including relaxation skills training (eg, deep breathing ⁶⁰ or guided imagery ⁶⁰).
Patient’s self-monitoring of daily sleeping patterns ⁶⁰	Daily self-monitoring of time in bed, sleep onset latency, time spent awake after sleep onset, and total sleep time. ⁶⁰

sleep pressure, or sleep drive, at sleep onset. If sleep efficiency increases to 85% or more, then the upward titration is initiated by adding 15 minutes to the prescribed time in bed.⁵ While people with insomnia knowingly often have lower levels of sleep pressure (ie, sleep onset) and increased levels of wake drives (ie, maintenance difficulties and hypervigilance),⁷³ interventions involving sleep and bedtime restrictions are aimed at improving sleep homeostasis, which occurs with the expansion of sleep opportunity, not only in response to sleep loss but also as a means of managing pain.⁵

Sleep hygiene should be improved alongside with improved stimulus control in order to reestablish an enhanced and optimal associations between bedroom and sleep, allowing for sleep to solely occur in association with the bedroom and related controlled parameters (eg, complete darkness, stable and sufficiently low room temperature, and lowest levels of sensory stimuli).⁵ Cognitive therapy entails changing negative thoughts about sleep. It includes “decatastrophization” to address the perception of dire consequences of

sleep loss.⁵ It is essential for clinicians to acknowledge that the barriers for achieving adequate sleep can present with substantial interindividual variabilities.¹¹ Such interventions are not limited to a specific discipline but available for any caregiver willing to learn more about the role of sleep and how to improve it. Regarding administration modalities, conservative sleep treatment are effective when delivered face-to-face^{4,5} or through the Internet.^{74,75}

Other interventions not directly targeting sleep may have a positive effect on sleep quality and quantity. For instance, stress management⁷⁶ and exercise therapy⁷⁶⁻⁷⁹ improve sleep in patients with cancer and noncancer pain.

Role of the Funding Source

This work was funded by the Applied Biomedical Research Program, Research Foundation Flanders (Fonds voor Wetenschappelijk Onderzoek Vlaanderen), Belgium (FWO-TBM project no. T001117N entitled “Cognitive behavioural therapy for insomnia in people with chronic spinal pain: a multi-center randomized controlled tri-

al”). The funder played no role in the conduct of this study.

Conclusion

Insomnia among individuals with chronic pain is highly prevalent and closely related to mechanisms of central sensitization. It is characterized by low-grade neuroinflammation, associated with stress/anxiety, and does not respond effectively to pharmacological treatment. CBT-I is effective for improvements of sleep initiation, efficiency and maintenance, perceived sleep quality and pain interference with daily activities in people with chronic pain. In addition, there is relatively little evidence about optimal administration procedures (timing, simultaneous priming, or hierarchy) for the treatment of sleep disorders in contexts of chronic pain. A recent systematic review concluded with high quality evidence that with additional training, physical therapist-led cognitive-behavioral interventions are effective for low back pain, allowing implementation within the field. CBT-I typically includes sleep (or bedtime) restriction recommendations, stimulus control instructions, psychoeducation regarding sleep and sleep hygiene, and

Downloaded from https://academic.oup.com/ptj/article/98/5/325/4841863 by U.S. Department of Justice user on 16 August 2022

cognitive therapy. CBT-I may not be a stand-alone treatment for people with chronic pain and comorbid insomnia, but may preferentially be combined with other effective treatment interventions, such as exercise therapy and CBT for pain.

Author Contributions

Concept/idea/research design: J. Nijs, O. Mairesse, D. Neu, L. Danneels, B. Cagnie, K. Ickmans, D. Goubert

Writing: J. Nijs, O. Mairesse, D. Neu, L. Danneels, B. Cagnie, M. Meeus, M. Moens, K. Ickmans

Project management: J. Nijs, L. Danneels
Providing institutional liaisons: D. Neu, M. Meeus

Consultation (including review of manuscript before submitting): O. Mairesse, D. Neu, L. Leysen, B. Cagnie, M. Meeus, M. Moens, K. Ickmans, D. Goubert

Funding

This work was funded by the Applied Biomedical Research Program, Research Foundation Flanders (Fonds voor Wetenschappelijk Onderzoek Vlaanderen), Belgium (FWO-TBM project no. T001117N, "Cognitive behavioural therapy for insomnia in people with chronic spinal pain: a multi-center randomized controlled trial"). K. Ickmans is a postdoctoral researcher funded by the Applied Biomedical Research Program, Institute for the Agency for Innovation by Science and Technology, Belgium (IWT-TBM project no. 150,180). D. Goubert is a postdoctoral researcher funded by the Applied Biomedical Research Program, Institute for the Agency for Innovation by Science and Technology, Belgium (IWT-TBM project no. IWT-TBM 130,246).

Disclosures

The authors completed the ICMJE Form for Disclosure of Potential Conflicts of Interest. M. Moens reported an independent research grant paid to his institution from Medtronic, Nevro, and money paid to him as a consulting fee or honorarium from Medtronic and Pfizer. D. Goubert reported a grant paid to her institution from the Applied Biomedical Research Program, Research Foundation Flanders (Fonds voor Wetenschappelijk Onderzoek Vlaanderen), Belgium (grant no. 130,246). There are no other conflict-of-interest disclosures from the authors.

J. Nijs is holder of the Chair "Exercise immunology and chronic fatigue in health and disease," funded by the Berekuy Academy/European College for Decongestive Lymphatic Therapy, the Netherlands. M. Moens

received the Lyrica Independent Investigator Research Award (LIIRA).

DOI: 10.1093/ptj/pzy020

References

- Henschke N, Ostelo RW, van Tulder MW, et al. Behavioural treatment for chronic low-back pain. *Cochrane Database Syst Rev*. 2010;(7):CD002014.
- Williams AC, Eccleston C, Morley S. Psychological therapies for the management of chronic pain (excluding headache) in adults. *Cochrane Database Syst Rev*. 2012;(11):CD007407.
- Cheung CW, Qiu Q, Choi SW, Moore B, Goucke R, Irwin M. Chronic opioid therapy for chronic non-cancer pain: a review and comparison of treatment guidelines. *Pain Physician*. 2014;17:401-414.
- Tang NK, Lereya ST, Boulton H, Miller MA, Wolke D, Cappuccio FP. Nonpharmacological treatments of insomnia for long-term painful conditions: a systematic review and meta-analysis of patient-reported outcomes in randomized controlled trials. *Sleep*. 2015;38:1751-1764.
- Jungquist CR, O'Brien C, Matteson-Rusby S, et al. The efficacy of cognitive-behavioral therapy for insomnia in patients with chronic pain. *Sleep Med*. 2010;11:302-309.
- Pigeon WR, Moynihan J, Matteson-Rusby S, et al. Comparative effectiveness of CBT interventions for co-morbid chronic pain & insomnia: a pilot study. *Behav Res Ther*. 2012;50:685-689.
- American Academy of Sleep Medicine. *International Classification of Sleep Disorders (ICSD)*. 3rd ed. Darien, IL: American Academy of Sleep Medicine; 2014.
- Becker N, Bondegaard Thomsen A, Olsen AK, Sjogren P, Bech P, Eriksen J. Pain epidemiology and health related quality of life in chronic non-malignant pain patients referred to a Danish multidisciplinary pain center. *Pain*. 1997;73:393-400.
- McCracken LM, Iverson GL. Disrupted sleep patterns and daily functioning in patients with chronic pain. *Pain Res Manag*. 2002;7:75-79.
- Tang NK, Wright KJ, Salkovskis PM. Prevalence and correlates of clinical insomnia co-occurring with chronic back pain. *J Sleep Res*. 2007;16:85-95.
- Daly-Eichenhardt A, Scott W, Howard-Jones M, Nicolaou T, McCracken LM. Changes in sleep problems and psychological flexibility following interdisciplinary acceptance and commitment therapy for chronic pain: an observational cohort study. *Front Psychol*. 2016;7:1326.
- Meeus M, Van Oosterwijck J, Ickmans K, et al. Interrelationships between pain processing, cortisol and cognitive performance in chronic whiplash-associated disorders. *Clin Rheumatol*. 2015;34:545-553.
- Finan PH, Goodin BR, Smith MT. The association of sleep and pain: an update and a path forward. *J Pain*. 2013;14:1539-1552.
- Tang NK, Sanborn AN. Better quality sleep promotes daytime physical activity in patients with chronic pain? A multilevel analysis of the within-person relationship. *PLoS One*. 2014;9:e92158.
- Holbrook AM, Crowther R, Lotter A, Cheng C, King D. Meta-analysis of benzodiazepine use in the treatment of insomnia. *CMAJ*. 2000;162:225-233.
- Johnson LC, Chernik DA. Sedative-hypnotics and human performance. *Psychopharmacology*. 1982;76:101-113.
- Alviar MJ, Hale T, Dungca M. Pharmacologic interventions for treating phantom limb pain. *Cochrane Database Syst Rev*. 2016;(10):CD006380.
- Walitt B, Klose P, Fitzcharles MA, Phillips T, Hauser W. Cannabinoids for fibromyalgia. *Cochrane Database Syst Rev*. 2016;(7):CD011694.
- Walitt B, Urrutia G, Nishishinya MB, Cantrell SE, Hauser W. Selective serotonin reuptake inhibitors for fibromyalgia syndrome. *Cochrane Database Syst Rev*. 2015;(6):CD011735.
- Saragiotto BT, Machado GC, Ferreira ML, Pinheiro MB, Abdel Shaheed C, Maher CG. Paracetamol for low back pain. *Cochrane Database Syst Rev*. 2016;(6):CD012230.
- Doufas AG, Panagiotou OA, Ioannidis JP. Concordance of sleep and pain outcomes of diverse interventions: an umbrella review. *PLoS One*. 2012;7:e40891.
- Breivik H, Collett B, Ventafridda V, Cohen R, Gallacher D. Survey of chronic pain in Europe: prevalence, impact on daily life, and treatment. *Eur J Pain*. 2006;10:287-333.
- International Association for the Study of Pain. Part III: pain terms: a current list with definitions and notes on usage. In: *Classification of Chronic Pain*. 2nd ed. Seattle, Washington: IASP Press; 1994:209-214.
- Woolf CJ. Central sensitization: implications for the diagnosis and treatment of pain. *Pain*. 2011;152(3 suppl):S2-S15.
- Meyer RA, Campbell IT, Raja SN. Peripheral neural mechanisms of nociception. In: Wall PD, Melzack R, eds. *Textbook of Pain*. 3rd ed. Edinburgh, Scotland: Churchill Livingstone; 1995:13-44.
- Nijs J, Van Houdenhove B, Oostendorp RA. Recognition of central sensitization in patients with musculoskeletal pain: application of pain neurophysiology in manual therapy practice. *Man Ther*. 2010;15:135-141.
- Nijs J, Malfliet A, Ickmans K, Baert I, Meeus M. Treatment of central sensitization in patients with 'unexplained' chronic pain: an update. *Expert Opin Pharmacother*. 2014;15:1671-1683.
- Woolf CJ. What to call the amplification of nociceptive signals in the central nervous system that contribute to widespread pain? *Pain*. 2014;155:1911-1912.

- 29 Onen SH, Alloui A, Gross A, Eschallier A, Dubray C. The effects of total sleep deprivation, selective sleep interruption and sleep recovery on pain tolerance thresholds in healthy subjects. *J Sleep Res.* 2001;10:35–42.
- 30 Schuh-Hofer S, Wodarski R, Pfau DB, et al. One night of total sleep deprivation promotes a state of generalized hyperalgesia: a surrogate pain model to study the relationship of insomnia and pain. *Pain.* 2013;154:1613–1621.
- 31 Smith MT, Edwards RR, McCann UD, Haythornthwaite JA. The effects of sleep deprivation on pain inhibition and spontaneous pain in women. *Sleep.* 2007;30:494–505.
- 32 Campbell CM, Buenaver LF, Finan P, et al. Sleep, pain catastrophizing, and central sensitization in knee osteoarthritis patients with and without insomnia. *Arthritis Care Res.* 2015;67:1387–1396.
- 33 Curatolo M, Muller M, Ashraf A, et al. Pain hypersensitivity and spinal nociceptive hypersensitivity in chronic pain: prevalence and associated factors. *Pain.* 2015;156:2373–2382.
- 34 Smith MT, Wickwire EM, Grace EG, et al. Sleep disorders and their association with laboratory pain sensitivity in temporomandibular joint disorder. *Sleep.* 2009;32:779–790.
- 35 Finan PH, Smith MT. The comorbidity of insomnia, chronic pain, and depression: dopamine as a putative mechanism. *Sleep Med Rev.* 2013;17:173–183.
- 36 Millan MJ. Descending control of pain. *Prog Neurobiol.* 2002;66:355–474.
- 37 Whitney MS, Shemery AM, Yaw AM, Donovan LJ, Glass JD, Deneris ES. Adult brain serotonin deficiency causes hyperactivity, circadian disruption, and elimination of siestas. *J Neurosci.* 2016;36:9828–9842.
- 38 Wei H, Pertovaara A. 5-HT_{1A} receptors in endogenous regulation of neuropathic hypersensitivity in the rat. *Eur J Pharmacol.* 2006;535:157–165.
- 39 Lopez-Canul M, Palazzo E, Dominguez-Lopez S, et al. Selective melatonin MT₂ receptor ligands relieve neuropathic pain through modulation of brainstem descending antinociceptive pathways. *Pain.* 2015;156:305–317.
- 40 Lopez-Canul M, Comai S, Dominguez-Lopez S, Granados-Soto V, Gobbi G. Antinociceptive properties of selective MT₂ melatonin receptor partial agonists. *Eur J Pharmacol.* 2015;764:424–432.
- 41 Greco MA, Fuller PM, Jhou TC, et al. Opioidergic projections to sleep-active neurons in the ventrolateral preoptic nucleus. *Brain Res.* 2008;1245:96–107.
- 42 Wang Q, Yue XF, Qu WM, et al. Morphine inhibits sleep-promoting neurons in the ventrolateral preoptic area via mu receptors and induces wakefulness in rats. *Neuropsychopharmacology.* 2013;38:791–801.
- 43 Mullington JM, Simpson NS, Meier-Ewert HK, Haack M. Sleep loss and inflammation. *Best Pract Res Clin Endocrinol Metab.* 2010;24:775–784.
- 44 Haack M, Lee E, Cohen DA, Mullington JM. Activation of the prostaglandin system in response to sleep loss in healthy humans: potential mediator of increased spontaneous pain. *Pain.* 2009;145:136–141.
- 45 Haack M, Sanchez E, Mullington JM. Elevated inflammatory markers in response to prolonged sleep restriction are associated with increased pain experience in healthy volunteers. *Sleep.* 2007;30:1145–1152.
- 46 Wisor JP, Schmidt MA, Clegern WC. Cerebral microglia mediate sleep/wake and neuroinflammatory effects of methamphetamine. *Brain Behav Immun.* 2011;25:767–776.
- 47 Pollmacher T, Haack M, Schuld A, Reichenberg A, Yirmiya R. Low levels of circulating inflammatory cytokines: do they affect human brain functions? *Brain Behav Immun.* 2002;16:525–532.
- 48 Neu D, Mairesse O, Montana X, et al. Dimensions of pure chronic fatigue: psychophysical, cognitive and biological correlates in the chronic fatigue syndrome. *Eur J Appl Physiol.* 2014;114:1841–1851.
- 49 Wodarski R, Schuh-Hofer S, Yurek DA, et al. Development and pharmacological characterization of a model of sleep disruption-induced hypersensitivity in the rat. *Eur J Pain.* 2015;19:554–566.
- 50 Nijs J, Loggia ML, Polli A, et al. Sleep disturbances and severe stress as glial activators: key targets for treating central sensitization in chronic pain patients? *Expert Opin Ther Targets.* 2017;21:817–826.
- 51 Denis D, Akhtar R, Holding BC, et al. Externalizing behaviors and callous-unemotional traits: different associations with sleep quality. *Sleep.* 2017;40:zxx070.
- 52 Nakamura M, Nagamine T. Neuroendocrine, autonomic, and metabolic responses to an orexin antagonist, suvorexant, in psychiatric patients with insomnia. *Innov Clin Neurosci.* 2017;14:30–37.
- 53 Kim EJ, Dimsdale JE. The effect of psychosocial stress on sleep: a review of polysomnographic evidence. *Behav Sleep Med.* 2007;5:256–278.
- 54 Ojeda B, Salazar A, Calahorra MJ, et al. Understanding the different relationships between mood and sleep disorders in several groups of non-oncological patients with chronic pain. *Curr Med Res Opin.* 2018;34:669–676.
- 55 Cigarán-Méndez M, Fernández-Muñoz JJ, Navarro-Pardo E, et al. Gender differences in variables associated with sleep quality in chronic tension type headache. *Women Health.* 2017 September 18 [Epub ahead of print]. doi: 10.1080/03630242.2017.1372845.
- 56 Morin CM, Hauri PJ, Espie CA, Spielman AJ, Buysse DJ, Bootzin RR. Nonpharmacologic treatment of chronic insomnia: an American Academy of Sleep Medicine review. *Sleep.* 1999;22:1134–1156.
- 57 Currie SR, Wilson KG, Curran D. Clinical significance and predictors of treatment response to cognitive-behavior therapy for insomnia secondary to chronic pain. *J Behav Med.* 2002;25:135–153.
- 58 Bastien CH, Vallieres A, Morin CM. Validation of the Insomnia Severity Index as an outcome measure for insomnia research. *Sleep Med.* 2001;2:297–307.
- 59 Morin CM, Belleville G, Belanger L, Ivers H. The Insomnia Severity Index: psychometric indicators to detect insomnia cases and evaluate treatment response. *Sleep.* 2011;34:601–608.
- 60 Finan PH, Buenaver LF, Coryell VT, Smith MT. Cognitive-behavioral therapy for comorbid insomnia and chronic pain. *Sleep Med Clin.* 2014;9:261–274.
- 61 Johns MW. A new method for measuring daytime sleepiness: the Epworth sleepiness scale. *Sleep.* 1991;14:540–545.
- 62 Kendzerska TB, Smith PM, Brignardello-Petersen R, Leung RS, Tomlinson GA. Evaluation of the measurement properties of the Epworth sleepiness scale: a systematic review. *Sleep Med Rev.* 2014;18:321–331.
- 63 Alda M, Luciano JV, Andrés E, et al. Effectiveness of cognitive behaviour therapy for the treatment of catastrophisation in patients with fibromyalgia: a randomised controlled trial. *Arthritis Res Ther.* 2011;13:R173.
- 64 Ehde DM, Dillworth TM, Turner JA. Cognitive-behavioral therapy for individuals with chronic pain: efficacy, innovations, and directions for research. *Am Psychol.* 2014;69:153–166.
- 65 van Dessel N, den Boeft M, van der Wouden JC, et al. Non-pharmacological interventions for somatoform disorders and medically unexplained physical symptoms (MUPS) in adults. *Cochrane Database Syst Rev.* 2014;(11):CD011142.
- 66 Hall A, Richmond H, Copsey B, et al. Physiotherapist-delivered cognitive-behavioural interventions are effective for low back pain, but can they be replicated in clinical practice? A systematic review. *Disabil Rehabil.* 2018;40:1–9.
- 67 Main CJ, George SZ. Psychologically informed practice for management of low back pain: future directions in practice and research. *Phys Ther.* 2011;91:820–824.
- 68 Currie SR, Wilson KG, Pontefract AJ, deLaplante L. Cognitive-behavioral treatment of insomnia secondary to chronic pain. *J Consult Clin Psychol.* 2000;68:407–416.
- 69 Smith MT, Finan PH, Buenaver LF, et al. Cognitive-behavioral therapy for insomnia in knee osteoarthritis: a randomized, double-blind, active placebo-controlled clinical trial. *Arthritis Rheum.* 2015;67:1221–1233.
- 70 Tang NK, Goodchild CE, Salkovskis PM. Hybrid cognitive-behaviour therapy for individuals with insomnia and chronic pain: a pilot randomised controlled trial. *Behav Res Ther.* 2012;50:814–821.

- 71 Ritterband LM, Bailey ET, Thorndike FP, Lord HR, Farrell-Carnahan L, Baum LD. Initial evaluation of an Internet intervention to improve the sleep of cancer survivors with insomnia. *Psychooncology*. 2012;21:695–705.
- 72 Perlis ML, Jungquist C, Smith MT, Posner D. *Cognitive Behavioral Treatment of Insomnia: A Session-by-Session Guide*. New York, NY: Springer-Verlag; 2005.
- 73 Neu D, Mairesse O, Verbanck P, Le Bon O. Slow wave sleep in the chronically fatigued: power spectra distribution patterns in chronic fatigue syndrome and primary insomnia. *Clin Neurophysiol*. 2126:1926–1933.
- 74 Cheng SK, Dizon J. Computerised cognitive behavioural therapy for insomnia: a systematic review and meta-analysis. *Psychother Psychosom*. 2012;81:206–216.
- 75 Seyffert M, Lagisetty P, Landgraf J, et al. Internet-delivered cognitive behavioral therapy to treat insomnia: a systematic review and meta-analysis. *PLoS One*. 2016;11:e0149139.
- 76 Davis MP, Goforth HW. Long-term and short-term effects of insomnia in cancer and effective interventions. *Cancer J*. 2014;20:330–344.
- 77 Mishra SI, Scherer RW, Geigle PM, et al. Exercise interventions on health-related quality of life for cancer survivors. *Cochrane Database Syst Rev*. 2012;(8):CD007566.
- 78 Buman MP, Hekler EB, Bliwise DL, King AC. Moderators and mediators of exercise-induced objective sleep improvements in midlife and older adults with sleep complaints. *Health Psychol*. 2011;30:579–587.
- 79 Driver HS, Taylor SR. Exercise and sleep. *Sleep Med Rev*. 2000;4:387–402.