

Does Using a Chair Backrest or Reducing Seated Hip Flexion Influence Trunk Muscle Activity and Discomfort? A Systematic Review

Máire Curran, Leonard O'Sullivan, University of Limerick, Limerick, Ireland, Peter O'Sullivan, Curtin University, Perth, Australia, Wim Dankaerts, University of Leuven, Leuven, Belgium, and Kieran O'Sullivan, University of Limerick, Limerick, Ireland

Objective: This paper systematically reviews the effect of chair backrests and reducing seated hip flexion on low back discomfort (LBD) and trunk muscle activation.

Background: Prolonged sitting commonly exacerbates low back pain (LBP). Several modifications to seated posture and chair design have been recommended, including using chairs with backrests and chairs that reduce hip flexion.

Method: Electronic databases were searched by two independent assessors. Part 1 of this review includes 26 studies comparing the effect of sitting with at least two different hip angles. In Part 2, seven studies that compared the effect of sitting with and without a backrest were eligible. Study quality was assessed using the PEDro scale.

Results: Significant confounding variables and a relatively small number of randomized controlled trials (RCTs) involving people with LBP complicates analysis of the results. There was moderate evidence that chair backrests reduce paraspinal muscle activation, and limited evidence that chair backrests reduce LBD. There was no evidence that chairs involving less hip flexion reduce LBP or LBD, or consistently alter trunk muscle activation. However, participants in several studies subjectively preferred the modified chairs involving less hip flexion.

Conclusion: The limited evidence to support the use of chairs involving less seated hip flexion, or the effect of a backrest, is consistent with the limited evidence that other isolated chair design features can reduce LBP.

Application: LBP management is likely to require consideration of several factors in addition to sitting position. Larger RCTs involving people with LBP are required.

Keywords: sitting, back pain, discomfort, systematic review

INTRODUCTION

Low back pain (LBP) is one of the most common musculoskeletal disorders (e.g., Woolf & Pfleger, 2003). Prolonged sitting by itself is not linked to the onset of LBP (Roffey, Wai, Bishop, Kwon, & Dagenais, 2010). However, sitting is commonly reported to increase the symptoms of people with LBP (e.g., Womersley & May, 2006). Consequently, there has been an increasing amount of research investigating sitting posture (e.g., Dankaerts, O'Sullivan, Burnett, & Straker, 2006), and the effect of seating modifications (e.g., Lengsfeld, Konig, Schmelter, & Ziegler, 2007), among people with LBP. Some studies have examined these factors among people with LBP (e.g., O'Sullivan et al., 2006), whereas other studies have examined the onset of low back discomfort (LBD) among pain-free subjects (e.g., Gadge & Innes, 2007).

Using lumbar supports (Williams, Hawley, McKenzie, & Van Wijmen, 1991) or devices (Reinecke, Hazard, & Coleman, 1994) to increase lumbar lordosis have been advocated in the management of LBP for many years. This reflects the fact that both health care professionals (O'Sullivan, O'Sullivan, O'Sullivan, & Dankaerts, 2012) and ergonomic recommendations (Corlett, 2006) favor lordotic sitting postures for the lumbar spine. However, although increasing seated lordosis may reduce LBP in the short term for some individuals (Williams et al., 1991), many patients with LBP already assume lordotic postures that are associated with their symptoms (Dankaerts et al., 2006, O'Sullivan, 2006). Furthermore, lordotic sitting postures are associated with high levels of trunk muscle activation (O'Sullivan, 2006), which may lead to compressive loading, fatigue, and discomfort (Carcone & Keir, 2007; Lander, Korbon, DeGood, & Rowlingson, 1987).

Address correspondence to Kieran O'Sullivan, Department of Clinical Therapies, University of Limerick, Ireland; kieran.osullivan@ul.ie.

HUMAN FACTORS

Vol. XX, No. X, Month XXXX, pp. 1–34

DOI: 10.1177/0018720815591905

Copyright © 2015, Human Factors and Ergonomics Society.

Sitting involves more lumbo-pelvic flexion than standing (Claus, Hides, Moseley, & Hodges, 2009; De Carvalho, Soave, Ross, & Callaghan, 2010; Dunk, Kedgley, Jenkyn, & Callaghan, 2009), due to the greater hip flexion causing posterior pelvic tilt. Sitting postures that maintain lumbar lordosis in sitting are commonly considered advantageous among health care professionals (O'Sullivan, O'Sullivan, et al., 2012; Pynt, Higgs, & Mackay, 2001) and the public (O'Sullivan et al., 2013). Consequently, attempting to reduce seated hip flexion, with a view to maintaining lumbar lordosis, has been proposed (Mandal, 1983). It has been demonstrated that reducing seated hip flexion helps maintain seated lumbar lordosis (Keegan, 1953; Saarni, Nygård, Rimpelä, Nummi, & Kaukainen, 2007) and that people may prefer to sit with less hip flexion than that usually recommended (Mandal, 1987). However, the effect of reducing hip flexion on trunk muscle activation is unclear with both reduced (e.g., Koskelo, Vuorikari, & Hänninen, 2007) and increased (e.g., Lander et al., 1987) muscle activation reported. In addition, it is unclear how such changes in muscle activation translate into clinically meaningful improvements in LBP. For example, schoolchildren have been shown to prefer chairs involving less hip flexion without such chairs necessarily decreasing pain prevalence or symptoms (Cardon, De Clercq, De Bourdeaudhuij, & Breithecker, 2004; Troussier, 1999).

Another commonly advocated chair modification is the use of a backrest (Carcone & Keir, 2007). Chair backrests have been proposed to promote good spinal posture, while also reducing trunk muscle activation and LBD (Anderson, Jonsson, & Ortengren, 1974; Carcone & Keir, 2007). In addition, using a backrest, especially increasing backrest inclination, has been proposed to reduce intradiscal pressure (Anderson, Ortengren, Nachemson, Elfström, & Broman, 1975). As a result, backrests have become standard in many workplaces. However, it appears that backrests are not always used optimally during common occupational tasks (Vergara & Page, 2000b), and a recent review cast significant doubt on the strength of the relationship between sitting and parameters such as

intradiscal pressure (Claus, Hides, Moseley, & Hodges, 2008). Furthermore, although much research has examined the effect of different types of backrest configurations (Ellegast et al., 2012; Groenesteijn et al., 2012), there has been no systematic review on whether using chair backrests actually reduces LBP/LBD, or how they influence trunk muscle activation.

Recent systematic reviews examining factors such as occupational seating have reported mixed results. A systematic review (Driessen et al., 2010) of randomized controlled trials (RCTs) concluded that there was little evidence to support the use of physical ergonomics interventions, including changes in seat design, for LBP or neck pain. In contrast to this, a more recent systematic review (Van Niekerk, Louw, & Hillier, 2012) of seated occupational interventions, which included more varied study designs and examined a broader range of body regions, suggested that there was some evidence of effectiveness. However, many of the included studies in this review (Van Niekerk et al., 2012) were at high risk of bias. Therefore, this systematic review specifically examined the effect of reducing seated hip flexion (Part 1) or providing a chair backrest (Part 2) on trunk muscle activation and/or LBP/LBD.

METHOD

Overview

The Cochrane and MEDLINE databases were initially searched, revealing no systematic reviews investigating the effect of using a chair backrest or hip angle on trunk muscle activation and/or LBP/LBD. These reviews were registered on the PROSPERO database (registration numbers CRD42012002343 and CRD42012002378; PROSPERO, 2012), and have been reported in accordance with the PRISMA statement (Moher, Liberati, Tetzlaff, & Altman, 2009).

Search Strategy and Inclusion Criteria

Two assessors (KO, MC) independently searched for the presence of an agreed range of keywords in the following databases; MEDLINE, SPORTDiscus, CINAHL, AMED, Academic Search Complete, and Biomedical




Figure 1. Four keyword groups used in search strategy examining chairs which reduce seated hip flexion.

Reference Collection. The search strategy used four keyword groups relating to (a) sitting, (b) the low back region, (c) muscle activation or discomfort, and (d) either changing hip angle in sitting (Figure 1) or backrests (Figure 2). The specific keywords for each group are detailed in Figures 1 and 2. The abstract had to contain at least one keyword from each group to be considered for these reviews. The four groups of keywords were combined using “AND.” Articles were limited to those involving humans and published in English, with no year limits applied. Only peer-reviewed journal articles were considered. Conference proceedings were excluded because they are not consistently peer reviewed, and often lack sufficient information to adequately assess methodological quality. After removing duplicates, the titles and abstracts that met these criteria were screened for suitability. If it was unclear whether a study was eligible, full-text articles were retrieved. Studies were included if they compared at least two sitting conditions with different seated hip flexion angles for Part 1, or if they compared sitting with a backrest to at least one other sitting

condition without a backrest for Part 2. In both reviews, included studies must have measured either LBP, LBD, or the activation of at least one trunk muscle. Studies had to either report quantitative values for muscle activation, LBD, or LBP, or perform statistical comparisons between the sitting conditions to be eligible for inclusion. Studies involving either pain-free participants or people with LBP were eligible. No minimum follow-up period was required, such that single-session comparisons of sitting conditions were eligible. The reference list of each article was also screened for further relevant articles.

Assessment of Methodological Quality

Methodological quality was rated independently by two assessors (KO, MC) using the Physiotherapy Evidence Database (PEDro) scale (<http://www.pedro.org.au>). The PEDro scale is a reliable (Maher, Sherrington, Herbert, Moseley, & Elkins, 2003) and valid (de Morton, 2009) method of assessing trial quality, based on the Delphi criteria (Verhagen et al., 1998). This scale investigates the inter-




Figure 2. Four keyword groups used in search strategy examining chairs using backrests.

nal validity of a study and contains 11 criteria, 10 of which are scored. When it was unclear whether a study met the PEDro criteria, the original authors were emailed for clarity, to enhance the accuracy and rigor of the reviews. Regarding the “baseline comparability” criterion, studies must typically describe a measure of disorder severity. Although this cannot be applied among pain-free participants, points for this criterion were automatically awarded for crossover design studies, despite there being no measure of “severity” available, in line with a recent systematic review (O’Sullivan, O’Keeffe, O’Sullivan, O’Sullivan, & Dankaerts, 2012). Regarding the “point estimates and variability” criterion, points were awarded if such data were provided for either muscle activation and/or LBP/LBD. In the event of disagreement between the two raters for an individual study, a consensus decision was reached. The quality was classified as “high” ($\geq 6/10$), “fair” (4–5/10), or “poor” ($< 4/10$), according to PEDro scores (Ye, Kalichman, Spittle, Dobson, &

Bennell, 2011), to aid interpretation of study quality in the event of inconsistent findings. An a priori decision was made to exclude studies rated as “poor,” in line with a recent systematic review (O’Sullivan, O’Keeffe, et al., 2012). In addition, the overall quality of the studies was evaluated under the headings of bias, confounding factors, strength of the results, and clinical applicability, similar to a recent systematic review (O’Sullivan, O’Keeffe, et al., 2012).

Data Extraction and Synthesis

Data regarding each study were extracted and cross-checked by two assessors (KO, MC). This included data on (a) sample size, (b) participant sex, (c) participant age, (d) sitting conditions, (e) study protocol, (f) muscles analyzed, (g) pain or discomfort measurement, (h) inclusion/exclusion criteria, and (i) the main results (Table 1). Significant differences in the outcome measures used for both muscle activation and LBP/LBD, as well as the comparison sitting conditions used, did not allow for pooled analysis of the data.

RESULTS

Study 1: Effect of Reducing Seated Hip Flexion

Identification of studies. The results of the search strategy are outlined in Figure 3. The electronic search returned 1,080 potentially relevant studies, which was reduced to 849 after removing duplicates. After screening these titles and abstracts, 19 potentially relevant studies were identified. After reviewing the full-text of these studies, 14 studies met the inclusion and exclusion criteria. Searching the reference lists of these articles added another 15 studies. Three studies were rated as “poor” quality on the PEDro scale and were excluded. Therefore, the final number of articles included in this review was 26 (Table 1). Five studies were clinical trials with a follow-up period. The remaining 21 studies were crossover studies, or used similar designs such as a single system repeated baseline study (Gadge & Innes, 2007) and studies comparing “old” and “new” seating designs (Aagaard-Hansen & Storr-Paulsen, 1995). In all, 21 studies examined LBP or LBD, 11 examined trunk muscle activation, with 6 studies examining both (Table 1). Only 7 studies included participants with LBP.

Crossover studies. Eight studies compared forward inclined kneeler chairs to at least one office chair with a flat seat pan (hip angle approximately 90°). Six of these eight studies used the same brand of armless kneeler chair (Balans kneeler chair), whereas three of them compared sitting to a flat seat pan that also had a backrest. Lander et al. (1987) reported that cervical erector spinae (CES) and lumbar erector spinae (LES) activation were significantly higher on the Balans kneeler chair than on an armless seat with a flat seat pan and backrest. In addition, the chair with the flat seat pan and backrest was reported as more comfortable on a 0 to 10 visual analogue scale (VAS). Bennett, Gillis, Portney, Romanow, and Sanchez (1989) reported no significant differences in LES activation while sitting on two armless seats with a flat seat pan and a backrest with the Balans kneeler chair. Cram and Vinitzky (1995) compared paraspinal activation (CES, thoracic erector spinae [TES], LES) on three different chairs: the Balans kneeler chair, a chair with a flat

seat pan and backrest, and using the Back Up, which is a chair accessory providing pelvic support. LES activation was again significantly higher on the Balans chair than on the flat seat pan with a backrest. However, this study did not specify if the chairs featured an armrest or whether participants were able to lean on the desk while writing for 10 minutes, which could confound results. Two other studies analyzed only discomfort and not muscle activation. Bendix, Jensen, and Bendix (1988) reported no statistically significant differences between sitting on a Balans chair and an armless tiltable chair. Similarly, this study did not state if participants leaned on the desk during the tasks. Bishu, Hallbeck, Riley, and Stentz (1991) reported that the Balans kneeler chair was less comfortable (measured by a discomfort questionnaire rated 0–5) than two other conventional sitting conditions with a flat seat pan. Although one of the conventional chairs in this study had armrests, it was not stated if participants were instructed to use them. Finally, and in contrast to the other five studies examining the same kneeler chair, Soderberg, Blanco, Cosentino, and Kurdelmeier (1986) reported that a kneeler chair at two different forward inclinations was preferred to a conventional chair with a flat seat pan. Furthermore, paraspinal activation (CES, TES, LES) was significantly lower on the kneeler chair. The conventional chair had armrests but participants could not use them and rested their wrists lightly on the computer keyboard to assist standardization. The other two studies of kneeler chairs used different brands, and combined the kneeler chair with sloping desks. Marschall, Harrington, and Steele (1995) compared a traditional workstation with a kneeler chair combined with sloping desks among schoolchildren. Neither chair featured armrests. Latissimus dorsi (LD) activity was significantly decreased on the kneeler chair and participants preferred this workstation. Finally, Bridger (1988) reported greater comfort when sitting for 20 minutes using a kneeler chair compared to a conventional office chair, especially when the kneeler chair was combined with a sloping desk. This study did not state whether chairs featured armrests which could confound results.

Thirteen studies examined the effect of varying seat pan inclination, or seat height, or both on

(text continues on p. 22)

TABLE 1: Description of Studies Included in the Review

Study	Year	Sample Size	Sex	Age (Years)	Sitting Conditions	Study Protocol	Muscles Measured	Discomfort Measurement	Inclusion/Exclusion Criteria	Results
Reducing hip flexion studies										
Aagaard-Hansen and Storr-Paulsen	1995	156	Mixed M/F (exact numbers NS)	7–11 years (range)	Conventional chair with flat seat pan and flat desk Higher, forward inclined (unspecified degree) chair with sloping desk Conventional chair with flat seat pan and sloping desk	Students sat on usual chair and desk at baseline. Then allocated in 3 clusters to one of the 3 study seats for 2–3 months.	N/A	Compared frequency of back pain between the children in the 3 different groups.	Children in forms 2, 3, and 5, both with and without LBP	No significant difference in frequency of back pain between groups, despite combination of higher, forward inclined chair with sloping desk receiving more positive overall ratings.
Bendix and Biering-Sørensen	1983	10	4M/6F	31	Horizontal seat Forward inclined seat pan (5°) Forward inclined seat pan (10°) Forward inclined seat pan (15°) The height of the seat increased in line with increasing forward inclination	Subjects sat on each seat for 1 hour.	N/A	Comfort scale from 1 to 5.	Healthy subjects No back trouble or back abnormalities All familiarized with tiltable chair for at least 2 weeks	No significant difference. Subjects showed a nonsignificant preference for the 0° and 5° forward inclinations over the 10° and 15° forward inclinations.

(continued)

TABLE 1: (continued)

Study	Year	Sample Size	Sex	Age (Years)	Sitting Conditions	Study Protocol	Muscles Measured	Discomfort Measurement	Inclusion/Exclusion Criteria	Results
Bendix	1984	10	2M/8F	28 (median) 20-39 (range)	Low seat, with backward inclined (-5°) seat pan High seat, with forward inclined (5°) seat pan Self-selected seat height, with freely tiltable (-10° to 30°) seat pan These 3 seating options were used with 4 different table heights	Subjects sat reading at each arrangement for 15 minutes on 3 different days.	N/A	Subjective acceptability on a scale from 1 to 5.	Healthy subjects All familiarized with tiltable chair for at least 2 weeks	No significant difference. Subjects showed a nonsignificant preference for the tiltable seat over the fixed forward and backward inclined seats.
Bendix et al.	1986	10	4M/6F	30 (median) 23-42 (range)	Low seat, with backward inclined (-5°) seat pan High seat, with forward inclined (5°) seat pan These 2 seating options were used with 3 different backrest positions and during 2 different seated tasks (deskwork and typing)	Subjects sat doing seated tasks (deskwork or typing) on each seat for 15 minutes each on 4 different days.	N/A	Subjective acceptability on a scale from 1 to 5.	Healthy subjects All familiar with using a typewriter, but not employed as typist All familiarized with tiltable chair for at least 2 weeks	Significantly greater acceptability of higher seat.

(continued)

TABLE 1: (continued)

Study	Year	Sample Size	Sex	Age (Years)	Sitting Conditions	Study Protocol	Muscles Measured	Discomfort Measurement	Inclusion/Exclusion Criteria	Results
Bendix et al.	1985	Total 24: Field study (n = 12) Laboratory study (n = 12)	0M/12F 0M/12F	31 (median) 22–47 (range) 34 (median) 22–41 (range)	Low seat, with backward inclined (–5°) seat pan High seat, with forward inclined (10°) seat pan High seat, with freely tiltable (–8° to 19.5°) seat pan	For the field study, subjects spent one full working day on each sitting condition. For the laboratory study, subjects sat for 1 hour on each sitting condition.	For the laboratory study only, LES (L3) was measured.	Subjective acceptability on a scale from 1 to 5.	Healthy subjects Most were professionals secretaries All familiarized with each chair setting for 2 weeks	No significant difference in muscular activity for any of the seat adjustments. There was a statistically significant preference for the tiltable chair in the laboratory study, but this trend did not reach statistical significance in the longer field study.
Bendix et al.	1988	12	6M/6F	28.2 23–39 (range)	Kneeler chair (Balans) with average forward inclination of 28.5° Tiltable seat, with average forward inclination of 11.5°	Subjects sat for 1 hour on each chair while performing office work and an assembly task	N/A	Subjective acceptability on a scale from 1 to 5.	Healthy subjects All familiarized with both chairs for at least 3 weeks	No statistically significant difference for acceptability of either chair. Nonsignificant trend for reduced acceptability and increased fatigue on the kneeler chair.

(continued)

TABLE 1: (continued)

Study	Year	Sample Size	Sex	Age (Years)	Sitting Conditions	Study Protocol	Muscles Measured	Discomfort Measurement	Inclusion/Exclusion Criteria	Results
Bishu et al.	1991	6	3M/3F	18-40 (range)	Conference room chair with flat seat pan Plastic classroom chair with flat seat pan Kneeler chair (Balans; unspecified degree)	Subjects sat in each chair for 30 minutes while discomfort ratings were measured at intervals.	N/A	Questionnaire assessing discomfort from 0-5.	Pain-free subjects considered to have normal posture	Significantly more back discomfort was reported for the forward inclined kneeler chair.
Bridger	1988	18	11M/7F	26.4	Office chairs with flat seat pan and backrest Kneeler chair with forward inclination of 25° and no backrest Both chairs were tested with both a flat table and a sloping (15°) desk	Subjects performed handwriting task for 20 minutes on both chairs used with both a flat table and a sloping (15°) table.	N/A	Comfort scale (rated 1-7).	Pain-free subjects with a sedentary occupation which involved writing No history of musculoskeletal disorders Not obese	Comfort was significantly greater when using kneeler chair, especially when combined with the sloping desk.
Gadge and Innes	2007	4	1M/3F	22.25 21-26 (range)	Standard office chair with flat seat pan Saddle stool (Bambach) with forward inclination of 15°	Measured discomfort every 5 minutes while carrying out typing exercises over a 30-minute period.	N/A	100 mm VAS to record discomfort.	No previous history of musculoskeletal pain or disorder that affected sitting or typing performance Able to type Able to participate on two separate occasions Proficiency in written and spoken English	Less overall body discomfort, and LBD, on forward inclined seat. However, greater hip and buttock discomfort on forward inclined seat.

TABLE 1: (continued)

Study	Year	Sample Size	Sex	Age (Years)	Sitting Conditions	Study Protocol	Muscles Measured	Discomfort Measurement	Inclusion/Exclusion Criteria	Results
Jensen and Bendix	1992	10	5M/5F	20-36 (range)	Low seat, with backward inclined (-5°) seat pan High seat, with forward inclined (10°) seat pan High seat, with freely tiltable (unspecified degree) seat pan High seat, with freely tiltable (unspecified degree) seat pan with sloping desk	Subjects sat reading in each condition for 60 minutes.	N/A	Subjective preference on a scale from 1 to 5.	Healthy volunteers All accustomed to use of tiltable chair for at least 2 weeks	No significant difference in subjective preference between sitting conditions.
Koskelo et al.	2007	30	14M/16F	16	Low, nonadjustable seat with backrest with flat seat pan Higher, adjustable seat with hip angle of 45° and no backrest with adjustable sloping desk also	Compared the effect of 2 years use of adjustable with nonadjustable school desks and chairs on sitting and standing postures, trunk muscle strength, and tension and pain levels.	ST LES (L1/2)	Presence of LBP, neck-shoulder pain, and headache assessed at baseline and follow-up. Questionnaires to determine satisfaction with both chair and table.	Volunteer students, both with and without LBP	Incidence of LBP and headache not different between the groups at follow-up. Neck-shoulder pain incidence decreased in both groups, and to a greater extent in the group who sat in less hip flexion. Trapezius and LES muscle activation lower among the group who sat in less hip flexion. Better satisfaction scores among the group who sat in less hip flexion.

(continued)

TABLE 1: (continued)

Study	Year	Sample Size	Sex	Age (Years)	Sitting Conditions	Study Protocol	Muscles Measured	Discomfort Measurement	Inclusion/Exclusion Criteria	Results
Linton et al.	1994	67	35M/32F	9.9	Higher desks and chairs, with forward inclined seat (unspecified degree) Conventional desks and chairs	Compared the effect of 6 months use of each type of furniture on posture, satisfaction, and comfort, and the presence of pain in back, neck, or head. Also completed 5 month follow-up.	N/A	Presence of pain in back, neck, or head assessed using yes/no question. If pain present, VAS used for frequency of pain in each area.	Children in fourth grade of school in Sweden, both with and without LBP	Significantly greater comfort using intervention desks and chairs. Significantly less incidence of back pain among the intervention group. Nonsignificant trend for lower incidence of headache, neck, and overall pain among the intervention group. Nonsignificant trend for lower frequency of back pain among the intervention group. No significant differences in posture, but increased perceived awareness of correct sitting among intervention group.

(continued)

TABLE 1: (continued)

Study	Year	Sample Size	Sex	Age (Years)	Sitting Conditions	Study Protocol	Muscles Measured	Discomfort Measurement	Inclusion/Exclusion Criteria	Results
Marschall et al.	1995	10	5M/5F	4.7	Standard chair with 5° backward inclined seat pan, backrest, and flat desk Kneeler chair (forward inclination of 15°) with backrest and sloping desk (15°)	Compared muscle activity and comfort on each chair for ten minutes.	LES (L4/5) ST LD	Subjectively rated which workstation they preferred in terms of comfort.	Children familiar with both workstations Children without postural deviations Sloping chair and desk had been present in classroom for 12 months in advance, to allow familiarization	Significantly less LD muscle activity at the workstation involving less hip flexion. Similar trend of decreased activity for LES activity however this was not statistically significant. No difference in ST activation between the sitting conditions. All subjects subjectively reported they preferred the workstation involving less hip flexion.

(continued)

TABLE 1: (continued)

Study	Year	Sample Size	Sex	Age (Years)	Sitting Conditions	Study Protocol	Muscles Measured	Discomfort Measurement	Inclusion/Exclusion Criteria	Results
Michel and Helander	1994	Total 16: 5 (young healthy) 6 (old healthy) 5 (old herniated)	3M/2F 6M/0F 5M/0F	24.6 23–26 (range) 40.2 30–44 (range) 40 30–47 (range)	Conventional chair (5° backward inclination) Sit-stand chair (140° hip angle, 20° forward inclination)	Measured discomfort while sitting on two different chairs while subjects performed a computer task.	N/A	Comfort (GCR) and discomfort (BPD) rated 1–5 for each sitting condition.	One healthy student population One elderly group without LBP One elderly group with diagnosis of lumbar disc herniation whose previous occupation involved heavy manual tasks	Both young and old healthy subjects were significantly more comfortable on the conventional chair. However, subjects with herniated discs were significantly less comfortable in the conventional chair. Significantly more buttock discomfort on the sit-stand chair.

(continued)

TABLE 1: (continued)

Study	Year	Sample Size	Sex	Age (Years)	Sitting Conditions	Study Protocol	Muscles Measured	Discomfort Measurement	Inclusion/Exclusion Criteria	Results
Naqvi	1994	25	25M/0F	22	Horizontal seat Forward inclined seat pan (5°) Forward inclined seat pan (10°) Forward inclined seat pan (15°) The height of the seat increased in line with increasing forward inclination	Measured comfort (neck, lower back, and overall) at each seat inclination with subjects sitting at a computer workstation for 15 minutes.	N/A	11-point subjective comfort scale.	No history or present orthopedic symptoms including the neck, back, and spine or any other painful syndrome	LBD increased significantly with forward inclination beyond 5°. However, because both neck discomfort and overall discomfort were lower at 10° than 5°, they recommended different angles of inclination for each spinal region.
Saarni et al.	2009	98	41M/57F	13	Conventional school chairs with flat seat pan (hip angle 100°) and adjustable sloping desks Saddle-type chairs (hip angle 125°) and adjustable sloping desks	Students attended school as normal for 26 months, with students allocated to one of the sitting conditions. 2 phases to study—first 14 months, and then final 12 months.	N/A	Subjectively rated how much they liked their workstation (0–100).	Schoolchildren in sixth or eighth grade, both with and without LBP without LBP	Despite a short-term preference for the intervention sitting condition in the first 14 months, this was lost at 26 month follow-up.

(continued)

TABLE 1: (continued)

Study	Year	Sample Size	Sex	Age (Years)	Sitting Conditions	Study Protocol	Muscles Measured	Discomfort Measurement	Inclusion/Exclusion Criteria	Results
Soderberg et al.	1986	20	10M/10F	24.4 22-33 (range)	Conventional chair (flat seat pan) without using backrest Kneeler chair (Balans) with forward inclination of 10° Kneeler chair (Balans) with forward inclination of 20°	All subjects had CES (C6) to complete a computer task while sitting on each seat for 15 minutes. A subsample (n = 10) completed the task for longer (30 minutes) on each chair.	CES (C6) TES (T10) LES (L3)	Rank ordering of preference between the 3 chairs/seats.	Healthy volunteers No history of treatment for back pain	All 3 muscles were significantly less active with the 10° inclination compared to the flat seat pan. CES and LES were significantly less active with the 20° inclination compared to the flat seat pan. 19/20 subjects preferred the kneeler chairs, especially the 20° deg option.

(continued)

TABLE 1: (continued)

Study	Year	Sample Size	Sex	Age (Years)	Sitting Conditions	Study Protocol	Muscles Measured	Discomfort Measurement	Inclusion/Exclusion Criteria	Results
van der Heide et al.	2003	Total 39: 29 children 10 young adults	10M/19F NS	6.3 23.6	Horizontal seat (flat seat pan) Forward inclined seat pan (15°) Backward inclined seat pan (15°)	Subjects reached for a BB toy with their NF dominant hand under the 3 sitting conditions while EMG was recorded. BF They also reached for a toy from flat seat pan with small weighted bracelet on wrist.	DE BB NF CES (C7) RA TES (T10) LES (L5) RF	N/A	Healthy children with normal motor development Healthy adults with no physical complaints	Mean TES activation was significantly increased with backward inclined seat pan. TES activation was significantly delayed with forward inclined seat pan (in adults and children aged 2-4). LES onset activation was significantly delayed with backward inclined seat pan. No other significant differences.

(continued)

TABLE 1: (continued)

Study	Year	Sample Size	Sex	Age (Years)	Sitting Conditions	Study Protocol	Muscles Measured	Discomfort Measurement	Inclusion/Exclusion Criteria	Results
Wang et al.	2008	293	114M/179F	38 18–68 (range)	Chair with flat seat pan Chair with curved seat pan, forward inclined approximately 15° Control group	Compared whether a flat or a curved seat pan could improve back and hip pain in sewing machine operators over a 4-month period.	N/A	Questionnaires to assess musculoskeletal symptom frequency and intensity (rated from 1–5 if present) in 6 body regions.	Employees that performed sewing machine work for more than 20 hours a week Subjects who were not in a probationary period Had worked for at least 3 months and did not have an active workers compensation claim. Subjects who reported LBP or hip pain in the previous month	Both intervention chairs reduced LBP over a 4-month period compared to the control chair. However, the chair with less hip flexion did not reduce back and hip pain as much as the flat seat pan. This effect was particularly evident among men and people with a high BMI.
Winkel and Bendix	1986	10	0M/10F	33 22–41 (range)	Low seat, with backward inclined (–5°) seat pan High seat, with forward inclined (10°) seat pan High seat, with freely tiltable (–8° to 19.5°) seat pan	Subjects performed seated tasks (deskwork and typing) for 60 minutes.	LES (L3) SOL	N/A	Healthy volunteers All familiarized with each chair setting for 2 weeks	No significant differences in muscle activation.

(continued)

TABLE 1: (continued)

Study	Year	Sample Size	Sex	Age (Years)	Sitting Conditions	Study Protocol	Muscles Measured	Discomfort Measurement	Inclusion/Exclusion Criteria	Results
Wu et al.	1998	Total 32 Experiment 1: 20 Experiment 2: 6 Experiment 3: 6	Experiment 1: 10M/ 10F Experiment 2: 3M/3F Experiment 3: 3M/3F	20-25 (range)	Experiment 1 Conventional chair with wedge (100° hip) Experiment 2 Conventional chair with wedge (100° hip) Experiment 3 Conventional chair with no wedge (90° hip) Conventional chair with 10° wedge (100° hip) Conventional chair with 20° wedge (110° hip) Conventional chair with 30° wedge (120° hip)	Experiment 1: Subjects sat still on a chair for 5 minutes. Experiment 2: Subjects performed a word processing task for 60 minutes. Experiment 3: Subjects performed a typing task for 45 minutes.	N/A	Experiments 1, 2, and 3: Questionnaire with 10 items on spinal loading and comfort (5-point rating scale). Experiment 3: Body discomfort and fatigue was also measured using a VAS for each body region.	Pain-free healthy volunteers	Participants rated the addition of a pelvic support wedge somewhat positively in terms of providing stability and changing load. However, a 100° angle was no different to a flat seat pan, and reducing seated hip flexion to 110° or 120° actually increased discomfort.
Yu et al.	1988	2	0M/2F	40	Adjusted 7 independent variables on an adjustable chair intermittently, including seat height (hip angles of 135° and 105°) seat inclination (parallel to thigh, and 7.5° forward inclined)	Subjects completed a sewing task for 6 hours (with regular rest periods) on 8 separate days.	LES (L3) GA ST DE	Overall body discomfort scale (rated 0-7) Sum of localized body discomfort scale (rated 0-7).	Subjects experienced in industrial sewing tasks. Specifically chosen to represent women of tall and short stature No history of health problems	Significantly less overall and localized body discomfort with a lower seat height (105°). Nonsignificant trend for less localized body discomfort with forward inclined (7.5°) seat. No difference in LES activation or LES fatigue with varying seat height or seat inclination.

(continued)

TABLE 1: (continued)

Study	Year	Sample Size	Sex	Age (Years)	Sitting Conditions	Study Protocol	Muscles Measured	Discomfort Measurement	Inclusion/Exclusion Criteria	Results
Backrest studies										
Gregory et al.	2006	14	7M/7F	23.8	Standard office chair with backrest Exercise ball	Compared emg and discomfort during a range of office tasks (typing, computer-aided design, combined typing/mouse work, and reading) for 15 minutes each (total 1 hour)	TES (T9) LES (L3) RA EO	100 mm VAS for discomfort for multiple body regions (head/neck, shoulders, upper back, lower back, and overall)	Free of LBP for at least the previous year	Reduced TES activation with backrest. No significant differences for LES, RA, or EO. Reduced LBD and OBD with backrest.
Kingma and van Dieen	2009	10	0M/10F	21.7 1.6 (SD)	Standard office chair with backrest Exercise ball	Compared emg on each chair during a 1-hour typing task. Assessed discomfort during pilot testing only.	TES (T10) LES (L3) Trapezius	Not specifically measured—assessed discomfort during pilot testing only.	Pain-free and working with a computer regularly	Reduced mean activation, reduced activation at rest, reduced variation in activation, and reduced fatigue for LES with the backrest. No significant differences for TES or trapezius. Reduced LBD and OBD with backrest.

(continued)

TABLE 1: (continued)

Study	Year	Sample Size	Sex	Age (Years)	Sitting Conditions	Study Protocol	Muscles Measured	Discomfort Measurement	Inclusion/Exclusion Criteria	Results
Yoo et al.	2008	23	12M/11F	29.5 26-32 (range)	Standard office chair, with backrest provided at 3 different heights (T4, T10, L3)	Compared emg on each chair during a 5 minute typing task	CES (C4) LES (L5) Trapezius SA IO	N/A	Computer workers with forward head posture No back or neck pain in last year No neck or upper limb pathologies No rheumatological or neurological conditions	CES activation reduced, whereas LES and trapezius activation increased, with T4 backrest. IO activation increased with T10 backrest. LES activation reduced with L5 backrest. SA activation unchanged.
Combined studies Bennett et al.	1989	20	8M/12F	25 22-37 (range)	Two chairs with backrests Kneeler chair without backrest (Balans)	Compared emg during relaxed and erect sitting postures, as well as during writing and typing tasks. Duration unclear. Data recorded for 10 seconds.	LES (L3) LES (L5)	N/A	No history of back or knee pain Unfamiliar with Balans chair Able to touch type	No significant difference in muscle activation between chairs.
Cram and Vinitzky	1995	24	9M/15F	38.4 25-45 (range)	Standard flat office chair with backrest Kneeler chair without backrest (Balans) Pelvic support accessory (Back Up)	Compared emg on each chair during a ten minute writing task.	CES (C1) TES (T6) TES (T10) LES (L3)	N/A	No LBP (n = 12), and LBP (n = 12) with history of recurring LBP for at least one year	Increased LES activation on kneeler chair compared to both other chairs, especially among LBP group. No significant differences for CES or TES.

(continued)

TABLE 1: (continued)

Study	Year	Sample Size	Sex	Age (Years)	Sitting Conditions	Study Protocol	Muscles Measured	Discomfort Measurement	Inclusion/Exclusion Criteria	Results
Hardage et al.	1983	20	20M/0F	NS	Dentist's chair at 3 different hip angles (105°, 90°, and 75°), done with and without backrest	Compared emg TES (T5) in each sitting LES (L3) condition while performing a one minute simulated dental task.	TES (T5) N/A	N/A	Pain-free participants	No statistically significant differences between different hip angles. Reduced LES & TES activation with backrest.
Lander et al.	1987	20	13M/7F	33 23–46 (range)	Standard flat (nonoffice) chair with backrest Kneeler chair without backrest (Balans)	Compared emg on each chair while watching a videotape for 30 minutes.	CES (C6) LES (L4)	10 mm VAS for comfort preference in multiple body regions (overall, low back, neck, and leg)	Healthy volunteers No history of neck or back pain No other medical complaints	Reduced LES & CES activation with standard chair with backrest. Increased overall and low back comfort using standard chair with backrest.

Note. Age expressed as mean unless otherwise stated. BB, biceps brachii; BF, biceps femoris; BPD, body part discomfort scale; CES, cervical erector spinae; C1/4/6, cervical vertebral levels; DE, deltoid; emg, electromyography; EO, external oblique; F, female; GA, gastrocnemius; GCR, general comfort rating scale; IO, internal oblique; LBD, low back discomfort; LBP, low back pain; LD, latissimus dorsi; LES, lumbar erector spinae; L1/2/3/4/5, lumbar vertebral levels; M, male; N/A, not assessed; NF, neck flexor (Sternocleidomastoid); NS, not stated; RA, rectus abdominis; RCT, randomized controlled trial; RF, rectus femoris; SOL, soleus; ST, superior trapezius; TA, tibialis anterior; TES, thoracic erector spinae; T4/5/6/9/10, thoracic vertebral levels; VAS, visual analogue scale.




Figure 3 Flow chart indicating study selection procedure for chairs which reduce seated hip flexion.

trunk muscle activation, LBP or LBD. Bendix and Biering-Sørensen (1983) reported no difference in discomfort on four different seats—a flat seat pan and three forward inclined seats whose height increased in line with increasing forward seat pan inclination. Bendix (1984) reported no significant difference in acceptability between three seats of varying inclination and mobility whose participants were instructed to rest their

elbows on the table. In contrast, Bendix, Jessen, and Winkel (1986) reported significantly greater acceptability for an armless, forward inclined higher seat compared to an armless, low, backward inclined seat. Bendix, Winkel, and Jessen (1985) observed that LES activation was not significantly different between three armless seats of varying inclination and mobility in a two-part study. Although the tiltable chair was significantly

more acceptable in a short laboratory study, this difference was not statistically significant in the longer field study. Winkel and Bendix (1986) reported no difference in LES or calf muscle activation between a low, backward inclined seat, and two higher seats that were forward inclined or freely tiltable. This study did not state whether chairs featured armrests or whether participants were instructed to lean on the table while completing the tasks. Another study by the same research group (Jensen & Bendix, 1992) also reported no significant difference in preference between three armless seats of varying inclination and mobility. Subjects in this study were instructed to sit as they pleased. Naqvi (1994) assessed discomfort every five minutes during 15 minutes of sitting on four different armless seats—a flat seat pan and three forward inclined seats. LBD increased significantly with increasing forward inclination beyond 5° and after 10 minutes of sitting. However, because both neck discomfort and overall discomfort were significantly lower at 10° than 5°, they recommended different angles of inclination for each spinal region. Van der Heide, Otten, van Eykern, and Hadders-Algra (2003) demonstrated that mean TES activation was significantly increased with a backward inclined seat pan, whereas LES onset activation was significantly delayed with a backward inclined seat pan. The authors did not specify if the seats featured armrests. Inclining the seat pan forward to varying degrees by sitting on three different wedges (10°, 20°, and 30°) was investigated in one study (Wu, Miyamoto, & Noro, 1998). In this series of three experiments, pain-free participants rated the use of a wedge positively in terms of the stability it provided, and the way they perceived it placed load on their spine and pelvis. However, the level of discomfort (rated on a 5-point scale) was increased with wedges which reduced hip flexion by 20° or 30°, whereas the level of discomfort using a 10° wedge was no better than using no wedge. Similarly, it was not stated if the seat featured armrests. Yu, Keyserling, and Chaffin (1988) modified seven sitting variables including seat height and seat inclination. They reported significantly less overall and localized body discomfort with a lower seat than with a higher seat. There were no significant differences during seat forward incli-

nation, and no significant differences in LES activation or LES fatigue with varying seat height or seat inclination.

The final three crossover studies simply controlled the degree of hip flexion in sitting. Hardage, Gildersleeve, and Rugh (1983) compared TES and LES activation while sitting with and without a backrest at three different hip angles. Despite some evidence of an interaction between hip angle and the presence of a backrest in sitting, there was no significant difference in paraspinal muscle activation between seats with different hip flexion angles. Gadge and Innes (2007) compared discomfort levels between an armless conventional chair with a flat seat pan and a Bambach armless saddle seat. It was not stated if participants were instructed to rest their elbows or whether the assessor monitored use of the table. Both LBD and overall body discomfort (OBD) increased more slowly on the Bambach seat, however discomfort was higher for the hips and buttocks when sitting on the Bambach seat. Michel and Helander (1994) reported that people with LBP reported significantly less LBD using an armless sit-stand chair than a conventional armless chair. However, pain-free participants reported less LBD on the conventional chair. Overall, buttock discomfort was significantly greater in the sit-stand chair.

Intervention studies. Five studies examined the effect of these chairs over a period of time greater than one day. An RCT by Wang et al. (2008) compared the effect of a curved seat pan to both (a) a flat seat pan and (b) a placebo group receiving only miscellaneous items (footrest, small table-top storage box, side table, lamp, and reading glasses). Although both armless “intervention” chairs reduced back and hip pain over a four month period, the curved seat pan did not actually reduce back and hip pain as much as the flat seat pan. The final four studies examined the effect of incorporating these seats into an ergonomically designed workstation including a sloping desk. All used schoolchildren as the population of interest. All seats were armless. An RCT by Linton, Hellsing, Halme, and Åkerstedt (1994) compared the effect of a traditional workstation with an ergonomic workstation over a six month period. The intervention group reported a significant reduction

in the incidence of LBP on a dichotomous (yes/no) scale. Comfort was also rated higher among the intervention group. However, the actual frequency of LBP, headache, neck pain, and overall musculoskeletal pain was not significantly different between groups at follow-up. The reduction in LBP incidence occurred without any change in actual spinal posture. Another RCT (Aagaard-Hansen & Storr-Paulsen, 1995) examined the effect of changing seat height and inclination, desk slope, or both. Combining a forward inclined, higher seat, and adjustable sloping desk was subjectively preferred overall to (a) a traditional flat seat pan and desk and (b) a flat seat pan and an adjustable sloping desk. However, there was no significant difference in the frequency of reported LBP. A nonrandomized clinical trial (Koskelo et al., 2007) compared a forward inclined seat combined with an adjustable sloped desk to standard furniture. Reducing seated hip flexion decreased trapezius and LES muscle activation, and was associated with better satisfaction scores at 2-year follow-up. The incidence of neck and shoulder pain decreased in both groups, especially in the group who sat in less hip flexion. However, there was no significant difference in the incidence of LBP or headache between the groups at follow-up. Finally, another nonrandomized clinical trial (Saarni et al., 2009) compared the use of saddle-type chairs and adjustable desks to a control group still using their usual chairs and desks. The ergonomic workstations were preferred in the first year, but not in the second year. In addition, no significant differences in seated posture or trunk mobility were observed between the two groups.

In conclusion, the use of chairs that reduce seated hip flexion appears to be associated with increased paraspinal muscle activation and/or increased discomfort. However, this is likely to be partly explained by other differences between the sitting conditions studied. This includes the use of a backrest or a sloping desk, which appear to ameliorate the negatives of forward inclined chairs (Bridger, 1988; Marschall et al., 1995; Soderberg et al., 1986) to the extent that reducing seated hip flexion in itself does not influence LBP, LBD, or trunk muscle activation.

Study 2: Effect of Providing a Backrest

Identification of studies. The results of the search strategy are outlined in Figure 4. The electronic search returned 386 potentially relevant papers, which was reduced to 275 after removing duplicates. After screening these titles and abstracts, 16 potentially relevant articles were identified. After reviewing these 16 full-text articles, only 7 of them met the inclusion and exclusion criteria. Searching the reference lists of these articles did not add any further articles. All seven eligible studies were rated as “fair” to “high” quality and were included in the review. Therefore, the final number of articles included in this review was seven. All seven studies were crossover design studies, with one (Kingma & van Dieen, 2009) involving testing on two separate days. All seven studies examined trunk muscle activation, whereas two of them (Gregory, Dunk, & Callaghan, 2006; Lander et al., 1987) also examined LBD. Only one study (Cram & Vinitzky, 1995) included both pain-free participants and people with LBP, with the remainder examining only pain-free participants.

Description of results. Four studies have already been described in Part 1 of this review, as they met the inclusion criteria for both systematic reviews (Bennett et al., 1989; Cram & Vinitzky, 1995; Hardage et al., 1983; Lander et al., 1987).

Two studies compared sitting with a backrest to sitting on an exercise ball. In one study (Gregory et al., 2006), the chair with a backrest involved significantly lower activation of TES, and significantly lower LBD and OBD. In the other study comparing to sitting on an exercise ball (Kingma & van Dieen, 2009), using a backrest was associated with lower LES mean activation, lower variation in activation of LES, and lower LES muscle fatigue. This study did not assess discomfort, although they reported that participants experienced less LBD and upper back discomfort when using the backrest during pilot testing. Finally, using a ball-shaped backrest at three different heights was compared to not using a backrest (Yoo et al., 2008). The effect of the backrest varied according to its height, suggesting that it may be clinically relevant to choose backrest height based on an individual's




Figure 4. Flow chart indicating study selection procedure for chairs using backrests.

height (spinal length) and region(s) of discomfort. It is important to acknowledge that the results of these two studies could be related to differences in mobility between the chairs, and not just the presence of a backrest, although a recent systematic review suggests dynamic sitting has little effect on LBP/LBD (O'Sullivan, O'Keeffe, et al., 2012).

In conclusion, the use of a backrest appears to reduce paraspinal muscle activation according to the height it is positioned at, most commonly the lower lumbar spine. The decrease in paraspinal activation is likely due to external support being provided by the backrest, which spares the muscles from needing to provide active support. This also appears to lead to a decrease in LBD, though the lack of RCTs involving people with LBP limits strong conclusions on this point.

Critical Appraisal

The detailed methodology of each included study is provided in Table 1. Table 2 illustrates that all included studies were rated as fair (4–5/10) to high (≥ 6 –10/10) quality, with all studies except three (Koskelo et al., 2007; Saarni et al., 2009; Wu et al., 1998) scoring in the range of 5–7/10. This section briefly summarizes the key methodological concerns regarding all studies included.

Bias. Randomization was acceptable in most studies, although the precise method of randomization (e.g., computer-generated or by selecting allocation from an opaque envelope) was not always stated. Although several studies allocated participants in a counterbalanced order instead, this is not a major cause for concern in crossover or repeated measures design studies. Concealment of allocation was performed in most, but not all

studies. All crossover studies clearly had similar groups at baseline, however one study (Wang et al., 2008) did not demonstrate comparability of groups at baseline. Only two studies reported blinded assessors (Linton et al., 1994; Wang et al., 2008). Blinding of therapists and participants is difficult to achieve in such studies, although some studies (Aagaard-Hansen & Storr-Paulsen, 1995; Wang et al., 2008) used several intervention chairs to partially address participant bias. Limited information was provided about inclusion/exclusion criteria overall, increasing the potential for selection bias. No study mentioned using a random recruitment strategy which decreases the generalizability of results. A range of participant ages and sex were included.

Confounders. Some significant confounding variables were present in several studies. Considering the review on reducing seated hip flexion first, several studies did not control for the presence of a backrest (Koskelo et al., 2007; Saarni et al., 2009) or the height and slope of the desk used (Koskelo et al., 2007; Linton et al., 1994; Saarni et al., 2009) between sitting conditions. Furthermore, despite the fact that changing seat pan inclination alters the pressure on a backrest (Bendix et al., 1985), the vast majority of studies did not investigate if backrest pressure varied between sitting conditions. In some studies, the size of the seat pan (Wang et al., 2008) or the size and inclination of the backrest (Gadge & Innes, 2007) varied considerably between the flat and inclined seats. In some studies, neither the inclination of the intervention seat nor its effect on hip angle was reported, and was not available from the authors when contacted (Aagaard-Hansen & Storr-Paulsen, 1995; Linton et al., 1994). In fact, it was not always clear if the modified chair design would have a major effect on the seated hip flexion angle (Wang et al., 2008).

For the review on chair backrests, it is again clear that several studies did not control for other factors which differed between the sitting conditions (other than the presence of a backrest) and which could affect the findings. This includes comparing to an exercise ball that involved greater movement, and comparing to a kneeler chair that not alone had no backrest, but also

involved far less hip flexion. The presence of these confounders reflects the fact that the chair with the backrest was often the control sitting condition in these studies. No study reported any objective method of checking that participants actually used the backrest when sitting in the backrest condition, which could confound the results (Vergara & Page, 2000a).

In both reviews, the duration of sitting exposures, and the use of rest periods between them, varied considerably between studies (Table 1), although these were similar between sitting conditions in all studies that mentioned these details. Similarly, some studies performed testing on separate days to avoid fatigue, however greater error in calculating muscle activation occurs between different days (Dankaerts, O'Sullivan, Burnett, Straker, & Danneels, 2004). Although the tasks performed varied between studies, within each study tasks were standardized in each sitting condition, to control for task variation affecting the results (Groenesteijn et al., 2012).

In both reviews, several studies did not control for the presence of armrests. Only two studies (Bendix, 1984; Soderberg et al., 1986) specifically stated that participants had to rest their elbows in front of them. Seven studies (Bridger, 1988; Cram & Vinitzky, 1995; Hardage et al., 1983; van der Heide et al., 2003; Winkel & Bendix, 1986; Wu et al., 1998; Yu et al., 1988) did not state whether seats featured armrests which could confound results. All other studies featured armless seats, however they did not monitor or report if participants were forward leaning onto a surface such as a table, which could confound results (Soderberg et al., 1986).

Strength of results. None of the included studies in either of the two reviews calculated their sample size based on a-priori power calculations, and many may have been underpowered to detect differences between sitting conditions. Appropriate methods of statistical analysis were used in nearly all studies. Only a small number of studies included a considerable number of drop-outs, and suitable intention-to-treat analysis was used in most of these studies reducing the risk of attrition bias. The sample size among studies in the review of seated hip flexion angles ranged from two to 293, with an overall sample of 986

TABLE 2: PEDro Score for Each Study

Study	Random Allocation	Concealed Allocation	Baseline Comparability	Assessor Blinded	Participant Blinded	Therapist Blinded	Follow-up	Intention-to-Treat Analysis	Between-Group Analysis	Point Estimates and Variability	Total Score/10
Reducing hip flexion studies											
Aagaard-Hansen and Storr-Paulsen	✓	✓	✓	X	X	X	✓	X	✓	X	5
Bendix and Biering-Sørensen	✓	X	✓	X	X	X	✓	✓	✓	✓	6
Bendix	✓	X	✓	X	X	X	✓	✓	✓	✓	6
Bendix et al. (1986)	✓	X	✓	X	X	X	✓	✓	✓	X	5
Bendix et al. (1985)	✓	X	✓	X	X	X	✓	✓	✓	✓	6
Bendix et al. (1988)	X	X	✓	X	X	X	✓	✓	✓	✓	5
Bishu et al.	X	X	✓	X	X	X	✓	✓	✓	✓	5
Bridger	✓	X	✓	X	X	X	✓	✓	✓	X	5
Gadge and Innes	X	✓	✓	X	X	X	✓	✓	✓	✓	6
Jensen and Bendix	✓	X	✓	X	X	X	✓	✓	✓	X	5
Koskelo et al.	X	X	✓	X	X	X	✓	X	✓	✓	4
Linton et al.	✓	✓	✓	✓	X	X	✓	✓	✓	X	7
Marschall et al.	X	X	✓	X	X	X	✓	✓	✓	✓	6
Michel and Helander	X	✓	✓	X	X	X	✓	✓	✓	X	5
Naqvi	✓	✓	✓	X	X	X	✓	✓	✓	X	6
Saarni et al.	X	X	✓	X	X	X	X	✓	✓	✓	4
Soderberg et al.	✓	✓	✓	X	X	X	✓	✓	✓	✓	7
van der Heide et al.	✓	✓	✓	X	X	X	✓	✓	✓	✓	7
Wang et al.	✓	✓	X	✓	X	X	X	✓	✓	✓	6
Winkel and Bendix	✓	X	✓	X	X	X	✓	✓	✓	X	5
Wu et al.	X	X	✓	X	X	X	✓	✓	✓	X	4
Yu et al.	X	X	✓	X	X	X	✓	✓	✓	✓	5
Backrest studies											
Gregory et al.	✓	✓	✓	X	X	X	✓	✓	✓	✓	7
Kingma & van Dieen	X	X	✓	X	X	X	✓	✓	✓	✓	5
Yoo et al.	✓	✓	✓	X	X	X	✓	✓	✓	✓	7
Combined studies											
Bennett et al.	✓	✓	✓	X	X	X	✓	✓	✓	✓	7
Cram and Vinitzky	✓	X	✓	X	X	X	✓	✓	✓	✓	6
Hardage et al.	X	X	✓	X	X	X	✓	✓	✓	✓	5
Lander et al.	✓	✓	✓	X	X	X	✓	✓	✓	✓	7

participants included. The sample size among studies in the review of chair backrests ranged from 10 to 24, with an overall sample of 131 participants included. The review on chair backrests found no randomized controlled trial (RCT) or longitudinal studies with a follow-up period, preventing any attempt to identify if using a chair backrest reduces LBP incidence or severity in the medium term.

Clinical applicability. Only seven studies across both reviews included participants with LBP, with little detail provided on their level of functional disability which limits its clinical applicability. In laboratory-based studies, participants were usually asked to perform simulated functional or occupational tasks to enhance clinical applicability. Only five studies examined the effect of reducing seated hip flexion over a prolonged period of time. However, some of these studies (Koskelo et al., 2007; Saarni et al., 2009) had long follow-up periods of approximately 2 years. Four of these five follow-up studies examined schoolchildren, which may not reflect the nature, or intensity, of LBP reported by adult populations. Several different scales to analyze LBP and LBD were used, preventing pooled analysis of the data. Some studies simply asked participants about their subjective preference, which is less clinically relevant. There were also large variations in the trunk muscles analyzed between studies (Table 1). Even when studies analyzed the same muscles, variations regarding electrode placement and the type of muscle activation analysis performed did not allow for pooled analysis of the data.

DISCUSSION

The results of both systematic reviews will be discussed for muscle activation and discomfort separately.

Muscle Activation

Muscle activation on chairs that reduce seated hip flexion. The use of kneeler chairs was associated with increased paraspinal muscle activation and/or increased discomfort in five of the eight studies that investigated them. In these five studies the control sitting condition offered participants a backrest whereas the kneeler chair did not, which

is likely to partly explain the difference between the sitting conditions. This is further highlighted by the fact that the three studies that reported at least some benefit from kneeler chairs ensured that the presence or absence of a backrest was consistent between the sitting conditions compared (Bridger, 1988; Marschall et al., 1995; Soderberg et al., 1986). In addition, two of the three studies which reported some benefit from a kneeler chair, combined it with a sloping desk, which may also have contributed to the benefit (Bridger, 1988; Marschall et al., 1995). Therefore, it would appear that kneeler chairs actually increase paraspinal muscle activation and/or discomfort, unless they have a backrest or are integrated into a wider ergonomic workstation with sloping desks, when they may actually reduce paraspinal muscle activation and are preferred by participants. The increase in discomfort associated with these chairs when they do not have a backrest suggest that paraspinal muscle activation increases in the absence of external support being provided by the backrest, leading to unnecessarily high levels of muscle activation being needed to provide active support. No study examined the use of kneeler chairs prospectively.

The 13 crossover studies which reduced seated hip flexion through changing the seat height, using tilted seat pans, or combining these features, revealed more consistent results, reflecting the fact that 12 of these studies controlled for confounding variables such as the presence of a backrest and the use of sloping desks. None of these 13 studies demonstrated unequivocal advantages for reducing seated hip flexion.

Muscle activation on chairs with backrests. Five of the seven studies in the backrest review reported that using a backrest significantly reduced LES muscle activation (Cram & Vinitzky, 1995; Hardage et al., 1983; Kingma & van Dieen, 2009; Lander et al., 1987; Yoo et al., 2008). A backrest had minimal effects on the other trunk muscles studied. The effect of the backrest appears to differ according to its position, with the paraspinal muscles at the height of the backrest being the most likely to be decreased (Yoo et al., 2008). The decrease in LES activation is likely due to external support being provided by the backrest, which spares the muscles from needing to provide active support.

Sitting Discomfort

Discomfort on chairs that reduce seated hip flexion. One study (Gadge & Innes, 2007) reported less LBD and OBD among pain-free participants when using a saddle chair with a backrest, but this was offset by an increase in buttock discomfort. Another study reported reduced LBD when sitting in less hip flexion among people with LBP (Michel & Helander, 1994), however pain-free participants reported greater discomfort using this chair. Finally, some limited evidence to support the use of reducing seated hip flexion was observed in two studies that reported a reduction in the activation of some paraspinal muscles (van der Heide et al., 2003) and enhanced subjective acceptability compared to a freely tiltable chair (Bendix et al., 1986). However, most of the crossover design studies reported no difference in LBP or LBD when seated hip flexion was slightly reduced (Bendix, 1984; Bendix & Biering-Sørensen, 1983; Bendix et al., 1985; Hardage et al., 1983; Jensen & Bendix, 1992; Winkel & Bendix, 1986; Wu et al., 1998). In fact, several studies (Bendix & Biering-Sørensen, 1983; Bendix et al., 1988; Naqvi, 1994; Wu et al., 1998) noted that reducing hip flexion to a much larger degree actually increased discomfort. One study (Yu et al., 1988) actually reported greater discomfort when seated hip flexion was reduced. However, the hip angles compared were 105° and 135°, probably reflecting the previous point that very large decreases in hip flexion appear to increase discomfort. Finally, although using ergonomic workstations including saddle-type chairs was preferred during the first year of a 2-year study (Saarni et al., 2009), this preference was lost in the second year. It is clear therefore that although some studies subjectively preferred chairs with reduced hip flexion, the majority of studies that examined discomfort demonstrated increased discomfort levels, particularly if large changes in hip flexion occurred.

Discomfort on chairs with backrests. The only two studies (Gregory et al., 2006; Lander et al., 1987) which specifically examined the effect of a backrest on sitting discomfort favored using a backrest. One other study (Kingma & van Dieen, 2009) did not specifically measure

discomfort, but also reported increased participant comfort using a backrest during pilot testing. None of these three studies included people with LBP.

Consideration of Confounding Variables

As previously stated, the presence of several significant confounders in many studies warrants some caution in interpreting these findings. Nevertheless, it is possible to at least partly interpret the contribution of such confounders to the results. For example, a recent systematic review (O'Sullivan, O'Keefe, et al., 2012) suggests that dynamic sitting does not significantly influence LBP or LBD. Consequently, the most likely reason for differences observed in the two dynamic comparisons (Gregory et al., 2006; Kingma & van Dieen, 2009) was the presence of a backrest rather than differences in mobility. It is more difficult to determine the confounding influence of hip angle when interpreting the effect of using a backrest. The single study that did actually control for hip angle between chairs with and without a backrest (Hardage et al., 1983) reported no difference in muscle activation between three different hip angles. Similarly, although the kneeler chair design typically increased paraspinal activation (Bennett et al., 1989; Cram & Vinitzky, 1995; Lander et al., 1987), other studies (e.g., Koskelo et al., 2007) have reported decreased paraspinal activation using other chair designs with less seated hip flexion. Variations in the degree of forward trunk lean between studies may also be relevant, as this could also influence paraspinal muscle activation if not monitored closely. This may explain why the differences between chairs in one of the studies were smaller during forward lean tasks than during sitting (Bennett et al., 1989). This overlaps with the fact that none of the included studies measured how much pressure was placed on the backrest, or whether in fact it was used by participants (Vergara & Page, 2000a). Another common potential confounder was that studies (e.g., Bridger, 1988; Cram & Vinitzky, 1995; Hardage et al., 1983; van der Heide et al., 2003; Winkel & Bendix, 1986; Wu et al., 1998; Yu et al., 1988) did not always state whether seats featured armrests or not. Finally,

it appears the effect of a backrest is closely related to its location (Yoo et al., 2008), which may explain some of the variation between studies. Notwithstanding these confounding factors, the results demonstrate moderate evidence that chair backrests reduce paraspinal muscle activation, and limited evidence that chair backrests reduce LBD. For chairs involving less hip flexion, the increased LBP/LBD reported in many studies probably reflects the confounding effect of not providing a backrest in the intervention chair. Therefore, for chairs involving less seated hip flexion the results demonstrate no evidence that they reduce LBP or LBD, or consistently alter trunk muscle activation.

Implications

This relatively limited supporting evidence to support the use of chairs involving less seated hip flexion or backrests is consistent with the limited evidence that other chair design features in isolation reduce LBP (Driessen et al., 2010, O'Sullivan, O'Keeffe, et al., 2012). This reflects the multidimensional nature of LBP where not just biomechanical and ergonomics factors, but also psychosocial (Carroll, Cassidy, & Côté, 2004; Jarvik et al., 2005; Main, Foster, & Buchbinder, 2010; Mitchell et al., 2010; Ramond et al., 2011), lifestyle (Chiu et al., 2005; Onen, Alloui, Gross, Eschallier, & Dubray, 2001), genetic (Battié, Videman, & Parent, 2004; Reichborn-Kjennerud et al., 2002), and neurophysiological (Apkarian, Baliki, & Geha, 2009; Wand et al., 2011) factors may be involved (O'Sullivan, 2012). Therefore, any unidimensional biomechanical or ergonomic approach to managing LBP is likely to be of limited effectiveness in isolation. Considering that people with LBP have previously been shown to have greater difficulty relaxing their trunk muscles than people without LBP (Geisser et al., 2005), using a backrest to reduce muscle activation may be of benefit to people with LBP. Another important consideration is the need to discriminate between participant ratings of preference in the short term compared to clinically meaningful changes in LBP/LBD in the long term. Participants in several studies subjectively preferred the chairs involving less hip flexion, but these chairs typically did not result in reductions

in LBP/LBD. It may be that simply changing spinal loading, and/or the novelty associated with a new chair design, explains these short-term positive perceptions without actually having a meaningful impact on LBP. All of the studies included in these reviews prescribe the same change in sitting to all participants. This is not consistent with clinical practice, and does not reflect well documented individual variations in posture and movement patterns among people with LBP (Dankaerts et al., 2006; Dankaerts et al., 2009). For example, two recent studies published after short listing for these systematic reviews was complete suggest that matching chair prescription to the individual presentations of people with LBP may be more effective than generic prescriptions that adopt a one-size-fits-all approach (Curran, Dankaerts, O'Sullivan, O'Sullivan, & O'Sullivan, 2014; O'Keeffe, Dankaerts, O'Sullivan, O'Sullivan, & O'Sullivan, 2013). We believe that if changes in chair design are to have a more significant impact on LBP it will be through matching the seating prescription to the individual, their specific demands at work or home, and/or linking it with other barriers to recovery from LBP across the biopsychosocial spectrum.

Limitations

The main limitations of these reviews are that significant differences in the outcome measure used, the participants included, and the comparison sitting conditions did not allow for pooled analysis of the data. There were very few RCT designs including people with LBP, and most tended to involve only very short-term follow-up of participants. Larger RCTs involving people with LBP, with suitable blinding, less confounding variables, and involving a long-term follow-up period, are required to fully confirm the findings of this review. Given the tendency for novel chair designs to differ in several ways (e.g., backrest presence and/or orientation, seat pan angle and motion, lower limb position; O'Sullivan et al., 2012a, 2012b), studies that identify the specific utility of each design feature may be useful. Many of the included studies were quite old, with specific design features which were unclear. The uncertainty as to whether armrests were used or not

in several of the studies may have influenced the findings. Further research is required to shed light on whether the acceptability of specific chair designs is affected by the duration of sitting studied.

CONCLUSION

These two reviews included a total of 29 studies investigating the effect of backrests and reduced seated hip flexion on sitting discomfort or trunk muscle activation. Significant confounding variables existed, and only a small number of RCTs involving people with LBP were found. There was moderate evidence that chair backrests reduce paraspinal muscle activation, and limited evidence that chair backrests reduce LBD. Despite participants in several studies subjectively preferring chairs involving less hip flexion, there was no evidence that these chairs reduce LBP or LBD, or consistently alter trunk muscle activation.

The limited evidence to support the use of chairs involving less seated hip flexion, or the effect of a backrest, is consistent with the limited evidence that other isolated chair design features can reduce LBP.

ACKNOWLEDGMENTS

KO was supported by a fellowship from the Health Research Board of Ireland. Thanks to Mary O’Keeffe and John Hurley for comments on the draft manuscript. Some of the authors are currently involved in studies using a chair that involves sitting in less hip flexion (www.backapp.eu), and that is used in the management of low back pain. Those studies are partially funded by the manufacturers of the chair. The authors do not feel this has in any way affected this systematic review, but are disclosing this fact to enhance transparency. None of the authors have any stake/shares in this, or any other specific chair design.

KEY POINTS

- The effect of a chair backrest and reducing seated hip flexion on low back pain (LBP)/low back discomfort (LBD) and trunk muscle activation was systematically reviewed.
- Many studies included several confounding variables, and only a small number of RCTs involving

people with LBP were found, which complicated analysis of the results.

- There was moderate evidence from seven studies that chair backrests reduce paraspinal muscle activation, and limited evidence that chair backrests reduce LBD.
- Despite participants in several studies subjectively preferring chairs involving less hip flexion, there was no evidence that these chairs reduce LBP or LBD, or consistently alter trunk muscle activation.
- The limited evidence to support the use of chairs involving less seated hip flexion, or the effect of a backrest, is consistent with the limited evidence that other isolated chair design features can reduce LBP.

REFERENCES

- Aagaard-Hansen, J., & Storr-Paulsen, A. (1995). A comparative study of three different kinds of school furniture. *Ergonomics*, 38(5), 1025–1035.
- Andersson, B., Ortengren, R., Nachemson, A., Elfström, G., & Broman, H. (1975). The sitting posture: An electromyographic and discometric study. *Orthopedic Clinics of North America*, 6(1), 105–120.
- Andersson, E. A., Jonsson, B., & Ortengren, R. (1974). Myoelectronic activity in individual lumbar erector spinae muscles in sitting. *Scandinavian Journal of Rehabilitation Medicine*, 3, 91–108.
- Apkarian, A. V., Baliki, M. N., & Geha, P. Y. (2009). Towards a theory of chronic pain. *Progress in Neurobiology*, 87(2), 81–97.
- Battié, M. C., Videman, T., & Parent, E. (2004). Lumbar disc degeneration: Epidemiology and genetic influences. *Spine*, 29(23), 2679–2690.
- Bendix, A., Jensen, C. V., & Bendix, T. (1988). Posture, acceptability and energy consumption on a tiltable and a knee-support chair. *Clinical Biomechanics*, 3(2), 66–73.
- Bendix, T. (1984). Seated trunk posture at various seat inclinations, seat heights, and table heights. *Human Factors*, 26, 695–703.
- Bendix, T., & Biering-Sørensen, F. (1983). Posture of the trunk when sitting on forward inclining seats. *Scandinavian Journal of Rehabilitation Medicine*, 15(4), 197–203.
- Bendix, T., Jessen, F. B., & Winkel, J. (1986). An evaluation of a tiltable office chair with respect to seat height, backrest position and task. *European Journal of Applied Physiology and Occupational Physiology*, 55(1), 30–36.
- Bendix, T., Winkel, J., & Jessen, F. (1985). Comparison of office chairs with fixed forwards or backwards inclining, or tiltable seats. *European Journal of Applied Physiology and Occupational Physiology*, 54(4), 378–385.
- Bennett, D. L., Gillis, D. K., Portney, L. G., Romanow, M., & Sanchez, A. S. (1989). Comparison of integrated electromyographic activity and lumbar curvature during standing and during sitting in three chairs. *Physical Therapy*, 69(11), 902–913.
- Bishu, R. R., Hallbeck, M. S., Riley, M. W., & Stentz, T. L. (1991). Seating comfort and its relationship to spinal profile: A pilot study. *International Journal of Industrial Ergonomics*, 8(1), 89–101.
- Bridger, R. S. (1988). Postural adaptations to a sloping chair and work surface. *Human Factors*, 30, 237–247.

- Carcone, S., & Keir, P. (2007). Effects of backrest design on biomechanics and comfort during seated work. *Applied Ergonomics*, 38(6), 755–764.
- Cardon, G., De Clercq, D., De Bourdeaudhuij, I., & Breithecker, D. (2004). Sitting habits in elementary schoolchildren: A traditional versus a "moving school." *Patient Education and Counseling*, 54(2), 133–142.
- Carroll, L. J., Cassidy, J. D., & Côté, P. (2004). Depression as a risk factor for onset of an episode of troublesome neck and low back pain. *Pain*, 107(1), 134–139.
- Chiu, Y., Silman, A., Macfarlane, G., Ray, D., Gupta, A., Dickens, C., Morriss, R., & McBeth, J. (2005). Poor sleep and depression are independently associated with a reduced pain threshold. Results of a population based study. *Pain*, 115(3), 316–321.
- Claus, A., Hides, J., Moseley, G. L., & Hodges, P. (2008). Sitting versus standing: Does the intradiscal pressure cause disc degeneration or low back pain? *Journal of Electromyography and Kinesiology*, 18(4), 550–558.
- Claus, A., Hides, J., Moseley, G., & Hodges, P. (2009). Is "ideal" sitting real? Measurement of spinal curves in four sitting postures. *Manual Therapy*, 14, 404–408.
- Corlett, E. N. (2006). Background to sitting at work: Research-based requirements for the design of work seats. *Ergonomics*, 49(14), 1538–1546.
- Cram, J. R., & Vinitzky, I. (1995). Effects of chair design on back muscle fatigue. *Journal of Occupational Rehabilitation*, 5(2), 101–113.
- Curran, M., Dankaerts, W., O'Sullivan, P., O'Sullivan, L., & O'Sullivan, K. (2014). The effect of a backrest and seatpan inclination on sitting discomfort and trunk muscle activation in subjects with extension-related low back pain. *Ergonomics*, 57(5), 733–743.
- Dankaerts, W., O'Sullivan, P., Burnett, A., & Straker, L. (2006). Differences in sitting postures are associated with nonspecific chronic low back pain disorders when patients are subclassified. *Spine*, 31(6), 698–704.
- Dankaerts, W., O'Sullivan, P. B., Burnett, A. F., Straker, L. M., & Danneels, L. A. (2004). Reliability of EMG measurements for trunk muscles during maximal and sub-maximal voluntary isometric contractions in healthy controls and CLBP patients. *Journal of Electromyography & Kinesiology*, 14(3), 333.
- Dankaerts, W., O'Sullivan, P., Burnett, A., Straker, L., Davey, P., & Gupta, R. (2009). Discriminating healthy controls and two clinical subgroups of nonspecific chronic low back pain patients using trunk muscle activation and lumbosacral kinematics of postures and movements: A statistical classification model. *Spine*, 34(15), 1610–1618.
- De Carvalho, D., Soave, D., Ross, K., & Callaghan, J. (2010). Lumbar spine and pelvic posture between standing and sitting: A radiologic investigation including reliability and repeatability of the lumbar lordosis measure. *Journal of Manipulative and Physiological Therapeutics*, 33(1), 48–55.
- de Morton, N. A. (2009). The PEDro scale is a valid measure of the methodological quality of clinical trials: A demographic study. *Australian Journal of Physiotherapy*, 55(2), 129–133.
- Driessen, M. T., Proper, K. I., Van Tulder, M. W., Anema, J. R., Bongers, P. M., & Van Der Beek, A. J. (2010). The effectiveness of physical and organisational ergonomic interventions on low back pain and neck pain: A systematic review. *Occupational and Environmental Medicine*, 67(4), 277–285.
- Dunk, N., Kedgley, A., Jenkyn, T., & Callaghan, J. (2009). Evidence of a pelvis-driven flexion pattern: Are the joints of the lower lumbar spine fully flexed in seated postures? *Clinical Biomechanics*, 24(2), 164–168.
- Ellegast, R. P., Kraft, K., Groenesteijn, L., Krause, F., Berger, H., & Vink, P. (2012). Comparison of four specific dynamic office chairs with a conventional office chair: Impact upon muscle activation, physical activity and posture. *Applied Ergonomics*, 43(2), 296–307.
- Gadge, K., & Innes, E. (2007). An investigation into the immediate effects on comfort, productivity and posture of the Bambach™ saddle seat and a standard office chair. *Work*, 29(3), 189–203.
- Geisser, M. E., Ranavaya, M., Haig, A. J., Roth, R. S., Zucker, R., Ambroz, C., & Caruso, M. (2005). A meta-analytic review of surface electromyography among persons with low back pain and normal, healthy controls. *Journal of Pain*, 6(11), 711–726.
- Gregory, D., Dunk, N., & Callaghan, J. (2006). Stability ball versus office chair: Comparison of muscle activation and lumbar spine posture during prolonged sitting. *Human Factors*, 48, 142–153.
- Groenesteijn, L., Ellegast, R. P., Keller, K., Krause, F., Berger, H., & de Looze, M. P. (2012). Office task effects on comfort and body dynamics in five dynamic office chairs. *Applied Ergonomics*, 43(2), 320–328.
- Hardage, J. L., Gildersleeve, J., & Rugh, J. (1983). Clinical work posture for the dentist: An electromyographic study. *Journal of the American Dental Association*, 107(6), 937–939.
- Jarvik, J. G., Hollingworth, W., Heagerty, P. J., Haynor, D. R., Boyko, E. J., & Deyo, R. A. (2005). Three-year incidence of low back pain in an initially asymptomatic cohort: Clinical and imaging risk factors. *Spine*, 30(13), 1541–1548.
- Jensen, C. V., & Bendix, T. (1992). Spontaneous movements with various seated-workplace adjustments. *Clinical Biomechanics*, 7(2), 87–90.
- Keegan, J. J. (1953). Alterations of the lumbar curve related to posture and seating. *Journal of Bone & Joint Surgery*, 35(3), 589–603.
- Kingma, I., & van Dieen, J. H. (2009). Static and dynamic postural loadings during computer work in females: Sitting on an office chair versus sitting on an exercise ball. *Applied Ergonomics*, 40(2), 199–205.
- Koskelo, R., Vuorikari, K., & Hänninen, O. (2007). Sitting and standing postures are corrected by adjustable furniture with lowered muscle tension in high-school students. *Ergonomics*, 50(10), 1643–1656.
- Lander, C., Korbon, G., DeGood, D., & Rowlingson, J. (1987). The Balans chair and its semi-kneeling position: An ergonomic comparison with the conventional sitting position. *Spine*, 12(3), 269–272.
- Lengsfeld, M., König, I., Schmelter, J., & Ziegler, A. (2007). Passive rotary dynamic sitting at the workplace by office-workers with lumbar pain: A randomized multicenter study. *Spine Journal*, 7, 531–540.
- Linton, S. J., Hellsing, A.-L., Halme, T., & Åkerstedt, K. (1994). The effects of ergonomically designed school furniture on pupils' attitudes, symptoms and behaviour. *Applied Ergonomics*, 25(5), 299–304.
- Maher, C. G., Sherrington, C., Herbert, R. D., Moseley, A. M., & Elkins, M. (2003). Reliability of the PEDro scale for rating quality of randomized controlled trials. *Physical Therapy*, 83(8), 713–721.
- Main, C. J., Foster, N., & Buchbinder, R. (2010). How important are back pain beliefs and expectations for satisfactory recovery from back pain? *Best Practice & Research Clinical Rheumatology*, 24(2), 205–217.
- Mandal, A. C. (1983). The correct height of school furniture. *Physiotherapy Canada*, 35(4), 212–218.

- Mandal, A. (1987). The influence of furniture height on backpain. *Behaviour & Information Technology*, 6(3), 347–352.
- Marschall, M., Harrington, A. C., & Steele, J. R. (1995). Effect of work station design on sitting posture in young children. *Ergonomics*, 38(9), 1932–1940.
- Michel, D. P., & Helander, M. G. (1994). Effects of two types of chairs on stature change and comfort for individuals with healthy and herniated discs. *Ergonomics*, 37(7), 1231–1244.
- Mitchell, T., O'Sullivan, P. B., Burnett, A., Straker, L., Smith, A., Thornton, J., & Rudd, C. J. (2010). Identification of modifiable personal factors that predict new-onset low back pain: A prospective study of female nursing students. *Clinical Journal of Pain*, 26(4), 275–283.
- Moher, D., Liberati, A., Tetzlaff, J., & Altman, D. G. (2009). Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *PLOS Medicine*, 6(7), e1000097.
- Naqvi, S. (1994). Study of forward sloping seats for VDT workstations. *Journal of Human Ergology*, 23(1), 41–49.
- O'Keefe, M., Dankaerts, W., O'Sullivan, P., O'Sullivan, L., & O'Sullivan, K. (2013). Specific flexion-related low back pain and sitting: Comparison of seated discomfort on two different chairs. *Ergonomics*, 56(4), 650–658.
- Onen, S. H., Alloui, A., Gross, A., Eschallier, A., & Dubray, C. (2001). The effects of total sleep deprivation, selective sleep interruption and sleep recovery on pain tolerance thresholds in healthy subjects. *Journal of Sleep Research*, 10(1), 35–42.
- O'Sullivan, K., McCarthy, R., White, A., O'Sullivan, L., & Dankaerts, W. (2012a). Can we reduce the effort of maintaining a neutral sitting posture? A pilot study. *Manual Therapy*, 17(6), 566–571.
- O'Sullivan, K., McCarthy, R., White, A., O'Sullivan, L., & Dankaerts, W. (2012b). Lumbar posture and trunk muscle activation during a typing task when sitting on a novel dynamic ergonomic chair. *Ergonomics*, 55(12), 1586–1595.
- O'Sullivan, K., O'Keefe, M., O'Sullivan, L., O'Sullivan, P., & Dankaerts, W. (2012). The effect of dynamic sitting on the prevention and management of low back pain and low back discomfort: A systematic review. *Ergonomics*, 55(8), 898–908.
- O'Sullivan, K., O'Keefe, M., O'Sullivan, L., O'Sullivan, P., & Dankaerts, W. (2013). Perceptions of sitting posture among members of the community, both with and without non-specific chronic low back pain. *Manual Therapy*, 18(6), 551–556.
- O'Sullivan, K., O'Sullivan, P., O'Sullivan, L., & Dankaerts, W. (2012). What do physiotherapists consider to be the best sitting spinal posture? *Manual Therapy*, 17(5), 432–437.
- O'Sullivan, P. (2006). Effect of different upright sitting postures on spinal-pelvic curvature and trunk muscle activation in a pain-free population. *Spine*, 31(19), 707–712.
- O'Sullivan, P. (2012). It's time for change with the management of non-specific chronic low back pain. *British Journal of Sports Medicine*, 46(4), 224–227.
- O'Sullivan, P., Dankaerts, W., Burnett, A. F., Farrell, G. T., Jefford, E., Naylor, C. S., & O'Sullivan, K. J. (2006). Effect of different upright sitting postures on spinal-pelvic curvature and trunk muscle activation in a pain-free population. *Spine*, 31(19), E707–E712.
- PROSPERO. (2012). *Centre for reviews and dissemination*. Retrieved from <http://www.york.ac.uk>
- Pynt, J., Higgs, J., & Mackay, M. (2001). Seeking the optimal posture of the seated lumbar spine. *Physiotherapy Theory and Practice*, 17, 5–21.
- Ramond, A., Bouton, C., Richard, I., Roquelaure, Y., Baufreton, C., Legrand, E., & Huez, J.-F. (2011). Psychosocial risk factors for chronic low back pain in primary care—A systematic review. *Family Practice*, 28, 12–21.
- Reichborn-Kjennerud, T., Stoltenberg, C., Tambs, K., Roysamb, E., Kringlen, E., Torgersen, S., & Harris, J. (2002). Back-neck pain and symptoms of anxiety and depression: A population-based twin study. *Psychological Medicine*, 32(6), 1009–1020.
- Reinecke, S., Hazard, R., & Coleman, K. (1994). Continuous passive motion in seating: A new strategy against low back pain. *Journal of Spinal Disorders & Techniques*, 7, 29–35.
- Roffey, D., Wai, E., Bishop, P., Kwon, B., & Dagenais, S. (2010). Causal assessment of occupational sitting and low back pain: Results of a systematic review. *Spine Journal*, 10(3), 252–261.
- Saarni, L., Nygård, C.-H., Nummi, T., Kaukiainen, A., & Rimpelä, A. (2009). Comparing the effects of two school workstations on spine positions and mobility, and opinions on the workstations—A 2-year controlled intervention. *International Journal of Industrial Ergonomics*, 39(6), 981–987.
- Saarni, L., Nygård, C.-H., Rimpelä, A., Nummi, T., & Kaukiainen, A. (2007). The working postures among schoolchildren—Controlled intervention study on the effects of newly designed workstations. *Journal of School Health*, 77(5), 240–247.
- Soderberg, G. L., Blanco, M. K., Cosentino, T. L., & Kurldelmeier, K. A. (1986). An EMG analysis of posterior trunk musculature during flat and anteriorly inclined sitting. *Human Factors*, 28, 483–491.
- Troussier, B. (1999). Comparative study of two different kinds of school furniture among children. *Ergonomics*, 42(3), 516–526.
- van der Heide, J. C., Otten, B., van Eykern, L. A., & Hadders-Algra, M. (2003). Development of postural adjustments during reaching in sitting children. *Experimental Brain Research*, 151(1), 32–45.
- Van Niekerk, S. M. S., Louw, Q. Q. A., & Hillier, S. S. (2012). The effectiveness of a chair intervention in the workplace to reduce musculoskeletal symptoms. A systematic review. *BMC Musculoskeletal Disorders*, 13(1), 145.
- Vergara, M., & Page, Á. (2000a). System to measure the use of the backrest in sitting-posture office tasks. *Applied Ergonomics*, 31(3), 247–254.
- Vergara, M., & Page, Á. (2000b). Technique to measure lumbar curvature in the ergonomic evaluation of chairs: Description and validation. *Clinical Biomechanics*, 15(10), 786–789.
- Verhagen, A. P., de Vet, H. C. W., de Bie, R. A., Kessels, A. G. H., Boers, M., Bouter, L. M., & Knipschild, P. G. (1998). The Delphi list: A criteria list for quality assessment of randomized clinical trials for conducting systematic reviews developed by Delphi consensus. *Journal of Clinical Epidemiology*, 51(12), 1235–1241.
- Wand, B., Parkitny, L., O'Connell, N., Luomajoki, L., McAuley, J., Thacker, M., & Moseley, G. L. (2011). Cortical changes in chronic low back pain: Current state of the art and implications for clinical practice. *Manual Therapy*, 16(1), 15–20.
- Wang, P.-C., Ritz, B. R., Janowitz, I., Harrison, R. J., Yu, F., Chan, J., & Rempel, D. M. (2008). A randomized controlled trial of chair interventions on back and hip pain among sewing machine operators: The Los Angeles garment study. *Journal of Occupational and Environmental Medicine*, 50(3), 255–262.
- Williams, M. M., Hawley, J. A., McKenzie, R. A., & Van Wijmen, P. M. (1991). A comparison of the effects of two sitting postures on back and referred pain. *Spine*, 16, 1185–1191.
- Winkel, J., & Bendix, T. (1986). Muscular performance during seated work evaluated by two different EMG methods. *European Journal of Applied Physiology and Occupational Physiology*, 55(2), 167–173.

- Womersley, L., & May, S. (2006). Sitting posture of subjects with postural backache. *Journal of Manipulative and Physiological Therapeutics*, 29(3), 213–218.
- Woolf, A., & Pfleger, B. (2003). Burden of major musculoskeletal conditions. *Bulletin of the World Health Organisation*, 81(9), 646–656.
- Wu, C.-S., Miyamoto, H., & Noro, K. (1998). Research on pelvic angle variation when using a pelvic support. *Ergonomics*, 41(3), 317–327.
- Ye, L., Kalichman, L., Spittle, A., Dobson, F., & Bennell, K. (2011). Effects of rehabilitative interventions on pain, function and physical impairments in people with hand osteoarthritis: A systematic review. *Arthritis Research & Therapy*, 13(1), R28.
- Yoo, W. G., Yi, C. H., Cho, S. H., Jeon, H., Cynn, H., & Choi, H. (2008). Effects of the height of ball-backrest on head and shoulder posture and trunk muscle activity in VDT workers. *Industrial Health*, 46(3), 289–297.
- Yu, C.-Y., Keyserling, W. M., & Chaffin, D. B. (1988). Development of a work seat for industrial sewing operations: Results of a laboratory study. *Ergonomics*, 31(12), 1765–1786.

Máire Curran is a physiotherapist working in clinical practice. She earned her BSc (Hons) in physiotherapy from the University of Limerick, Ireland in 2013.

Leonard O’Sullivan is a senior lecturer in the Department of Design and Manufacturing Technology at the University of Limerick, Ireland. He earned his PhD in ergonomics from the University of Limerick in 2002.

Peter O’Sullivan is professor of musculoskeletal physiotherapy at Curtin University, Perth, Australia. He earned his PhD in physiotherapy from Curtin University in 1998.

Wim Dankaerts is a professor at the University of Leuven, Belgium. He earned his PhD in physiotherapy from Curtin University in 2005.

Kieran O’Sullivan is a physiotherapy lecturer in the Department of Clinical Therapies at the University of Limerick, Ireland. He earned his PhD in physiotherapy from the University of Limerick in 2012.

Date received: September 20, 2014

Date accepted: May 15, 2015