

Small exotic 4–manifolds

ANAR AKHMEDOV

In this article, we construct the first example of a simply-connected minimal symplectic 4–manifold that is homeomorphic but not diffeomorphic to $3\mathbb{C}\mathbb{P}^2 \# 7\overline{\mathbb{C}\mathbb{P}^2}$. We also construct the first exotic minimal *symplectic* $\mathbb{C}\mathbb{P}^2 \# 5\overline{\mathbb{C}\mathbb{P}^2}$.

57N65, 57N13; 57M50

1 Introduction

Over the past several years, there has been a considerable amount of progress in the discovery of exotic smooth structures on simply-connected 4–manifolds with small Euler characteristic. In early 2004, Jongil Park [15] has constructed the first example of exotic smooth structure on $\mathbb{C}\mathbb{P}^2 \# 7\overline{\mathbb{C}\mathbb{P}^2}$, ie 4–manifold homeomorphic but not diffeomorphic to $\mathbb{C}\mathbb{P}^2 \# 7\overline{\mathbb{C}\mathbb{P}^2}$. Later that year, András Stipsicz and Zoltán Szabó used a similar technique to construct an exotic smooth structure on $\mathbb{C}\mathbb{P}^2 \# 6\overline{\mathbb{C}\mathbb{P}^2}$ [18]. Then Fintushel and Stern [5] introduced a new technique, the double node surgery, which demonstrated that in fact $\mathbb{C}\mathbb{P}^2 \# k\overline{\mathbb{C}\mathbb{P}^2}$, $k = 6, 7$ and 8 have infinitely many distinct smooth structures. Using the double node surgery technique [5], Park, Stipsicz and Szabó constructed infinitely many smooth structures on $\mathbb{C}\mathbb{P}^2 \# 5\overline{\mathbb{C}\mathbb{P}^2}$ [17]. The examples in [17] are not known if symplectic. Based on similar ideas, Stipsicz and Szabó constructed the exotic smooth structures on $3\mathbb{C}\mathbb{P}^2 \# k\overline{\mathbb{C}\mathbb{P}^2}$ for $k = 9$ [19] and Park for $k = 8$ [16]. In this article, we construct an exotic smooth structure on $3\mathbb{C}\mathbb{P}^2 \# 7\overline{\mathbb{C}\mathbb{P}^2}$. We also construct an exotic *symplectic* $\mathbb{C}\mathbb{P}^2 \# 5\overline{\mathbb{C}\mathbb{P}^2}$, the first known such symplectic example.

Our approach is different from the above constructions in the sense that we do not use any rational-blowdown surgery (Fintushel and Stern [3], Jongil [14]). Also, in contrary to the previous constructions, we use non-simply connected building blocks (Akhmedov [1], Matsumoto [11]) to produce the simply-connected examples. The main surgery technique used in our construction is the symplectic fiber sum operation (Gompf [7], McCarthy and Wolfson [12]) along the genus two surfaces. Our results can be stated as follows.

Theorem 1.1 *There exist a smooth closed simply-connected minimal symplectic 4–manifold X that is homeomorphic but not diffeomorphic to $3\mathbb{C}\mathbb{P}^2\#7\overline{\mathbb{C}\mathbb{P}^2}$.*

Theorem 1.2 *There exist a smooth closed simply-connected minimal symplectic 4–manifold Y which is homeomorphic but not diffeomorphic to the rational surface $\mathbb{C}\mathbb{P}^2\#5\overline{\mathbb{C}\mathbb{P}^2}$.*

This article is organized as follows. The first two sections give a quick introduction to Seiberg–Witten invariants and a fiber sum operation. In [Section 4](#), we review the symplectic building blocks for our construction. Finally, in [Section 5](#) and [Section 6](#), we construct minimal symplectic 4–manifolds X and Y homeomorphic but not diffeomorphic to $3\mathbb{C}\mathbb{P}^2\#7\overline{\mathbb{C}\mathbb{P}^2}$ and $\mathbb{C}\mathbb{P}^2\#5\overline{\mathbb{C}\mathbb{P}^2}$, respectively.

Acknowledgments I would like to thank John Etnyre, Ron Stern and András Stipsicz for their interest in this work and for their encouragement. Also, I am grateful to B Doug Park for the comments on the first draft of this article, kindly pointing out some errors in the fundamental group computations and for the corrections. Finally, I wish to thank the referee for many helpful suggestions which improved the exposition of this article. This work is partially supported by NSF grant FRG-0244663.

Dedication Dedicated to Professor Ronald J Stern on the occasion of his sixtieth birthday.

2 Seiberg–Witten Invariants

In this section, we briefly recall the basics of Seiberg–Witten invariants introduced by Seiberg and Witten. Seiberg–Witten invariant of a smooth closed oriented 4–manifold X with $b_2^+(X) > 1$ is an integer valued function which is defined on the set of spin^c structures over X (Witten [\[23\]](#)). For simplicity, we assume that $H_1(X, \mathbf{Z})$ has no 2–torsion. Then there is a one-to-one correspondence between the set of spin^c structures over X and the set of characteristic elements of $H^2(X, \mathbf{Z})$.

In this set up, we can view the Seiberg–Witten invariant as an integer valued function

$$\text{SW}_X: \{k \in H^2(X, \mathbf{Z}) \mid k \equiv w_2(TX) \pmod{2}\} \longrightarrow \mathbf{Z}.$$

The Seiberg–Witten invariant SW_X is a diffeomorphism invariant. We call β a *basic class* of X if $\text{SW}_X(\beta) \neq 0$. It is a fundamental fact that the set of basic classes is finite. Also, if β is a basic class, then so is $-\beta$ with

$$\text{SW}_X(-\beta) = (-1)^{(e+\sigma)(X)/4} \text{SW}_X(\beta)$$

where $e(X)$ is the Euler characteristic and $\sigma(X)$ is the signature of X .

Theorem 2.1 (Taubes [20]) *Suppose that (X, ω) is a closed symplectic 4-manifold with $b_2^+(X) > 1$ and the canonical class K_X . Then $SW_X(\pm K_X) = \pm 1$.*

3 Fiber Sum

Definition 3.1 Let X and Y be closed, oriented, smooth 4-manifolds each containing a smoothly embedded surface Σ of genus $g \geq 1$. Assume Σ represents a homology class of infinite order and has self-intersection zero in X and Y , so that there exist a tubular neighborhood, say $\nu\Sigma \cong \Sigma \times D^2$, in both X and Y . Using an orientation-reversing and fiber-preserving diffeomorphism $\psi: S^1 \times \Sigma \rightarrow S^1 \times \Sigma$, we can glue $X \setminus \nu\Sigma$ and $Y \setminus \nu\Sigma$ along the boundary $\partial(\nu\Sigma) \cong \Sigma \times S^1$. This new oriented smooth 4-manifold $X \#_\psi Y$ is called a *generalized fiber sum* of X and Y along Σ , determined by ψ .

Definition 3.2 Let $e(X)$ and $\sigma(X)$ denote the Euler characteristic and the signature of a closed oriented smooth 4-manifold X , respectively. We define

$$c_1^2(X) := 2e(X) + 3\sigma(X), \quad \chi_h(X) := \frac{e(X) + \sigma(X)}{4}.$$

In the case that X is a complex surface, then $c_1^2(X)$ and $\chi_h(X)$ are the self-intersection of the first Chern class $c_1(X)$ and the holomorphic Euler characteristic, respectively.

Lemma 3.3 *Let X and Y be closed, oriented, smooth 4-manifolds containing an embedded surface Σ of self-intersection 0. Then*

$$\begin{aligned} c_1^2(X \#_\psi Y) &= c_1^2(X) + c_1^2(Y) + 8(g - 1), \\ \chi_h(X \#_\psi Y) &= \chi_h(X) + \chi_h(Y) + (g - 1), \end{aligned}$$

where g is the genus of the surface Σ .

Proof The above simply follows from the well-known formulas

$$e(X \#_\psi Y) = e(X) + e(Y) - 2e(\Sigma), \quad \sigma(X \#_\psi Y) = \sigma(X) + \sigma(Y). \quad \square$$

If X, Y are symplectic manifolds and Σ is an embedded symplectic submanifold in X and Y , then according to theorem of Gompf [7] $X \#_\psi Y$ admits a symplectic structure.

We will use the following theorem of M Usher [21] to show that the symplectic manifolds constructed in Section 5 and Section 6 are minimal. Here we slightly abuse the above notation for the fiber sum.

Theorem 3.4 (Usher [21], Minimality of Symplectic Sums) *Let $X = X_1 \#_{F_1=F_2} X_2$ be symplectic fiber sum of manifolds X_1 and X_2 .*

- (i) *If either $X_1 \setminus F_1$ or $X_2 \setminus F_2$ contains an embedded symplectic sphere of square -1 , then X is not minimal.*
- (ii) *If one of the summands X_i (say X_1) admits the structure of an S^2 -bundle over a surface of genus g such that F_i is a section of this fiber bundle, then X is minimal if and only if X_2 is minimal.*
- (iii) *In all other cases, X is minimal.*

4 Building blocks

The building blocks for our construction will be as follows.

- (i) The manifold $T^2 \times S^2 \# 4\overline{\mathbb{C}\mathbb{P}^2}$ equipped with the genus two Lefschetz fibration of Matsumoto [11].
- (ii) The symplectic manifolds X_K and Y_K [1]. For the convenience of the reader, we recall the construction in [1].

4.1 Matsumoto fibration

First, recall that the manifold $Z = T^2 \times S^2 \# 4\overline{\mathbb{C}\mathbb{P}^2}$ can be described as the double branched cover of $S^2 \times T^2$ where the branch set $B_{2,2}$ is the union of two disjoint copies of $S^2 \times \{\text{pt}\}$ and two disjoint copies of $\{\text{pt}\} \times T^2$. The branch cover has 4 singular points, corresponding to the number of the intersections points of the horizontal lines and the vertical tori in the branch set $B_{2,2}$. After desingularizing the above singular manifold, one obtains $T^2 \times S^2 \# 4\overline{\mathbb{C}\mathbb{P}^2}$. The vertical fibration of $S^2 \times T^2$ pulls back to give a fibration of $T^2 \times S^2 \# 4\overline{\mathbb{C}\mathbb{P}^2}$ over S^2 . A generic fiber of the vertical fibration is the double cover of T^2 , branched over 2 points. Thus a generic fiber will be a genus two surface. According to Matsumoto [11], this fibration can be perturbed to be a Lefschetz fibration over S^2 with the global monodromy $(\beta_1\beta_2\beta_3\beta_4)^2 = 1$, where the curves $\beta_1, \beta_2, \beta_3$ and β_4 are shown in Figure 1.

Let us denote the regular fiber by Σ'_2 and the images of standard generators of the fundamental group of Σ'_2 as a_1, b_1, a_2 and b_2 . Using the homotopy exact sequence for a Lefschetz fibration,

$$\pi_1(\Sigma'_2) \longrightarrow \pi_1(Z) \longrightarrow \pi_1(S^2)$$




Figure 1: Dehn Twists for Matsumoto Fibration

we have the following identification of the fundamental group of Z [13]:

$$\pi_1(Z) = \pi_1(\Sigma'_2) / \langle \beta_1, \beta_2, \beta_3, \beta_4 \rangle.$$

- (1) $\beta_1 = b_1 b_2,$
- (2) $\beta_2 = a_1 b_1 a_1^{-1} b_1^{-1} = a_2 b_2 a_2^{-1} b_2^{-1},$
- (3) $\beta_3 = b_2 a_2 b_2^{-1} a_1,$
- (4) $\beta_4 = b_2 a_2 a_1 b_1.$

Hence $\pi_1(Z) = \langle a_1, b_1, a_2, b_2 \mid b_1 b_2 = [a_1, b_1] = [a_2, b_2] = a_1 a_2 = 1 \rangle.$

Note that the fundamental group of $T^2 \times S^2 \# 4\overline{\mathbb{C}\mathbb{P}^2}$ is $\mathbf{Z} \oplus \mathbf{Z}$, generated by two of these standard generators (say a_1 and b_1). The other two generators a_2 and b_2 are the inverses of a_1 and b_1 in the fundamental group. Also, the fundamental group of the complement of $\nu\Sigma'_2$ is $\mathbf{Z} \oplus \mathbf{Z}$. It is generated by a_1 and b_1 . The normal circle $\lambda' = pt \times \partial D^2$ to Σ'_2 can be deformed using one of the exceptional spheres, thus is trivial in $\pi_1(T^2 \times S^2 \# 4\overline{\mathbb{C}\mathbb{P}^2} \setminus \nu\Sigma'_2) = \mathbf{Z} \oplus \mathbf{Z}$.

Lemma 4.1 $c_1^2(Z) = -4, \sigma(Z) = -4$ and $\chi_h(Z) = 0$.

Proof We have $c_1^2(Z) = c_1^2(T^2 \times S^2) - 4 = -4, \sigma(Z) = \sigma(T^2 \times S^2) - 4 = -4$ and $\chi_h(Z) = \chi_h(T^2 \times S^2) = 0$. □

Note that this Lefschetz fibration can be given a symplectic structure. This means that Z admits a symplectic structure such that the regular fibers are symplectic submanifolds. We consider such a symplectic structure on Z .

4.2 Symplectic 4-manifolds cohomology equivalent to $S^2 \times S^2$

Our second building block will be X_K , the symplectic cohomology $S^2 \times S^2$ [1], or the symplectic manifold Y_K , an intermediate building block in that construction [1], (see also Fintushel and Stern [4]). For the sake of completeness, the details of this construction are included below. We refer the reader to [1] for more details and for the generalization of these symplectic building blocks.

Let K be a fibered knot of genus one (ie, the trefoil or the figure eight knot) in S^3 and m be a meridional circle to K . We perform 0-framed surgery on K and denote the resulting 3-manifold by M_K . Since K is fibered and has genus one, it follows the 3-manifold M_K is a torus bundle over S^1 ; hence the 4-manifold $M_K \times S^1$ is a torus bundle over a torus. Furthermore, $M_K \times S^1$ admits a symplectic structure, and both the torus fiber and the torus section $T_m = m \times S^1 = m \times x$ are symplectically embedded and have a self-intersection zero. The first homology of $M_K \times S^1$ is generated by the standard first homology generators m and x of the torus section. On the other hand, the classes of circles γ_1 and γ_2 of the fiber F , coming from the Seifert surface, are trivial in homology. In addition, $M_K \times S^1$ is minimal symplectic, ie, it does not contain symplectic -1 sphere.

We form a twisted fiber sum of two copies of the manifold $M_K \times S^1$, we identify the fiber F of one fibration to the section T_m of other. Let Y_K denote the mentioned twisted fiber sum $Y_K = M_K \times S^1 \#_{F=T_m} M_K \times S^1$. It follows from Gompf's theorem [7] that Y_K is symplectic and by Usher's Theorem 3.4 that Y_K is minimal symplectic.

Let T_1 be the section of the first copy of $M_K \times S^1$ and T_2 be the fiber in the second copy. Then the genus two surface $\Sigma_2 = T_1 \# T_2$ symplectically embeds into Y_K and has self-intersection zero. Let X_K be a symplectic 4-manifold constructed as follows: Take two copies of Y_K and form the fiber sum along the genus two surface Σ_2 using the special gluing diffeomorphism ϕ , the vertical involution of Σ_2 with two fixed points. Thus $X_K := Y_K \#_{\phi} Y_K$. Let m, x, γ_1 and γ_2 denote the generators of $\pi_1(\Sigma_2)$ under the inclusion. The diffeomorphism $\phi: T_1 \# T_2 \rightarrow T_1 \# T_2$ of Σ_2 maps on the generators as follows: $\phi_*(m') = \gamma_1$, $\phi_*(x') = \gamma_2$, $\phi_*(\gamma_1') = m$ and $\phi_*(\gamma_2') = x$. In [1] we show that the manifold X_K has first Betti number zero and has the integral cohomology of $S^2 \times S^2$. Furthermore, $H_2(X_K, \mathbf{Z}) = \mathbf{Z} \oplus \mathbf{Z}$, where the basis for the second homology are the classes of $\Sigma_2 = S$ and the new genus two surface T resulting from the last fiber sum operation (two punctured genus one surfaces glues to form a genus two surface). Also, $S^2 = T^2 = 0$ and $S \cdot T = 1$. Furthermore, $c_1^2(X_K) = 8$, $\sigma(X_K) = 0$ and $\chi_h(X_K) = 1$. Since Y_K is minimal symplectic, it follows from Theorem 3.4 that X_K is minimal symplectic as well.

4.2.1 Fundamental Group of $M_K \times S^1$ We will assume that K is the trefoil knot. Let a, b and c denote the Wirtinger generators of the trefoil. The knot group of the trefoil has the following presentations: $\pi_1(K) = \langle a, b, c \mid ab = bc, ca = ab \rangle = \langle a, b \mid aba = bab \rangle = \langle u, v \mid u^2 = v^3 \rangle$ where $u = bab$ and $v = ab$. The homotopy classes of the meridian and the longitude of the trefoil are given as follows: $m = uv^{-1} = b$ and $l = u^2(uv^{-1})^{-6} = ab^2ab^{-4}$ (Burde and Zieschang [2]). Also, the homotopy classes of γ_1 and γ_2 are given as follows: $\gamma_1 = a^{-1}b$ and $\gamma_2 = b^{-1}aba^{-1}$. Notice that the fundamental group of M_K , 0-surgery on the trefoil, is obtained from the knot group of the trefoil by adjoining the relation $l = u^2(uv^{-1})^{-6} = ab^2ab^{-4} = 1$. Thus, we have $\pi_1(M_K) = \langle u, v \mid u^2 = v^3, u^2(uv^{-1})^{-6} = 1 \rangle = \langle a, b \mid aba = bab, ab^2a = b^4 \rangle$ and $\pi_1(M_K \times S^1) = \langle a, b, x \mid aba = bab, ab^2a = b^4, [x, a] = [x, b] = 1 \rangle$.

4.2.2 Fundamental Group of Y_K The next step is to take two copies of the manifold $M_K \times S^1$ and perform the fiber sum along symplectic tori. In the first copy of $M_K \times S^1$, we take a tubular neighborhood of the torus section T_m , remove it from $M_K \times S^1$ and denote the resulting manifold by C_S . In the second copy, we remove a tubular neighborhood of the fiber F and denote it by C_F . Notice that $C_S = M_K \times S^1 \setminus \nu T_m = (M_K \setminus \nu(m)) \times S^1$. We have $\pi_1(C_S) = \pi_1(K) \oplus \langle x \rangle$ where x is the generator corresponding to the S^1 copy. Also using the above computation, we easily derive: $H_1(C_S) = H_1(M_K) = \langle m \rangle \oplus \langle x \rangle$.

To compute the fundamental group of the C_F , we will use the following observation: νF is the preimage of the small disk on $T_{m'} = m' \times y$. The elements y and $m' = d$ of the $\pi_1(C_F)$ do not commute anymore, but y still commutes with generators γ'_1 and γ'_2 . The fundamental group and the first homology of the C_F will be isomorphic to the following: $\pi_1(C_F) = \langle d, y, \gamma'_1, \gamma'_2 \mid [y, \gamma'_1] = [y, \gamma'_2] = [\gamma'_1, \gamma'_2] = 1, d\gamma'_1d^{-1} = \gamma'_1\gamma'_2, d\gamma'_2d^{-1} = (\gamma'_1)^{-1} \rangle$ and $H_1(C_F) = \langle d \rangle \oplus \langle y \rangle$.

We use the Van Kampen's Theorem to compute the fundamental group of Y_K .

$$\begin{aligned} \pi_1(Y_K) &= \pi_1(C_F) *_{\pi_1(T^3)} \pi_1(C_S) \\ &= \langle d, y, \gamma'_1, \gamma'_2 \mid [y, \gamma'_1] = [y, \gamma'_2] = [\gamma'_1, \gamma'_2] = 1, d\gamma'_1d^{-1} = \gamma'_1\gamma'_2, d\gamma'_2d^{-1} \\ &= (\gamma'_1)^{-1} \rangle_{\langle \gamma'_1=x, \gamma'_2=b, \lambda'=\lambda \rangle} \langle a, b, x \mid aba = bab, [x, a] = [x, b] = 1 \rangle \\ &= \langle a, b, x, \gamma'_1, \gamma'_2, d, y \mid aba = bab, [x, a] = [x, b][y, \gamma'_1] = [y, \gamma'_2] \\ &= [\gamma'_1, \gamma'_2] = 1, d\gamma'_1d^{-1} = \gamma'_1\gamma'_2, d\gamma'_2d^{-1} = (\gamma'_1)^{-1}, \gamma'_1 = x, \\ &\gamma'_2 = b, [\gamma'_1, \gamma'_2] = [d, y] \rangle. \end{aligned}$$

Inside Y_K , we can find a genus 2 symplectic submanifold Σ_2 which is the internal sum of a punctured fiber in C_S and a punctured section in C_F . The inclusion-induced

homomorphism maps the standard generators of $\pi_1(\Sigma_2)$ to $a^{-1}b, b^{-1}aba^{-1}, d$ and y in $\pi_1(Y_K)$.

Lemma 4.2 ([1]) *There are nonnegative integers m and n such that*

$$\begin{aligned} \pi_1(Y_K \setminus \nu\Sigma_2) = \langle a, b, x, d, y; g_1, \dots, g_m \mid &aba = bab, \\ [y, x] = [y, b] = 1, dxd^{-1} = xb, dbd^{-1} = x^{-1}, & \\ ab^2ab^{-4} = [d, y], r_1 = \dots = r_n = 1, r_{n+1} = 1 \rangle, & \end{aligned}$$

where the generators g_1, \dots, g_m and relators r_1, \dots, r_n all lie in the normal subgroup N generated by the element $[x, b]$ and the relator r_{n+1} is a word in x, a and elements of N . Moreover, if we add an extra relation $[x, b] = 1$, then the relation $r_{n+1} = 1$ simplifies to $[x, a] = 1$.

Proof This follows from Van Kampen’s Theorem. Note that $[x, b]$ is a meridian of Σ_2 in Y_K . Hence setting $[x, b] = 1$ should turn $\pi_1(Y_K \setminus \nu\Sigma_2)$ into $\pi_1(Y_K)$. Also note that $[x, a]$ is the boundary of a punctured section in $C_S \setminus \nu(\text{fiber})$ and is no longer trivial in $\pi_1(Y_K \setminus \nu\Sigma_2)$. By setting $[x, b] = 1$, the relation $r_{n+1} = 1$ is to turn into $[x, a] = 1$.

It remains to check that the relations in $\pi_1(Y_K)$ other than $[x, a] = [x, b] = 1$ remain the same in $\pi_1(Y_K \setminus \nu\Sigma_2)$. By choosing a suitable point $\theta \in S^1$ away from the image of the fiber that forms part of Σ_2 , we obtain an embedding of the knot complement $(S^3 \setminus \nu K) \times \{\theta\} \hookrightarrow C_S \setminus \nu(\text{fiber})$. This means that $aba = bab$ holds in $\pi_1(Y_K \setminus \nu\Sigma_2)$. Since $[\Sigma_2]^2 = 0$, there exists a parallel copy of Σ_2 outside $\nu\Sigma_2$, wherein the identity $ab^2ab^{-4} = [d, y]$ still holds. The other remaining relations in $\pi_1(Y_K)$ are coming from the monodromy of the torus bundle over a torus. Since these relations will now describe the monodromy of the punctured torus bundle over a punctured torus, they hold true in $\pi_1(Y_K \setminus \nu\Sigma_2)$. □

4.2.3 Fundamental Group of X_K Finally, we carry out the computations of the fundamental group and the first homology of X_K . Suppose that e, f, z, s and t are the generators of the fundamental group in the second copy of Y_K corresponding to the generators a, b, x, d and y as in above discussion. Our gluing map ϕ maps the generators of $\pi_1(\Sigma_2)$ as follows:

$$\phi_*(a^{-1}b) = s, \phi_*(b^{-1}aba^{-1}) = t, \phi_*(d) = e^{-1}f, \phi_*(y) = f^{-1}efe^{-1}.$$

By Van Kampen’s Theorem and Lemma 4.2, we have

$$\begin{aligned} \pi_1(X_K) = \langle a, b, x, d, y; e, f, z, s, t; g_1, \dots, g_m; h_1, \dots, h_m \mid \\ aba = bab, [y, x] = [y, b] = 1, \\ dx d^{-1} = xb, db d^{-1} = x^{-1}, ab^2 ab^{-4} = [d, y], \\ r_1 = \dots = r_{n+1} = 1, r'_1 = \dots = r'_{n+1} = 1, \\ efe = fef, [t, z] = [t, f] = 1, \\ szs^{-1} = zf, sfs^{-1} = z^{-1}, ef^2 ef^{-4} = [s, t], \\ d = e^{-1} f, y = f^{-1} efe^{-1}, a^{-1} b = s, b^{-1} aba^{-1} = t, \\ [x, b] = [z, f] \rangle, \end{aligned}$$

where the elements g_i, h_i ($i = 1, \dots, m$) and r_j, r'_j ($j = 1, \dots, n + 1$) all are in the normal subgroup generated by $[x, b] = [z, f]$.

Notice that it follows from our gluing that the images of standard generators of the fundamental group of Σ_2 are $a^{-1} b, b^{-1} aba^{-1}, d$ and y in $\pi_1(X_K)$. By abelianizing $\pi_1(X_K)$, we easily see that $H_1(X_K, \mathbf{Z}) = 0$.

5 Construction of an exotic $3\mathbb{C}P^2 \# 7\overline{\mathbb{C}P^2}$

In this section, we construct a simply-connected minimal symplectic 4-manifold X homeomorphic but not diffeomorphic to $3\mathbb{C}P^2 \# 7\overline{\mathbb{C}P^2}$. Using Seiberg–Witten invariants, we will distinguish X from $3\mathbb{C}P^2 \# 7\overline{\mathbb{C}P^2}$.

Our manifold X will be the symplectic fiber sum of X_K and $Z = T^2 \times S^2 \# 4\overline{\mathbb{C}P^2}$ along the genus two surfaces Σ_2 and Σ'_2 . Recall that $a^{-1} b, b^{-1} aba^{-1}, d, y$ and $\lambda = \{pt\} \times S^1 = [x, b][z, f]^{-1}$ generate the inclusion-induced image of $\pi_1(\Sigma_2 \times S^1)$ inside $\pi_1(X_K \setminus \nu\Sigma_2)$. Let a_1, b_1, a_2, b_2 and $\lambda' = 1$ be generators of $\pi_1(Z \setminus \nu\Sigma'_2)$ as in Section 4.1. We choose the gluing diffeomorphism $\psi: \Sigma_2 \times S^1 \rightarrow \Sigma'_2 \times S^1$ that maps the fundamental group generators as follows:

$$\psi_*(a^{-1} b) = a_2, \psi_*(b^{-1} aba^{-1}) = b_2, \psi_*(d) = a_1, \psi_*(y) = b_1, \psi_*(\lambda) = \lambda'.$$

λ and λ' above denote the meridians of Σ and Σ'_2 in X_K and Z , respectively.

It follows from Gompf’s theorem [7] that $X = X_K \#_{\psi} (T^2 \times S^2 \# 4\overline{\mathbb{C}P^2})$ is symplectic.

Lemma 5.1 X is simply connected.

Proof By Van Kampen’s theorem, we have

$$\pi_1(X) = \frac{\pi_1(X_K \setminus \nu\Sigma_2) * \pi_1(Z \setminus \nu\Sigma'_2)}{\langle a^{-1}b = a_2, b^{-1}aba^{-1} = b_2, d = a_1, y = b_1, \lambda = 1 \rangle}.$$

Since λ' is nullhomotopic in $Z \setminus \nu\Sigma'_2$, the normal circle λ of $\pi_1(X_K \setminus \nu\Sigma_2)$ becomes trivial in $\pi_1(X)$. Also, using the relations $b_1b_2 = [a_1, b_1] = [a_2, b_2] = b_2a_2b_2^{-1}a_1 = a_1a_2 = 1$ in $\pi_1(Z \setminus \nu\Sigma'_2)$, we get the following relations in the fundamental group of X : $a^{-1}bd = [a^{-1}b, b^{-1}aba^{-1}] = [d, y] = [d, b^{-1}aba^{-1}] = yb^{-1}aba^{-1} = 1$. Note that the fundamental group of Z is an abelian group of rank two. In addition, we have the following relations in $\pi_1(X)$ coming from the fundamental group of $\pi_1(X_K \setminus \nu\Sigma_2)$: $aba = bab, efe = fef, [y, b] = [t, f] = 1, dbd^{-1} = x^{-1}, dxd^{-1} = xb, sfs^{-1} = z^{-1}, szs^{-1} = zf, a^{-1}b = s, b^{-1}aba^{-1} = t, y = f^{-1}efe^{-1}$ and $e^{-1}f = d$. These set of relations give rise to the following identities:

- (5) $yab = ba,$
- (6) $a = bd,$
- (7) $yb = by,$
- (8) $aba = bab.$

Next, multiply the relation (5) by a from the right and use $aba = bab$. We have $yaba = ba^2 \implies ybab = ba^2$. By cancelling the element b , we obtain $yab = a^2$. Finally, applying the relation (5) again, we have $ba = a^2$. The latter implies that $b = a$. Since $a = bd, dbd^{-1} = x^{-1}, dxd^{-1} = xb, aba = bab$ and $yb^{-1}aba^{-1} = 1$, we obtain $d = y = x = b = a = 1$. Furthermore, using the relations $a^{-1}b = s, b^{-1}aba^{-1} = t, efe = fef, e^{-1}f = d, sfs^{-1} = z^{-1}$ and $szs^{-1} = zf$, we similarly have $s = t = z = f = e = 1$. Thus, we can conclude that the elements $a, b, x, d, y, e, f, z, s$ and t are all trivial in the fundamental group of X . Since we identified $a^{-1}b$ and $b^{-1}aba^{-1}$ with generators a_2 and b_2 of the group $\pi_1(Z \setminus \nu\Sigma'_2) = \mathbf{Z} \oplus \mathbf{Z}$, it follows that a_2 and b_2 are trivial in the fundamental group of X as well. This proves that X is simply connected. □

Lemma 5.2 $c_1^2(X) = 12, \sigma(X) = -4$ and $\chi_h(X) = 2$.

Proof We have $c_1^2(X) = c_1^2(X_K) + c_1^2(T^2 \times S^2 \# 4\overline{\mathbb{C}\mathbb{P}^2}) + 8, \sigma(X) = \sigma(X_K) + \sigma(T^2 \times S^2 \# 4\overline{\mathbb{C}\mathbb{P}^2})$ and $\chi_h(X) = \chi_h(X_K) + \chi_h(T^2 \times S^2 \# 4\overline{\mathbb{C}\mathbb{P}^2}) + 1$. Since $c_1^2(X_K) = 8, \sigma(X_K) = 0$ and $\chi_h(X_K) = 1$, the result follows from [Lemma 3.3](#) and [Lemma 4.1](#). □

By Freedman’s theorem [6], Lemma 5.1 and Lemma 5.2, X is homeomorphic to $3\mathbb{C}\mathbb{P}^2\#7\overline{\mathbb{C}\mathbb{P}^2}$. It follows from Taubes Theorem 2.1 that $\text{SW}_X(K_X) = \pm 1$. Next we apply the connected sum theorem for the Seiberg–Witten invariant and show that SW function is trivial for $3\mathbb{C}\mathbb{P}^2\#7\overline{\mathbb{C}\mathbb{P}^2}$. Since the Seiberg–Witten invariants are diffeomorphism invariants, we conclude that X is not diffeomorphic to $3\mathbb{C}\mathbb{P}^2\#7\overline{\mathbb{C}\mathbb{P}^2}$. Notice that case (i) of Theorem 3.4 does not apply and X_K is a minimal symplectic manifold. Thus, we can conclude that X is minimal. Since symplectic minimality implies irreducibility for simply-connected 4-manifolds with $b_2^+ > 1$ (Kotschick [9]), it follows that X is also smoothly irreducible.

6 Construction of an exotic symplectic $\mathbb{C}\mathbb{P}^2\#5\overline{\mathbb{C}\mathbb{P}^2}$

In this section, we construct a simply-connected minimal symplectic 4-manifold Y homeomorphic but not diffeomorphic to $\mathbb{C}\mathbb{P}^2\#5\overline{\mathbb{C}\mathbb{P}^2}$. Using Usher’s Theorem [21], we will distinguish Y from $\mathbb{C}\mathbb{P}^2\#5\overline{\mathbb{C}\mathbb{P}^2}$.

The manifold Y will be the symplectic fiber sum of Y_K and $T^2 \times S^2 \# 4\overline{\mathbb{C}\mathbb{P}^2}$ along the genus two surfaces Σ_2 and Σ'_2 . Let us choose the gluing diffeomorphism $\varphi: \Sigma_2 \times S^1 \rightarrow \Sigma'_2 \times S^1$ that maps the generators $a^{-1}b, b^{-1}aba^{-1}, d, y$ and μ of $\pi_1(Y_K \setminus \nu\Sigma_2)$ to the generators a_1, b_1, a_2, b_2 and μ' of $\pi_1(Z \setminus \nu\Sigma'_2)$ according to the following rule:

$$\varphi_*(a^{-1}b) = a_2, \varphi_*(b^{-1}aba^{-1}) = b_2, \varphi_*(d) = a_1, \varphi_*(y) = b_1, \varphi_*(\mu) = \mu'.$$

Here, μ and μ' denote the meridians of Σ and Σ'_2 .

Again, by Gompf’s theorem [7], $Y = Y_K \#_{\varphi} (T^2 \times S^2 \# 4\overline{\mathbb{C}\mathbb{P}^2})$ is symplectic.

Lemma 6.1 *Y is simply connected.*

Proof By Van Kampen’s theorem, we have

$$\pi_1(Y) = \frac{\pi_1(Y_K \setminus \nu\Sigma_2) * \pi_1(Z \setminus \nu\Sigma'_2)}{\langle a^{-1}b = a_2, b^{-1}aba^{-1} = b_2, d = a_1, y = b_1, \lambda = 1 \rangle}.$$

The following set of relations hold in $\pi_1(Y)$.

- (9) $a = bd,$
- (10) $yb = by,$
- (11) $aba = bab,$
- (12) $yab = ba.$

Using the same argument as in proof of [Lemma 5.1](#), we have $a = b = x = d = y = 1$. Thus $\pi_1(Y) = 0$. \square

Lemma 6.2 $c_1^2(Y) = 4$, $\sigma(Y) = -4$ and $\chi_h(Y) = 1$.

Proof We have $c_1^2(Y) = c_1^2(Y_K) + c_1^2(T^2 \times S^2 \# 4\overline{\mathbb{C}\mathbb{P}^2}) + 8$, $\sigma(Y) = \sigma(Y_K) + \sigma(T^2 \times S^2 \# 4\overline{\mathbb{C}\mathbb{P}^2})$ and $\chi_h(Y) = \chi_h(Y_K) + \chi_h(T^2 \times S^2 \# 4\overline{\mathbb{C}\mathbb{P}^2}) + 1$. Since $c_1^2(Y_K) = 0$, $\sigma(Y_K) = 0$ and $\chi_h(Y_K) = 0$, the result follows from [Lemma 3.3](#) and [Lemma 4.1](#). \square

By Freedman's classification theorem [\[6\]](#), [Lemma 6.1](#) and [Lemma 6.2](#) above, Y is homeomorphic to $\mathbb{C}\mathbb{P}^2 \# 5\overline{\mathbb{C}\mathbb{P}^2}$. Notice that Y is a fiber sum of the non-minimal manifold $Z = T^2 \times S^2 \# 4\overline{\mathbb{C}\mathbb{P}^2}$ with the minimal manifold Y_K . All 4 exceptional spheres E_1, E_2, E_3 and E_4 in Z meet with the genus two fiber $2T + S - E_1 - E_2 - E_3 - E_4$. Also, any embedded symplectic -1 sphere in $T^2 \times S^2 \# 4\overline{\mathbb{C}\mathbb{P}^2}$ is of the form $mS \pm E_i$, thus intersect non-trivially with the fiber class $2T + S - E_1 - E_2 - E_3 - E_4$. It follows from [Theorem 3.4](#) that Y is a minimal symplectic manifold. Since symplectic minimality implies irreducibility for simply-connected 4-manifolds for $b_2^+ = 1$ [\[8\]](#), it follows that Y is also smoothly irreducible. We conclude that Y is not diffeomorphic to $\mathbb{C}\mathbb{P}^2 \# 5\overline{\mathbb{C}\mathbb{P}^2}$.

Remark Alternatively, one can apply the concept of symplectic Kodaira dimension to prove the exoticness of X and Y . We refer the reader to the articles by Li and Yau [\[10\]](#) and Usher [\[22\]](#) for a detailed treatment of how the Kodaira dimension behaves under the symplectic fiber sum.

References

- [1] **A Akhmedov**, *Construction of symplectic cohomology $S^2 \times S^2$* , Gökova Geometry-Topology Proc 14 (2007) 36–48
- [2] **G Burde, H Zieschang**, *Knots*, de Gruyter Studies in Mathematics 5, Walter de Gruyter & Co., Berlin (1985) [MR808776](#)
- [3] **R Fintushel, R J Stern**, *Rational blowdowns of smooth 4-manifolds*, J. Differential Geom. 46 (1997) 181–235 [MR1484044](#)
- [4] **R Fintushel, R J Stern**, *Families of simply connected 4-manifolds with the same Seiberg–Witten invariants*, Topology 43 (2004) 1449–1467 [MR2081432](#)
- [5] **R Fintushel, R J Stern**, *Double node neighborhoods and families of simply connected 4-manifolds with $b^+ = 1$* , J. Amer. Math. Soc. 19 (2006) 171–180 [MR2169045](#)

- [6] **MH Freedman**, *The topology of four-dimensional manifolds*, J. Differential Geom. 17 (1982) 357–453 [MR679066](#)
- [7] **RE Gompf**, *A new construction of symplectic manifolds*, Ann. of Math. (2) 142 (1995) 527–595 [MR1356781](#)
- [8] **MJD Hamilton**, **D Kotschick**, *Minimality and irreducibility of symplectic four-manifolds*, Int. Math. Res. Not. (2006) Art. ID 35032, 13 [MR2211144](#)
- [9] **D Kotschick**, *The Seiberg–Witten invariants of symplectic four-manifolds (after CH Taubes)*, Astérisque (1997) Exp. No. 812, 4, 195–220 [MR1472540](#) Séminaire Bourbaki, Vol. 1995/96
- [10] **T-J Li**, **ST Yau**, *Embedded surfaces and Kodaira Dimension*, preprint
- [11] **Y Matsumoto**, *Lefschetz fibrations of genus two—a topological approach*, from: “Topology and Teichmüller spaces (Katinkulta, 1995)”, World Sci. Publ., River Edge, NJ (1996) 123–148 [MR1659687](#)
- [12] **JD McCarthy**, **JG Wolfson**, *Symplectic normal connect sum*, Topology 33 (1994) 729–764 [MR1293308](#)
- [13] **B Ozbagci**, **AI Stipsicz**, *Noncomplex smooth 4-manifolds with genus-2 Lefschetz fibrations*, Proc. Amer. Math. Soc. 128 (2000) 3125–3128 [MR1670411](#)
- [14] **J Park**, *Seiberg–Witten invariants of generalised rational blow-downs*, Bull. Austral. Math. Soc. 56 (1997) 363–384 [MR1490654](#)
- [15] **J Park**, *Simply connected symplectic 4-manifolds with $b_2^+ = 1$ and $c_1^2 = 2$* , Invent. Math. 159 (2005) 657–667 [MR2125736](#)
- [16] **J Park**, *Exotic smooth structures on $3\mathbb{C}P^2\#8\overline{\mathbb{C}P^2}$* , Bull. Lond. Math. Soc. 39 (2007) 95–102 [MR2303524](#)
- [17] **J Park**, **AI Stipsicz**, **Z Szabó**, *Exotic smooth structures on $\mathbb{C}P^2\#5\overline{\mathbb{C}P^2}$* , Math. Res. Lett. 12 (2005) 701–712 [MR2189231](#)
- [18] **AI Stipsicz**, **Z Szabó**, *An exotic smooth structure on $\mathbb{C}P^2\#6\overline{\mathbb{C}P^2}$* , Geom. Topol. 9 (2005) 813–832 [MR2140993](#)
- [19] **AI Stipsicz**, **Z Szabó**, *Small exotic 4-manifolds with $b_2^+ = 3$* , Bull. London Math. Soc. 38 (2006) 501–506 [MR2239045](#)
- [20] **CH Taubes**, *The Seiberg–Witten invariants and symplectic forms*, Math. Res. Lett. 1 (1994) 809–822 [MR1306023](#)
- [21] **M Usher**, *Minimality and symplectic sums*, Int. Math. Res. Not. (2006) Art. ID 49857, 17 [MR2250015](#)
- [22] **M Usher**, *Kodaira dimension and Symplectic Sums*, to appear in Commentarii Mathematici Helvetici
- [23] **E Witten**, *Monopoles and four-manifolds*, Math. Res. Lett. 1 (1994) 769–796 [MR1306021](#)

Mathematics Department, Columbia University, New York, NY 10027, USA

anar@math.columbia.edu

Received: 12 March 2007 Revised: 7 July 2008