

Small gaps between products of two primes

D. A. Goldston, S. W. Graham, J. Pintz and C. Y. Yıldırım

ABSTRACT

Let q_n denote the n th number that is a product of exactly two distinct primes. We prove that $q_{n+1} - q_n \leq 6$ infinitely often. This sharpens an earlier result of the authors, which had 26 in place of 6. More generally, we prove that if ν is any positive integer, then $(q_{n+\nu} - q_n) \leq \nu e^{\nu-\gamma}(1 + o(1))$ infinitely often. We also prove several other related results on the representation of numbers with exactly two prime factors by linear forms.

1. Introduction

As an approximation to the twin prime conjecture it was proved in [13] that

$$\liminf_{n \rightarrow \infty} \frac{p_{n+1} - p_n}{\log p_n} = 0. \quad (1.1)$$

The strongest approximation for the twin prime conjecture in another direction was proved in the celebrated work of Chen [4][†] (see also [15, Chapter 11]), where he showed that there are infinitely many primes p such that $p + 2 \in \mathcal{P}_2$, where

$$\mathcal{P}_2 := \{n : \Omega(n) \leq 2\}. \quad (1.2)$$

If \mathcal{P} denotes the set of primes, then Chen's theorem asserts that at least one of the relations

$$p + 2 = p' \in \mathcal{P} \quad (1.3)$$

or

$$p + 2 = p_1 p_2, \quad p_1, p_2 \in \mathcal{P} \quad (1.4)$$

holds for infinitely many primes p .

The phenomenon that we cannot specify which one of the two equations (1.3) and (1.4) has infinitely many solutions (in reality most probably both, naturally) is the most significant particular case of the parity problem, a heuristic principle stating that sieve methods cannot differentiate between integers with an even and an odd number of prime factors. This principle is based on some extremal examples of Selberg (see [14, Chapter 4; 28, p. 204]). Accordingly, until very recently, problems involving numbers that are products of two distinct prime factors (which we called E_2 -numbers in [11]) seemed to be as difficult as problems involving primes, since sieve methods seemed to be not suitable to attack these problems due to the parity problem. For example, the analog of (1.1),

$$\liminf_{n \rightarrow \infty} \frac{q_{n+1} - q_n}{\log q_n / \log \log q_n} = 0 \quad (1.5)$$

(where $q_1 < q_2 < \dots$ denotes the sequence of E_2 -numbers) was, similar to (1.1), not known.

Received 20 December 2007; published online 5 November 2008.

2000 *Mathematics Subject Classification* 11N25 (primary), 11N36 (secondary).

The first and second authors were supported in part by NSF Grants, the third author by OTKA grants No. 43623, 49693, 67676, and the Balaton program, and the fourth author by TÜBITAK.

[†]Chen's result was announced in 1966 [3]. However, due to the Cultural Revolution, the complete proof was not published until 1973.

The present authors observed that the method, a variant of Selberg’s sieve, which led to the proof of (1.1) in [13], can be used even more successfully for E_2 -numbers. In our preceding work [11] we gave an alternative proof of (1.1); further we showed that E_2 -numbers are infinitely often a bounded distance apart, more precisely,

$$\liminf_{n \rightarrow \infty} (q_{n+1} - q_n) \leq 26. \tag{1.6}$$

The relation (1.6) was actually a simple consequence of a more general result, according to which every admissible (see the definition below) 8-tuple contains at least two E_2 -numbers infinitely often. The following far reaching generalization of the twin prime conjecture was formulated qualitatively 100 years ago by Dickson [6], and two decades later in a quantitative form by Hardy and Littlewood [16]. In order to formulate the conjecture we define a set

$$\mathcal{H} = \{h_i\}_{i=1}^k \quad h_i \in \mathbb{Z}^+ \cup \{0\} \tag{1.7}$$

to be admissible if for every prime number p the set \mathcal{H} does not cover all residue classes mod p . Prime-tuple conjecture. Given any admissible set \mathcal{H} , there are infinitely many integers n such that all numbers of the form $n + h_i$ ($1 \leq i \leq k$) are primes. The number of such ns below N is asymptotically equal to

$$\frac{N}{\log^k N} \mathfrak{S}(\mathcal{H}) = \frac{N}{\log^k N} \prod_p \left(1 - \frac{\nu_p(\mathcal{H})}{p}\right) \left(1 - \frac{1}{p}\right)^{-k}, \tag{1.8}$$

where $\nu_p(\mathcal{H})$ denotes the number of residue classes mod p covered by \mathcal{H} .

The above conjecture includes (as the case $k = 2$) the generalized twin prime conjecture, which states that every even number can be written as the difference of two primes in infinitely many ways. This was formulated by de Polignac [26] in 1849 in a qualitative way, and in the same work of Hardy and Littlewood [16] in a quantitative form.

If we substitute primes by almost primes of the form P_r (integers having at most $r \geq 2$ prime factors) then the qualitative form of the analogous conjecture is true for $k = 2$, as shown by Chen’s theorem (1.2), even for $r = 2$. This trivially implies that we have infinitely often at least two P_2 -numbers in any admissible k -tuple for any $k \geq 2$.

We will examine the problem whether for any ν we can guarantee that there are infinitely often at least ν P_2 -numbers (or at least ν P_r -numbers with a given fixed r , independent of ν) in any admissible k -tuple if k is sufficiently large, that is, $k \geq C_0(\nu)$.

Such a result seems to be unknown for any fixed value of r . The strongest result in this direction is due to Heath-Brown [18] who showed that if $\{h_i\}_{i=1}^k$ is an admissible k -tuple then there are infinitely many n such that

$$\max_{1 \leq i \leq k} \omega(n + h_i) < C \log k. \tag{1.9}$$

This improved an earlier result of Halberstam and Richert [15, Chapter 10], where the analog of (1.9) was proved with the max replaced by the average of $\omega(n + h_i)$.

In the case of the primes it was shown in [13] that if the level ϑ of distribution of primes (see Definition 1.17) is any fixed number in $(1/2, 1]$, then

$$\liminf_{n \rightarrow \infty} (p_{n+1} - p_n) < \infty. \tag{1.10}$$

On the other hand, we needed the Elliott–Halberstam conjecture (EH) (see [7]) in its full strength to obtain

$$\liminf_{n \rightarrow \infty} \frac{p_{n+2} - p_n}{\log p_n} = 0. \tag{1.11}$$

For $p_{n+3} - p_n$ the best result we were able to prove on EH in [12] was

$$\liminf_{n \rightarrow \infty} \frac{p_{n+3} - p_n}{\log p_n} \leq e^{-\gamma} (\sqrt{3} - \sqrt{2})^2. \tag{1.12}$$

The incredible depth of the assumption EH in (1.11) and (1.12) suggests that it might be very difficult to prove

$$\liminf_{n \rightarrow \infty} (q_{n+\nu} - q_n) < \infty, \tag{1.13}$$

already for $\nu = 2$ or 3 . We will show, however, that our method can be applied very efficiently to this problem.

In the present work we will show the existence of at least ν E_2 -numbers in any admissible k -tuple if $k \geq C_1(\nu)$. We will also show that $C_1(2) = 3$ is permitted, that is, every admissible triplet contains at least two E_2 -numbers infinitely often.

The mentioned work of Heath-Brown [18] is based on a method of Selberg [28]. Selberg considered only the case $k = 2$ and showed that there are infinitely many pairs $n, n + 2$ such that one of them is a P_2 -number, the other a P_3 -number.

Our method, a modified form of the above mentioned methods of Selberg and Heath-Brown, also shows that $C_0(2) = 2$. Therefore, we have

$$n, n + 2 \in \mathcal{P}_2, \tag{1.14}$$

infinitely often, improving Selberg’s result but falling short of (1.2).

We will, in fact, prove the above results in the following more general form, similar to Heath-Brown [18]. Let

$$L_i(x) = a_i x + b_i \quad (1 \leq i \leq k) \quad a_i, b_i \in \mathbb{Z}, \quad a_i > 0 \tag{1.15}$$

be an admissible k -tuple of distinct linear forms. In other words, we suppose that for every prime p there exists $x_p \in \mathbb{Z}$ such that

$$p \nmid \prod_{i=1}^k (a_i x_p + b_i). \tag{1.16}$$

In order to formulate the results we will introduce the level ϑ of distribution of primes in arithmetic progressions. We say that the primes have level of distribution ϑ if for any positive A there exists a constant $C = C(A)$ such that

$$\sum_{q \leq N^\vartheta} \max_{\substack{a \\ (a,q)=1}} \left(\sum_{\substack{p \equiv a \\ (\text{mod } q)}} 1 - \frac{\text{li}(N)}{\varphi(q)} \right) \ll_A \frac{N}{(\log N)^A}. \tag{1.17}$$

The Bombieri–Vinogradov theorem states that $\vartheta = 1/2$ is admissible. Elliott and Halberstam [7] conjectured that (1.17) is true for any $\vartheta < 1$. Friedlander and Granville [10] proved that (1.17) is not true with $\vartheta = 1$, but it is possible that it still holds for any fixed $\vartheta < 1$.

In the following we suppose that an analog of (1.17) is true for E_2 -numbers with the same value of ϑ . This is true with $\vartheta = 1/2$ unconditionally, as shown by Motohashi [23]. Motohashi gives a more general result; he proves that if two functions satisfy analogs of the Bombieri–Vinogradov theorem, then under certain reasonable conditions, the convolution of the two functions also satisfies an analog of Bombieri and Vinogradov. This may also be proved using a slight variation of the argument of Bombieri [1, Theorem 22].

In the formulation of the theorems below, we assume that ϑ ($1/2 \leq \vartheta < 1$) is a common level of distribution for primes and E_2 -numbers. We then define

$$B = \frac{2}{\vartheta}. \tag{1.18}$$

Unconditionally, we may take $B = 4$. The Elliott–Halberstam conjecture for primes and E_2 -numbers is equivalent to taking $B = 2 + \epsilon$.

THEOREM 1. *Let D be any constant and let $L_i(x)$ ($1 \leq i \leq k$) be an admissible k -tuple of distinct linear forms. Then there are $\nu + 1$ forms among them, which take simultaneously E_2 -numbers as values with both prime factors above D if[†]*

$$k \geq C_1(\nu) := \frac{4e^{-\gamma}(1 + o(1))}{B} e^{B\nu/4}. \quad (1.19)$$

THEOREM 2. *Let $\{L_1(n), L_2(n), L_3(n)\}$ be an admissible triplet of linear forms. Among these, exist two forms L_i, L_j such that for infinitely many n , $L_i(n), L_j(n)$ are both E_2 -numbers, all the prime factors of which exceed $n^{1/144}$.*

THEOREM 3. *Let $\{L_1(n), L_2(n)\}$ be an admissible pair of linear forms. Then there exist infinitely many n such that both $L_1(n)$ and $L_2(n)$ are P_2 -numbers, and the prime factors of $L_1(n)L_2(n)$ all exceed $n^{1/10}$. In particular, there are infinitely many integers n such that*

$$n, n - d \in \mathcal{P}_2 \quad (1.20)$$

for any even integer d .

Theorem 1 shows that, in contrast to the case of primes, we can really prove the existence of infinitely many blocks of ν consecutive E_2 -numbers with a bounded diameter (depending on ν) for any given ν .

COROLLARY 1. *We have for any $\nu > 0$,*

$$\liminf_{n \rightarrow \infty} (q_{n+\nu} - q_n) \leq C_2(\nu) = e^{-\gamma} \nu e^{B\nu/4} (1 + o(1)). \quad (1.21)$$

Taking the admissible triplet $\{n, n + 2, n + 6\}$, we see that Theorem 2 implies an improvement of (1.6), namely

COROLLARY 2. $\liminf_{n \rightarrow \infty} (q_{n+1} - q_n) \leq 6$.

The question arises: why is our present method much more successful for E_2 -numbers than for primes, as indicated by (1.11), (1.12) and (1.21)?

Usually sieve methods are at any rate unable to detect E_r -numbers for any given r due to the parity problem, and even in the case of P_r -numbers (r fixed) they produce only numbers with all prime factors larger than

$$X^{1/w}, \quad w > 0 \text{ fixed}, \quad (1.22)$$

where $\text{card } \mathcal{A} \sim X$, where \mathcal{A} is a starting set containing almost primes, as emphasized by Kan [20, 21]. In these cases the number of almost primes considered below N is $O_w(N/\log N)$ (the same as the number of primes), whereas the true order of magnitude of the number of P_r -numbers (or E_r -numbers) is

$$c(r) \frac{N(\log_2 N)^{r-1}}{(\log N)}. \quad (1.23)$$

[†]For clarity, we remark that here and in subsequent results (Theorems 4, 5, 6, Corollaries 1, 3, 4, 5) the notation $o(1)$ denotes a function $g(\nu)$ such that $g(\nu) \rightarrow 0$ as $\nu \rightarrow \infty$.

Differently from almost all other applications of sieve methods (for exceptions see the mentioned works [20, 21] of Kan), our method is able to make use of E_2 -numbers that satisfy

$$n = p_1 p_2, \quad p_1 < n^\varepsilon, \quad p_2 > n^{1-\varepsilon}, \tag{1.24}$$

for any given small $\varepsilon > 0$. In the proof of Theorem 8, we allow E_2 -numbers with prime factors of any size.

This phenomenon (the larger density of E_2 -numbers over primes) is crucial in our method. A careful consideration of the proof of Theorem 1 reveals that without taking into account the contribution of E_2 -numbers with (1.24) for all $\varepsilon > 0$, our method would fail to prove Theorem 1. If we exclude numbers of type (1.24) for $\varepsilon < c_0$, then we would be unable to show Theorem 1, and so Corollary 1 for any $\nu > \nu_0(c_0) \asymp c_0^{-1}$.

As we have seen in (1.10) and (1.11), the level ϑ of distribution of primes has dramatic consequences for the strength of the result we can show about the existence of primes in tuples. On the other hand, the value of ϑ , that is, of B , is much less important in the distribution of E_2 -numbers; only the quantitative value $C_1(\nu)$ depends on the value of B , that is, of ϑ . The dependence of $C_1(\nu)$ in (1.19) on ϑ is not too strong: we have in the exponent of $C_1(\nu)$

$$\frac{B\nu}{4} = \frac{\nu}{(2\vartheta)} \in \left[\frac{\nu}{2}, \nu \right] \quad \text{for } \vartheta \in \left[\frac{1}{2}, 1 \right]. \tag{1.25}$$

This observation has theoretical importance, for we do not need the full strength of the Bombieri–Vinogradov theorem. Moreover, it can be used to generalize the results of Theorem 1 for a situation when $\vartheta = \varepsilon \Leftrightarrow B = 2/\varepsilon$, for example. We remark that, contrary to this, the proof of (1.1) would break down if we had just a fixed $\vartheta < 1/2$ at our disposal, even if these values were very close to $1/2$.

The case of ϑ being small occurs when we would like to find blocks of bounded length of E_2 -numbers in short intervals of type

$$[N, N + N^{7/12+\varepsilon}], \quad \varepsilon > 0 \text{ fixed.} \tag{1.26}$$

In this case it was proved by Perelli, Pintz and Salerno [24] in 1985 that one has a short interval version of Bombieri–Vinogradov theorem for intervals of type (1.26), where (surprisingly) we can choose ϑ as a fixed positive constant, $\vartheta = 1/40$ for any $\varepsilon > 0$. This was improved two years later by Timofeev [30] to

$$\vartheta = \frac{1}{30} \iff B = 60. \tag{1.27}$$

The result proved in [24] reads as

$$\sum_{q \leq Q} \max_{(a,q)=1} \max_{h \leq y} \max_{x/2 < z \leq x} \left| \sum_{\substack{p \equiv a \pmod{q} \\ z < p \leq z+h}} \log p - \frac{h}{\varphi(q)} \right| \ll \frac{y}{(\log x)^A}, \tag{1.28}$$

where $y = x^{7/12+\varepsilon}$, $Q = x^\vartheta (\log x)^{-D}$, A arbitrary and $D = D(A)$.

The method of proof of both works [24, 30] uses Heath-Brown’s identity, therefore, the analog of (1.28) can be proved *mutatis mutandis* for E_2 -numbers as well. Accordingly, we will prove the following theorem.

THEOREM 4. *Under the conditions of Theorem 1 we can find $\nu + 1$ linear forms $L_{i_j}(n)$ which take E_2 -numbers with both prime factors above D and for any $\varepsilon > 0$ we can require*

$$n \in [N, N + N^{7/12+\varepsilon}] \tag{1.29}$$

if

$$k \geq C_3(\nu) := \frac{e^{-\gamma}(1 + o(1))}{15} e^{15\nu}, \quad N > N_0(k, \varepsilon, D). \tag{1.30}$$

COROLLARY 3. *Let $\nu \in \mathbb{Z}^+$, $\varepsilon > 0$. If $N > N_1(\nu, \varepsilon)$, then there exist $n, m \in \mathbb{Z}^+$ such that*

$$n \in [N, N + N^{7/12+\varepsilon}] \tag{1.31}$$

$$n < q_m < q_{m+1} < \dots < q_{m+\nu} < n + e^{-\gamma} \nu e^{15\nu} (1 + o(1)). \tag{1.32}$$

We can further restrict our E_2 -numbers $p_1 p_2$ to be of the form

$$p_1 p_2 = x^2 + y^2 \quad (x, y \in \mathbb{Z}) \iff p_1, p_2 \equiv 1 \pmod{4}. \tag{1.33}$$

It is relatively easy to show the existence of infinitely many families of triplets of consecutive integers that are sums of two squares, see [5, 19] for a more general result. A modification of our proof of Theorem 1 shows that Theorem 1 remains essentially valid for E_2 -numbers which are sums of two squares.

THEOREM 5. *Under the conditions of Theorem 1, we have infinitely many n such that at least $\nu + 1$ linear forms $L_{i_j}(n)$ take simultaneously E_2 -values (with both prime factors above D) which are sums of two squares, if*

$$k \geq C_4(\nu) := \frac{4e^{-\gamma}(1 + o(1))}{B} e^{B\nu}. \tag{1.34}$$

COROLLARY 4. *Under the conditions of Theorem 5 we have at least ν E_2 -numbers which are sums of two squares infinitely often in intervals of length K if*

$$K \geq C_5(\nu) := 4e^{-\gamma} \nu e^{B\nu} (1 + o(1)). \tag{1.35}$$

Finally we can combine the results of Theorems 4 and 5 to have ν E_2 -numbers which are sums of two squares if the value n is localized in a short interval of type (1.29).

THEOREM 6. *In Theorem 4 we may require that the ν E_2 -values of the linear forms should be sums of two squares if we have in place of (1.30) the restriction*

$$k \geq C_6(\nu) := \frac{e^{-\gamma}(1 + o(1))}{15} e^{60\nu}. \tag{1.36}$$

COROLLARY 5. *Let $\{q'_n\}_1^\infty$ denote the set of E_2 -numbers which can be written as sums of two squares. Then Corollary 3 is true if (1.32) is replaced by*

$$n < q'_m < q'_{m+1} < \dots < q'_{m+\nu} < n + 4e^{-\gamma} \nu e^{60\nu} (1 + o(1)). \tag{1.37}$$

It will be clear from the proofs that all of the above theorems and corollaries remain true if we require that all of the constructed E_2 -numbers have both of their prime factors exceeding some specific constant. Indeed, this holds more generally if both prime factors exceed some $Y(N)$ with $\log Y(N)/\log N \rightarrow 0$ as $N \rightarrow \infty$.

Our methods open the way toward a new, simpler, and unified treatment of some conjectures of Erdős [8] on consecutive integers, the most well known of them is the Erdős–Mirsky [9] conjecture, which states that

$$d(n) = d(n + 1) \text{ infinitely often}; \tag{1.38}$$

the two others are the analogous conjectures with $d(n)$ replaced by the functions $\omega(n)$ or $\Omega(n)$ (number of prime divisors of n counted without and with multiplicity, respectively). Similar to

(1.3) and (1.4), the parity problem seemed to prevent a solution of these conjectures. However, as observed by Spiro [29] and Heath-Brown [17], the parity problem can be “sidestepped,” and it is possible to prove the conjectures without specifying the common value of $f(n) = f(n + 1)$ (or even its parity) for the relevant functions $f = d$ or Ω . Recently, the same was shown for $f = \omega$ by Schlage-Puchta [27].

In a future paper, we will show these conjectures in the stronger form, where we can specify the common value $f(n) = f(n + 1)$, thereby overcoming the parity problem in these cases. Among other things, we will prove that there are infinitely many positive integers n such that the relations

$$d(n) = d(n + 1) = 24, \quad \Omega(n) = \Omega(n + 1) = 5, \quad \text{and} \quad \omega(n) = \omega(n + 1) = 4$$

simultaneously hold. Furthermore, we can prove similar statements for an arbitrary shift b in place of 1. This generalization was proved for every b by Pinner [25] in 1997 (without specifying the common value of f) for $f = d$ and Ω ; however, Schlage-Puchta’s method for ω does not work for general b . On the other hand, Buttkewitz [2] has recently proved an analogous result for an infinite set \mathcal{B} of possible shifts b .

2. Notation and Preliminary Lemmas

Most of our notation will be introduced as needed, but it is useful to make the following comments here. Throughout this paper, we use k to denote an integer $k \geq 2$, \mathcal{L} to denote an admissible k -tuple of linear forms, and P to denote a polynomial. The constants implied by “ O ” and “ \ll ” may depend on k, \mathcal{L} , and P . Let $\tau_k(n)$ denote the number of ways of writing n as product of k factors, and let $\omega(n)$ denote the number of distinct prime factors of n . Let $\phi(n)$ and $\mu(n)$ be the usual functions of Euler and Möbius, respectively. The letters N and R denote real numbers regarded as tending to infinity, and we always assume that $R \leq N^{1/2}$.

To count E_2 -numbers, we introduce the following function β . Let Y be a real number with $1 \leq Y \leq N^{1/4}$, and define

$$\beta(n) = \begin{cases} 1 & \text{if } n = p_1 p_2, \quad Y < p_1 \leq N^{1/2} < p_2, \\ 0 & \text{otherwise.} \end{cases} \tag{2.1}$$

The notation $\pi(x)$ is commonly used to denote the number of primes up to x , and $\pi(x; q, a)$ denotes the number of primes up to x that are congruent to $a \pmod{q}$. For our purposes, it is convenient to define the following related quantities:

$$\begin{aligned} \pi^b(x) &= \sum_{x < p \leq 2x} 1 = \pi(2x) - \pi(x). \\ \pi^b(x; q, a) &= \sum_{\substack{x < p \leq 2x \\ p \equiv a \pmod{q}}} 1 = \pi(2x; q, a) - \pi(x; q, a). \\ \pi_\beta(x) &= \sum_{x < n \leq 2x} \beta(n). \\ \pi_{\beta,u}(x) &= \sum_{\substack{x < n \leq 2x \\ (n,u)=1}} \beta(n). \\ \pi_\beta(x; q, a) &= \sum_{\substack{x < n \leq 2x \\ n \equiv a \pmod{q}}} \beta(n). \end{aligned}$$

As mentioned in the introduction, we will employ results on the level of distribution for both prime numbers and E_2 -numbers. For primes, define

$$\Delta(x; q, a) = \pi^b(x; q, a) - \frac{1}{\phi(q)}\pi^b(x)$$

and

$$\Delta^*(x; q) = \max_{y \leq x} \max_{a; (a, q)=1} |\Delta(y; q, a)|.$$

LEMMA 1. *Assume that the primes have level of distribution ϑ , $\vartheta \leq 1$. For every $A > 0$ and for every fixed integer $h \geq 0$, there exists $C = C(A, h)$ such that if $Q \leq x^\vartheta (\log x)^{-C}$, then*

$$\sum_{q \leq Q} \mu^2(q) h^{\omega(q)} \Delta^*(x; q) \ll_A x (\log x)^{-A}.$$

By the Bombieri–Vinogradov Theorem, this lemma is unconditional for $\vartheta \leq 1/2$. The incorporation of the factor $h^{\omega(q)}$ is familiar feature in sieve applications; see [15, Lemma 3.5], for example.

For the function β , we define

$$\begin{aligned} \Delta_\beta(x; q, a) &= \sum_{\substack{x < n \leq 2x \\ n \equiv a \pmod{q}}} \beta(n) - \frac{1}{\phi(q)} \sum_{\substack{x < n \leq 2x \\ (n, q)=1}} \beta(n) = \pi_\beta(x; q, a) - \frac{1}{\phi(q)} \pi_{\beta, q}(x), \\ \Delta_\beta^*(x; q) &= \max_{y \leq x} \max_{a; (a, q)=1} |\Delta_\beta(y; q, a)|. \end{aligned}$$

LEMMA 2. *Assume that E_2 -numbers have a level of distribution ϑ , $\vartheta \leq 1$. For every $A > 0$ and for every fixed integer $h \geq 0$, there exists $C = C(A, h)$ such that if $Q \leq x^\vartheta (\log x)^{-C}$, then*

$$\sum_{q \leq Q} \mu^2(q) h^{\omega(q)} \Delta_\beta^*(x; q) \ll_A x (\log x)^{-A}. \tag{2.2}$$

This follows from a general result of Motohashi [23] when $\vartheta \leq 1/2$. When $1/2 < \vartheta \leq 1$, Lemmas 1 and 2 are both hypothetical.

Our next lemma is central to the estimation of the sums that arise in Selberg’s sieve.

LEMMA 3. *Suppose that γ is a multiplicative function, and suppose that there are positive real numbers κ, A_1, A_2, L such that*

$$0 \leq \frac{\gamma(p)}{p} \leq 1 - \frac{1}{A_1}, \tag{2.3}$$

and

$$-L \leq \sum_{w \leq p < z} \frac{\gamma(p) \log p}{p} - \kappa \log \frac{z}{w} \leq A_2 \tag{2.4}$$

if $2 \leq w \leq z$. Let g be the multiplicative function defined by

$$g(d) = \prod_{p|d} \frac{\gamma(p)}{p - \gamma(p)}. \tag{2.5}$$

Then

$$\sum_{d < z} \mu^2(d) g(d) = c_\gamma \frac{(\log z)^\kappa}{\Gamma(\kappa + 1)} \left\{ 1 + O\left(\frac{L}{\log z}\right) \right\},$$

where

$$c_\gamma = \prod_p \left(1 - \frac{\gamma(p)}{p}\right)^{-1} \left(1 - \frac{1}{p}\right)^\kappa.$$

This is a combination of Lemmas 5.3 and 5.4 of Halberstam and Richert’s book [15]. In [15], the hypothesis (2.3) is denoted by (Ω_1) , and hypothesis (2.4) is denoted by $(\Omega_2(\kappa, L))$. The constant implied by “ O ” may depend on A_1, A_2, κ , but it is independent of L .

Our next lemma is a variant of the previous one with the terms $g(d)$ weighted by an appropriate function.

LEMMA 4. Assume the hypotheses of Lemma 3. Assume also that $F : [0, 1] \rightarrow \mathbb{R}$ is a piecewise differentiable function. Then

$$\sum_{d < z} \mu^2(d)g(d)F\left(\frac{\log z/d}{\log z}\right) = c_\gamma \frac{(\log z)^\kappa}{\Gamma(\kappa)} \int_0^1 F(1-x)x^{\kappa-1}dx \tag{2.6}$$

$$+ O(c_\gamma LM(F)(\log z)^{\kappa-1}),$$

where $M(F) = \sup\{|F(x)| + |F'(x)| : 0 \leq x \leq 1\}$. The constant implied by “ O ” may depend on A_1, A_2 , and κ , but it is independent of L and F .

Proof. The left-hand side of the proposed conclusion is

$$\int_{1^-}^z F\left(\frac{\log z/u}{\log z}\right) dG(u),$$

where

$$G(u) = \sum_{d < u} \mu^2(d)g(d) = c_\gamma \frac{(\log u)^\kappa}{\Gamma(\kappa + 1)} + E(u),$$

and $E(u) \ll c_\gamma L(\log 2u)^{\kappa-1}$ by the previous lemma. Consequently, the sum in (2.6) may be written as

$$\int_1^z F\left(\frac{\log z/u}{\log z}\right) dc_\gamma \frac{(\log u)^\kappa}{\Gamma(\kappa + 1)} + \int_{1^-}^z F\left(\frac{\log z/u}{\log z}\right) dE(u).$$

In the first integral, we make the change of variables $u = z^x$; this gives the first term. We use integration by parts on the second integral to obtain

$$\int_{1^-}^z F\left(\frac{\log z/u}{\log z}\right) dE(u) = F\left(\frac{\log z/u}{\log z}\right) E(u) \Big|_{1^-}^z + \int_1^z E(u)F'\left(\frac{\log z/u}{\log z}\right) \frac{du}{u \log z}.$$

The desired result now follows by using the above-mentioned bound for $E(u)$. □

3. Initial Considerations

Let $\mathcal{L} = \{L_1, L_2, \dots, L_k\}$ be an admissible k -tuple of linear forms; that is, a set of linear forms satisfying the conditions of (1.15) and (1.16). Admissibility can also be defined in terms of solutions of congruences. Define

$$P_{\mathcal{L}}(n) = \prod_{i=1}^k L_i(n) = (a_1n + b_1) \cdots (a_kn + b_k),$$

and for each prime p define

$$\begin{aligned} \Omega_p(\mathcal{L}) &= \{n : 1 \leq n \leq p \text{ and } P_{\mathcal{L}}(n) \equiv 0 \pmod{p}\}, \\ \nu_p(\mathcal{L}) &= \#\Omega_p(\mathcal{L}). \end{aligned}$$

The condition that \mathcal{L} is admissible is equivalent to requiring that

$$\nu_p(\mathcal{L}) < p \tag{3.1}$$

for all primes p . We always have $\nu_p(\mathcal{L}) \leq k$, so the above condition is automatic for any prime $p > k$.

The singular series connected to \mathcal{L} is defined as

$$\mathfrak{S}(\mathcal{L}) = \prod_p \left(1 - \frac{\nu_p(\mathcal{L})}{p}\right) \left(1 - \frac{1}{p}\right)^{-k}.$$

The product converges because $\nu_p = k$ for all but finitely many primes p , and \mathcal{L} is admissible if and only if $\mathfrak{S}(\mathcal{L}) \neq 0$.

Next, we examine those primes p satisfying $\nu_p < k$. First of all, if $p|a_i$ for some i , then the congruence $a_i n + b_i \equiv 0 \pmod{p}$ will have no solutions, and so $\nu_p < k$ in this case. Now suppose that $p \nmid a_i$ for all i . In this instance, $\nu_p < k$ if and only if there are indices i, j ($i \neq j$) such that

$$\bar{a}_i b_i \equiv \bar{a}_j b_j \pmod{p},$$

where \bar{a}_i denotes the multiplicative inverse of $a_i \pmod{p}$. We, therefore, see that $\nu_p < k$ if and only if $p|A$, where

$$A = A(\mathcal{L}) = \prod_{i=1}^k a_i \prod_{1 \leq i < j \leq k} |a_i b_j - a_j b_i|. \tag{3.2}$$

For technical reasons, it is useful to adopt the normalization introduced by Heath-Brown [18]. For each prime $p|A$, there is an integer n_p such that $p \nmid P_{\mathcal{L}}(n_p)$. By the Chinese remainder theorem, there is an integer B such that $B \equiv n_p \pmod{p}$ for all $p|A$. For $i = 1, \dots, k$, we define

$$L'_i(n) = L_i(An + B) = a'_i n + b'_i,$$

where $a'_i = a_i A$ and $b'_i = L_i(B) = a_i B + b_i$. Set

$$\mathcal{L}' = \{L'_1, \dots, L'_k\}.$$

We claim that

$$\nu_p(\mathcal{L}') = \begin{cases} k & \text{if } p \nmid A, \text{ and} \\ 0 & \text{if } p | A. \end{cases} \tag{3.3}$$

To justify this claim, we assume first that $p|A$. Then

$$L'_i(n) \equiv L_i(B) \equiv L_i(n_p) \not\equiv 0 \pmod{p}$$

for all integers n , and so $\nu_p(\mathcal{L}') = 0$. Next assume that $p \nmid A$. As noted before, $\nu_p(\mathcal{L}') < k$ if and only if $p|(a'_i b'_j - a'_j b'_i)$ for some choice of i, j with $1 \leq i < j \leq k$. However,

$$a'_i b'_j - a'_j b'_i = \det \begin{vmatrix} a_i A & a_j A \\ a_i B + b_i & a_j B + b_j \end{vmatrix} = A \det \begin{vmatrix} a_i & a_j \\ b_i & b_j \end{vmatrix},$$

and this is not divisible by p .

For brevity, it is useful to relabel L'_i as L_i and to assume the following hypothesis.

HYPOTHESIS 1. Let $\mathcal{L} = \{L_1, \dots, L_k\}$ be an admissible k -tuple of linear forms. The functions $L_i(n) = a_i n + b_i (1 \leq i \leq k)$ have integer coefficients with $a_i > 0$. Each of the coefficients a_i is composed of the same primes, none of which divides any of the b_i . If $i \neq j$, then any prime factor of $a_i b_j - a_j b_i$ divides each of the a_i .

For sets of linear forms \mathcal{L} satisfying Hypothesis 1, we redefine

$$A = \prod_i a_i.$$

In this case,

$$\mathfrak{S}(\mathcal{L}) = \prod_{p|A} \left(1 - \frac{1}{p}\right)^{-k} \prod_{p \nmid A} \left(1 - \frac{k}{p}\right) \left(1 - \frac{1}{p}\right)^{-k}.$$

Note that by (3.1) and (3.3),

$$p \leq k \implies p|A, \tag{3.4}$$

so $\mathfrak{S}(\mathcal{L})$ is indeed positive.

The primary tool for proving Theorems 1–3 is an adaptation of the basic construction of Goldston, Pintz, and Yıldırım. Let $\mathcal{L} = \{L_1, L_2, \dots, L_k\}$ be a set of linear forms satisfying Hypothesis 1, and let β be as defined in (2.1). For the proof of Theorem 1, we seek an asymptotic formula for the sum

$$\mathcal{S} = \sum_{N < n \leq 2N} \left\{ \sum_{j=1}^k \beta(L_j(n)) - \nu \right\} \left(\sum_{d|P_{\mathcal{L}}(n)} \lambda_d \right)^2, \tag{3.5}$$

where the λ_d are real numbers to be chosen in due course. The significance of \mathcal{S} is that a value of n contributes a positive amount only if at least $\nu + 1$ elements of the set $\{L_1(n), \dots, L_k(n)\}$ are E_2 -numbers.

We immediately decompose \mathcal{S} as

$$\mathcal{S} = \sum_{j=1}^k \mathcal{S}_{1,j} - \nu \mathcal{S}_0, \tag{3.6}$$

where

$$\mathcal{S}_{1,j} = \sum_{N < n \leq 2N} \beta(L_j(n)) \left(\sum_{d|P_{\mathcal{L}}(n)} \lambda_d \right)^2,$$

and

$$\mathcal{S}_0 = \sum_{N < n \leq 2N} \left(\sum_{d|P_{\mathcal{L}}(n)} \lambda_d \right)^2.$$

The motivation for the use of the coefficient λ_d comes from the realm of the Selberg sieve. More specifically, consider the problem of bounding the number of n for which all of $L_1(n), \dots, L_k(n)$ are prime. Start from the observation that if $\lambda_1 = 1$ and $\lambda_d = 0$ for $d > N$, then

$$\begin{aligned} \sum_{\substack{N < n \leq 2N \\ \text{all } L_i(n) \text{ prime}}} 1 &\leq \sum_{N < n \leq 2N} \left(\sum_{d|P_{\mathcal{L}}(n)} \lambda_d \right)^2 \\ &= \sum_{d,e} \lambda_d \lambda_e \sum_{\substack{N < n \leq 2N \\ [d,e]|P_{\mathcal{L}}(n)}} 1. \end{aligned} \tag{3.7}$$

As we noted in the introduction, we take $\nu_p(\mathcal{L})$ to be the number of solutions of $P_{\mathcal{L}}(n) \equiv 0 \pmod p$. We extend this definition to arbitrary squarefree d by multiplicativity. Consequently,

$$\sum_{\substack{N < n \leq 2N \\ d|P_{\mathcal{L}}(n)}} 1 = N \frac{\nu_d(\mathcal{L})}{d} + O\left(k^{\omega(d)}\right)$$

for squarefree d . Returning to (3.7), we find that the expression there is

$$N \sum_{d,e} \frac{\lambda_d \lambda_e \nu_{[d,e]}(\mathcal{L})}{[d,e]} + O\left(\sum_{d,e} |\lambda_d \lambda_e| k^{\omega([d,e])}\right).$$

We control the size of the error term by specifying that $\lambda_d = 0$ if $d \geq R$, where R will be chosen later. Moreover, the terms with $([d,e], A) > 1$ make no contribution since $\nu_{[d,e]}(\mathcal{L}) = 0$ for these terms. Accordingly, we restrict the sum to terms with $(d, A) = (e, A) = 1$. It is also convenient to specify that

$$\lambda_d = 0 \text{ if } d \text{ is not squarefree.}$$

The coefficient of N in the main term may be rewritten as

$$\sum'_{d,e} \frac{\lambda_d \lambda_e}{f([d,e])}, \tag{3.8}$$

where \sum' denotes that the sum is over all values of the indices that are relatively prime to A , and

$$f(d) = \frac{d}{\nu_d(\mathcal{L})} = \frac{d}{\tau_k(d)} = \prod_{p|d} \frac{p}{k} \tag{3.9}$$

for squarefree d with $(d, A) = 1$.

The typical approach in the Selberg sieve is to choose the λ_d to minimize the form in (3.8). To make this problem feasible, one needs to diagonalize this bilinear form. This can be done by making a change of variables

$$y_r = \mu(r) f_1(r) \sum'_d \frac{\lambda_{dr}}{f(dr)}, \tag{3.10}$$

where f_1 is the multiplicative function defined by $f_1 = f * \mu$. In other words,

$$f_1(d) = \prod_{p|d} \frac{p-k}{k}, \tag{3.11}$$

whenever d is squarefree and $(d, A) = 1$. (Note that the sum in (3.10) is finite because $\lambda_d = 0$ for $d > R$. Note also that there is an implicit condition $(d, r) = 1$ because $\lambda_{dr} = 0$ if dr is not squarefree.) The sum in (3.8) is then transformed into

$$\sum'_r \frac{y_r^2}{f_1(r)},$$

and the bilinear form is minimized by taking

$$y_r = \mu^2(r) \frac{\lambda_1}{V} \tag{3.12}$$

when $r < R$ and $(r, A) = 1$, where

$$V = \sum'_{r < R} \frac{\mu^2(r)}{f_1(r)}.$$

The minimum of the form in (3.8) is then seen to be

$$\frac{\lambda_1^2}{V}.$$

One usually assumes that $\lambda_1 = 1$, but this is not an essential element of the Selberg sieve, and it is sometimes useful to assign some other nonzero value to λ_1 .

Using Möbius inversion, one can easily show that

$$\lambda_d = \mu(d)f(d) \sum_r' \frac{y_{rd}}{f_1(rd)}. \tag{3.13}$$

Consequently, specifying a choice for λ_d is equivalent to specifying a choice for y_r . Our choice of λ_d is different from the choice implied by (3.12), and it is most easily described in terms of y_r . We will take

$$y_r = \begin{cases} \mu^2(r)\mathfrak{S}(\mathcal{L})P\left(\frac{\log R/r}{\log R}\right) & \text{if } r < R \text{ and } (r, A) = 1, \\ 0 & \text{otherwise.} \end{cases} \tag{3.14}$$

Here, P is a polynomial to be determined later.

Our estimate for \mathcal{S} follows from the following two results.

THEOREM 7. *Suppose that \mathcal{L} is a set of linear forms satisfying Hypothesis 1. Suppose that λ_d is given by (3.13) and (3.14). There is some constant C such that if $R \leq N^{1/2}(\log N)^{-C}$, then*

$$\mathcal{S}_0 = \frac{\mathfrak{S}(\mathcal{L})N(\log R)^k}{(k-1)!} J_0 + O(N(\log N)^{k-1}),$$

where

$$J_0 = \int_0^1 P(1-x)^2 x^{k-1} dx.$$

THEOREM 8. *Suppose that the primes and the E_2 -numbers have a common level of distribution $\vartheta \leq 1$, and let \mathcal{L} be a set of linear forms satisfying Hypothesis 1. Suppose that λ_d is given by (3.13) and (3.14), and let the polynomial \tilde{P} be defined as*

$$\tilde{P}(x) = \int_0^x P(t) dt.$$

There is some constant C such that if $R = N^{\vartheta/2}(\log N)^{-C}$, then

$$\mathcal{S}_{1,j} = \frac{\mathfrak{S}(\mathcal{L})N(\log R)^{k+1}}{(k-2)!(\log N)} (J_1 + J_2 + J_3) + O(N(\log \log N)(\log N)^{k-1}),$$

where $Y = N^\eta$, $B = 2/\vartheta$, and

$$\begin{aligned} J_1 &= \int_{B\eta}^1 \frac{B}{y(B-y)} \int_0^{1-y} \left(\tilde{P}(1-x) - \tilde{P}(1-x-y) \right)^2 x^{k-2} dx dy, \\ J_2 &= \int_{B\eta}^1 \frac{B}{y(B-y)} \int_{1-y}^1 \tilde{P}(1-x)^2 x^{k-2} dx dy, \\ J_3 &= \int_1^{B/2} \frac{B}{y(B-y)} \int_0^1 \tilde{P}(1-x)^2 x^{k-2} dx dy. \end{aligned}$$

Finally, we mention the following result, which is needed for the proof of Theorem 3.

THEOREM 9. Assume the hypotheses of Theorem 8. Let ϖ denote the characteristic function of the primes; that is, $\varpi(p) = 1$ if p is a prime and $\varpi(p) = 0$ otherwise. There is some constant C such that if $R \leq N^{\theta/2}(\log N)^{-C}$, then

$$\sum_{N < n \leq 2N} \varpi(L_j(n)) \left(\sum_{d|P_{\mathcal{L}}(n)} \lambda_d \right)^2 = \frac{\mathfrak{S}(\mathcal{L})N(\log R)^{k+1}}{(k-2)!(\log N)} J_{\varpi} + O(N(\log \log N)(\log N)^{k-1}),$$

where

$$J_{\varpi} = \int_0^1 \tilde{P}(1-x)^2 x^{k-2} dx.$$

This result is very similar to Theorem 1.6 of [11] and to Theorem 8; consequently, we will give just a short sketch of the proof in Section 7.

4. Proof of Theorem 7

From the definition of S_0 , we see that

$$\begin{aligned} S_0 &= \sum'_{d,e} \lambda_d \lambda_e \sum_{\substack{N < n \leq 2N \\ [d,e]|P_{\mathcal{L}}(n)}} 1 = N \sum'_{d,e} \frac{\lambda_d \lambda_e}{f([d,e])} + O\left(\sum'_{d,e} |\lambda_d \lambda_e r_{[d,e]}|\right) \\ &= NS_{01} + O(S_{02}), \end{aligned} \tag{4.1}$$

say, where

$$r_d = \sum_{\substack{N < n \leq 2N \\ d|P_{\mathcal{L}}(n)}} 1 - \frac{N}{f(d)}. \tag{4.2}$$

Now

$$S_{01} = \sum'_{d,e} \frac{\lambda_d \lambda_e}{f(d)f(e)} \sum'_{\substack{r|d \\ r|e}} f_1(r) = \sum'_r f_1(r) \left(\sum'_d \frac{\lambda_{dr}}{f(dr)} \right)^2 = \sum'_r \frac{\mu^2(r)y_r^2}{f_1(r)}.$$

We use Lemma 4 with

$$\gamma(p) = \begin{cases} k & \text{if } p \nmid A, \\ 0 & \text{if } p|A \end{cases}$$

and $\kappa = k, F(x) = P(x)^2$. We deduce that

$$S_{01} = \frac{\mathfrak{S}(\mathcal{L})(\log R)^k}{(k-1)!} \int_0^1 P(1-x)^2 x^{k-1} dx + O((\log R)^{k-1}).$$

For S_{02} , we first note the bound

$$|r_{[d,e]}| \leq k^{\omega([d,e])}$$

that follows from (4.2). We will later establish the bound

$$|\lambda_d| \ll (\log R)^k \tag{4.3}$$

whenever $d \leq R$ and d is squarefree. Assuming this momentarily, we find that

$$\begin{aligned} S_{02} &\ll (\log R)^{2k} \sum_{d,e < R} \mu^2(d)\mu^2(e)k^{\omega([d,e])} \ll (\log R)^{2k} \sum_{r < R^2} \mu^2(r)(3k)^{\omega(r)} \\ &\ll R^2(\log R)^{2k} \sum_{r < R^2} \frac{\mu^2(r)(3k)^{\omega(r)}}{r} \ll R^2(\log R)^{2k} \prod_{p < R^2} \left(1 + \frac{3k}{p}\right) \\ &\ll R^2(\log R)^{5k}. \end{aligned}$$

Therefore $S_{02} \ll N$ if $R \leq N^{1/2}(\log N)^{-3k}$.

To finish, we need to establish the bound (4.3) on $|\lambda_d|$. From (3.13), we see that if $d \leq R$ and d is squarefree, then

$$\begin{aligned} |\lambda_d| &= \mathfrak{S}(\mathcal{L}) \frac{f(d)}{f_1(d)} \sum'_{\substack{r < R/d \\ (r,d)=1}} \frac{\mu^2(r)}{f_1(r)} P\left(\frac{\log R/dr}{\log R}\right) \\ &\leq \mathfrak{S}(\mathcal{L}) \sup_{0 \leq u \leq 1} |P(u)| \sum_{\delta|d} \frac{1}{f_1(\delta)} \sum'_{\substack{r < R/\delta \\ (r,d)=1}} \frac{\mu^2(r)}{f_1(r)} \\ &\ll \sum'_{r < R} \frac{\mu^2(r)}{f_1(r)} \ll (\log R)^k, \end{aligned}$$

where we have used Lemma 3 in the last line.

5. Proof of Theorem 8, Initial Steps

From the definition of $\mathcal{S}_{1,j}$, we see that

$$\mathcal{S}_{1,j} = \sum_{d,e} \lambda_d \lambda_e \sum_{\substack{N < n \leq 2N \\ [d,e]|P_{\mathcal{L}}(n)}} \beta(L_j(n)). \tag{5.1}$$

We analyze the inner sum in the next lemma.

LEMMA 5. *Suppose that u is squarefree, $(u, A) = 1$, and all prime divisors of u are less than R . Define*

$$M_j(u) = \sum_{\substack{N < n \leq 2N \\ u|P_{\mathcal{L}}(n)}} \beta(L_j(n)).$$

Then

$$\begin{aligned} M_j(u) &= \frac{\tau_{k-1}(u)}{\phi(a_j u)} \sum_{\substack{Y < p \leq N^{1/2} \\ p \nmid u}} \pi^b(a_j N/p) + \sum_{\substack{Y < p < R \\ p|u}} \frac{\tau_{k-1}(u/p)}{\phi(a_j u/p)} \pi^b(a_j N/p) + O(\tau_k(u)) \\ &\quad + O(\tau_{k-1}(u)\Delta_{\beta}^*(AN; a_j u)) + O\left(\tau_{k-1}(u) \sum_{p|u} \Delta^*(AN/p; a_j u/p)\right). \end{aligned} \tag{5.2}$$

Proof. Assume that $u|P_{\mathcal{L}}(n)$ and let $u_i = (P_{\mathcal{L}}(n), u)$. Then $u = u_1 \dots u_k$, where each $u_i|P_i(n)$. Moreover, we claim that this decomposition is unique because \mathcal{L} satisfies Hypothesis 1. To justify this, assume that the decomposition is not unique. Then there must be some prime

p such that $p|L_i(n)$ and $p|L_j(n)$ for distinct values of i and j . We conclude that $p|(a_i b_j - a_j b_i)$; this, however, contradicts Hypothesis 1.

Therefore,

$$\sum_{\substack{N < n \leq 2N \\ u|P_{\mathcal{L}}(n)}} \beta(L_j(n)) = \sum_{u_1 \dots u_k = u} \sum_{\substack{N < n \leq 2N \\ u_i | L_i(n) \\ i=1, \dots, k}} \beta(L_j(n)). \tag{5.3}$$

Set $L_j(n) = m$. Then $a_j N + b_j < m \leq 2a_j N + b_j$. Moreover, $u_j | m$. Now when $\beta(m) \neq 0$, m has exactly one prime divisor p with $p \leq N^{1/2}$, and all prime divisors of u are less than $R \leq N^{\theta/2} \leq N^{1/2}$. Consequently, we may assume that either $u_j = 1$ or $u_j = p$ for some prime $p < R$. In the latter case, our definition of β implies that we may also assume $p > Y$.

From our definition of m , we also have

$$m \equiv b_j \pmod{a_j} \quad \text{and} \quad a_i m \equiv a_i b_j - a_j b_i \pmod{u_i} \quad \text{for } i \neq j.$$

We use the Chinese remainder theorem to combine these into one congruence

$$m \equiv m_0 \pmod{a_j u / u_j}.$$

Observe that m_0 is relatively prime to $a_j u / u_j$ by Hypothesis 1; the condition $(u, A) = 1$ implies that u is coprime to a_i, a_j , and $a_i b_j - a_j b_i$.

Now we fix values of u_1, \dots, u_k . The inner sum of (5.3) is

$$\sum_{\substack{a_j N < m \leq 2a_j N \\ m \equiv m_0 \pmod{a_j u / u_j}}} \beta(m) + O(1).$$

Summing the error term over all values of u_1, \dots, u_k gives the first error term in (5.2).

Next, we consider the effect of different values of u_j . First, we assume that $u_j = 1$. Then

$$\begin{aligned} \sum_{\substack{a_j N < m \leq 2a_j N \\ m \equiv m_0 \pmod{a_j u}}} \beta(m) &= \frac{\pi_{\beta, a_j u}(a_j N)}{\phi(a_j u)} + \Delta_{\beta}(a_j N; a_j u, m_0) \\ &= \frac{1}{\phi(a_j u)} \sum_{\substack{Y < p \leq N^{1/2} \\ p \nmid u}} \pi^b(a_j N/p) + \Delta_{\beta}(a_j N; a_j u, m_0). \end{aligned} \tag{5.4}$$

Now, assume that $u_j = p$ for some prime $p, Y < p < R$. Let \bar{p} be the inverse of $p \pmod{a_j u / p}$. Then

$$\begin{aligned} \sum_{\substack{a_j N < m \leq 2a_j N \\ m \equiv m_0 \pmod{a_j u / u_j}}} \beta(m) &= \sum_{\substack{\frac{a_j N}{p} < p_2 \leq \frac{2a_j N}{p} \\ p_2 \equiv m_0 \bar{p} \pmod{a_j u / p}}} \beta(pp_2) \\ &= \pi^b(a_j N/p; a_j u/p, m_0 \bar{p}) \\ &= \frac{\pi^b(a_j N/p)}{\phi(a_j u/p)} + \Delta(a_j N/p; a_j u/p, \bar{p} m_0). \end{aligned} \tag{5.5}$$

We now sum (5.4) and (5.5) over all choices of u_1, u_2, \dots, u_k with $u_1 u_2 \dots u_k = u$ to finish the proof of the lemma. □

Returning to the estimate of $\mathcal{S}_{1,j}$, we inject Lemma 5 into (5.1). The first two error terms contribute

$$\ll \sum_{d, e < R} |\lambda_d \lambda_e| \tau_k([d, e]) \{1 + \Delta_{\beta}^*(AN, a_j[d, e])\}.$$

Using (4.3), we find that this is

$$\ll (\log R)^{2k} \sum_{r < R^2} \mu^2(r)(3k + 3)^{\omega(r)} \{1 + \Delta_\beta^*(AN, a_j r)\}.$$

By Lemma 2, there is some constant C such that if $R \leq N^{\theta/2}(\log N)^{-C}$, then the above is $\ll N$.

The contribution of the third error term requires a slightly more elaborate analysis. After using (4.3), we find that this contribution is

$$\begin{aligned} &\ll (\log R)^{2k} \sum_{r < R^2} \mu^2(r)(3k)^{\omega(r)} \sum_{p|r; p < R} \Delta^*(AN/p, a_j r/p) \\ &\ll (\log R)^{2k} \sum_{p < R} \sum_{m < R^2/p} \mu^2(m)(3k)^{\omega(m)} \Delta^*(AN/p, a_j m). \end{aligned}$$

We use Lemma 1 to bind the innermost sum. If $R \leq N^{\theta/2}(\log N)^{-C}$ for some sufficiently large C , then the above is

$$\sum_{p < R} \frac{N}{p(\log N/p)} \ll N.$$

We close this section by updating our progress on $\mathcal{S}_{1,j}$. Till now, we have

$$\mathcal{S}_{1,j} = \frac{1}{\phi(a_j)} \sum'_{Y < p \leq N^{1/2}} \pi^b \left(\frac{a_j N}{p} \right) T_p + O(N),$$

where we define

$$T_p = \sum_{\substack{d,e \\ p \nmid [d,e]}} \frac{\lambda_d \lambda_e \tau_{k-1}([d,e])}{\phi([d,e])} + \sum_{\substack{d,e \\ p \mid [d,e]}} \frac{\lambda_d \lambda_e \tau_{k-1}([d,e]/p)}{\phi([d,e]/p)}. \tag{5.6}$$

Note that in the sum defining T_p , we implicitly have the conditions $d < R$ and $e < R$ because we are assuming that $\lambda_d = 0$ if $d \geq R$. Therefore, if $p \geq R$, the second sum in (5.6) is empty, and the condition that $p \nmid [d,e]$ is vacuous. In other words, if $R \leq p < N^{1/2}$, then

$$T_p = \sum_{d,e} \frac{\lambda_d \lambda_e \tau_{k-1}([d,e])}{\phi([d,e])}. \tag{5.7}$$

However, when $p < R$, the sum T_p is more complicated, and we will analyze this case in more detail in the next section.

Before closing this section, we use the prime number theorem to write

$$\pi^b \left(\frac{a_j N}{p} \right) = \frac{a_j N}{p \log N} \alpha(p) + O \left(\frac{N}{p(\log N)^2} \right),$$

where

$$\alpha(p) = \frac{\log N}{\log(N/p)}.$$

Note that by Hypothesis 1, a_j and A have exactly the same prime divisors. Consequently, $a_j/\phi(a_j) = A/\phi(A)$, and

$$\mathcal{S}_{1,j} = \frac{A}{\phi(A)} \frac{N}{\log N} \sum'_{Y < p \leq N^{1/2}} \frac{\alpha(p)}{p} T_p + O \left(N + \frac{N}{(\log N)^2} \sum'_{Y < p \leq N^{1/2}} \frac{T_p}{p} \right). \tag{5.8}$$

6. Evaluation of T_p

Analogous to the function f defined in (3.9), we define

$$f^*(d) = \frac{\phi(d)}{\tau_{k-1}(d)} \tag{6.1}$$

whenever d is squarefree and relatively prime to A . We use this to define

$$T_\delta = \sum'_{d,e} \frac{\lambda_d \lambda_e}{f^*([d, e, \delta]/\delta)}. \tag{6.2}$$

When $\delta = p$, (6.2) reduces to the earlier definition of T_p . We will analyze the more general quantity T_δ ; this provides additional insight at the cost of little extra complication of detail.

An expression similar to T_δ occurs in Selberg’s $\Lambda^2 \Lambda^-$ sieve. See, for example, the last displayed equation in [28, p. 85] or in [14, equation (1.9)]. In our notation, those results can be stated as

$$\sum_{d,e} \frac{\lambda_d \lambda_e}{f([d, e, \delta]/\delta)} = \sum'_{(r,\delta)=1} \frac{\mu^2(r)}{f_1(r)} \left(\sum_{s|\delta} \mu(s) y_{rs} \right)^2.$$

Our next lemma is an analog of this result with f replaced by f^* .

LEMMA 6. *If δ is squarefree and relatively prime to A , then*

$$T_\delta = \sum'_{(r,\delta)=1} \frac{\mu^2(r)}{f_1^*(r)} \left(\sum_{s|\delta} \mu(s) y_{rs}^* \right)^2,$$

where

$$f_1^*(d) = \mu * f^*(d) = \prod_{p|d} \frac{p-k}{k-1}$$

whenever d is squarefree and $(d, A) = 1$, and

$$y_r^* = \frac{\mu^2(r)r}{\phi(r)} \sum'_m \frac{y_{mr}}{\phi(m)}. \tag{6.3}$$

Proof. Define $g_\delta^*(d) = g^*(d)$ by the relation

$$g^*(d) = f^* \left(\frac{d}{(d, \delta)} \right).$$

If p is a prime, $p \nmid A$, then

$$g^*(p) = \begin{cases} f^*(p) & \text{if } p \nmid \delta \\ 1 & \text{if } p|\delta. \end{cases}$$

With this notation, we may write

$$T_\delta = \sum'_{d,e} \frac{\lambda_d \lambda_e g^*((d, e))}{g^*(d)g^*(e)} = \sum'_{d,e} \frac{\lambda_d \lambda_e}{g^*(d)g^*(e)} \sum_{\substack{r|d \\ r|e}} g_1^*(r),$$

where $g_1^* = g^* * \mu$. Note that

$$g_1^*(p) = \begin{cases} f_1^*(p) & \text{if } p \nmid \delta, \\ 0 & \text{if } p|\delta. \end{cases}$$

After changing the order of summation in the last sum, we find that

$$T_\delta = \sum'_{(r,\delta)=1} g_1^*(r) \left(\sum'_{r|d} \frac{\lambda_d}{g^*(d)} \right)^2.$$

The condition that $(r, \delta) = 1$ may be inserted because $g_1^*(r) = 0$ if $(r, \delta) \neq 1$.

Define

$$w_r^* = \mu(r)g_1^*(r) \sum'_d \frac{\lambda_{dr}}{g^*(dr)}.$$

Then

$$T_\delta = \sum'_{(r,\delta)=1} \frac{\mu^2(r)}{g_1^*(r)} (w_r^*)^2. \tag{6.4}$$

Assume henceforth that $(r, \delta) = 1$. Then

$$\begin{aligned} w_r^* &= \frac{\mu(r)g_1^*(r)}{g^*(r)} \sum'_d \frac{\lambda_{dr}}{g^*(d)} \\ &= \frac{\mu(r)g_1^*(r)}{g^*(r)} \sum'_d \frac{\mu(dr)f(dr)}{g^*(d)} \sum'_t \frac{y_{drt}}{f_1(drt)} \\ &= \frac{\mu^2(r)g_1^*(r)f(r)}{g^*(r)f_1(r)} \sum'_m \frac{y_{mr}}{f_1(m)} \sum'_{d|m} \frac{\mu(d)f(d)}{g^*(d)}. \end{aligned}$$

We note that $g_1^*(r) = f_1^*(r)$ and $g^*(r) = f^*(r)$ because of our hypothesis $(r, \delta) = 1$. Thus

$$\frac{g_1^*(r)f(r)}{g^*(r)f_1(r)} = \frac{f_1^*(r)f(r)}{f^*(r)f_1(r)} = \frac{r}{\phi(r)}.$$

Next, we consider the sum

$$\sum'_{d|m} \frac{\mu(d)f(d)}{g^*(d)} = \sum'_{d|m} \frac{\mu(d)f(d)}{f^*\left(\frac{d}{(d,\delta)}\right)}.$$

We write $d = d_1d_2$, with $(d_1, \delta) = 1$ and $d_2|\delta$. The above sum is then

$$\sum'_{\substack{d_1|m \\ (d_1,\delta)=1}} \frac{\mu(d_1)f(d_1)}{f^*(d_1)} \sum'_{\substack{d_2|m \\ d_2|\delta}} \mu(d_2)f(d_2). \tag{6.5}$$

The first factor in (6.5) is

$$\sum'_{\substack{d_1|m \\ (d_1,\delta)=1}} \frac{\mu(d_1)f(d_1)}{f^*(d_1)} = \prod_{\substack{p|m \\ p \nmid \delta}} \left(1 - \frac{p(k-1)}{(p-1)k} \right) = \frac{f_1(m/(m,\delta))}{\phi(m/(m,\delta))}.$$

The second factor in (6.5) is

$$\sum_{d_2|(m,\delta)} \mu(d_2)f(d_2) = \prod_{p|(m,\delta)} (1 - f(p)) = \mu((m,\delta))f_1((m,\delta)).$$

We conclude that the expression in (6.5) is

$$\frac{f_1(m)}{\phi(m)} \mu((m,\delta))\phi((m,\delta)).$$

Now

$$\mu((m, \delta))\phi((m, \delta)) = \sum_{\substack{s|m \\ s|\delta}} \mu(s)s,$$

so

$$w_r^* = \frac{\mu^2(r)r}{\phi(r)} \sum_m \frac{y_{mr}}{\phi(m)} \sum_{\substack{s|m \\ s|\delta}} \mu(s)s.$$

The definition of w_r^* depends on r as well as δ . Using the definition of y_r^* given in (6.3), we find that

$$w_r^* = \sum_{s|\delta} \mu(s)y_{rs}^*.$$

Inserting this into (6.4) completes the proof of the lemma. □

When $\delta = p$, Lemma 6 becomes

$$T_p = \sum'_{\substack{r \\ (r,p)=1}} \frac{\mu^2(r)}{f_1^*(r)} (y_r^* - y_{rp}^*)^2. \tag{6.6}$$

Now $y_{rp}^* = 0$ if $r \geq R/p$, so

$$T_p = \sum'_{\substack{r < R/p \\ (r,p)=1}} \frac{\mu^2(r)}{f_1^*(r)} (y_r^* - y_{rp}^*)^2 + \sum'_{\substack{R/p \leq r < R \\ (r,p)=1}} \frac{\mu^2(r)}{f_1^*(r)} (y_r^*)^2.$$

When $p \geq R$, the second sum above is empty, and the condition $(p, r) = 1$ in the first sum is vacuous. In other words, if $p \geq R$, then

$$T_p = \sum'_{r < R} \frac{\mu^2(r)}{f_1^*(r)} (y_r^*)^2 = T_1.$$

This is equivalent to the observation that we made earlier in (5.7).

Now we turn our attention to the sum

$$\sum'_{Y < p \leq N^{1/2}} \frac{\alpha(p)}{p} T_p$$

that appears in the main term of (5.8). Using the above observations on T_p , we find that

$$\sum'_{Y < p \leq N^{1/2}} \frac{\alpha(p)}{p} T_p = S_1 + S_2 + S_3, \tag{6.7}$$

where

$$S_1 = \sum'_{Y < p < R} \frac{\alpha(p)}{p} \sum'_{\substack{r < R/p \\ (r,p)=1}} \frac{\mu^2(r)}{f_1^*(r)} (y_r^* - y_{rp}^*)^2, \tag{6.8}$$

$$S_2 = \sum'_{Y < p < R} \frac{\alpha(p)}{p} \sum'_{\substack{R/p \leq r < R \\ (r,p)=1}} \frac{\mu^2(r)}{f_1^*(r)} (y_r^*)^2, \tag{6.9}$$

$$S_3 = \sum'_{R \leq p < N^{1/2}} \frac{\alpha(p)}{p} \sum'_{r < R} \frac{\mu^2(r)}{f_1^*(r)} (y_r^*)^2. \tag{6.10}$$

LEMMA 7. Assume that $r < R$, $(r, A) = 1$, and r is squarefree. Let y_r^* be as defined in (6.3). Then

$$y_r^* = \frac{\phi(A)}{A} \mathfrak{S}(\mathcal{L})(\log R) \tilde{P}\left(\frac{\log R/r}{\log R}\right) + O(L(r)),$$

where

$$L(r) = 1 + \sum_{p|r} \frac{\log p}{p}.$$

Proof. From (6.3) and (3.14), we see that

$$y_r^* = \mu^2(r) \frac{r}{\phi(r)} \mathfrak{S}(\mathcal{L}) \sum_{\substack{m \leq R/r \\ (m, rA)=1}} \frac{\mu^2(m)}{\phi(m)} P\left(\frac{\log R/rm}{\log R}\right). \tag{6.11}$$

We apply Lemma 4 with

$$\gamma(p) = \begin{cases} 1 & \text{if } p \nmid rA, \\ 0 & \text{if } p|rA. \end{cases}$$

Then $c_\gamma = \phi(rA)/rA$ and condition (2.4) is satisfied with $\kappa = 1$ and

$$L = \sum_{p|rA} \frac{\log p}{p} + O(1).$$

We are regarding A as fixed, so $L \ll L(r)$. Using Lemma 4 with

$$F(x) = P\left(x \frac{\log R/r}{\log R}\right),$$

we obtain

$$\begin{aligned} \sum_{\substack{m < R/r \\ (m, rA)=1}} \frac{\mu^2(m)}{\phi(m)} P\left(\frac{\log R/rm}{\log R}\right) &= \frac{\phi(rA)}{rA} (\log(R/r)) \int_0^1 P\left(\frac{\log R/r}{\log R} (1-x)\right) dx \\ &\quad + O\left(\frac{\phi(r)}{r} L(r)\right). \end{aligned}$$

The desired results follows by making an appropriate change of variables in the integral on the right-hand side. □

LEMMA 8. For $u \geq 1$, define

$$G^*(u) := \sum'_{r < u} \frac{\mu^2(r)}{f_1^*(r)}.$$

Then

$$G^*(u) = \frac{A}{\phi(A)} \frac{(\log u)^{k-1}}{\mathfrak{S}(\mathcal{L})(k-1)!} + E^*(u), \tag{6.12}$$

where $E^*(u) \ll (\log(2u))^{k-2}$.

Proof. We apply Lemma 3 with

$$\gamma(p) = \begin{cases} \frac{p(k-1)}{p-1} & \text{if } p \nmid A, \\ 0 & \text{if } p|A, \end{cases} \tag{6.13}$$

and $\kappa = k - 1$. As noted in (3.4), every prime $p \leq k$ divides A , so

$$\frac{\gamma(p)}{p} \leq 1 - \frac{1}{k}$$

for $p \nmid A$. Therefore, (2.3) is satisfied with $A_1 = k$. We are treating A as fixed, so (2.4) is satisfied with $L \ll 1$. Moreover,

$$c_\gamma = \prod_{p|A} \left(1 - \frac{1}{p}\right)^{k-1} \prod_{p \nmid A} \left(1 - \frac{k-1}{p-1}\right)^{-1} \left(1 - \frac{1}{p}\right)^{k-1} = \frac{A}{\phi(A)} \frac{1}{\mathfrak{S}(\mathcal{L})},$$

and (6.12) follows from Lemma 3. □

LEMMA 9. *If p is prime, then $T_p \ll (\log R)^{k+1}$.*

Proof. From (6.6),

$$T_p = \sum'_{\substack{r \\ (r,p)=1}} \frac{\mu^2(r)}{f_1^*(r)} (y_r^* - y_{rp}^*)^2.$$

We are regarding P as fixed, so Lemma 7 implies that $y_j^* \ll \log R$ for any $j < R$, and $y_j^* = 0$ if $j \geq R$. Therefore

$$T_p \ll (\log R)^2 \sum'_{r < R} \frac{\mu^2(r)}{f_1^*(r)},$$

and the lemma now follows by using (6.12) with $u = R$. □

Using the above lemma, we see that the second error term in (5.8) is

$$\frac{N}{(\log N)^2} (\log R)^{k+1} \sum_{p \leq N^{1/2}} \frac{1}{p} \ll N (\log \log N) (\log N)^{k-1}. \tag{6.14}$$

Combining (5.8), (6.14), and (6.7), we now have

$$\mathcal{S}_{1,j} = \frac{A}{\phi(A)} \frac{N}{\log N} (S_1 + S_2 + S_3) + O(N (\log \log N) (\log N)^{k-1}). \tag{6.15}$$

To finish the proof of Theorem 8, we will show that when $i = 1, 2$, or 3 ,

$$S_i = \frac{\phi(A)}{A} \frac{\mathfrak{S}(\mathcal{L})(\log R)^{k+1}}{(k-2)!} J_i + O((\log \log R)(\log R)^k),$$

where J_1, J_2, J_3 are as defined in the statement of Theorem 8.

7. Completion of proof of Theorem 8

LEMMA 10. *Let S_1 be as defined in (6.8), and let J_1 be as defined in the statement of Theorem 8. Then*

$$S_1 = \frac{\phi(A)}{A} \frac{\mathfrak{S}(\mathcal{L})(\log R)^{k+1}}{(k-2)!} J_1 + O((\log \log R)(\log R)^k).$$

Proof. Assume that $r < R$, r is squarefree, p is a prime with $p < R/r$, $(p, r) = 1$, and $(pr, A) = 1$. By Lemma 7,

$$y_r^* - y_{rp}^* = \frac{\phi(A)}{A} \mathfrak{S}(\mathcal{L})(\log R) \int_{1-\log pr/\log R}^{1-\log r/\log R} P(x) dx + O(L(r)).$$

In the above, we have used the simple observation that

$$L(rp) = L(r) + \frac{\log p}{p} \ll L(r) + 1 \ll L(r).$$

Note also that

$$(\log R) \int_{1-\log pr/\log R}^{1-\log r/\log R} P(x)dx \ll \log p,$$

and

$$\begin{aligned} L(r) &\leq 1 + \sum_{p \leq \log R} \frac{\log p}{p} + \sum_{\substack{p|r \\ p > \log R}} \frac{\log p}{p} \\ &\ll 1 + \log \log R + \frac{\log \log R}{\log R} \frac{\log R}{\log \log R} \\ &\ll \log \log R. \end{aligned} \tag{7.1}$$

In particular, $L(r) \ll \log p$ when $p > Y$. Therefore

$$\begin{aligned} (y_r^* - y_{rp}^*)^2 &= \frac{\phi(A)^2}{A^2} \mathfrak{S}(\mathcal{L})^2 (\log R)^2 \left(\tilde{P} \left(\frac{\log R/r}{\log R} \right) - \tilde{P} \left(\frac{\log R/rp}{\log R} \right) \right)^2 \\ &\quad + O((\log p)L(r)). \end{aligned}$$

We use this in the definition of S_1 to obtain

$$\begin{aligned} S_1 &= \frac{\phi(A)^2}{A^2} \mathfrak{S}(\mathcal{L})^2 (\log R)^2 \sum'_{Y < p < R} \frac{\alpha(p)}{p} \sum'_{\substack{r < R/p \\ (r,p)=1}} \frac{\mu^2(r)}{f_1^*(r)} \left(\tilde{P} \left(\frac{\log R/r}{\log R} \right) - \tilde{P} \left(\frac{\log R/rp}{\log R} \right) \right)^2 \\ &\quad + O \left(\sum_{p < R} \frac{1}{p} \sum'_{r < R/p} \frac{\mu^2(r)}{f_1^*(r)} (\log p)L(r) \right) \\ &= S_{11} + O(S_{12}), \end{aligned}$$

say.

For S_{12} , we reverse the order of summation and use (6.12) to obtain

$$\begin{aligned} S_{12} &= \sum'_{r < R} \frac{\mu^2(r)}{f_1^*(r)} L(r) \sum_{p < R/r} \frac{\log p}{p} \ll (\log \log R)(\log R) \sum'_{r < R} \frac{\mu^2(r)}{f_1^*(r)} \\ &\ll (\log \log R)(\log R)^k. \end{aligned} \tag{7.2}$$

Now we consider S_{11} . We write this as $S_{13} - S_{14}$, where

$$S_{13} = \frac{\phi(A)^2}{A^2} \mathfrak{S}(\mathcal{L})^2 (\log R)^2 \sum'_{Y < p < R} \frac{\alpha(p)}{p} \sum'_{r < R/p} \frac{\mu^2(r)}{f_1^*(r)} \left(\tilde{P} \left(\frac{\log R/r}{\log R} \right) - \tilde{P} \left(\frac{\log R/rp}{\log R} \right) \right)^2, \tag{7.3}$$

and S_{14} is the same sum with the extra condition that $p|r$.

For S_{14} , we note that

$$\tilde{P} \left(\frac{\log R/r}{\log R} \right) - \tilde{P} \left(\frac{\log R/rp}{\log R} \right) = \int_{\log(R/rp)/\log R}^{\log(R/r)/\log R} P(t)dt \ll \frac{\log p}{\log R}.$$

We also note that $f_1^*(p) = (p - k)/(k - 1) \gg p$. Making the change of variables $r = mp$, we get

$$S_{14} \ll (\log R)^2 \sum'_{Y < p < R} \frac{1}{p^2} \sum'_{m < R/p} \frac{\mu^2(m)}{f_1^*(m)} \left(\frac{\log p}{\log R} \right)^2$$

$$\ll (\log R)^{k-1} \sum_{p < R} \frac{(\log p)^2}{p^2}$$

by Lemma 8. The last sum converges, so

$$S_{14} \ll (\log R)^{k-1}. \tag{7.4}$$

For S_{13} , we evaluate the inner sum using Lemma 4 with $z = R$, $g(d) = 1/f_1^*(d)$, $\kappa = k - 1$, γ as defined in (6.13), and

$$F\left(\frac{\log R/r}{\log R}\right) = \begin{cases} \left\{ \tilde{P}\left(\frac{\log R/r}{\log R}\right) - \tilde{P}\left(\frac{\log R/rp}{\log R}\right) \right\}^2 & \text{if } r < R/p, \\ 0 & \text{if } R/p \leq r < R. \end{cases}$$

If we set $y = \log p / \log R$ and $x = \log r / \log R$, then the last is equivalent to

$$F(1 - x) = F_p(1 - x) = \begin{cases} \left(\tilde{P}(1 - x) - \tilde{P}(1 - x - y) \right)^2 & \text{if } x < 1 - y, \\ 0 & \text{if } 1 - y \leq x < 1. \end{cases}$$

Making the substitution $w = 1 - x$, we see that this is the same as

$$F(w) = F_p(w) = \begin{cases} \left(\tilde{P}(w) - \tilde{P}(w - y) \right)^2 & \text{if } y \leq w \leq 1, \\ 0 & \text{if } 0 \leq w < y. \end{cases}$$

From Lemma 4, we find that

$$\sum'_{r < R/p} \frac{\mu^2(r)}{f_1^*(r)} \left(\tilde{P}\left(\frac{\log R/r}{\log R}\right) - \tilde{P}\left(\frac{\log R/rp}{\log R}\right) \right)^2 \tag{7.5}$$

$$= \frac{A}{\phi(A)} \frac{(\log R)^{k-1}}{\mathfrak{S}(\mathcal{L})(k-2)!} V_1\left(\frac{\log p}{\log R}\right) + O(M(F_p)(\log R)^{k-2}).$$

where

$$V_1(y) = \int_0^{1-y} \left\{ \tilde{P}(1-x) - \tilde{P}(1-x-y) \right\}^2 x^{k-2} dx.$$

Observe that if $y \leq x \leq 1$, then

$$|F_p(x)| = \left(\int_{x-y}^x P(t) dt \right)^2 \leq y^2 \sup_{t \in [0,1]} |P(t)| \ll 1,$$

where the implied constant depends on P but not on p . Similarly, $F'_p(x) \ll 1$, and therefore $M(F_p) \ll 1$ uniformly in p . The error term in (7.5) thus contributes

$$\ll (\log R)^k \sum_{p < R} \frac{1}{p} \ll (\log \log R)(\log R)^k \tag{7.6}$$

to S_{13} . Incorporating the contribution of the main term from (7.5), we now have

$$S_{13} = \frac{\phi(A)}{A} \frac{\mathfrak{S}(\mathcal{L})(\log R)^{k+1}}{(k-2)!} \sum'_{Y < p < R} \frac{\alpha(p)}{p} V_1\left(\frac{\log p}{\log R}\right) + O((\log \log R)(\log R)^k). \tag{7.7}$$

Now let $Z(u)$ be defined by the relation

$$\sum_{p \leq u} \log p = u + Z(u). \tag{7.8}$$

From the classical form of the prime number theorem, we know that

$$Z(u) \ll u \exp(-c\sqrt{\log u})$$

for some absolute constant c . Therefore the sum in (7.7) is

$$\int_Y^R \alpha(u) V_1 \left(\frac{\log u}{\log R} \right) \frac{du}{u \log u} + \int_Y^R \alpha(u) V_1 \left(\frac{\log u}{\log R} \right) \frac{dZ(u)}{u \log u}. \tag{7.9}$$

In the first integral, we make the change of variable $u = R^y$, and we set

$$b = \frac{\log N}{\log R} \tag{7.10}$$

to obtain

$$\int_Y^R \alpha(u) V_1 \left(\frac{\log u}{\log R} \right) \frac{du}{u \log u} = \int_{b\eta}^1 \frac{b}{y(b-y)} V_1(y) dy = J_1. \tag{7.11}$$

Note that we have used the fact that $\log Y / \log N = \eta$. Comparing the definitions of b and B (see (1.18)), we see that

$$b = \left(\frac{\vartheta}{2} - \frac{C \log \log N}{\log N} \right)^{-1} = B + O(\log \log R / \log R).$$

We may, therefore, replace b by B on the right-hand side of (7.11) at the cost of an error term $O(\log \log R / \log R)$. The first integral in (7.9) is thus

$$= \int_{B\eta}^1 \frac{B}{B-y} V_1(y) dy + O \left(\frac{\log \log R}{\log R} \right) = J_1 + O \left(\frac{\log \log R}{\log R} \right). \tag{7.12}$$

We write the second integral in (7.9) as

$$\int_Y^R F_1(u) dZ(u) = F_1(R)Z(R) - F_1(Y)Z(Y) - \int_Y^R Z(u) F_1'(u) du,$$

where

$$F_1(u) = \frac{\alpha(u)}{u \log u} V_1 \left(\frac{\log u}{\log R} \right).$$

Now $V_1(y) \ll y$, so

$$F_1(u) \ll \frac{1}{u \log R}.$$

Moreover,

$$\begin{aligned} F_1'(u) &= F_1(u) \frac{d}{du} \log F_1(u) \\ &= F_1(u) \left\{ \frac{1}{u \log(N/u)} - \frac{V_1' \left(\frac{\log u}{\log R} \right)}{V_1 \left(\frac{\log u}{\log R} \right)} \frac{1}{u \log R} - \frac{1}{u} - \frac{1}{u \log u} \right\}, \end{aligned}$$

so

$$|F_1'(u)| \ll \frac{|F_1(u)|}{u} \ll \frac{1}{u^2 \log R}.$$

Therefore,

$$\int_Y^R Z(u) F_1'(u) du \ll \int_Y^R \frac{\exp(-c\sqrt{\log u})}{u \log R} du \ll (\log R)^{-1}.$$

We also note that

$$|F_1(R)Z(R)| + |F_1(Y)Z(Y)| \ll (\log R)^{-1}.$$

From the last two estimates, (7.7) and (7.12), we conclude that

$$S_{13} = \frac{\phi(A)}{A} \frac{\mathfrak{S}(\mathcal{L})(\log R)^{k+1}}{(k-2)!} J_1 + O((\log \log R)(\log R)^k).$$

We combine this with (7.2) and (7.4) to complete the proof. □

LEMMA 11. *Let S_2 be as defined in (6.9), and let J_2 be as defined in the statement of Theorem 8. Then*

$$S_2 = \frac{\phi(A)}{A} \frac{\mathfrak{S}(\mathcal{L})(\log R)^{k+1}}{(k-2)!} J_2 + O((\log \log R)(\log R)^k).$$

Proof. From Lemma 7 and (7.1), we see that

$$(y_r^*)^2 = \frac{\phi(A)^2}{A^2} \mathfrak{S}(\mathcal{L})^2 (\log R)^2 \tilde{P} \left(\frac{\log R/r}{\log R} \right)^2 + O(L(r) \log R). \tag{7.13}$$

Therefore

$$\begin{aligned} S_2 &= \frac{\phi(A)^2}{A^2} \mathfrak{S}(\mathcal{L})^2 (\log R)^2 \sum'_{Y < p < R} \frac{\alpha(p)}{p} \sum'_{\substack{R/p \leq r < R \\ (r,p)=1}} \frac{\mu^2(r)}{f_1^*(r)} \tilde{P} \left(\frac{\log R/r}{\log R} \right)^2 \\ &\quad + O \left((\log R) \sum'_{Y < p < R} \frac{1}{p} \sum'_{R/p \leq r < R} \frac{\mu^2(r)}{f_1^*(r)} L(r) \right) \\ &= S_{21} + O(S_{22}), \end{aligned} \tag{7.14}$$

say.

We first consider S_{22} . From the above definition, we see that

$$S_{22} \ll (\log \log R)(\log R) \sum'_{r < R} \frac{\mu^2(r)}{f_1^*(r)} L(r).$$

Now

$$\sum'_{r < R} \frac{\mu^2(r)}{f_1^*(r)} L(r) = \sum'_{r < R} \frac{\mu^2(r)}{f_1^*(r)} + \sum'_{r < R} \frac{\mu^2(r)}{f_1^*(r)} \sum_{p|r} \frac{\log p}{p},$$

and

$$\begin{aligned} \sum'_{r < R} \frac{\mu^2(r)}{f_1^*(r)} \sum_{p|r} \frac{\log p}{p} &= \sum'_{p < R} \frac{\log p}{p f_1^*(p)} \sum'_{t < R/p; (t,p)=1} \frac{\mu^2(t)}{f_1^*(t)} \\ &\ll \sum'_{p < R} \frac{\log p}{p^2} (\log R)^{k-1} \ll (\log R)^{k-1}. \end{aligned}$$

Therefore,

$$\sum'_{r < R} \frac{\mu^2(r)}{f_1^*(r)} L(r) \ll (\log R)^{k-1}, \tag{7.15}$$

and

$$S_{22} \ll (\log \log R)(\log R)^k. \tag{7.16}$$

Now $S_{21} = S_{23} - S_{24}$, where

$$S_{23} = \frac{\phi(A)^2}{A^2} \mathfrak{S}(\mathcal{L})^2 (\log R)^2 \sum'_{Y < p < R} \frac{\alpha(p)}{p} \sum'_{R/p \leq r < R} \frac{\mu^2(r)}{f_1^*(r)} \tilde{P} \left(\frac{\log R/r}{\log R} \right)^2, \tag{7.17}$$

and S_{24} is the same sum with the extra condition that $p|r$.

For S_{24} , we begin by noting that $\tilde{P}(y) \ll y$. Therefore, if $R/p \leq r < R$, then

$$\tilde{P} \left(\frac{\log R/r}{\log R} \right) \ll \left(\frac{\log p}{\log R} \right)^2.$$

Consequently,

$$S_{24} \ll (\log R)^2 \sum_{p < R} \frac{(\log p)^2}{p^2} \sum'_{t < R} \frac{\mu^2(t)}{f_1^*(t)} \ll (\log R)^{k-1}. \tag{7.18}$$

Using Lemma 4, we find that the innermost sum in S_{23} is

$$\frac{A}{\phi(A)} \frac{(\log R)^{k-1}}{\mathfrak{S}(\mathcal{L})(k-2)!} V_2 \left(\frac{\log p}{\log R} \right) + O((\log R)^{k-2}),$$

where

$$V_2(y) = \int_{1-y}^1 \tilde{P}(1-x)^2 x^{k-2} dx.$$

Inserting this into (7.17), we find that

$$S_{23} = \frac{\phi(A)}{A} \frac{\mathfrak{S}(\mathcal{L})(\log R)^{k+1}}{(k-2)!} \sum'_{Y < p < R} \frac{\alpha(p)}{p} V_2 \left(\frac{\log p}{\log R} \right) + O((\log \log R)(\log R)^k). \tag{7.19}$$

The sum in the main term is

$$\int_Y^R \alpha(u) V_2 \left(\frac{\log u}{\log R} \right) \frac{du}{u \log u} + \int_Y^R \alpha(u) V_2 \left(\frac{\log u}{\log R} \right) \frac{dZ(u)}{u \log u}, \tag{7.20}$$

where $Z(u)$ was defined in (7.8). In the first integral, we let $u = R^y$ to obtain

$$\int_Y^R \alpha(u) V_2 \left(\frac{\log u}{\log R} \right) \frac{du}{u \log u} = \int_{b\eta}^1 \frac{b}{y(b-y)} V_2(y) dy,$$

where $b = \log N / \log R$, as defined in (7.10). As in the proof of Lemma 10, we may replace b by B at the cost of an error term $O(\log \log R / \log R)$; therefore,

$$\int_Y^R \alpha(u) V_2 \left(\frac{\log u}{\log R} \right) \frac{du}{u \log u} = J_2 + O \left(\frac{\log \log R}{\log R} \right).$$

The second integral in (7.20) may be written as

$$\int_Y^R F_2(u) dZ(u),$$

where

$$F_2(u) = \frac{\alpha(u)}{u \log u} V_2 \left(\frac{\log u}{\log R} \right).$$

We estimate this by using the argument following (7.12), but with F_1 and V_1 replaced by F_2 and V_2 . Note that $V_2(y) \ll y$, so $F_2(u) \ll (u \log R)^{-1}$. The end result is that

$$\int_Y^R F_2(u) dZ(u) \ll (\log R)^{-1}.$$

We combine the above estimates to get

$$\sum'_{Y < p < R} \frac{\alpha(p)}{p} V_2 \left(\frac{\log p}{\log R} \right) = J_2 + O \left(\frac{\log \log R}{\log R} \right).$$

The proof of the lemma is completed by combining this with (7.14), (7.16), (7.18), and (7.19). □

LEMMA 12. *Let S_3 be as defined in (6.10), and let J_3 be as defined in the statement of Theorem 8. Then*

$$S_3 = \frac{\phi(A)}{A} \frac{\mathfrak{S}(\mathcal{L})(\log R)^{k+1}}{(k-2)!} J_3 + O((\log \log R)(\log R)^k).$$

Proof. Let S_3 is a product of two sums. Using (7.13), we see that the second sum is

$$\begin{aligned} \sum'_{r < R} \frac{\mu^2(r)}{f_1^*(r)} (y_r^*)^2 &= \frac{\phi(A)^2}{A^2} \mathfrak{S}(\mathcal{L})^2 (\log R)^2 \sum'_{r < R} \frac{\mu^2(r)}{f_1^*(r)} \tilde{P} \left(\frac{\log R/r}{\log R} \right)^2 \\ &\quad + O \left((\log R) \sum'_{r < R} \frac{\mu^2(r)}{f_1^*(r)} L(r) \right). \end{aligned}$$

We use Lemma 4 for the main term and (7.15) for the error term. Therefore,

$$\sum'_{r < R} \frac{\mu^2(r)}{f_1^*(r)} (y_r^*)^2 = \frac{\phi(A)}{A} \frac{\mathfrak{S}(\mathcal{L})(\log R)^{k+1}}{(k-2)!} \int_0^1 \tilde{P}(1-x)^2 x^{k-2} dx + O((\log R)^k). \tag{7.21}$$

The first sum in the definition of S_3 is

$$\int_R^{N^{1/2}} \alpha(u) \frac{du}{u \log u} + \int_R^{N^{1/2}} \alpha(u) \frac{dZ(u)}{u \log u}. \tag{7.22}$$

In the first integral, we set $u = R^y$ to get

$$\int_R^{N^{1/2}} \alpha(u) \frac{du}{u \log u} = \int_1^{b/2} \frac{b}{y(b-y)} dy.$$

As in the proofs of Lemma 10 and Lemma 11, we may replace b by B at the cost of a small error term, and therefore

$$\int_R^{N^{1/2}} \alpha(u) \frac{du}{u \log u} = \int_1^{B/2} \frac{B}{y(B-y)} dy + O \left(\frac{\log \log R}{\log R} \right).$$

Letting $F_3(u) = \alpha(u)/(u \log u)$, we see that the second integral in (7.22) is

$$\begin{aligned} \int_R^{N^{1/2}} F_3(u) dZ(u) &\ll |F_3(R)Z(R)| + |F_3(N^{1/2})Z(N^{1/2})| + \int_R^{N^{1/2}} |F_3'(u)| \exp(-c\sqrt{\log u}) du \\ &\ll (\log R)^{-1}. \end{aligned}$$

Therefore,

$$\sum'_{R \leq p < N^{1/2}} \frac{\alpha(p)}{p} = \int_1^{B/2} \frac{B}{y(B-y)} dy + O((\log R)^{-1}).$$

We combine this with (7.21) to complete the proof. □

Theorem 8 now follows by combining the previous three lemmas and (6.15).

We close this section by giving, as promised earlier, a short sketch of the proof of Theorem 9. The left-hand side of the conclusion is

$$= \sum_{d,e} \lambda_d \lambda_e Q_j([d, e]), \tag{7.23}$$

where

$$Q_j(u) = \sum_{\substack{N < n \leq 2N \\ u|P_{\mathcal{L}}(n)}} \varpi(L_j(n)).$$

This last sum can be evaluated in the same way as the related sum $M_j(u)$ considered in Lemma 5. The evaluation is simpler because only the case $u_j = 1$ occurs in this instance. The final result is

$$Q_j(u) = \frac{\tau_{k-1}(u)}{\phi(a_j u)} \pi^b(a_j N) + O(\tau_{k-1}(u) \Delta^*(AN, u)).$$

We insert this into (7.23) and use the Bombieri–Vinogradov theorem to handle the error terms. The main term is

$$\frac{\pi^b(a_j N)}{\phi(a_j)} T_1 = \frac{A}{\phi(A)} \frac{N}{\log N} T_1 + O(T_1 N (\log N)^{-2}),$$

where T_1 is given by (6.2) with $\delta = 1$. By Lemma 6, T_1 is equal to the sum considered in (7.21), and the proof is completed by appealing to the formula there.

8. Proofs of Theorems 1–3 and Corollaries

For the proof of Theorem 1, we use (3.5). For our choice of P , we take $\ell = \lceil \sqrt{k} \rceil$, and

$$P(x) = \frac{x^\ell}{\ell!}, \quad \tilde{P}(x) = \frac{x^{\ell+1}}{(\ell + 1)!}.$$

We take $Y = 1$ in the definition of β ; therefore $\eta = 0$.

From Theorems 7 and 8, we see that (cf. (3.6))

$$\mathcal{S} \sim \mathfrak{S}(\mathcal{L}) N (\log R)^k J,$$

where

$$J = \left\{ \frac{k}{B} \frac{(J_1 + J_2 + J_3)}{(k - 2)!} - \nu \frac{J_0}{(k - 1)!} \right\}. \tag{8.1}$$

Next, we write

$$\frac{J_1 + J_2}{(k - 2)!} = J_4 + J_5 + J_6, \tag{8.2}$$

where

$$\begin{aligned} J_4 &= \int_0^1 \left(\frac{1}{y} + \frac{1}{B - y} \right) \int_0^1 \tilde{P}(1 - x)^2 \frac{x^{k-2}}{(k - 2)!} dx dy, \\ J_5 &= -2 \int_0^1 \left(\frac{1}{y} + \frac{1}{B - y} \right) \int_0^{1-y} \tilde{P}(1 - x) \tilde{P}(1 - y - x) \frac{x^{k-2}}{(k - 2)!} dx dy, \\ J_6 &= \int_0^1 \left(\frac{1}{y} + \frac{1}{B - y} \right) \int_0^{1-y} \tilde{P}(1 - x - y)^2 \frac{x^{k-2}}{(k - 2)!} dx dy. \end{aligned} \tag{8.3}$$

In J_5 , we can write

$$(1 - x)^{\ell+1} = (1 - y - x)^{\ell+1} + \sum_{j=1}^{\ell+1} \binom{\ell+1}{j} y^j (1 - y - x)^{\ell+1-j} \tag{8.4}$$

and denote the corresponding integrals by $J_5^{(0)}$ and $J_5^{(1)}$, respectively. The terms J_4 , $J_5^{(0)}$, and J_6 will contribute to the main term, $J_5^{(1)}$ to the secondary term.

We will often use the evaluation ($m, n \in \mathbb{Z}^+$)

$$\int_0^1 x^m (1 - x)^n dx = \frac{m!n!}{(m + n + 1)!}, \tag{8.5}$$

which is a special case of a standard formula for the Euler beta function (see, for example, Karacuba [22, p. 46]). For later convenience, we define $A(k, \ell)$ should not be confused with the quantity A defined in (3.2).

$$A(k, \ell) = \binom{2\ell + 2}{\ell + 1} \frac{1}{(k + 2\ell + 1)!}.$$

Using (8.5) we obtain

$$\begin{aligned} J_5^{(0)} + J_6 &= -J_6 = -\int_0^1 \left(\frac{1}{y} + \frac{1}{B - y}\right) \binom{2\ell + 2}{\ell + 1} \frac{(1 - y)^{k+2\ell+1}}{(k + 2\ell + 1)!} dy \\ &= -A(k, \ell) \int_0^1 \left(\frac{1}{y} + \frac{1}{B - y}\right) (1 - y)^{k+2\ell+1} dy, \end{aligned} \tag{8.6}$$

$$J_4 = A(k, \ell) \int_0^1 \left(\frac{1}{y} + \frac{1}{B - y}\right) dy, \tag{8.7}$$

$$\begin{aligned} J_7 &= J_4 + J_5^{(0)} + J_6 = A(k, \ell) \int_0^1 \left(\frac{1}{y} + \frac{1}{B - y}\right) (1 - (1 - y)^{k+2\ell+1}) dy \\ &= A(k, \ell) \int_0^1 \left(\frac{1}{y} \sum_{j=0}^{k+2\ell} (1 - y)^j + \frac{1}{B - y} - \frac{(1 - y)^{k+2\ell+1}}{B - y}\right) dy. \end{aligned} \tag{8.8}$$

With the notation

$$L(n) = \sum_{i=1}^n \frac{1}{i} = \log n + \gamma + O\left(\frac{1}{n}\right), \tag{8.9}$$

we obtain by $B \geq 2$,

$$\begin{aligned} J_7 &= A(k, \ell) \left(\sum_{j=0}^{k+2\ell} \int_0^1 (1 - y)^j dy + \int_0^1 \frac{dy}{B - y} - \int_0^1 \frac{(1 - y)^{k+2\ell+1}}{B - y} dy \right) \\ &= A(k, \ell) \left(L(k + 2\ell + 1) + \log \frac{B}{B - 1} + O(1/k) \right) \\ &= A(k, \ell) \left(\log k + \gamma + \log \frac{B}{B - 1} + O(1/\sqrt{k}) \right). \end{aligned} \tag{8.10}$$

Since in the term $J_5^{(1)}$ the factor y appears, we can directly work with $J_5^{(1)}$, and we get

$$\begin{aligned}
 -\frac{J_5^{(1)}}{2} &= \int_0^1 \sum_{j=1}^{\ell+1} \frac{y^{j-1} + \frac{y^j}{B-y}}{(\ell+1)!(\ell+1-j)!j!} \int_0^{1-y} \frac{(1-y-x)^{2\ell+2-j} x^{k-2}}{(k-2)!} dx dy \tag{8.11} \\
 &= \sum_{j=1}^{\ell+1} \int_0^1 \frac{(y^{j-1} + O(y^j))(1-y)^{k+2\ell+1-j} (2\ell+2-j)!}{(\ell+1)!(\ell+1-j)!j!(k+2\ell+1-j)!} dy \\
 &= A(k, \ell) \sum_{j=1}^{\ell+1} \frac{(\ell+1) \dots (\ell+1-(j-1))}{(2\ell+2) \dots (2\ell+2-(j-1))} \left(\frac{1}{j} + O\left(\frac{1}{k+2\ell+2}\right) \right) \\
 &= A(k, \ell) (C(\ell) + O(1/k)).
 \end{aligned}$$

By $\log(1-x) = -\sum_{j=1}^{\infty} x^j/j$ we have

$$C(\ell) = \sum_{j=1}^{\infty} \frac{1}{2^j} \cdot \frac{1}{j} + o(1) = -\log\left(1 - \frac{1}{2}\right) + o(1) = \log 2 + o(1) \tag{8.12}$$

as $\ell \rightarrow \infty$, so

$$J_5^{(1)} = -A(k, \ell)(\log 4 + o(1)).$$

Finally,

$$\frac{J_3}{(k-2)!} = \int_1^{B/2} \left(\frac{1}{y} + \frac{1}{B-y} \right) \cdot \frac{(2\ell+2)!}{((\ell+1)!)^2} \frac{dy}{(k+2\ell+1)!} = A(k, \ell) \log(B-1). \tag{8.13}$$

Summarizing (8.2)–(8.13), we obtain

$$\begin{aligned}
 \frac{J_1 + J_2 + J_3}{(k-2)!} &= J_7 + J_5^{(1)} + \frac{J_3}{(k-2)!} + o(1) \tag{8.14} \\
 &= A(k, \ell) (\log k + \gamma + \log B - \log 4 + o(1)) \\
 &= A(k, \ell) \left(\log \frac{Be^\gamma k}{4} + o(1) \right).
 \end{aligned}$$

From (8.5), we deduce that

$$\frac{J_0}{(k-1)!} = \binom{2\ell}{\ell} \frac{1}{(k+2\ell)!} = A_0(k, \ell),$$

say.

Returning to (8.1), we find that

$$\begin{aligned}
 J &= \frac{k}{B} A(k, \ell) \log\left(\frac{Be^\gamma k}{4}\right) - \nu A_0(k, \ell) + o(kA(k, \ell)) \\
 &= A_0(k, \ell) \left(\frac{2\left(2 - \frac{1}{\ell+1}\right)k}{B(k+2\ell+1)} \log\left(\frac{Be^\gamma k}{4}\right) - \nu + o(1) \right) \\
 &= A_0(k, \ell) \left(\frac{4}{B} \log\left(\frac{Be^\gamma k}{4}\right) - \nu + o(1) \right).
 \end{aligned}$$

This is positive if

$$\frac{Be^\gamma k}{4} \geq e^{B\nu/4}(1 + o(1)), \tag{8.15}$$

and this proves Theorem 1.

We remark that in the above proof, we are finding “unsifted” E_2 -numbers; that is, the E_2 -numbers found in the proof can have small prime factors. However, it should be clear from the

argument that if one desires, one may take Y in the definition of β to be any function of N such that $\log Y/\log N \rightarrow 0$ as $N \rightarrow \infty$, and the same argument goes through.

In order to show Corollary 1, we have only to note that if $p_1 < p_2 < \dots$ are the consecutive primes, then

$$\mathcal{H} = \{p_{\pi(k)+1} \dots p_{\pi(k)+k}\} \tag{8.16}$$

forms an admissible k -tuple and $p_{\pi(k)+k} \sim k \log k$.

Now we consider Theorem 2. Let \mathcal{S} be as defined in (3.5) with $\nu = 1$. By Theorems 7 and 8, we see that

$$\mathcal{S} \sim \frac{\mathfrak{S}(\mathcal{L})N(\log R)^k}{(k-1)!}J,$$

where

$$J = \frac{k(k-1)}{B}(J_1 + J_2 + J_3) - J_0.$$

We take $k = 3, B = 4, \eta = 1/144$, and $P(x) = 1 + 6x$. Straightforward computations show that

$$\begin{aligned} J_0 &= \frac{38}{15} = 2.5333, \dots, \\ J_1 &= 4824 \log\left(\frac{143}{108}\right) - \frac{13641020155}{10077696} = 0.57625 \dots, \\ J_2 &= -\frac{77824}{15} \log\left(\frac{143}{108}\right) + \frac{14680965985}{10077696} = 0.36202 \dots, \\ J_3 &= \frac{41}{60} \log 3 = 0.75071 \dots, \\ J &= \frac{41}{40} \log 3 - \frac{2732}{5} \log\left(\frac{143}{108}\right) + \frac{852438101}{5598720} = 0.00016493 \dots \end{aligned}$$

For Theorem 3, we take $k = 2, B = 4, \eta = 1/10$, and consider the sum

$$\tilde{\mathcal{S}}(\mathcal{L}) = \sum_{N < n \leq 2N} \left\{ \sum_{j=1}^2 (\beta(L_j(n)) + \varpi(L_j(n)) - 1) \right\} \left(\sum_{d|P_{\mathcal{L}}(n)} \lambda_d \right)^2.$$

From Theorems 7, 8, and 9, we see that

$$\tilde{\mathcal{S}}(\mathcal{L}) \sim N\mathfrak{S}(\mathcal{L})(\log R)^2J,$$

where

$$J = \frac{1}{2}(J_1 + J_2 + J_3 + J_{\varpi}) - J_0.$$

With $P(x) = 1 + x$, we see that

$$\begin{aligned} J_0 &= \frac{11}{12} = 0.91667 \dots, \\ J_1 &= -144 \log(6/5) + \frac{66363}{2500} = 0.29089 \dots, \\ J_2 &= \frac{2048}{15} \log(6/5) - \frac{308429}{12500} = 0.21864 \dots, \end{aligned}$$

$$J_3 = \frac{19}{30} \log 3 = 0.69578 \dots,$$

$$J_\infty = \frac{19}{30} = 0.63333 \dots,$$

$$J = \frac{19}{60} \log 3 - \frac{56}{15} \log(6/5) + \frac{4193}{12500} = 0.00266 \dots,$$

and the theorem follows. The result of (1.20) follows by taking $\mathcal{L} = \{n, n - d\}$.

Now we mention the slight changes which lead to the proofs of Theorems 4–6.

Theorem 4 follows from the proof of Theorem 1 by taking $B = 60$ in view of (1.27)–(1.28). For Theorem 5 we have to restrict p and q to primes of the form $4m + 1$. This means that the density of both p and q is half of that of all primes; therefore, we obtain finally for all S_i and J_i ($i = 2, 4, 5, 6$) a quantity which is $1/4$ of that in the proof of Theorem 1, which has the same effect as to writing 4ν in place of ν .

Finally, the proof of Theorem 6 is just a combination of the proofs of Theorems 4 and 5. The result is that we have to take $B = 60$ as in Theorem 4 and to replace ν by 4ν as in Theorem 5. This leads finally to (1.36).

Corollaries 3–5 follow from Theorems 4–6 in the same way as Corollary 1 follows from Theorem 1 (see (8.16)).

References

1. E. BOMBIERI, *Le grand crible dans la théorie analytique des nombres*, Asterisque 18, 2nd edn (Société Mathématique de Paris, 1987) 103.
2. Y. BUTTKEWITZ, *Aufeinanderfolgende Werte zahlentheoretischer Funktionen*, Master's Thesis, University of Freiburg, 2003.
3. J. -R. CHEN, 'On the representation of a large even integer as the sum of a prime and the product of at most two primes', *Kexue Tongbao* 17 (1966) 385–386 (Chinese).
4. J. -R. CHEN, 'On the representation of a large even integer as the sum of a prime and the product of at most two primes', *Sci. Sinica* 16 (1973) 157–176.
5. T. COCHRANE and R. E. DRESSLER, 'Consecutive triples of sums of two squares', *Arch. Math.* (Basel) 49 (1987) 301–304.
6. L. E. DICKSON, 'A new extension of Dirichlet's theorem on prime numbers', *Messenger of Math.* (2) 33 (1904) 155–161.
7. P. D. T. A. ELLIOTT and H. HALBERSTAM, 'A conjecture in prime number theory', 1970 Symposia Mathematica 4, INDAM, Rome, 1968/69 (Academic Press, London, 1970) 59–72.
8. P. ERDŐS, 'Some problems on number theory', *Analytic and elementary number theory* (Marseille, 1983), Publications Mathématiques d'Orsay 86-1, 53–67, University of Paris, Orsay 1986.
9. P. ERDŐS and L. MIRSKY, 'The distribution of values of the divisor function $d(n)$ ', *Proc. London Math. Soc.* (3) 2 (1952) 257–271.
10. J. FRIEDLANDER and A. GRANVILLE, 'Limitations to the equi-distribution of primes I', *Ann. of Math.* (2) 129 (1989) 363–382.
11. D. A. GOLDSTON, S. W. GRAHAM, J. PINTZ and C. Y. YILDIRIM, 'Small gaps between primes and almost primes', *Trans. Amer. Math. Soc.*, to appear.
12. D. A. GOLDSTON, J. PINTZ and C. Y. YILDIRIM, 'On the difference $p_{n+\nu} - p_n$ ', *Funct. Approx. Comment. Math.* XXXV (2006) 79–89.
13. D. A. GOLDSTON, J. PINTZ and C. Y. YILDIRIM, 'Primes in tuples I', *Ann. of Math.*, to appear.
14. G. GREAVES, *Sieves in number theory* (Springer, Berlin, 2001).
15. H. HALBERSTAM and H. -E. RICHERT, *Sieve methods* (Academic Press, London, New York, 1974).
16. G. H. HARDY and J. E. LITTLEWOOD, 'Some problems of 'Partitio Numerorum': III on the expression of a number as a sum of primes', *Acta Math.* 44 (1923) 1–70.
17. D. R. HEATH-BROWN, 'The divisor function at consecutive integers', *Mathematika* 31 (1984) 141–149.
18. D. R. HEATH-BROWN, 'Almost-prime k -tuples', *Mathematika* 44 (1997) 245–266.
19. C. HOOLEY, 'On the intervals between numbers that are sums of two squares: II', *J. Number Theory* 5 (1973) 215–217.
20. J. KAN, 'On the lower bound sieve', *Mathematika* 37 (1990) 273–286.
21. J. KAN, 'A theorem in multiplicative number theory', *J. Number Theory* 38 (1991) 1–17.
22. A. KARACUBA, *Basic Analytic Number Theory*, Translated from the second (1983) Russian edition by M. B. Nathanson (Springer, Berlin, 1993).

23. Y. MOTOHASHI, 'An induction principle for the generalization of Bombieri's prime number theorem', *Proc. Japan Acad.* 52 (1976) 273–275.
24. A. PERELLI, J. PINTZ and S. SALERNO, 'Bombieri's theorem in short intervals. II', *Invent. Math.* (1) 79 (1985) 1–9.
25. CH. G. PINNER, 'Repeated values of the divisor function', *Quart. J. Math. Oxford Ser. 2* (192) 48 (1997) 499–502.
26. A. DE POLIGNAC, 'Six propositions arithmologiques sur les nombres premiers', *Nouv. Ann. Math.* 7 (1849) 423–429.
27. J. -C. SCHLAGE-PUCHTA, 'The equation $\omega(n) = \omega(n + 1)$ ', *Mathematika* (1–2) 50 (2003) (2005) 99–101.
28. A. SELBERG, *Lectures on Sieves*, Collected Papers II (Springer, Berlin, 1991) 65–247.
29. C. A. SPIRO, 'The frequency with which an integer-valued, prime-independent, multiplicative or additive function of n divides a polynomial function of n ', PhD Dissertation, University of Illinois at Urbana-Champaign, 1981.
30. N. M. TIMOFEEV, 'Distribution of arithmetic functions in short intervals in the mean with respect to progressions (Russian)', *Izv. Akad. Nauk SSSR Ser. Mat.* 51 (1987) 341–362; English translation *Math. USSR Izv.* 30 (1988) 315–335.

D. A. Goldston
 Department of Mathematics
 San Jose State University
 San Jose, CA 95192
 USA

goldston@math.sjsu.edu

J. Pintz
 Rényi Mathematical Institute of the
 Hungarian Academy of Sciences
 P.O.B. 127
 H-1364 Budapest
 Hungary

pintz@renyi.hu

S. W. Graham
 Department of Mathematics
 Central Michigan University
 Mount Pleasant, MI 48859
 USA

graha1sw@cmich.edu

C. Y. Yıldırım
 Department of Mathematics
 Boğaziçi University,
 Istanbul 34342
 Turkey

and

Feza Gürsey Enstitüsü
 Çengelköy
 Istanbul
 P.K. 6, 81220
 Turkey

yalciny@boun.edu.tr