

26. Delcourt, H. R. & Harris, W. F. Carbon budget of the southeastern U.S. biota: analysis of historical change in trend from source to sink. *Science* **210**, 321–323 (1980).
27. Brown, S. L. & Schroeder, P. E. Spatial patterns of aboveground production and mortality of woody biomass for eastern U.S. forests. *Ecol. Appl.* **9**, 968–980 (1999).
28. Schimel, D. *et al.* Contribution of increasing CO<sub>2</sub> and climate to carbon storage by ecosystems in the United States. *Science* **287**, 2004–2006 (2000).
29. Casperson, J. P. *et al.* Contributions of land-use history to carbon accumulation in U.S. forests. *Science* **290**, 1148–1151 (2000).
30. Weisberg, S. *Applied Linear Regression* 114–117 (Wiley, New York, 1985).

**Acknowledgements**

We thank K. Harrison and J. Andrews for collecting pretreatment samples for covariance analysis; H. Hemric, R. Gill, M. Lavine and A. Mace for technical assistance; L. Giles for mass spectrometry analyses; and J. Clark, E. Davidson, L. Dellwo, A. Hirsch and D. Schimel for critical reviews of the manuscript. Operated in cooperation with Brookhaven National Laboratory, the Duke Forest FACE project is supported by the US Department of Energy.

Correspondence and requests for materials should be addressed to W.H.S. (e-mail: schlesin@duke.edu).

**Soil fertility limits carbon sequestration by forest ecosystems in a CO<sub>2</sub>-enriched atmosphere**

Ram Oren\*, David S. Ellsworth†‡, Kurt H. Johnsen§, Nathan Phillips||, Brent E. Ewers\*, Chris Maier§, Karina V.R. Schäfer\*, Heather McCarthy\*, George Hendrey‡, Steven G. McNulty¶ & Gabriel G. Katul\*

\* Nicholas School of the Environment and Earth Sciences, Duke University, Durham, North Carolina 27708, USA

† School of Natural Resources and Environment, University of Michigan, Ann Arbor, Michigan 48109, USA

‡ Department of Environmental Sciences, Brookhaven National Laboratory, Upton, New York 11973-5000, USA

§ Southern Research Station, US Forest Service, Research Triangle Park, North Carolina 27709, USA

|| Department of Geography, Boston University, Boston, Massachusetts 02215, USA


¶ Southern Global Climate Change Program, US Forest Service, Raleigh, North Carolina 27606, USA

Northern mid-latitude forests are a large terrestrial carbon sink<sup>1–4</sup>. Ignoring nutrient limitations, large increases in carbon sequestration from carbon dioxide (CO<sub>2</sub>) fertilization are expected in these forests<sup>5</sup>. Yet, forests are usually relegated to sites of moderate to poor fertility, where tree growth is often limited by nutrient supply, in particular nitrogen<sup>6,7</sup>. Here we present evidence that estimates of increases in carbon sequestration of forests, which is expected to partially compensate for increasing CO<sub>2</sub> in the atmosphere, are unduly optimistic<sup>8</sup>. In two forest experiments on maturing pines exposed to elevated atmospheric CO<sub>2</sub>, the CO<sub>2</sub>-induced biomass carbon increment without added nutrients was undetectable at a nutritionally poor site, and the stimulation at a nutritionally moderate site was transient, stabilizing at a marginal gain after three years. However, a large synergistic gain from higher CO<sub>2</sub> and nutrients was detected with nutrients added. This gain was even larger at the poor site (threefold higher than the expected additive effect) than at the moderate site (twofold higher). Thus, fertility can restrain the response of wood carbon sequestration to increased atmospheric CO<sub>2</sub>. Assessment of future carbon sequestration should consider the limitations imposed by soil fertility, as well as interactions with nitrogen deposition.

By burning fossil fuel and forests, and converting land to intensive agriculture use, humans have elevated the atmospheric concentration of CO<sub>2</sub> (ref. 3) and the deposition of atmospheric nitrogen (N)<sup>9,10</sup>. Growth of many tree species is enhanced with provisions of both CO<sub>2</sub> and N in the suboptimal range<sup>11</sup>, making it difficult to assess the effect of increased availability of either one when the supply of both increases concurrently<sup>10</sup>. Experiments indicate decreased nutrient availability<sup>12</sup>, owing to increased carbon (C)/N ratios in elevated CO<sub>2</sub>-grown foliage and litter<sup>13</sup>, but increased nutrient uptake with elevated CO<sub>2</sub>-induced growth enhancement of fine roots<sup>14</sup>. Because most forests occur on low-nutrient soils, the uncertain effects of elevated CO<sub>2</sub> on nutrient supply hinders our ability to estimate forest C sequestration for future global C budgets.

If tree nutrient uptake does not increase in proportion to the growth response to elevated CO<sub>2</sub>, then maturing pine trees should show less elevated CO<sub>2</sub>-induced growth enhancements on low-fertility sites than on moderate sites, and thus should respond more to elevated CO<sub>2</sub> under improved nutrition. To evaluate these predictions, we used two field experiments with related loblolly pine (*Pinus taeda* L.) genotypes: the longest running forest-based free-air CO<sub>2</sub> enrichment (FACE) experiment conducted on a moderately fertile site; and a whole-tree chamber CO<sub>2</sub> enrichment experiment on an infertile site. The large difference in fertility of the two sites is indicated by a much greater increase in growth at the infertile site (130%) than at the moderate site (15%) when added nutrients augmented the sites to the same optimal fertility<sup>7</sup>. Additional site and soil characteristics that reflect fertility are provided in Methods.

The FACE prototype (FACE<sub>p</sub>) was built in 1993 in a 10-year-old, 8.5-m tall loblolly pine plantation in the lower Piedmont plateau, growing on moderately low fertility, acidic clay-loam, at the Duke Forest of Duke University, North Carolina (35° 58' N, 79° 08' W; elevation 130 m). Currently the pines are about 15-m tall, and comprise 98% of the basal area<sup>15</sup>. FACE<sub>p</sub> has enriched (550 p.p.m.v. CO<sub>2</sub>) a 30-m diameter circular patch in the forest since 1994, during daylight hours of the growing season<sup>16</sup>. Before CO<sub>2</sub> enrichment in FACE<sub>p</sub> commenced, growth (that is, the amount of C sequestered in woody biomass increment) was similar in FACE<sub>p</sub> and an adjacent,




**Figure 1** A comparison of annual carbon increment under elevated atmospheric CO<sub>2</sub> concentration (initiated in 1994) and ambient concentration without nutrient addition. **a**, Plot-level comparison between the free-air CO<sub>2</sub> enrichment prototype (FACE<sub>p</sub>) and a nearby untreated, ambient CO<sub>2</sub> plot (in the past 2 yr, the number of untreated plots was increased to five). **b**, Individual tree comparison between trees in FACE<sub>p</sub> and trees selected at random from the entire stand. Data for 1993 are shown as means for the first 10 yr of the stand's life.

untreated plot (Fig. 1a), as were foliar N concentration and physiological capacity<sup>16</sup>.

Averaged over the first 3 yr, the elevated CO<sub>2</sub> plot had a 34% increase in growth relative to the ambient CO<sub>2</sub> plot, reaching a maximum of 40%, but only a 6% increase over the next 4 yr. After four growing seasons of enrichment, we cored all pine trees in FACE<sub>p</sub> (*n* = 78) and trees selected at random from the entire stand (*n* = 56; none in the untreated plot), and calculated annual individual tree growth for every year since 1993. The temporal pattern in individual tree growth was fundamentally the same as that of the stand with lower growth of FACE<sub>p</sub> trees before the commencement of CO<sub>2</sub> enrichment, a large initial relative response to elevated CO<sub>2</sub>, and a subsequent decrease in the response (Fig. 1b).

The CO<sub>2</sub>-induced growth enhancement over the first 2 yr in the adjacent replicated FACE study (*n* = 3)<sup>15</sup>, which began operation in 1996, averaged about 24%, and ranged from 7 to 57% (1 s.d. = 15%). The initial growth enhancement in the FACE<sub>p</sub> plot was not different from that at replicated FACE plots, after matching annual growth for the same year since enrichment commenced in both studies (2 yr, *t*-test, *P* > 0.1; see Methods for statistical approach). Thus, the early growth responses to elevated CO<sub>2</sub> were consistent in both experiments, but these responses are transient (Fig. 1).

The decrease in response to elevated CO<sub>2</sub> (Fig. 1) might be caused by nutrient limitation that can develop fairly rapidly in this moderate-fertility site with a limited rooting depth (< 0.3 m)<sup>17</sup>. The January foliar N concentration was similarly low in FACE<sub>p</sub> and in the surrounding stand growing under ambient CO<sub>2</sub> (0.97 ± 0.04% by weight; mean ± s.e., *n* = 16). Plant-soil models predicted responses to an experimental stepwise increase in elevated CO<sub>2</sub> in nutrient-limited forests<sup>18</sup>, similar to that shown in Fig. 1. To test whether nutrient limitations reduced the tree response to elevated CO<sub>2</sub>, we partitioned FACE<sub>p</sub> and its unenriched counterpart in 1998, and added a balanced fertilizer to half the area (> 265 m<sup>2</sup>)




**Figure 2** Effect of atmospheric CO<sub>2</sub> concentration and nutrient supply on annual carbon increment in woody tissue of genotypically similar loblolly pine. **a**, A moderate-fertility Duke Forest site (averaged for 1999 and 2000); **b**, an infertile SETRES site (1997 and 1998). *P* values represent test results between ambient and elevated CO<sub>2</sub> in each fertility level. In both sites, elevating only CO<sub>2</sub> had no significant effect (in **a**, data without nutrient addition are the mean of the past 2 yr shown in Fig. 1a); fertilizing in ambient CO<sub>2</sub> had a significant effect; and fertilizing under elevated CO<sub>2</sub> had significantly higher effects than the sum of the single CO<sub>2</sub> and N effects.

at the end of the growing season, aiming to attain optimal nutrition (1.4% N)<sup>7,19</sup>. We also established four plot pairs, randomly positioned in the stand, with one member of each fertilized similarly, thus producing five plot pairs under ambient CO<sub>2</sub>. After 1 yr, fertilization increased foliar N concentration under ambient CO<sub>2</sub> to 1.18 ± 0.08% (mean ± s.e.), but foliar N remained low under elevated CO<sub>2</sub> (1.02%), similar to unfertilized plots under both ambient and elevated CO<sub>2</sub> (1.05 ± 0.07%).

Averaged over 1999 and 2000, about 378 ± 16.0 g C m<sup>-2</sup> yr<sup>-1</sup> (mean ± s.e.) was incorporated into woody tissue under ambient CO<sub>2</sub> and without fertilization (Fig. 2a). Annual woody growth under elevated CO<sub>2</sub> without nutrient addition sequestered only 7% more C (*P* > 0.05), and improved nutrient supply in ambient CO<sub>2</sub> atmosphere increased growth by 15% (*P* = 0.024). However, the combination of improved nutrition and elevated CO<sub>2</sub> increased growth by 47% (~175 g C m<sup>-2</sup> yr<sup>-1</sup>; *P* = 0.007)—more than twice the sum of the separate responses to each factor (total ~80 g C m<sup>-2</sup> yr<sup>-1</sup>). This clearly indicates a synergistic effect of CO<sub>2</sub> and nutrient supply that raised the relative CO<sub>2</sub>-induced growth response to the maximum previously seen in FACE<sub>p</sub> (~40%).

We assessed whether this synergism increases with decreasing site fertility by evaluating the response of trees (*n* = 3, whole-tree open-top chambers) in another North Carolina loblolly pine plantation of a similar age and genetic stock growing on an infertile sandy soil (USDA Forest Service Southeast Tree Research and Education Site, SETRES; 34° 48' N, 79° 12' W). At this infertile site, foliar N was 1.02 ± 0.04% (mean ± s.e.) in both CO<sub>2</sub> environments, increased with fertilization to 1.29 ± 0.05%, and decreased to 1.15 ± 0.04% when fertilized trees were subjected to elevated CO<sub>2</sub> for two growing seasons (550 p.p.m.v.). Without added nutrients, annual growth did not respond to elevated CO<sub>2</sub> in the 2-yr period (*P* > 0.05; Fig. 2b). Under optimal nutrition<sup>19</sup> and ambient CO<sub>2</sub>, the growth of these trees increased by 21% (*P* < 0.05). However, the synergistic enhancement from improved nutrition and higher atmospheric CO<sub>2</sub> was 74% (*P* = 0.008)—more than three times the sum of the separate responses. This suggests that CO<sub>2</sub> responses of growth in pine forests will be highly variable and depend greatly on site fertility, perhaps to the point of not responding at all on the nutritionally poorest sites.

In addition to nutrient limitations, water deficits could also limit forest response to elevated CO<sub>2</sub>. The early part of the 1999 growing season was unusually dry along the eastern seaboard of the USA,



**Figure 3** Cumulative carbon increment in woody tissue under ambient and elevated atmospheric CO<sub>2</sub> concentration with and without the addition of nutrients. The carbon increment under elevated CO<sub>2</sub> without nutrient addition was never significantly greater than under ambient CO<sub>2</sub> (*P* > 0.05). Arrows depict the earliest time in each growing season in which the carbon increment under elevated CO<sub>2</sub> with nutrient addition was significantly higher than the indicated treatment (*P* < 0.05). The period of CO<sub>2</sub> enrichment in each year is indicated as a horizontal bar at the bottom. In 1999, drought-breaking rains (indicated) raised soil moisture in the root zone from 0.14 to 0.32 m<sup>3</sup> m<sup>-3</sup>. Error bars represent s.e.

with the May–July rainfall 84% below the 30-yr site average. In contrast, rain was ample and evenly distributed during the 2000 growing season. Weekly measurements of stem diameter at the Duke Forest site permitted us to quantify drought-induced interaction effects with the CO<sub>2</sub> and nutrition treatments on the seasonal dynamics of C investment in woody tissue. A fraction of the stem increment that occurred after each rainfall event (such as that in 1999; Fig. 3) was due to actual wood growth and a fraction was due to stem swelling with hydration; we estimated the latter, reversible ‘increment’ as  $\leq 17.4 \text{ g C m}^{-2}$  after a given event. Without drought in 2000, growth enhancement in fertilized elevated CO<sub>2</sub> conditions became significant earlier (Fig. 3), and sequestered  $45 \text{ g C m}^{-2} \text{ yr}^{-1}$  (27%) more than with early season drought in 1999. Hence, the synergistic effect of CO<sub>2</sub> and nutrients is especially strong when seasonal water limitations are alleviated.

Treatments imposed in a stepwise manner may invoke different responses than gradual changes<sup>8</sup>. Where site fertility is very low, responses to stepwise CO<sub>2</sub> enrichment experiments may be muted (Fig. 2b). Where fertility is moderate, responses may be transient and decrease when soil nutrients are depleted by the initial increase in growth rate<sup>19</sup> (Fig. 1). Thus, the short-term dynamics of the elevated CO<sub>2</sub> response<sup>15</sup> may not portray the longer-term, multi-year responses shown in this study, particularly where dynamic changes in nutrient cycles are involved<sup>12</sup>. There are many reasons to expect the long- and short-term responses to be different<sup>18,20</sup>, and this study highlights the need for caution in broadly extrapolating from short-term results.

Along a gradient of decreasing native-site fertility, nutrient amendments will become increasingly necessary if forest ecosystems (planted or natural) are intended to both take advantage of elevated atmospheric CO<sub>2</sub> and aid in reducing the rate at which atmospheric CO<sub>2</sub> concentrations increase. Under current atmospheric CO<sub>2</sub>, nitrogen deposition has been estimated to contribute little to enhanced C sequestration in temperate forests<sup>10,21,22</sup>, and N deposition may ultimately cause a growth reduction by causing nutritional disharmony<sup>23</sup>. Nevertheless, a model comparison has concluded that interaction between CO<sub>2</sub> and N deposition may be involved in the response of terrestrial ecosystems to future levels of atmospheric CO<sub>2</sub> (ref. 24).

Our amendment of  $\sim 11 \text{ g N m}^{-2} \text{ yr}^{-1}$  is more than 10 times the current deposition rates of  $\sim 0.65 \text{ g N m}^{-2} \text{ yr}^{-1}$  at both sites. At the moderate-fertility site, C sequestration was enhanced by  $\sim 14.1$  and  $\sim 17.3 \text{ g C g}^{-1} \text{ N}$  addition under elevated atmospheric CO<sub>2</sub> in the dry and wet year, respectively (that is, an average of  $15.7 \text{ g C g}^{-1} \text{ N}$ ). At the nutritionally poor site, the enhancement under elevated CO<sub>2</sub> was  $19.1 \text{ g C g}^{-1} \text{ N}$  addition. This provides a first estimate of the effect of N supply on CO<sub>2</sub>-induced C sequestration in woody biomass increment at the scale of forest stands. We note, however, that industrial N fertilizer production and use already has an impact on the global N cycle<sup>25</sup>, and any plan to enhance C sequestration that relies on a significant increase in N fertilization must be carefully evaluated for both local and global effects. □

## Methods

### Setting

The FACE and SETRES sites are mid-rotation plantations without density-related mortality. The summers are warm and humid, and winters are moderate. Annual precipitation (1,100–1,200 mm) is evenly distributed throughout the year, but occasional deficits in growing-season water occur. The sites are considered moderate (FACE) and low (SETRES) fertility, as reflected in the heights of dominant *P. taeda* of 15.7 and 10.4 m at age 25 yr, respectively. Soil incubations for net N mineralization at these sites yielded roughly 2.7 (upper 7.5 cm) and 1.3 (15 cm)  $\text{g N m}^{-2} \text{ yr}^{-1}$ , respectively, and cation exchange capacity was about 12 (upper 15 cm) and 2 (20 cm)  $\text{cmol}(+) \text{ g}^{-1}$ . Seasonal CO<sub>2</sub> enrichment in FACE<sub>p</sub>, typically from late April to late October, was achieved in 1994–2000 by the free-air CO<sub>2</sub> enrichment technique<sup>26</sup>.

Annual growth of trees at the FACE<sub>p</sub> and its paired untreated plot was measured from 1993 onwards, at the end of each growing season. In 1998, an impermeable barrier to 1-m depth, about three times the depth of fine roots in this forest<sup>17</sup>, partitioned both plots into two sections containing roughly the same mass of above-ground biomass per unit ground. Concurrently, four pairs of 10 m × 10 m plots were established nearby with a minimum

distance of 20 m between plots. One randomly selected section (FACE<sub>p</sub> and its counterpart) and one randomly chosen plot of each newly established plot pair were fertilized to meet optimal values<sup>7,19</sup>. An application rate of  $11.2 \text{ g N m}^{-2} \text{ yr}^{-1}$  in the form of urea began in July 1998.

The SETRES stand was planted in 1985 in the Sandhills of North Carolina<sup>19</sup>, a site with sandy, siliceous, well-drained soil. Annual nutrient additions to meet optimal values commenced in March 1992, with annual applications averaging  $11.2 \text{ g N m}^{-2} \text{ yr}^{-1}$  over the 7 yr (ref. 18). Six fertilized trees and six unfertilized trees of a similar diameter range and mean height of  $8.6 \pm 1 \text{ m}$  were enclosed in open-top chambers<sup>27</sup> in 1996. CO<sub>2</sub> enrichment to  $\sim 550 \text{ p.p.m.v.}$  in three chambers per treatment commenced in August 1996, and lasted for two complete growing seasons.

### Carbon sequestration

Woody biomass (that is, in branches, bole and roots down to 5 mm in diameter) was estimated at both sites from diameter and allometric equations for ambient CO<sub>2</sub> conditions<sup>19,28</sup>. At Duke Forest, estimates included trees with diameters > 50 mm at 1.4 m above ground, representing 99% of the biomass, less than 5% of which is in hardwood species. At SETRES, equations were available for trees with and without fertilization<sup>19</sup>. From SETRES, the density of wood decreased from 0.51 to 0.47  $\text{g cm}^{-3}$  with fertilization. From cores taken at FACE<sub>p</sub>, the density of wood decreased from 0.52 to 0.48  $\text{g cm}^{-3}$  with elevated CO<sub>2</sub>. Thus, when growth estimated from diameter was increased by either treatment, the values were adjusted to reflect these reductions. Woody biomass above ground was converted to C content by multiplying by 0.48, and below ground by 0.44, as determined from combustion (CHN Analyzer; Perkin Elmer). In SETRES, biomass was converted to a ground-area value on the basis of 1,260 trees  $\text{ha}^{-1}$ . At Duke Forest, diameter was measured weekly during the growing season (Fig. 3) with stainless steel dendrometer bands<sup>15</sup> installed 1.4 m above ground on all trees. Edge conditions created when a 3-m wide strip was cleared around FACE<sub>p</sub> did not affect growth: there was no pattern in relative growth rate with distance from the edge, nor was there a difference between the population of trees positioned less than one-third of the radius from the periphery and that composed of the rest of the trees ( $P > 0.5$ ).

### Statistical considerations

We analysed the SETRES data by paired *t*-test after pairing trees on the basis of initial biomass. Data from FACE<sub>p</sub> and its counterpart were not compared statistically (Fig. 1a); however, data from FACE<sub>p</sub> and its unenriched counterpart were compared with data from the nearby replicated FACE ( $n = 3$ ), and the 1999 and 2000 data from FACE<sub>p</sub> were compared with data from all nearby unenriched plots ( $n = 5$ ) by a *t*-test between a single observation and a population mean<sup>29</sup>. This approach permits testing whether the response of FACE<sub>p</sub> is different from that of the replicated unenriched plots, but without replications it is not possible to estimate the mean response to elevated supplies of both CO<sub>2</sub> and N. Furthermore, we note that the statistical scope of inference of most controlled experiments, including the replicated FACE and SETRES, is limited<sup>30</sup>.

Received 6 February; accepted 22 March 2001.

1. Tans, P. P. & White, J. W. C. In balance, with a little help from the plants. *Science* **281**, 183–184 (1998).
2. Ciais, P., Tans, P. P., Trolier, M., White, J. W. C. & Francey, R. J. A large Northern Hemisphere terrestrial CO<sub>2</sub> sink indicated by <sup>13</sup>C/<sup>12</sup>C of atmospheric CO<sub>2</sub>. *Science* **269**, 1098–1102 (1995).
3. Schimel, D. S. Terrestrial ecosystems and the carbon cycle. *Global Change Biol.* **1**, 77–91 (1995).
4. Houghton, R. A., Davidson, E. A. & Woodwell, G. M. Missing sinks, feedbacks, and understanding the role of terrestrial ecosystems in the global carbon balance. *Global Biogeochem. Cycles* **12**, 25–34 (1998).
5. Idso, S. B. & Kimball, B. A. Tree growth in carbon-dioxide enriched air and its implications for global carbon cycling and maximum levels of atmospheric CO<sub>2</sub>. *Global Biogeochem. Cycles* **7**, 537–555 (1993).
6. Vitousek, P. M. & Howarth, R. W. Nitrogen limitations on land and in the sea: How can it occur? *Biogeochemistry* **13**, 87–115 (1991).
7. Linder, S. Foliar analysis for detecting and correcting nutrient imbalances in Norway spruce. *Ecol. Bull.* **44**, 178–190 (1995).
8. Körner, C. Towards a better experimental basis for upscaling plant responses to elevated CO<sub>2</sub> and climate warming. *Plant Cell Env.* **19**, 1101–1110 (1995).
9. Aber, J. et al. Nitrogen saturation in temperate forest ecosystems. *Bioscience* **48**, 921–934 (1998).
10. Nadelhoffer, K. J. et al. Nitrogen deposition makes a minor contribution to carbon sequestration in temperate forests. *Nature* **398**, 145–148 (1999).
11. Wullschlegel, S. D., Post, W. M. & King, A. W. in *Biotic Feedbacks in the Global Climatic System* (eds Woodwell, G. M. & Mackenzie F. T.) 85–107 (Oxford Univ. Press, New York, 1995).
12. Hu, S., Chapin III, F. S., Firestone, M. K., Field, C. B. & Chiarello, N. R. Nitrogen limitation of microbial decomposition in a grassland under elevated CO<sub>2</sub>. *Nature* **409**, 188–191 (2001).
13. Norby, R. J. & Cotrufo, M. F. A question of litter quality. *Nature* **396**, 17–18 (1998).
14. Berntson, G. M. & Bazzaz, F. A. Belowground positive and negative feedbacks on CO<sub>2</sub> growth enhancement. *Plant Soil* **187**, 119–131 (1996).
15. DeLucia, E. H. et al. Net primary production of a forest ecosystem with experimental CO<sub>2</sub> enrichment. *Science* **284**, 1177–1179 (1999).
16. Ellsworth, D., Oren, R., Ce, H., Phillips, N. & Hendrey, G. Leaf and canopy responses to elevated CO<sub>2</sub> in a pine forest using a free-air CO<sub>2</sub> enrichment technique. *Oecologia* **104**, 139–146 (1995).
17. Oren, R., Ewers, B. E., Todd, P., Phillips, N. & Katul, G. Water balance delineates the layer in which soil moisture affects canopy conductance. *Ecol. Appl.* **8**, 990–1002 (1998).
18. Comins, H. N. & McMurtrie, R. E. Long-term responses of nutrient-limited forests to CO<sub>2</sub> enrichment: Equilibrium behavior of plant-soil models. *Ecol. Appl.* **3**, 666–681 (1993).
19. Albaugh, T. J., Allen, H. L., Dougherty, P. M., Kress, L. W. & King, J. S. Leaf area and above- and belowground growth responses of loblolly pine to nutrient and water additions. *For. Sci.* **44**, 1–12 (1998).
20. Bazzaz, F. A., Miao, S. L. & Wayne, P. M. CO<sub>2</sub>-induced enhancements of co-occurring tree species decline at different rates. *Oecologia* **96**, 478–482 (1994).
21. Houghton, R. A., Hackler, J. L. & Lawrence, K. T. The U.S. carbon budget: Contributions from land-

- use change. *Science* **285**, 574–578 (1999).
22. Field, C. B. & Fung, I. Y. The not-so-big U.S. carbon sink. *Science* **285**, 544–545 (1999).
23. Oren, R. in *Plant Response to Air Pollution* (eds Iqbal, M. & Yunus, M.) 75–98 (John Wiley & Sons, Chichester, UK, 1996).
24. Kicklighter, D. W. *et al.* A first-order analysis of the potential role of CO<sub>2</sub> fertilization to affect global carbon budget: a comparison of four terrestrial biosphere models. *Tellus B* **51**, 343–366 (1999).
25. Vitousek, P. M. Beyond global warming: Ecology and global change. *Ecology* **75**, 1861–1876 (1994).
26. Hendrey, G. R., Ellsworth, D. S., Lewin, K. F. & Nagy, J. A free-air enrichment system for exposing tall forest vegetation to elevated atmospheric CO<sub>2</sub>. *Global Change Biol.* **5**, 293–309 (1999).
27. Tissue, D. T., Thomas, R. B. & Strain, B. R. Growth and photosynthesis of loblolly pine (*Pinus taeda*) after exposure to elevated CO<sub>2</sub> for 19 months in the field. *Tree Physiol.* **16**, 49–60 (1996).
28. Naidu, S. L., DeLucia, E. H. & Thomas, R. B. Contrasting patterns of biomass allocation in dominant and suppressed loblolly pine. *Can. J. For. Res.* **28**, 1116–1124 (1998).
29. Sokal, R. R. & Rohlf, F. J. *Biometry* (W. H. Freeman and Co., San Francisco, 1969).
30. Bolin, B., Canadel, J., Moore, B., Noble, I. & Steffen, W. Effect on the biosphere of elevated atmospheric CO<sub>2</sub>. *Science* **285**, 1851–1852 (1999).

## Acknowledgements

This study was supported by the Department of Energy through both the Office of Biological and Environmental Research and the National Institute for Global Environmental Change, Southeast Regional Center at the University of Alabama, and by the US Forest Service through both the Southern Global Climate Change Program and the Southern Research Station. This work contributes to the Global Change and Terrestrial Ecosystems (GCTE) core project of the International Geosphere–Biosphere Programme (IGBP). We thank D. E. Pataki for helping with data acquisition; T. Albaugh for descriptive site data; and C. Körner, C. Field and R. Norby for helpful comments on an early version.

Correspondence and requests for materials should be addressed to R.O. (e-mail: ramoren@duke.edu).

## An Early Cambrian tunicate from China

D.-G. Shu, L. Chen, J. Han & X.-L. Zhang

Early Life Institute & Department of Geology, Northwest University, Xi'an, 710069, China

Like the Burgess Shales of Canada, the Chengjiang Lagerstätte from the Lower Cambrian of China is renowned for the detailed preservation as fossils of delicate, soft-bodied creatures<sup>1–9</sup>, providing an insight into the Cambrian explosion. The fossils of possible hemichordate chordates<sup>5–7</sup> and vertebrates<sup>9</sup> have attracted particular attention. Tunicates, or urochordates, comprise the most basal chordate clade<sup>10</sup>, and details of their evolution could be important in understanding the sequence of character acquisition that led to the emergence of chordates and vertebrates<sup>11–18</sup>. However, definitive fossils of tunicates from the Cambrian are scarce or debatable<sup>4,9,19–24</sup>. Here we report a probable tunicate *Cheungkongella ancestralis* from the Chengjiang fauna. It resembles the extant ascidian tunicate genus *Styela* whose morphology could be useful in understanding the origin of the vertebrates.

Phylum Chordata  
Subphylum Urochordata  
Class Ascidiacea  
*Cheungkongella ancestralis* gen. et sp. nov.

**Type species.** *Cheungkongella ancestralis*.

**Etymology.** Genetic name is a metaphor of China and is also in honour of the Cheungkong Scholars Programme that supports this work; the specific name is a reference to its possible primitive position.

**Holotype.** Early Life Institute (ELI), Northwest University, Xi'an: ELI-0000195.


**Stratigraphy and locality.** Qiongzhusi Formation, Yu'an-shan Member (*Eoredlichia* Zone); Lower Cambrian. The specimen was collected by L.C. and J.H. from the same locality and horizon as the

animal *Xidazoon*<sup>8</sup> and agnathan vertebrate *Myllokunmingia*<sup>9</sup>.

**Diagnosis.** The body is club-shaped, reminiscent of extant ascidian *Styela*, with two-fold division: an upper main body and a lower thick supporting stem attached to hard substratum (Fig. 1). The body is wholly enclosed within a structure interpreted as a secreted tunic. The stem tapers downward, and the main body is bucket-shaped in outline, bearing a large oral siphon with short tentacles on its top and a small cloacal one on the lower dorsal side. A pharynx occupies over two-thirds of the body volume.

**Description.** *Cheungkongella ancestralis*, new genus and species, is known from a single specimen, with a total length of about 25 mm. The whole body consists of two regions: a stout stem, which in life supported a sub-spherical main body. The stem (about 15 mm long) tapers distally, and is attached to the exterior surface of the left, free cheek of a trilobite *Eoredlichia intermedia*, an index fossil for the Lower Cambrian. The stem bears some transverse creases, consistent with an enclosing tunic, and prominent longitudinal 'ribs'. The distal section has a conspicuously coarse texture, and has several patches of agglutinated sediment including quartz grains.

The main body (roughly 10 mm long) was probably sub-spherical in life. Wrinkling of the compressed body on the ventral side (opposite to cloacal siphon) is consistent with folding of a tough



**Figure 1** The Lower Cambrian urochordate *Cheungkongella ancestralis* gen. et sp. nov. from Haikou, Kunming, Yunnan. Specimen ELI-0000195, viewed from the left. Scale bar, 1 mm. Bt, buccal tentacles; Cs, cloacal siphon; Dt?, degenerating tail; Os, oral siphon; En?, presumed endostyle; Es?, possible esophagus; Ph, pharynx; S?, presumed stomach; St, stem; T, tunic; Tf, tentacle-like fringe; Tfc, trilobite free cheek.