

Soil pipe distribution and hydrological functioning within the humid tropics: a synthesis

Nick A. Chappell*

Lancaster Environment Centre, Lancaster University, Lancaster LA1 4YQ, UK

Abstract:

Some of the most responsive runoff systems in the world are found within the humid tropics. These runoff responses are likely to be affected by the presence of *natural pipes* within the soil. This study provides a synthesis of the hydrological aspects of these phenomena within the humid tropics. Of the studies reporting the presence of soil piping within the humid tropics, most are associated with Ultisol soils, and, locally, most pipe outlets are observed on the lower sections of hillslopes. While the drainage role of pipes has been observed (providing faster and slower components of stream hydrographs), the mechanism of their recharge remains less clear. In part, this is because their spatial extent is poorly mapped within the hillslopes of the humid tropics. Further studies quantifying the length and recharge of soil pipe networks within the humid tropics are needed. Copyright © 2010 John Wiley & Sons, Ltd.

KEY WORDS Cicada burrow; humid tropics; preferential flow; soil pipe; Ultisol

Received 1 September 2009; Accepted 9 November 2009

INTRODUCTION

The humid tropics cover one-fifth of the land surface of the globe (Figure 1 in Bonell and Balek, 1993; Chang and Lau, 1993). They extend from parts of Mexico to Brazil within the Americas, Senegal to Madagascar within the African domain and India to Northern Australia within the Indo-Malayan domain. The extent is similar to that covered by climate type 'A' of the Köppen-Geiger system (Peel *et al.*, 2007). This global region produces the greatest amount of runoff (Fekete *et al.*, 2002), and has headwater basins noted for their flashy rainfall–runoff responses (e.g. Bonell and Gilmour, 1978; Chappell *et al.*, 1999b; Dykes and Thornes, 2000; Godsey *et al.*, 2004). Furthermore, the region's catchments are subject to the greatest intensity of land cover disturbance through forest cutting (FAO, 2005). Despite the extent of this hydroclimatic region, its hydrological significance and the current pace of disturbance, there are relatively few studies addressing rainfall–runoff processes within humid tropical hillslopes and basins (Bonell, 2004; Chappell *et al.*, 2007).

Intensive research within experimental basins in temperate climates indicate that *natural soil piping* can be a dominant pathway for routing rainfall to rivers (Jones *et al.*, 1997; Uchida *et al.*, 1999; Jones and Connelly, 2002). These hydrological features can be defined as water conduits that are typically several centimetres or tens of centimetres in diameter and their networks can extend several metres or tens of metres

within hillslopes. Their size and length make them a special type of macropore flow (cf. Jones, 1981, 1990; Scherrer *et al.*, 2007). Pipes can be enlarged by a combination of (a) sheer stresses exerted by flowing water on soil particles in pipe walls (a process formally called 'tunnel erosion'), and (b) by seepage forces within the soil that comprises pipe walls, which lead to Coulomb failure or liquefaction (so called 'true piping'; see Dunne, 1990; Bryan and Jones, 1997). In areas where piping is present, accurate simulation of subsurface flow within hillslopes may need the hydrological functioning of pipe systems to be parameterized explicitly (Jones and Connelly, 2002), or may necessitate modelling approaches alternative to those based on the Darcy–Richards Equation (Kirkby, 1988 p. 336). With much less experimental work on soil piping undertaken within the humid tropics (Bonell and Balek, 1993; Negishi *et al.*, 2007), their extent and hydrological role remains less clear. Studies such as Elsenbeer and Cassel (1990); Chappell *et al.* (1998); Elsenbeer and Vertessy (2000); Sayer *et al.* (2004); Chappell and Sherlock (2005) and Negishi *et al.* (2007) do, however, suggest that piping within humid tropical soils may have a significant impact on streamflow generation and the way that catchment models are parameterized and results interpreted.

Consequently, this study seeks to review and synthesize previous studies on soil piping within the humid tropics to consider: (i) the characteristics of their spatial distribution, (ii) their hydrological functioning, and (iii) key research needs. The review of published work is supplemented by further detailed observations from Danum (East Malaysia), the humid tropical location with possibly the greatest number of published pipe studies.

* Correspondence to: Nick A. Chappell, Lancaster Environment Centre, Lancaster University, Lancaster LA1 4YQ, UK.
E-mail: n.chappell@lancaster.ac.uk




Figure 1. A map of the tropics (23.4°N and 23.4°S shown with broken lines) showing the distribution of published studies reporting soil pipe incidence within humid tropical regions. Names of the numbered locations are given in Table I

This study focuses on piping within soil in the strictest sense of the solum (i.e. A and B soil horizons). Within the humid tropics, piping has also been observed deep within underlying *regolith* in Sierra Leone (Dixey, 1920 p. 219; Thomas, 1968), Guinea (de Chételat, 1938), Guyana (McBeath and Barron, 1954 p97), Uganda (Patz, 1965), Columbia (Feininger, 1969), and Brazil (Bremer, 1973). However, detailed hydrological studies have not been undertaken on these, what might be called, *regolith pipes*. In addition, soil piping may be seen as having some of the preferential flow characteristics of rock fracture systems (Zhao, 1998) and karstic phenomena within rock (Gunn, 1986). These deeper strata (*regolith* and rock) are not examined within this review.

SPATIAL DISTRIBUTION OF PIPING WITHIN HUMID TROPICAL SOILS

The dearth of experimental studies on soil piping within humid tropical soils makes understanding their spatial distribution important for extrapolation of findings (however tentatively) and in designing and locating new experimental studies. Three questions arise when considering an initial assessment of their spatial distribution:

1. Do certain soil types found within the humid tropics have a greater reported incidence of soil piping?
2. Are the pipes of the humid tropics preferentially located with certain topographic locations?
3. Where within vertical profiles of humid tropical soils are pipes located?

Reported incidence of soil piping in humid tropical soils

If piping is primarily associated with specific soil types, then there may be underlying physicochemical reasons for this. As a result, the soil type associated with each reported incidence of piping within humid tropics is presented in Table I and Figure 1. The relatively limited number of studies presented within Table I shows that there are fewer studies than reported for soils within other global regions, particularly those within temperate or semi-arid climates (see e.g. the reviews of Jones, 1981, 1994; Jones *et al.*, 1997). The majority of studies in the humid tropics are reported for soils of the Ultisol order (Table I; Soil Survey Staff,

1999), though a few studies also report piping within Alfisol and Andisol soils (Banerjee, 1972; Walsh, 1980; Bruijnzeel, 1983; Walsh and Howells, 1988). Ultisol soils are equivalent to the former Acrisol group (FAO-UNESCO, 1974) and the Acrisol-Alisol groups of the current system (e.g. FAO-UNESCO, 1988). Acrisol is the dominant soil group of tropical Asia, and is particularly important within Malaysia, southern China, Cambodia, Vietnam, parts of Thailand and Indonesia (e.g. Sumatra and Sulawesi). Within tropical Africa it dominates only in southern Ghana and Côte D'Ivoire, and within South America, only within northern Venezuela and the south-western headwaters of the Amazon (FAO-UNESCO, 2007; Driessen *et al.*, 2001). Outside of the humid tropics, the Acrisol group dominates in Georgia, Alabama and the Carolinas in the south-eastern USA. Indeed, piping within Acrisol soils is reported within this humid sub-tropical region of the USA (e.g. Uchida *et al.*, 2005). Alisol soils are associated with Acrisol soils but can be distinguished by cation exchange capacities exceeding $24 \text{ cmol}^+ \text{ kg}^{-1}$ clay (by $1 \text{ M NH}_4\text{O Ac}$ at pH 7). For example, the piped soils at Danum, East Malaysia (Table I) are classified as a Haplic Alisol (Chappell *et al.*, 1999a). Globally, the Acrisol group is estimated to cover 1000 million hectares, while the Alisol group covers approximately 100 million hectares (Driessen *et al.*, 2001). This combined area is equivalent to more than twice the size of Southeast Asia.

The limited number of reported studies even for Ultisol soils could imply that such humid tropical soils, while being influenced by macroporosity (e.g. van Noordwijk *et al.*, 1991; Schwartz *et al.*, 1998; Reichenberger *et al.*, 2002; Ciglasch *et al.*, 2005), do not have extensive soil piping. Alternatively, the limited number of published studies may result from a wider dearth of research on runoff processes within the humid tropics and the difficulty of observing pipe outlets and depression lines within the region's dense rainforest cover. Indeed, subsurface flow within some hillslopes can be strongly influenced by soil piping despite the only visible evidence of piping arising during the excavation of soil pits (Sherlock, 1997; Lancaster, 1999; Chappell and Sherlock, 2005).

Ultisol soil within the humid tropics may be more sensitive to piping due to the strong decline of saturated

Table I. Location (see also Figure 1) and characteristics of pipes reported for soils in the humid tropics, ordered by date of first publication reporting piping at each location

Location	Soil type	Detailed pipe study (Y/N)	Diameter(s) (cm)	Length(s) (m)	Outlet density (per channel length or ha)	Flow measured (Y/N)	Reference
Majeedi (W Malaysia)-1	Ultisol	N	—	—	—	N	Burton (1964)
Lalgarth (India)-2	Alfisol	Y	1.0	—	—	N	Banerjee (1972)
Garbeta (India)-2	Alfisol	Y	10	—	—	N	Banerjee (1972)
Kuala Lumpur (W Malaysia)-3	Ultisol	N	—	—	—	N	Morgan (1973)
Mt Duau (Papua New Guinea)-4	Ultisol	Y	3–30	10–15	—	N	Löffler (1974)
Niah (E Malaysia)-5	Ultisol	Y	20	>30	—	N	Baillie (1975)
Bkt Mersing (E Malaysia)-6	Ultisol	Y	20	>30	—	N	Baillie (1975)
Segan (E Malaysia)-7	Ultisol	Y	15–25	>90	—	N	Baillie (1975)
Palmas (Dominica)-8	Andisol	Y	30	1.5	Limited	Y	Walsh (1980)
Kali Mondo (Indonesia)-9	Andisol	N	—	—	—	N	Bruijnzeel (1983)
Pointe Baptiste (Dominica)-10	Alfisol	Y	Mostly <10	—	29/500-m channel	Y	Walsh and Howells (1988)
Bkt Timah (Singapore)-11	Ultisol	N	2–8	—	—	N	Chatterjea (1989)
Bkt Timah (Singapore)-11	Ultisol	N	10	—	—	Y	Sherlock (1997)
La Cuenca (E Peru)-12	Ultisol	N	—	—	6/0.75 ha	Y	Elsenbeer and Cassel (1990)
La Cuenca (E Peru)-12	Ultisol	Y	5	—	6/0.75 ha	Y	Elsenbeer and Lack (1996)
La Cuenca (E Peru)-12	Ultisol	Y	—	—	6/0.75 ha	Y	Elsenbeer and Vertessy (2000)
Danum (E Malaysia)-13	Ultisol	N	5	—	—	N	Sinun (1991)
Danum (E Malaysia)-13	Ultisol	Y	1–10	—	—	N	Bidin (1995)
Danum (E Malaysia)-13	Ultisol	Y	—	—	—	Y	Chappell <i>et al.</i> (1998)
Danum (E Malaysia)-13	Ultisol	Y	14	>30	—	Y	Sayer (2002)
Danum (E Malaysia)-13	Ultisol	N	10–60	—	—	Y	Chappell <i>et al.</i> (2004a)
Danum (E Malaysia)-13	Ultisol	Y	—	—	—	Y	Chappell and Sherlock (2005)
Danum (E Malaysia)-13	Ultisol	Y	2–30	>30	—	Y	Sayer <i>et al.</i> (2006)
Babinda (NE Australia)-14	Ultisol	N	—	—	Limited	N	Bonell and Balek (1993)
La Selva (Costa Rica)-15	—	N	—	—	—	N	Parker (1994)
Belalong (Brunei)-16	Ultisol	N	10–15	—	—	N	Dykes and Thornes (2000)
Talakaveri (India)-17	Ultisol	Y	Several centimetre	—	—	Y	Putty and Prasad (2000)
Bkt Tarek (Malaysia)-18	Ultisol	Y	3.0–7.5	0.07–0.80	—	Y	Negishi <i>et al.</i> (2007)

The soil type utilizes Soil Survey Staff (1999); FAO-UNESCO (2007) and the references listed.

hydraulic conductivity (K_s) with depth and associated periodic water-logging (cf. Dunne, 1990 p. 22; Driessen *et al.*, 2001). In addition, Ultisol soils (particularly those equivalent to Alisol soils) are considered to be highly erodible (Driessen *et al.*, 2001; Dudal, 2005).

Outlet and upslope distribution of soil pipes in humid tropical basins

Several studies within the humid tropics report the location of *pipe outlets* (with visible water-flow or signs



Figure 2. Photographs of two pipe outlets: (a) a 5-cm diameter pipe outlet showing a water-level within the pipe just at the ground surface (pipe location is at $4^{\circ}57'18''\text{N}$, $117^{\circ}48'12''\text{E}$ in the Danum area, East Malaysia), and (b) the emergence of a tracer slug (1 : 10 diluted, white emulsion paint) from a pipe outlet after its input to a former cicada burrow, 10-m upslope (see Figure 3b)

that it has taken place recently). Most pipe outlets in the humid tropics are observed on the lower sections of hillslopes, close to ephemeral or perennial stream channels (Burton, 1964; Banerjee, 1972; Löffler, 1974; Baillie, 1975; Walsh, 1988; Bruijnzeel, 1983; Walsh and Howells, 1980; Sayer, 2002; Sayer *et al.*, 2006; Putty and Prasad, 2000; Negishi *et al.*, 2007). From theoretical considerations (e.g. Terzaghi and Peck, 1948; Dunne, 1990; Jones, 1990), areas of a hillslope with greater seepage forces are more likely to generate pipes. Hillslope areas with greater seepage forces are likely to be downslope areas and areas of subsurface flow convergence—perhaps formed by depressions in the rock-head and its impact on the topography of overlying soil horizons. Once a pipe develops in an area of subsurface convergence, then drainage of this area would be increased and thereby further increase the degree of convergence to this area in a positive feedback (cf. theory of formation of solution dolines: Williams, 1985). The exact location of pipes within these zones of subsurface convergence is then likely to be a function of the local spatial variability in the saturated hydraulic conductivity, K_s (Dunne, 1990 p. 29; Vieira and Fernandes, 2004).

Within downslope areas of studied slopes in the humid tropics, the outlets can be: (i) an upwelling on the floor of the hillslope concavities or shallow side slopes (Figure 2a), (ii) a hole in the vertical headwall of small seeps (e.g. Elsenbeer and Lack, 1996), or (iii) a hole in the head or wall of incised channels (e.g. Löffler, 1974; Baillie, 1975; Walsh and Howells, 1988; Sayer, 2002; Sayer *et al.*, 2006; Negishi *et al.*, 2007). Indeed, some of the incised, ephemeral channels in the humid tropics are reported to be forming through: (1) retreat of headwalls accelerated by Coulomb failure and soil fall at pipe outlets (Löffler, 1974; Bidin, 1995), and/or

(2) collapse of short sections of pipe roof (Burton, 1964; Baillie, 1975; Sayer, 2002). Pipe outlets can be found on the lower sections of hillslopes both at the head of permanent stream channels (e.g. Negishi *et al.*, 2007) and on the sideslopes of permanent stream channels (e.g. Baillie, 1975; Walsh and Howells, 1988). In addition, within the Danum area of East Malaysia, 5 to 15-cm diameter pipe outlets can be observed to discharge water intermittently from the lower sideslopes of second- and third-order permanent channels (e.g. at $4^{\circ}58'39''\text{N}$, $117^{\circ}48'98''\text{E}$ and $4^{\circ}57'61''\text{N}$, $117^{\circ}48'17''\text{E}$; Figure 2b) but also through the bed of these permanent channels (e.g. at $4^{\circ}57'93''\text{N}$, $117^{\circ}48'00''\text{E}$).

While active pipe outlets are easily recognizable by their visible resurgence of water during rain-events (particularly during intense rain-events), the *upslope extent* of soil pipes is difficult to identify within soils of the humid tropics (Walsh and Howells, 1988; Negishi *et al.*, 2007). Table I shows that data on the upslope extent is rarely measured and reported within piping studies in the humid tropics. Löffler (1974); Sayer *et al.* (2006) and Baillie (1975) do, however, report pipes extending 15, 30, and 90 m within hillslopes, respectively. Pipes exposed by intermittent collapses are more readily identified. Figure 3 shows such a pipe at Segan (redrawn from Baillie, 1975) and at Danum (i.e. 'PT trail pipe' at $4^{\circ}57'68''\text{N}$, $117^{\circ}48'15''\text{E}$, redrawn from a survey by A. Freer) both in East Malaysia.

Further evidence for the presence of soil pipes extending away from near-channel outlets has come from their observation within man-made soil pits (Sinun, 1991; Elsenbeer and Lack, 1996; Sherlock, 1997) and slope cuts (Morgan, 1973; Parker, 1994; Putty and Prasad, 2000). However, with such data, only the presence of pipes within upslope areas can be established, not their length.




Figure 3. Lateral pipe cum gully systems at: (a) Segan, East Malaysia (redrawn from Baillie, 1975) and (b) the Palum Tambun trail area of Danum, East Malaysia ($4^{\circ}57'68''\text{N}$, $117^{\circ}48'15''\text{E}$; redrawn from a survey by A. Freer). Former cicada burrow X is shown to be connected hydrologically to the river-bank outlet (Y) during the tracer test illustrated in Figure 2b

Better experimental evidence for pipe length comes from observations at Danum. Here flowing pipe outlets in gully sidewalls are sometimes associated with a very shallow (<10-cm relief) linear depression extending away from the pipe outlet in the gully (e.g. pipe outlet 'B' in Figure 3 of Sayer, 2002). Recently, artificial addition of tracer (namely 1:10 diluted, emulsion paint of various colours) into several 3-cm diameter natural shafts (probably former cicada burrows) along the line of this pipe depression (at up to 10 m from the pipe outlet) was visible within minutes to tens of minutes at the outlet (personal observation). Such tracer tests showing upslope extent of individual pipes (and the connectivity with some former cicada burrows) have been undertaken on other pipes in the Danum area. Figure 2b shows emergence of a tracer slug from a pipe outlet (marked with a 'Y' in Figure 3b) after its input to a former cicada burrow, in this case 10-m upslope (marked with an 'X' in Figure 3b).

Other evidence for the presence of pipes extending away from channel areas may come from the behaviour of soil moisture probes. Figure 4 shows a time series of soil moisture content measured at 0.50-m depth on a steep slope section at $4^{\circ}58'52''\text{N}$, $117^{\circ}49'03''\text{E}$ (Danum, East Malaysia). After installation, it was noticed that the profile of soil moisture measurements at this location was immediately downslope of a shallow linear

depression that extends up towards the hillslope divide. The marked bimodal recessions shown in Figure 4 (with upper sections drying just as quickly as the soil wetted) may indicate that the soil at this depth is being saturated by a soil pipe beneath the depression during storm events. This unusual behaviour was not noted at other moisture measured profiles throughout the Baru basin (Suhaimi, in prep). This hillslope was also subject to an earlier tracer test using a steady-state application of water (followed by pulse of technicium-99m tracer). Water was applied to a trench upslope of the same linear depression, and at e.g. 19 m further downslope (in riparian piezometer No. 4) the peak in the water breakthrough curve was attained after only 185 min. Following the analysis of Radulovich *et al.* (1989), this rapid water migration would be consistent with the presence of a preferential pathway such as a pipe system, as inferred from the moisture dynamics.

Negishi *et al.* (2007) attempted to measure pipe lengths by inserting a straight stick into several pipe outlets at their Bukit Tarek site in West Malaysia. They were only able to insert the stick up to 80 cm from the outlet. Given that individual pipes can follow a tortuous route (see e.g. Figure 2a, b), insertion of straight stick will be soon stopped even where a pipe extends over many metres. Indeed, the general lack of measurements of pipe length within studies in the humid tropics (Table I) is no basis for assuming that individual pipes do not extend beyond the few tens of centimetres visible by looking into an outlet.

From personal observation, it should be emphasized that some flowing outlets near ephemeral or perennial streams are not associated with either a line of pipe collapses or a shallow linear depression upslope of the outlet (e.g. at $4^{\circ}58'39''\text{N}$, $117^{\circ}48'98''\text{E}$). In these circumstances, establishing pipe length would certainly require the progressive excavation of the whole hillslope or monitoring pipe outlets for tracers applied to upslope sections of pipe (Smart and Wilson, 1984; Anderson *et al.*, 2008). Indeed, these approaches would be beneficial on all humid tropical hillslopes with evidence of piping. Only with observations of: (i) lines of pipe collapses (Baillie, 1974), (ii) whole hillslope excavations (Anderson *et al.*, 2008) or (iii) tracing along pipe sections (Smart and Wilson, 1984) can the typical, upslope extent of piping be quantified in the humid tropics.

Location of lateral pipes within soil profiles in humid tropical basins

Within the humid tropics, the solum (or A and B soil horizons) of Ultisol soils can range from <0.5 m to >1.5-m deep (e.g. Acres *et al.*, 1975). Lateral soil pipes exposed by gully incision, soil pit excavation or slope cutting during road construction are found at any depth within this range and several pipes at different depths can be present in the same exposure of a soil profile (Bidin, 1995; Negishi *et al.*, 2007; Figure 5). As noted earlier, pipes also may be present in the regolith below the solum.




Figure 4. A time series of volumetric moisture content ($\text{cm}^3 \text{cm}^{-3}$) monitored at 0.5-m depth and immediately downslope of a shallow (<10-cm relief) linear depression (at $4^\circ 58' 52'' \text{N}$, $117^\circ 49' 03'' \text{E}$ within the 2M Baru basin, Danum, East Malaysia). This depression extends up towards the hillslope divide. The marked bimodal recessions, with upper sections drying as quickly as the soil wetted, may indicate that the soil matrix at this depth is being saturated by a lateral pipe during storm events. This behaviour was not noted at other moisture measurement profiles throughout the Baru basin (Suhaimi, in prep). The soil moisture values at the measurement profiles were observed using Campbell Scientific Ltd CS616 probes and automatically recorded every 5 min using a CR1000 data-logger

At Niah (East Malaysia), the greatest depth in the soil profile where pipes have been recorded is 1.5 m (Baillie, 1975), at Pointe Baptiste, Dominica, it is 1.3 m (Walsh and Howells, 1988), at Bukit Tarek, West Malaysia, 1.5 m (Negishi *et al.*, 2007) and at Danum, East Malaysia, 1.4 m (Figure 5a). For example, Figure 5a shows the location of active, lateral pipes (with evidence of water-flow) within a 1.5-m high gully sidewall ($4^\circ 57' 18'' \text{N}$, $117^\circ 48' 12'' \text{E}$), while Figure 5b shows them within a gully head 5 km away (redrawn from Bidin, 1995). The locations of small macropores only ca. 0.5 to 1-cm diameter are also shown within Figure 5a, b, but it is not known which of these regularly conduct significant amounts of water-flow (i.e. 'hydrologically active macropores') and which of the faunal burrows (associated with worms, termites etc.) rarely conduct water. Within gully walls that contain active lateral pipes at several levels, the largest pipes tend to be at the base of the gully wall (e.g. Figure 5a, b). The same pipes also tend to flow more frequently (personal observation at Danum).

Three factors may affect the location of lateral pipes at one or more depths within Ultisol soil profiles within the humid tropics. These factors are vertical changes in: (i) K_s , (ii) soil erodibility, and (iii) proximity to the local water-table.

1. *Saturated hydraulic conductivity (K_s) reduction with depth:* Research outside of the tropics has shown that large lateral pipes are often found immediately above layers of reduced K_s (Yasuhara, 1980; Jones, 1981 p81). As soils within the humid tropics often exhibit a predominantly monotonic decline of K_s with depth (Chappell and Ternan, 1992; Elsenbeer, 2001; Chappell and Sherlock, 2005), a lower K_s in the strata immediately below a lateral pipe is likely. Figure 6 shows the vertical distribution of K_s of soil profiles containing piping at La Cuenca (Elsenbeer and Lack, 1996; Elsenbeer and Vertessy, 2000), Bukit Tarek (Ziegler *et al.*, 2006) and Danum (Chappell *et al.*, 1998).

2. *Local increases in erodibility of piped strata:* Aggregate stability has been shown to be lower within piped soil horizons within temperate and semi-arid climates (Beckedahl, 1977; Jones, 1981). Such data are extremely limited for piped soils in the humid tropics. A small number of aggregate stability tests were, however, undertaken on a soil layer with active piping and on adjacent layers at one location in the Danum area (Chappell *et al.*, 1999a). The piped layer at 0.94-m depth (in a profile surveyed by Bidin, 1995; Figure 5b) had a lower aggregate stability (i.e. 60%) compared to higher strata (i.e. 77–100%).

Exchangeable Sodium Percentage (ESP) is inversely correlated with aggregate stability within some soil types (Goldberg *et al.*, 1988). Within Alfisol soils of dry tropical Zimbabwe, Stocking (1981) observed that soil horizons with pipes have a higher ESP compared to non-piped horizons. The study of Chappell *et al.* (1999a) conducted within the humid tropics, demonstrated a similar relationship, with an ESP of 7.43% for the piped layer, compared to 2.58–4.14% for the layers higher in the profile.

3. *Changing likelihood of water-flow in lateral pipes with depth:* Walsh and Howells (1988), Bidin (1995) and Figure 5a show that where lateral pipes at several levels are exposed by gully incision, the larger pipes that also flow more frequently, tend to be located at the base of channel banks. In some contrast, the study of Negishi *et al.* (2007) showed that while flow was correlated with the size of pipe outlet, neither was correlated with depth at their studied gully head in West Malaysia.

Tracer application at Danum (e.g. at pipes shown in Figure 5a and shown in Figure 3 of Sayer, 2002) would indicate that multilevel, lateral pipes may be connected. Where this is the case, it is expected that the deeper, lateral pipes would maintain local saturation or free-surface flow for longer (with higher pipes draining into




Figure 5. Soil profiles along incised gullies that contain small macropores (which may or may not transmit water) and soil pipes either with observed flow or with evidence of previous flow. Survey (a) is of a gully sidewall at 4°57.18'N, 117°48.12'E (redrawn from a survey by A Freer), and (b) a nearby gully head (redrawn from Bidin, 1995)




Figure 6. The median value of saturated hydraulic conductivity (K_s) at each depth in the soil profile representative of three locations with soil piping, namely: (i) La Cuenca, Peru (crosses; data from Elsenbeer and Lack, 1996 and Elsenbeer and Vertessy, 2000), (ii) Bukit Tarek, West Malaysia (closed circles; data from Ziegler *et al.*, 2006) and (iii) Danum, East Malaysia (open circles; data from Figure 4a in Chappell *et al.*, 1998)

these; cf. Jones and Connelly, 2002). With more water-flow, greater pipe enlargement would be expected, so making these pipes larger. Consequently, there may be a two-way relation between the local pipe distribution and hydrological functioning at some humid tropical sites.

Vertical pipes and cicada burrows

Cicadas, insects mostly of the family Cicadidae, when newly born as nymphs, excavate vertical burrows within humid tropical soils. After living on roots for 2–5 years, they emerge from the surface via 8 to 14-cm high domed

‘chimneys’ (Lander, 1895; Sinun, 1991). These chimneys erode away within a few months (Sinun, 1991), to leave an exposed vertical shaft some 3 cm in diameter. Some cicada burrows remain as isolated, dry shafts, e.g. the 60-cm deep vertical shaft shown in Figure 7 that was filled with Plaster of Paris (calcium sulphate hemihydrates, $CaSO_4 \cdot \frac{1}{2}H_2O$) and then excavated. Other cicada shafts have been shown to connect to large, lateral pipes. For example, cicada burrow marked with an ‘X’ in Figure 3b has been confirmed (using a diluted emulsion paint tracer) to connect to a large lateral pipe. Other cicada burrows have been shown to connect to lateral pipe networks by attempting, but failing to fill the burrows with a 6 l/s water pump (in a soil where the K_s reduces to <10 cm/hr at 0.45-cm depth; Figure 6c). Furthermore, in some downslope locations within the Danum area, water has been seen to spout from such holes during rain-events; holes, which look like the holes found in hillslope areas containing clusters of many cicada chimneys (personal observation). Where lateral pipe-flow erodes a connection with a vertical cicada ‘shaft’, and the piezometric surface (within the pipe system) goes above the local ground surface, pipe flow would be expected to emerge from the ground via these former cicada burrows. The pipe outlet shown in Figure 2a may have once been a cicada ‘shaft’. Where such vertical cicada ‘shafts’ have such hydrological connections with the large lateral pipes they should be considered as part of the pipe system.

A repeat survey of cicada ‘chimneys’ in six 20 m² plots at Danum by Sinun (1991) showed densities of 15 chimneys per m². Thus, density was as high as the 14 chimneys per m² found by the seminal study of Scully (1942) at sites within the USA. Given that cicadas are present throughout the humid tropics, the possibility that the vertical shafts that they create could act as outlets for pipe flow requires further investigation.



Figure 7. A vertical cicada burrow filled with Plaster of Paris and then excavated (undertaken by V. Grimsey). The shaft extends to a depth of 60 cm (undisturbed soil is observed beneath), and is located only 20 cm from the bank of a gully formed by pipe collapse ($4^{\circ}57'68''\text{N}$, $117^{\circ}48'15''\text{E}$; Figure 3b). The shaft diameter is mostly between 2 and 3 cm. The top 10-cm section of the shaft has been removed and is not visible in the photograph

Effect of hillslope disturbance on piping within the humid tropics

In the humid tropics, where forestry roads have been built across ephemeral stream channels without the incorporation of a drainage conduit (typically a hollow log culvert) new pipes can sometimes be observed to develop through the fill materials (personal observation, e.g. at $4^{\circ}58'51''\text{N}$, $117^{\circ}49'14''\text{E}$). However, there are no published studies for the humid tropics that have examined how such soil and hillslope disturbances may affect the incidence or hydrological functioning of such soil pipes, as was studied by Ziemer (1992) within a temperate region of the USA. This contrasts with the number of studies quantifying the loss of small macropores resulting from tropical rainforest disturbance (e.g. Spaans *et al.*, 1989; Balbino *et al.*, 2004; Schack-Kirchner *et al.*, 2007).

HYDROLOGICAL FUNCTIONING OF PIPES WITHIN HUMID TROPICAL SOILS

Lateral soil pipes within the humid tropics are an extreme form of preferential flow that can focus the drainage and subsequent outflow of a significant proportion of hillslope water (Löffler, 1974). At this hillslope scale, we might consider a continuum of increasing preferential flow from topographic convergence to topographic convergence with a *percoline* (i.e. linear zone of high K_s : Bunting, 1961) to a long lateral pipe. Within some instrumented microbasins in the humid tropics, a significant proportion




Figure 8. Lateral saturated hydraulic conductivity of the soil profile at Danum, East Malaysia after core-based permeametry values have been up-scaled (solid line), together with the *effective lateral K_s* values derived from inversion of a catchment model (broken line; data from Chappell *et al.*, 1998). The latter includes the effects of soil piping on catchment-scale runoff generation, and only minimal overland flow

of the streamflow generation can come via lateral soil pipes (Sayer *et al.*, 2004). The study of Putty and Prasad (2000) states 25%, Walsh and Howells (1988) state >20%, Negishi *et al.* (2007) state 48% from several pipes at the channel head, while Sayer *et al.* (2004, 2006) show up to 47% from a single pipe resurgence.

Impact of lateral pipes on effective K_s of humid tropical hillslopes

Considering the hillslope as a hydrological unit, the presence of large, lateral pipes within the soil will increase the *effective, lateral saturated hydraulic conductivity* of the soil profile. Figure 8 shows the measured K_s of the Danum soil profile after being up-scaled to a *lateral K_s* value (adapted from Chappell *et al.*, 1998, 2004b). The broken line in the same figure shows the *effective lateral K_s* values derived from inversion of a catchment model (TOPMODEL) applied to rainfall–runoff data from the Baru catchment (similarly adapted from Chappell *et al.*, 1998, 2004b). The latter implicitly includes the dominant runoff pathways (thus, potentially, the effects of soil piping) on catchment-scale runoff generation. Studies undertaken outside of the tropics (e.g. Brooks *et al.*, 2004) show a similar magnitude of difference between the K_s for a hillslope profile derived from measurements at a point (e.g. core-based or auger hole-based tests) and those inverted from the whole hillslope or catchment behaviour. For hillslopes containing these extreme preferential pathways, the actual patterns of subsurface flow may bear little relation to those indicated by small core-based (or auger hole-based) K_s tests. Rather than flow responding to the patterns in K_s linked to tropical soil horizons (Elsenbeer, 2001) or a tropical soil catena (Chappell and Ternan, 1992), patterns may be overwhelmingly a function of the spatial distribution of the lateral pipes network

(Kirkby, 1988; Bonell and Balek, 1993, p. 198; Bonell, 2004 p. 317). Indeed, the tropical tracer study of Chappell and Sherlock (2005) provides evidence that this may be the case with subsurface flow direction being little related to the vertical distribution of core-based K_s values.

Comparison of soil pipe outlet and stream time constants within humid tropical basins

Time series' of water discharge emerging from lateral soil pipes have been measured at several experimental sites within the humid tropics (Walsh and Howells, 1988; Sayer *et al.*, 2004, 2006; Negishi *et al.*, 2007). At other sites within the humid tropics, surface flows have been gauged a few metres downslope of pipe outlets (Elsenbeer and Lack, 1996; Elsenbeer and Vertessy, 2000; Putty and Prasad, 2000; Sayer *et al.*, 2004, 2006). The studies of Sayer *et al.* (2004) and Negishi *et al.* (2007) allow comparison of flow in an ephemeral stream channel with that proportion generated by the resurgence of pipe flow. The flow observed within the channels will be a mix of pipe flow, exfiltration of water from the soil matrix (including flow through small macropores), and precipitation falling onto the locally saturated soil. The gauged pipes in both the Negishi *et al.* (2007) and Sayer *et al.* (2004) studies cease flowing several hours after rainfall. The hydrograph recessions for the gauged pipes in these two studies can be faster than that of flow within the ephemeral stream channel (see e.g. pipe 3 in Figure 6 of Negishi *et al.*, 2007) or indeed slower (see e.g. W3 pipe in Figure 1 in Sayer *et al.*, 2004). Figure 9 shows an 11-day period of discharge gauged at the outlet of the W3 pipe, and at 30-m downstream in the ephemeral stream channel. The difference in the rate of recession and hence time constant of the response is apparent, with the pipe having longer recessions. The nearby gauged W8S5 third-order stream (4°57'68"N, 117°47'66"E) also has faster hydrograph recessions in comparison to the W3 pipe. The W8S5 stream has a time constant, TC (determined by a first-order transfer function, with or without a non-linear transform: Chappell *et al.*, 1999b) of approximately 1 h and hence is comparable to that of the nearby Baru stream (Chappell *et al.*, 1999b). In contrast, the W3 pipe has a time constant of approximately 40 h. The time constants relate to the slope of the recession of output (i.e. pipe flow or channel flow) relative to that of the input (i.e. hyetograph). The longer time constant for the W3 pipe outlet (gauging an area of <1 ha: Sayer *et al.*, 2006) implies that the dominant water pathway to this outlet is significantly slower than that of the 1.5 km² W8S5 basin, or indeed of the 1.32 ha microbasin gauged only 30-m downstream of the pipe outlet.

Faster paths observed in the streamflow records could be related to (i) saturated ground, (ii) a dominance of hillslopes shorter than those with extensive pipe systems, or (iii) lateral pipes higher within the soil profile that have a more flashy flow regime. Local observations of these three factors are available:

1. Only a small proportion of the streamside soils within the W8S5 or Baru basins are saturated (i.e. from




Figure 9. A 11-day time series of discharge gauged at the outlet of the W3 pipe, and 30-m downstream in the ephemeral channel (data collected by A Sayer and J Hanapi)

moisture measurements at 1197 points within 10 m of all perennial channel within the Baru, less than 10% are more than 95% saturated: S. Abu Bakar, personal communication).

2. As the resistance to flow within a large soil pipe is considerably less than the resistance to flow expected through the matrix (or even fine macropores) of the silt loam soil (Chappell *et al.*, 1999a), the slower response is more likely to relate to the mechanism of recharge of the W3 pipe. The W3 pipe basin has an elongated headwater area, which extends to more than 60-m upslope of the pipe outlet (see Figure 2 in Sayer *et al.*, 2006). Such basin shape effects can damp the delivery of water towards a main pipe (or stream; see Bras, 1990 p. 397).
3. An 8-cm diameter pipe which supplies all of the flow to weir E6 in the Baru catchment (and then drains into a 60-cm diameter pipe: Chappell *et al.*, 2004a) has a time constant of approximately 1 h and hence is comparable to that of the third-order Baru stream (or W8S5 stream). This pipe outlet only produces flow irregularly and is located towards the top of the gully head at a height of 1.2 m. This contrasts with the W3 pipe of Sayer *et al.* (2004) with its slower time constant, which discharges close to the base of the gully head, and hence close to the level which is regularly saturated (i.e. the local water-table along the channel: see Bidin *et al.*, 1993; Chappell *et al.*, 2006). Indeed, visual observations would suggest that the pipe above weir E6 may be discharging water on to an initially dry channel, as noticed by Putty and Prasad (2000) at monitored sites in India.

The gauged pipe within the 0.75 ha La Cuenca basin (East Peru) has an even shorter time constant of only a few minutes (Elsenbeer and Vertessy, 2000). This pipe outlet is less than 10 m from an ephemeral stream channel, but is 16 m above the height of the perennial stream ('P1' in Figure 3 of Elsenbeer and Vertessy, 2000), and hence the 'permanent' water-table.




Figure 10. A perceptual model of the hydrological function (notably drainage) of soil pipes within humid tropical soils. The figure shows two different transects or vertical slices through the soil profile, one in a channel head depression (b and c), and one traversing a gully sidewall (d and e); both are located on an ephemeral stream channel (a). Two periods are illustrated for each transect. First, a 'wet period' is given to show hydrological functioning during a typical rain-event or within a few hours of such an event, where the ephemeral stream channel maintains some saturation and channel flow (b and d). The second period is for an episode of either intense rainfall or a long duration event where the soil profile within near-channel areas has a water-table at or close to the ground surface (c and e). The lower pipe in d and e is located close to the base of the B soil horizon

Possible role of cicada 'shafts' as outlets for lateral soil pipes

The widespread occurrence of vertical shafts excavated by cicada nymphs is observed in the Danum area. Overland flow running into these shafts has not been observed. As noted earlier, holes (which have the appearance of slightly enlarged cicada shafts) have, however, been observed to spout water during intense rainfall events and many have been shown to be connected to large, lateral pipes. Thus, some of these cicada shafts are considered to function as pipe outlets (Figure 2a) while others remain as dry shafts, though none have been observed to behave as insurgencies within this particular humid tropical environment. This contrasts with some *regolith pipes* within Columbia, which have been shown to act as insurgencies for channel flow (Feininger, 1969). Given that cicadas are present throughout the humid tropics, the possibility that cicada 'shafts' can be modified by seepage forces from lateral pipes to act as pipe outlets needs to be examined throughout the region.

Recharge of lateral soil pipes

The mechanism for recharge of the lateral pipes remains unclear within the literature for piping within the humid tropics, and from wider observations at Danum. Given that the moisture content of the Danum subsoil remains close to saturation in so called *tension-saturated wedges* (Sherlock, 1997; Suhaimi, in prep) perhaps such

zones within piped slopes rapidly saturate during rain-events and thereby recharge pipe flow (cf. Dunne, 1990). With rapid saturation, the network of small macropores within these wedges may fill connecting near-surface macropores with those in the lower soil profile (cf. Sidle *et al.*, 2001), which are themselves connected with the soil pipes.

A perceptual model of the hydrological function of soil pipes in the humid tropics

Based on the preceding review of literature pertaining to the humid tropics and the additional observations included, Figure 10 illustrates a tentative *perceptual model* (using terminology of Beven, 1991, 2001) of the hydrological functioning of soil pipes within humid tropical soils. This model is based particularly on direct observations of several pipe systems within the Danum area, East Malaysia (e.g. at 4°57.18'N, 117°48.12'E), but seeks to be consistent with published findings from elsewhere in the humid tropics where the necessary observations of pipe extent have yet to be made. Figure 10 shows two transects or vertical slices through the soil profile, one in a channel head depression (where there is a gradual slope rather than a gully headwall, as in Figures 5b, 10a–c), and one traversing a gully sidewall (Figure 10a, d, e). Two periods are illustrated for each transect. First, a 'wet period' is given to show hydrological functioning during a typical rain-event or within a few hours of such an event, where the ephemeral channel (Figure 10a)

maintains some saturation and channel flow (Figure 10b, d). The second period illustrated is during either intense rainfall or a long-duration event where the soil profile within near-channel areas has a water-table at or close to the ground surface (Figure 10c, d; Bidin, 1995).

A large lateral pipe is shown running beneath the channel head (Figure 10b, c; following Baillie, 1975; Sayer, 2005). During typical rain-events, there may be no visible signs of these pipes (Figure 10b) except where the course of the lateral pipe intercepts the surface, collapses occur or a depression line is present (Burton, 1964; Baillie, 1975; Sayer, 2002, 2005; Figure 3). During intense or long-duration events, the water-table extends further up the depression so that it is above the ground surface at the locations of former cicada shafts connected to the lateral pipes (see Figure 2a). Under these conditions, the pipe flow then returns to the ground surface (to generate saturation overland flow) via the former cicada 'shafts' modified by pipe flow (Figure 10c).

In the sidewalls of incised gullies, those outlets of lateral pipes that are close to the channel floor (and close to the bedrock) are likely to maintain flow when the ephemeral stream channel has flow (Figure 10d). Such pipes are also likely to be only partially filled with water during these conditions. Furthermore, pipes higher within the profile may be dry (Figures 10d and 5a, b). When a large proportion of the riparian soil profile attains saturation during intense or long-duration events (Noguchi *et al.*, 1997; Sherlock, 1997; Chappell and Sherlock, 2005), then pipes higher within the profile may contain flow and also discharge into the ephemeral channel (Figure 10e; note that pipes higher within the profile of Figure 5a, b do show evidence of intermittent water-flow). Under these conditions, the deeper pipes may be completely filled with water and are under significant hydrostatic pressure.

This model illustrates some of the mechanisms of hillslope drainage by soil pipes. It needs to be extended to include the mechanisms that recharge the lateral pipes, though this needs more field survey and monitoring. Given the dearth of published work, this tentative model needs to be evaluated over a wide range of piped sites within the humid tropics.

CONCLUSIONS AND SUGGESTED RESEARCH PRIORITIES

The preceding review of the distribution and hydrological function of soil piping within the humid tropics was supplemented by additional information from the location stimulating the most published studies (Table I). The combined analysis has produced several findings, all needing to be explored with further research.

1. *Soil pipe distribution across the humid tropics* Most of the locations in the humid tropics with published reports of soil piping are within areas of Ultisol soil, though a few have either Alfisol or Andisol. Further

work is needed to determine why Ultisol soils (rather than Oxisols or other locally dominant soils) have a greater reported incidence of piping. While there appears to be a link between the Ultisol soil order and soil piping, it is not clear how widespread the phenomenon is within this extensive soil type. A more extensive field survey of the incidence of soil piping throughout the humid tropics is needed, particularly within areas of Ultisol soil.

2. *Upslope and vertical extent* Several studies report the incidence and location of pipe resurgences (or outlets) within humid tropical basins. The most frequently active of these resurgences are in the lower sections of hillslopes (Burton, 1964; Banerjee, 1972; Löffler, 1974; Baillie, 1975; Walsh, 1980; Walsh and Howells, 1980; Bruijnzeel, 1983; Putty and Prasad, 2000; Sayer, 2002; Sayer *et al.*, 2006; Negishi *et al.*, 2007). Very few studies have, however, attempted to measure the upslope extent of soil pipes in the humid tropics. Those studies that have measured lengths within humid tropical soils have shown that they can extend up to 15 m (Löffler, 1974), 30 m (Sayer *et al.*, 2006; Figure 3b) or 90 m (Baillie, 1975) from pipe outlets. Within deep regolith strata in humid tropical Columbia, individual pipes have been shown to extend over several hundred metres (Feininger, 1969). Such limited data cannot be generalized, so it is imperative that pipe length is determined at many more humid tropical sites, most particularly at those sites where pipe studies have already taken place. The success of experiments at the Danum humid tropical site using diluted paint to establish possible connections between the resurgences of large, lateral pipes and upslope pipe collapses may be the first step in this local pipe mapping. This work would then need to be supported by excavation of pipe systems upslope from the pipe outlets, perhaps following a similar approach to that of Anderson *et al.* (2008). Injecting Plaster of Paris or expanding foam into lateral pipes via upslope shafts or collapses could help to maintain the integrity and visibility of the pipes during the excavation phase. Indeed, systematic surveys to show how many of the vertical shafts produced by cicada nymphs have become connected (by piping processes?) to the lateral pipe systems would be helpful in this regard. Known volumes of low viscosity resin could be poured into these shafts to identify those shafts not capable of being filled and hence connected with the lateral pipe network. Of the few cicada burrows examined by this method at Danum (East Malaysia), those that were isolated could be filled with the addition of 300–700 cm³ of resin or dilute solutions of plaster.

Within humid tropical soils, lateral pipes are observed to be present at many different depths. Several studies do, however, report that those lateral pipes that are most often flowing occur at a depth of 1.3–1.5 m, which corresponds to the base of the soil, i.e. lower B soil horizon (Baillie, 1975; Walsh and Howells, 1988; Negishi *et al.*, 2007; this study). Several humid tropical

studies have also reported the incidence of piping beneath the soil profile within deep regolith (Dixey, 1920; de Chélat, 1938; McBeath and Barron, 1954; Patz, 1965; Thomas, 1968; Feininger, 1969; Bremer, 1973). Within humid tropical countries, these deeper *regolith pipes* are often reported to form beneath a duricrust (Goudie, 1973). These studies have not involved hydrological investigations and are not in areas of the humid tropics with published hydrological studies on piping within the solum (i.e. A and B soil horizons). New studies that investigate the relationship between these deeper pipe systems (i.e. 2 to >10-m depth) and those that occur within 2 m of the ground surface (i.e. soil) are needed.

3. Lumped response of pipe systems from monitoring pipe resurgences Active pipe resurgences within humid tropical basins can be gauged relatively easily using weir boxes (e.g. Figure 2.3 in Sayer, 2005). These devices do not significantly affect the natural flow from pipes or surrounding soils, unlike devices installed within excavated hillslope pits (Knapp, 1970). As such, they provide a reliable way of measuring (preferential) hillslope drainage prior to the formation of channel flow (Negishi *et al.*, 2007) or overland flow outside of channels (Elsenbeer and Vertessy, 2000). The resulting high quality time series of hillslope drainage (e.g. Sayer, 2005; Negishi *et al.*, 2007) unaffected by artificial conditions imposed by the measurement make them amenable to characterization, physical interpretation, and are potentially useful for rainfall–runoff modelling.

Time constants (or residence times of response) of measured pipe resurgences in the humid tropics vary from those faster to those slower than the channel flow into which they drain. From the few time series available, the majority of the gauged pipes with outlets just above the height of the channel flow have responses dominated by a time constants longer in comparison to the channel flows. This would imply that pathways with recessions faster than those of deeper pipes contribute significant volumes of flow to the channels. Such faster paths within humid tropical basins may be related to a combination of (i) overland flow and pressure-wave effects in areas of saturated ground, (ii) hillslopes shorter than those with deep soil pipes or (iii) lateral pipes higher within the soil profile that have a more flashy flow regime.

As large, lateral pipes will offer less resistance to flow in comparison with the soil matrix (or small macropores) through which the pipes are recharged, quantification of the recharge processes may be more important than the pipe geometry in the interpretation of the time constants of response. These recharge processes are very poorly understood within humid tropical basins and warrant detailed investigation.

4. Recharge of large, lateral pipe systems Models of the recharge pathways of lateral pipes within temperate soils have been developed (e.g. Jones and Connelly, 2002). Field investigations within humid tropical basins are needed to allow similar quantification. In piped hillslopes where pipe resurgences are gauged (by

existing or new structures), many automated measurements of soil moisture content and pressure potential are needed at multiple depths. With intensive spatial sampling these data would show which parts of the lateral pipe network are within saturated soil. One minute time steps may be needed given the flashy nature of some shallow lateral pipes (see Elsenbeer and Lack, 1996). Such data would also show the timing of the soil wetting (at different depths and slope positions) with respect of the time series of pipe outflow. These data could be analysed using a combination of transfer function approaches (cf. Fawcett *et al.*, 1997; Chappell *et al.*, 1999b), and physics-based models of soil moisture dynamics (Radulovich *et al.*, 1989; Germann *et al.*, 1997; Wollschläger *et al.*, 2009). These physics-based approaches would necessitate that hydraulic conductivity is sampled systematically throughout the monitored hillslopes. The saturated hydraulic conductivity (K_s) of typically fines-rich, Ultisol soils could be measured accurately with techniques such as ring permeametry or piezometer tests, as such methods are less sensitive to artificial smearing effects within fines-rich soils (Chappell and Lancaster, 2007). Ideally, the upper part of the unsaturated hydraulic conductivity curve would be measured with tension infiltrometry. These measurements may also show the presence of percolines and, where present, their relation to the lateral soil pipes. This destructive sampling would need to be undertaken after the soil moisture monitoring and perhaps at the same time as the excavation of the pipe networks discussed earlier.

Understanding the role of the small, but numerous macropores formed by tropical tree roots (van Noordwijk *et al.*, 1991) and soil fauna (Blanchart *et al.*, 1999) on these hydraulic conductivity measurements and the processes recharging lateral soil pipes is a fundamental research question (Beven and Germann, 1982) that has yet to be addressed within hillslopes of the humid tropics.

5. Application With detailed investigation of the spatial distribution of lateral pipes and surrounding hydraulic conductivity within specific hillslopes in the humid tropics, together with equally detailed monitoring of soil moisture dynamics, robust *conceptual models* (using the terminology of Beven, 1991, 2001) of piping within humid tropics may be capable of being developed. If a more extensive survey of the presence of piping within tropical Ultisols shows they are present within many hillslopes covered by Ultisols, then the availability of such models would allow better quantification of the hydrological processes over large areas of the humid tropics. This could allow improvement to rainfall–runoff models and hence a better understanding of why humid tropics basins may have very flashy responses. Secondly, this could allow improvement to models of ion transport that represent more accurately the migration of nutrients or contaminants within natural or disturbed hillslopes (Elsenbeer and Lack, 1996; Laabs *et al.*, 2002). Lastly, this would provide a better description of the flow

processes leading to gully development and sediment delivery within the humid tropics (Baillie, 1975; Sayer *et al.*, 2006). These geomorphological processes would also benefit from studies quantifying the impact of forestry or agricultural disturbances to tropical pipe systems. As expected for a region containing a dearth of studies, this review has highlighted many aspects of soil piping that warrant further investigation.

ACKNOWLEDGEMENTS

This synthesis of research findings benefits from past discussions with Keith Beven, Kawi Bidin, Mike Bonell, Adam Freer, Verity Grimsey, Jamil Hanapi, Paun Kukuon, Jeff McDonnell, Shoji Noguchi, Mark Sherlock, Roy Sidle, Waidi Sinun, and Rory Walsh, for which I am grateful. The constructive comments of the two reviewers are also appreciated.

REFERENCES

- Acres BD, Bower RP, Burrough PA, Folland CJ, Kalsi MS, Thomas P, Wright PS. 1975. The soils of Sabah. *Land Resource Study 20*, vol. 5. Ministry of Overseas Development: Surbiton; 1–147.
- Anderson AE, Weiler M, Alila Y, Hudson RO. 2008. Dye staining and excavation of a lateral preferential flow network. *Hydrology and Earth System Science Discussion 5*: 1043–1065.
- Baillie IC. 1975. Piping as an erosion process in the uplands of Sarawak. *Journal of Tropical Geography 41*: 9–15.
- Balbino LC, Bruand A, Cousin I, Brossard M, Quéting P, Grimaldi M. 2004. Change in the hydraulic properties of a Brazilian clay Ferralsol on clearing for pasture. *Geoderma 120*: 297–307.
- Banerjee AK. 1972. Morphology and genesis of 'pipe structure' in ferrallitic soils of Midnapore District, West Bengal. *Journal of the Indian Society of Soil Science 20*: 399–402.
- Beckedahl HR. 1977. Subsurface erosion near the Olivier-shoek pass, Drakensberg. *South African Geographical Journal 59*: 130–138.
- Beven KJ. 1991. Spatially distributed modelling: conceptual approach to runoff prediction. In *Recent Advances in the Modelling of Hydrologic Systems*, Bowles DS, O'Connell PE (eds). Kluwer: Dordrecht; 373–387.
- Beven KJ. 2001. *Rainfall-runoff Modelling: The Primer*. Wiley: Chichester; 360.
- Beven K, Germann P. 1982. Macropores and water flow in soils. *Water Resources Research 18*: 1311–1325.
- Bidin K. 1995. *Subsurface flow controls of runoff in a bornean natural rainforest*. MSc thesis, University of Manchester, Manchester, 128.
- Bidin K, Douglas I, Greer T. 1993. Dynamic response of subsurface water levels in a zero-order tropical rainforest basin, Sabah, Malaysia. In *Proceedings of the Hydrology of Warm Humid Regions*, Gladwell JS (ed.). IAHS Publication No. 216, IAHS Press: Wallingford; 491–496.
- Blanchart E, Albrecht A, Alegre J, Duboiset A, Gilot C, Pashanasi B, Lavelle P, Brussaard L. 1999. Effects of earthworms on soil structure and physical properties. In *Earthworm Management in Tropical Agroecosystems*, Lavelle P, Brussaard L, Hendrix P (eds). CAB International: Wallingford; 149–172.
- Bonell M. 2004. Runoff generation processes in tropical forests. In *Forests, Water and People in the Humid Tropics*, Bonell M, Bruijnzeel LA (eds). Cambridge University Press: Cambridge; 314–406.
- Bonell M, Balek J. 1993. Recent scientific developments and research needs in hydrological processes of the humid tropics. In *Hydrology and Water Management in the Humid Tropics*, Bonell M, Hufschmidt MM, Gladwell JS (eds). Cambridge University Press: Cambridge, 167–260.
- Bonell M, Gilmour DA. 1978. The development of overland flow in a tropical rainforest catchment. *Journal of Hydrology 39*: 365–382.
- Bras RL. 1990. *Hydrology: An Introduction to Hydrologic Science*. Addison-Wesley: Reading, MA; 643.
- Bremer H. 1973. Der Formungsmechanismus im tropischen Regenwald Amazoniens. *Zeitschrift Fur Geomorphologie Supplementband 17*: 195–222.
- Brooks ES, Boll J, McDaniel PA. 2004. A hillslope-scale experiment to measure lateral saturated hydraulic conductivity. *Water Resources Research 40*: W04208. DOI: 10.1029/2003WR002858.
- Bruijnzeel LA. 1983. Evaluation of runoff sources in a forested basin in a wet monsoonal environment: a combined hydrological and hydrochemical approach. In *Hydrology of Humid Tropical Regions with Particular Reference to the Hydrological Effects of Agriculture and Forestry Practice*, Keller R (ed.). IAHS Publication No. 140, IAHS Press: Wallingford; 165–174.
- Bryan RB, Jones JAA. 1997. The significance of soil piping process: inventory and prospect. *Geomorphology 20*: 209–218.
- Bunting BT. 1961. The role of seepage moisture in soil formation, slope development, and stream initiation. *American Journal of Science 259*: 503–518.
- Burton CK. 1964. The older alluvium of Johore and Singapore. *Journal of Tropical Geography 18*: 30–42.
- Chang JH, Lau LS. 1993. Definition of the humid tropics. In *Hydrology and Water Management in the Humid Tropics*, Bonell M, Hufschmidt MM, Gladwell JS (eds). Cambridge University Press: Cambridge; 571–575.
- Chappell NA, Franks SW, Larenus J. 1998. Multi-scale permeability estimation for a tropical catchment. *Hydrological Processes 12*: 1507–1523.
- Chappell NA, Lancaster JW. 2007. Comparison of methodological uncertainty within permeability measurements. *Hydrological Processes 21*: 2504–2514.
- Chappell NA, Sherlock MD. 2005. Contrasting flow pathways within tropical forest slopes of Ultisol soil. *Earth Surface Processes and Landforms 30*: 735–753.
- Chappell NA, Ternan JL. 1992. Flow-path dimensionality and hydrologic modelling. *Hydrological Processes 6*: 327–345.
- Chappell NA, Ternan JL, Bidin K. 1999a. Correlation of physicochemical properties and sub-erosional landforms with aggregate stability variations in a tropical Ultisol disturbed by forestry operations. *Soil & Tillage Research 50*: 55–71.
- Chappell NA, McKenna P, Bidin K, Douglas I, Walsh RPD. 1999b. Parsimonious modelling of water and suspended-sediment flux from nested-catchments affected by selective tropical forestry. *Philosophical Transactions of the Royal Society of London Series B 354*: 1831–1846.
- Chappell NA, Douglas I, Hanapi JM, Tych W. 2004a. Source of suspended-sediment within a tropical catchment recovering from selective logging. *Hydrological Processes 18*: 685–701.
- Chappell NA, Bidin K, Sherlock MD, Lancaster JW. 2004b. Parsimonious spatial representation of tropical soils within dynamic, rainfall-runoff models. In *Forests, Water and People in the Humid Tropics*, Bonell M, Bruijnzeel LA (eds). Cambridge University Press: Cambridge; 756–769.
- Chappell NA, Vongtanaboon S, Jiang Y, Tangtham N. 2006. Return-flow prediction and buffer designation in two rainforest headwaters. *Forest Ecology and Management 224*: 131–146.
- Chappell NA, Sherlock M, Bidin K, Macdonald R, Najman Y, Davies G. 2007. Runoff processes in Southeast Asia: role of soil, regolith and rock type. In *Forest Environments in the Mekong River Basin*, Sawada H, Araki M, Chappell NA, LaFrankie JV, Shimizu A (eds). Springer Verlag: Tokyo; 3–23.
- Chatterjea K. 1989. Surface wash—the dominant geomorphic process in the surviving rainforest of Singapore. *Singapore Journal of Tropical Geography 10*: 95–109.
- Ciglasch H, Amelung W, Totrakool S, Kaupenjohann M. 2005. Water flow patterns and pesticide fluxes in an upland soil in northern Thailand. *European Journal of Soil Science 56*: 765–777.
- de Chérelat E. 1938. Le modele laterique de l'ouest de la Guinee Francaise. *Revue de Geographie Physique et de Geologie Dynamique 11*: 5–120.
- Dixey F. 1920. Laterization in Sierra Leone. *Geological Magazine 57*: 211–220.
- Driessen PM, Deckers JA, Spaargaren OC, Nachtergaele FO. 2001. *Lecture Notes on the Major Soils of the World, World Soil Resources Reports 94*. Food and Agriculture Organization of the United Nations: Rome; 334.
- Dudal R. 2005. Soils of Southeast Asia. In *Physical Geography of Southeast Asia*, Gupta A (ed.). Oxford University Press: Oxford; 94–103.
- Dunne T. 1990. Hydrology, mechanics and geomorphic implications of erosion by subsurface flow. In *Groundwater Geomorphology*, Higgins CC, Coates DR (eds). Geological Society of America: Boulder, CO; 1–28.
- Dykes AP, Thornes JB. 2000. Hillslope hydrology in tropical rainforest steeplands in Brunei. *Hydrological Processes 14*: 215–235.

- Elsenbeer H. 2001. Hydrologic flowpaths in tropical rainforest soils— a review. *Hydrological Processes* **15**: 1751–1759.
- Elsenbeer H, Cassel DK. 1990. Surficial processes in the rainforest of western Amazonia. In *Research Needs and Applications to Reduce Erosion and Sedimentation in Tropical Steeplands*, Ziemer RR, O'Loughlin CL, Hamilton LS (eds). IAHS Publication No. 192, IAHS Press: Wallingford; 289–297.
- Elsenbeer H, Lack A. 1996. Hydrological pathways and water chemistry in Amazonian rain forests. In *Advances in Hillslope Processes*, Anderson MG, Brooks SM (eds). John Wiley and Sons: Chichester; 939–959.
- Elsenbeer H, Vertessy RA. 2000. Stormflow generation and flowpath characteristics in an Amazonian rainforest catchment. *Hydrological Processes* **14**: 2367–2381.
- FAO. 2005. *Global Forest Resources Assessment 2005 (FRA 2005)*. Food and Agriculture Organisation (FAO): Rome; 350.
- FAO-UNESCO. 1974. *Soil Map of the World, Volume 1 Legend*. UNESCO: Paris; 62.
- FAO-UNESCO. 1988. *Soil Map of the World, Revised Legend*. UNESCO: Paris; 119.
- FAO-UNESCO. 2007. *Digital Soil Map of the World v3-6*. FAO: Rome.
- Fawcett CP, Young PC, Feyen H. 1997. Validation of data-based mechanistic non-linear rainfall-flow model. In *Proceedings of the twelfth International Conference on Systems Engineering, ICSE'97*, Coventry, 9–11 September 1997. Coventry University: Coventry; 252–257.
- Feininger T. 1969. Pseudokarst on quartz diorite, Colombia. *Zeitschrift Fur Geomorphologie* **13**: 287–296.
- Fekete BM, Vorosmarty CJ, Grabs W. 2002. High-resolution fields of global runoff combining observed river discharge and simulated water balances. *Global Biogeochemical Cycles* **16**: 15–1 to 15–10.
- Germann PF, Di Pietro L, Singh VP. 1997. Momentum of flow in soils assessed with TDR-moisture readings. *Geoderma* **80**: 153–168.
- Godsey S, Elsenbeer H, Stallard R. 2004. Overland flow generation in two lithologically distinct rainforest catchments. *Journal of Hydrology* **295**: 276–290.
- Goldberg S, Suarez DL, Glaubig RA. 1988. Factors affecting clay dispersion and aggregate stability of arid zone soils. *Soil Science* **146**: 317–325.
- Goudie A. 1973. *Duricrusts in Tropical and Subtropical Landscapes*. Clarendon Press: Oxford; 174.
- Gunn J. 1986. A conceptual model for conduit flow dominated karst aquifers. In *Karst Water Resources*, Giinay G, Johnson AI (eds). IAHS Publication No. 161, IAHS Press: Wallingford; 587–596.
- Jones JAA. 1981. *The Nature of Soil Piping: A Review of Research*. Geo Books: Norwich; 301.
- Jones JAA. 1990. Piping effects in humid lands. In *Groundwater Geomorphology*, Higgins CG, Coates DR (eds). Geological Society of America: Boulder, CO; 111–137.
- Jones JAA. 1994. Soil piping and its hydrogeomorphic function. *Cuaternario y Geomorfologia* **8**: 77–102.
- Jones JAA, Connelly LJ. 2002. A semi-distributed simulation model for natural pipeflow. *Journal of Hydrology* **262**: 28–49.
- Jones JAA, Richardson JM, Jacob HJ. 1997. Factors controlling the distribution of piping in Britain: a reconnaissance. *Geomorphology* **20**: 289–306.
- Kirkby MJ. 1988. Hillslope runoff process and models. *Journal of Hydrology* **100**: 315–339.
- Knapp BJ. 1970. *Patterns of water movement on a steep Upland Hillside, Plynlimon, Central Wales*. PhD thesis. University of Reading, Reading, 213.
- Laabs V, Amelunga W, Pintob A, Zecha W. 2002. Fate of pesticides in tropical soils of Brazil under field conditions. *Journal of Environmental Quality* **31**: 256–268.
- Lancaster JW. 1999. *Multi-scale estimation of effective permeability within the greenholes beck catchment*. PhD thesis. Lancaster University, Lancaster.
- Lander B. 1895. Domed burrows of Cicada Septendecim. *Journal of the New York Entomological Society* **3**: 33–38.
- Löffler E. 1974. Piping and pseudokarst features in the tropical lowlands of New Guinea. *Erdkunde* **28**: 13–18.
- McBeath DM, Barron CN. 1954. Report on the Lateritic Ore Deposit at Iron and Wamara Mountain, Berbice. Report Geological Survey Department, British Guiana, for year 1953. Geological Survey Department: Georgetown; 87–100.
- Morgan RPC. 1973. Soil-slope relationships in the lowlands of Selangor and Negri Sembilan, West Malaysia. *Zeitschrift fur Geomorphologie* **17**: 139–155.
- Negishi JN, Noguchi S, Sidle RC, Ziegler AD, Nik AR. 2007. Stormflow generation involving pipe flow in a zero-order basin of Peninsular Malaysia. *Hydrological Processes* **21**: 789–806. DOI:10.1002/hyp.6271.
- Noguchi S, Abdul Rahim N, Baharuddin K, Sammori T, Tani M, Tsuboyama Y. 1997. Rainfall characteristics of tropical rain forest and temperate forest: comparison between Bukit Tarek in Peninsular Malaysia and Hitachi Ohta in Japan. *Journal of Tropical Forest Science* **9**: 206–220.
- Parker GG. 1994. Soil fertility, nutrient acquisition, and nutrient cycling. In *La Selva: Ecology and Natural History of a Neotropical Rainforest*, McDade LA, Bawa KS, Hespenheide HA, Hartshorn GS (eds). University of Chicago Press: Chicago; 54–63.
- Patz MJ. 1965. Hilltop hollows: further investigations. *Uganda Journal* **29**: 225–228.
- Peel MC, Finlayson BL, McMahon TA. 2007. Updated world map of the Köppen-Geiger climate classification. *Hydrology and Earth System Science* **11**: 1633–1644.
- Putty MRY, Prasad R. 2000. Runoff processes in headwater catchments—an experimental study in Western Ghats, South India. *Journal of Hydrology* **235**: 63–71.
- Radulovich R, Solorzano E, Sollins P. 1989. Soil macropore size distribution from water breakthrough curves. *Soil Science Society of America Journal* **53**: 556–559.
- Reichenberger S, Amelung W, Laabs V, Pinto A, Totsche KU, Zech W. 2002. Pesticide displacement along preferential flow pathways in a Brazilian Oxisol. *Geoderma* **110**: 63–68.
- Sayer A. 2002. Soil pipe dynamics and morphology in primary rainforest, Malaysian Borneo: a tentative study. *Swansea Geographer* **37**: 75–81.
- Sayer A. 2005. *The hydrogeomorphological role of soil pipes in small tropical rainforest catchments, Sabah, Malaysia*. PhD thesis. University of Wales Swansea, Swansea, 316.
- Sayer AM, Walsh RPD, Bidin K. 2004. Pipeflow as a storm runoff process in small catchments in Sabah, Malaysia. In *Proceedings of the Second Asia Pacific Association of Hydrology and Water Resources Conference*, Singapore; 160–167.
- Sayer AM, Walsh RPD, Bidin K. 2006. Pipeflow suspended sediment dynamics and their contribution to stream sediment budgets in small rainforest catchments, Sabah, Malaysia. *Forest Ecology and Management* **224**: 119–130.
- Schack-Kirchner H, Fenner PT, Hildebrand E. 2007. Different responses in bulk density and saturated hydraulic conductivity to soil deformation by logging machinery on a Ferralsol under native forest. *Soil Use and Management* **23**: 286–293.
- Scherrer S, Naef F, Faeh AO, Cordery I. 2007. Formation of runoff at the hillslope scale during intense precipitation. *Hydrology and Earth System Science* **11**: 907–922.
- Schwartz RC, Juo ASR, McInnes KJ, Cervantes CE. 1998. Anion transport in a fine-textured Ultisol in Costa Rica. In *Proceedings of the 16th World Congress of Soil Science*, Montpellier, France, 20–26 August, 1998; 1–11.
- Scully NJ. 1942. Root distribution and environment in a maple-oak forest. *Botanical Gazette* **103**: 492–517.
- Sherlock M. 1997. Plot-scale hydrometric and tracer characterisation of soil water flow in two tropical rain forest catchments in Southeast Asia. PhD thesis. Lancaster University, Lancaster, 346.
- Sidle RC, Noguchi S, Tsuboyama Y, Laursen K. 2001. A conceptual model of preferential flow systems in forested hillslopes: evidence of self-organization. *Hydrological Processes* **15**: 1675–1692.
- Sinun W. 1991. Hillslope hydrology, hydrogeomorphology and hydrochemistry of an equatorial lowland rainforest, Danum Valley, Sabah, Malaysia. MSc thesis. University of Manchester, Manchester.
- Smart PL, Wilson CM. 1984. Two methods for the tracing of pipe flow on hillslopes. *Catena* **11**: 159–168.
- Soil Survey Staff. 1999. *Soil Taxonomy: A Basic System of Soil Classification for Making and Interpreting Soil Surveys*, 2nd edn. US Government Printing Office: Washington, DC; 436.
- Spaans EJA, Baltissen GAM, Bouma J, Miedeme R, Lansu ALE, Schoonderbeek D, Wielemaker WG. 1989. Changes in physical properties of young and old volcanic surface soils in Costa Rica after clearing of tropical rain forest. *Hydrological Processes* **3**: 383–392.
- Stocking MA. 1981. Model of piping in soils. *Transactions, Japanese Geomorphological Union* **2**: 263–278.
- Suhaimi J. in prep. *Moisture control on evapotranspiration and streamflow generation in tropical forest*. PhD thesis. Lancaster University, Lancaster.
- Terzaghi K, Peck RB. 1948. *Soil Mechanics in Engineering Practice*. John Wiley and Sons: New York; 566.

- Thomas MF. 1968. Some Outstanding Problems in the Interpretation of the Geomorphology of Tropical Shields, Occasional Paper 5. British Geomorphological Research Group: Nottingham; 41–49.
- Uchida T, Kosugi K, Mizuyama T. 1999. Runoff characteristics of pipeflow and effects of pipeflow on rainfall–runoff phenomena in a mountainous watershed. *Journal of Hydrology* **222**: 18–36.
- Uchida T, Tromp-van Meerveld HJ, McDonnell JJ. 2005. The role of lateral pipe flow in hillslope runoff response: an intercomparison of non-linear hillslope response. *Journal of Hydrology* **311**: 117–133.
- van Noordwijk M, Widiyanto HM, Hairiah K. 1991. Old tree root channels in acid soils in the humid tropics: important for crop root penetration, water infiltration and nitrogen management. *Plant and Soil* **134**: 37–44.
- Vieira BC, Fernandes NF. 2004. Landslides in Rio de Janeiro: the role played by variations in soil hydraulic conductivity. *Hydrological Processes* **18**: 791–805.
- Walsh RPD. 1980. Runoff processes and models in the humid tropics. *Zeitschrift für Geomorphologie* **36**: 176–202.
- Walsh RPD, Howells KA. 1988. Soil pipes and their role in runoff generation and chemical denudation in a humid tropical catchment in Dominica. *Earth Surface Processes and Landforms* **13**: 9–17.
- Williams PW. 1985. Subcutaneous hydrology and the development of doline and cockpit karst. *Zeitschrift für Geomorphologie* **29**: 463–482.
- Wollschläger U, Pfaff T, Roth K. 2009. Field scale effective hydraulic parameterisation obtained from TDR time series and inverse modelling. *Hydrology and Earth System Science Discussion* **6**: 1489–1522.
- Yasuhara M. 1980. Streamflow generation in a small forested watershed. MSc thesis. University of Tsukuba, Tsukuba, 55.
- Zhao J. 1998. Rock mass hydraulic conductivity of the Bukit Timah granite, Singapore. *Engineering Geology* **50**: 211–216.
- Ziegler AD, Negishi JN, Sidle RC, Noguchi S, Nik AR. 2006. Impacts of logging disturbance on hillslope saturated hydraulic conductivity in a tropical forest in Peninsular Malaysia. *Catena* **67**: 89–104.
- Ziemer RR. 1992. Effect of logging on subsurface pipeflow and erosion: coastal northern California, USA. In *Erosion, Debris Flows and Environment in Mountain Regions*, Proceedings of the Chengdu Symposium, July 1992, Shamir U, Jiaqi C (eds). IAHS Publication No. 209, IAHS Press: Wallingford; 187–197.