

SoilTemp: a global database of near-surface temperature

Running title – SoilTemp: call for data

Jonas J. Lembrechts(1), Juha Aalto(2,3), Michael B. Ashcroft(4,5), Pieter De Frenne(6), Martin Kopecký(7,8), Jonathan Lenoir(9), Miska Luoto(3), Ilya M. D. Maclean(10), Olivier Roupsard(11,12), Eduardo Fuentes-Lillo(13,14,15,1), Rafael A. García(13,14), Loïc Pellissier(16,17), Camille Pitteloud(16,17), Juha M. Alatalo(18,19), Stuart W. Smith(20,21), Robert G. Björk(22,23), Lena Muffler(24,25), Simone Cesarz(26,27), Felix Gottschall(26,27), Amanda Ratier Backes(28,26), Joseph Okello(29,30), Josef Urban(31,32), Roman Plichta(31), Martin Svátek(31), Shyam S. Phartyal(33,34), Sonja Wipf(35,36), Nico Eisenhauer(26,27), Mihai Puşcaş(37), Pavel Dan Turtureanu(38), Andrej Varlagin(39), Romina D. Dimarco(40), Alistair S. Jump(41), Krystal Randall(42), Ellen Dorrepaal(43), Keith Larson(43), Josefina Walz(43), Luca Vitale(44), Miroslav Svoboda(8), Rebecca Finger Higgens(45), Aud H. Halbritter(46), Salvatore R. Curasi(47), Ian Klupar(47), Austin Koontz(48), William D. Pearse(48,49), Elizabeth Simpson(48), Michael Stemkovski(48), Bente Jessen Graae(20), Mia Vedel Sørensen(20), Toke T. Høye(50), M. Rosa Fernández Calzado(51), Juan Lorite(51), Michele Carbognani(52), Marcello Tomaselli(52), T'ai G. W. Forte(52), Alessandro Petraglia(52), Stef Haesen(53), Ben Somers(53), Koenraad Van Meerbeek(53), Mats P. Björkman(22,23), Kristoffer Hylander(54), Sonia Merinero(55), Mana Gharun(56), Nina Buchmann(56), Jiri Dolezal(7,57), Radim Matula(8), Andrew D. Thomas(58), Joseph J. Bailey(59), Dany Ghosn(60), George Kazakis(60), Miguel Angel de Pablo(61), Julia Kempainen(3), Pekka Niittynen(3), Lisa Rew(62), Tim Seipel(62), Christian Larson(62), James D. M. Speed(63), Jonas Ardö(64), Nicoletta Cannone(65), Mauro Guglielmin(66), Francesco Malfasi(66), Maaike Y. Bader(67), Rafaella Canessa(67), Angela Stanisci(68), Juergen Kreyling(24), Jonas Schmeddes(24), Laurenz Teuber(24), Valeria Aschero(69,70), Marek Čiliak(71), František Máliš(72), Pallieter De Smedt(6), Sanne Govaert(6), Camille Meeussen(6), Pieter Vangansbeke(6), Khatuna Gigauri(73), Andrea Lamprecht(74), Harald Pauli(74), Klaus Steinbauer(74), Manuela Winkler(74), Masahito Ueyama(75), Martin A. Nuñez(76), Tudor-Mihai Ursu(77), Sylvia Haider(28,26), Ronja E. M. Wedegärtner(20), Marko Smiljanic(78), Mario Trouillier(78), Martin Wilmking(78), Jan Altman(7), Josef Brůna(7), Lucia Hederová(7), Martin Macek(7), Matěj Man (7), Jan Wild(7), Pascal Vittoz(79), Meelis Pärtel(80), Peter Barančok(81), Róbert Kanka(81), Jozef Kollár(81), Andrej Palaj(81), Agustina Barros(70), Ana Clara Mazzolari(70), Marijn Bauters(29), Pascal Boeckx(29), José Luis Benito Alonso(82), Shengwei Zong(83), Valter Di Cecco(84), Zuzana Sitková(85), Katja Tielbörger(86), Liesbeth van den Brink(86), Robert Weigel(25), Jürgen Homeier(25), C. Johan Dahlberg(54,87), Sergiy Medinets(88), Volodymyr Medinets(88), Hans J. De Boeck(89), Miguel Portillo-Estrada(1), Lore T. Verryckt(1), Ann Milbau(90), Gergana N. Daskalova(91), Haydn J. D. Thomas(91), Isla H. Myers-Smith(91), Benjamin Blonder(92,93), Jörg G. Stephan(94), Patrice Descombes(16,17,95), Florian Zellweger(95), Esther R. Frei(95,35), Bernard Heinesch(96), Christopher Andrews(97), Jan Dick(97), Lukas Siebicke(98), Adrian Rocha(99), Rebecca A. Senior(100), Christian Rixen(35), Juan J. Jimenez(101), Julia Boike(102,103), Aníbal Pauchard(13,14), Thomas Scholten(104), Brett Scheffers(105), David Klinges(106), Edmund W. Basham(106), Jian Zhang(107), Zhaochen Zhang(107), Charly Géron(108), Fatih Fazlioglu(109), Onur Candan(109), Jhonatan Sallo Bravo(110), Filip Hrbacek(111), Kamil Laska(111), Edoardo Cremonese(112), Peter Haase(113,114), Fernando E. Moyano(98), Christian Rossi(115,116,36), Ivan Nijs(1)

Author contributions: JLL performed the analyses and wrote the manuscript , JLL, JA, MBA, PDF, MK, JL, ML, IMDM and IN lead the consortium and contributed to the writing; all authors contribute to the consortium and provided editorial advice.

*Corresponding author, OrcID = <https://orcid.org/0000-0002-1933-0750>,
Jonas.lembrechts@uantwerpen.be, +3232651727

OrcIDs (alphabetically ordered)

Juha Aalto <https://orcid.org/0000-0001-6819-4911>
Juha M. Alatalo <https://orcid.org/0000-0001-5084-850X>
Jan Altman <https://orcid.org/0000-0003-4879-5773>
Jonas Ardö <https://orcid.org/0000-0002-9318-0973>
Valeria Aschero <https://orcid.org/0000-0003-3865-4133>
Maaïke Y Bader <http://orcid.org/0000-0003-4300-7598>
Peter Barančok <https://orcid.org/0000-0003-1171-2524>
Edmund Basham <https://orcid.org/0000-0002-0167-7908>
José-Luis Benito-Alonso <https://orcid.org/0000-0003-1086-8834>
Robert G. Björk <https://orcid.org/0000-0001-7346-666X>
Mats P. Björkman <https://orcid.org/0000-0001-5768-1976>
Julia Boike <https://orcid.org/0000-0002-5875-2112>
Josef Brůna <https://orcid.org/0000-0002-4839-4593>
Nina Buchmann <https://orcid.org/0000-0003-0826-2980>
Onur Candan <https://orcid.org/0000-0002-9254-4122>
Rafaella Canessa <https://orcid.org/0000-0002-6979-9880>
Michele Carbognani <https://orcid.org/0000-0001-7701-9859>
Marek Čiliak <https://orcid.org/0000-0002-6720-9365>
Edoardo Cremonese <https://orcid.org/0000-0002-6708-8532>
Salvatore R. Curasi: <https://orcid.org/0000-0002-4534-3344>
C. Johan Dahlberg <https://orcid.org/0000-0003-0271-3306>
Gergana Daskalova <https://orcid.org/0000-0002-5674-5322>
Miguel Ángel de Pablo Hernández <https://orcid.org/0000-0002-4496-2741>
Pallier De Smedt <https://orcid.org/0000-0002-3073-6751>
Jiri Dolezal <https://orcid.org/0000-0002-5829-4051>
Nico Eisenhauer <https://orcid.org/0000-0002-0371-6720>
Fatih Fazlioglu <https://orcid.org/0000-0002-4723-3640>
T'ai G. W. Forte <https://orcid.org/0000-0002-8685-5872>
Esther R. Frei <https://orcid.org/0000-0003-1910-7900>
Charly Géron: <https://orcid.org/0000-0001-7912-4708>
Mana Gharun <https://orcid.org/0000-0003-0337-7367>
Dany Ghosn <https://orcid.org/0000-0003-1898-9681>
Felix Gottschall <https://orcid.org/0000-0002-1247-8728>
Sanne Govaert <https://orcid.org/0000-0002-8939-1305>
Peter Haase <https://orcid.org/0000-0002-9340-0438>
Stef Haesen <https://orcid.org/0000-0002-4491-4213>
Sylvia Haider <https://orcid.org/0000-0002-2966-0534>
Bernard Heinesch <https://orcid.org/0000-0001-7594-6341>
Filip Hrbacek <https://orcid.org/0000-0001-5032-9216>
Juan J. Jiménez <https://orcid.org/0000-0003-2398-0796>
Alistair S. Jump <https://orcid.org/0000-0002-2167-6451>
Róbert Kanka <https://orcid.org/0000-0002-7071-7280>
Julia Kemppinen <https://orcid.org/0000-0001-7521-7229>
Austin Koontz <https://orcid.org/0000-0002-6103-5894>
Andrea Lamprecht <https://orcid.org/0000-0002-8719-026X>
Christian Larson <https://orcid.org/0000-0002-7567-4953>
Kamil Laska <https://orcid.org/0000-0002-5199-9737>
Jonathan Lenoir <http://orcid.org/0000-0003-0638-9582>
Juan Lorite <https://orcid.org/0000-0003-4617-8069>
František Máliš <https://orcid.org/0000-0003-2760-6988>
Matěj Man <https://orcid.org/0000-0002-4557-8768>
Sergiy Medinets <http://orcid.org/0000-0001-5980-1054>
Volodymyr Medinets <https://orcid.org/0000-0001-7543-7504>
Camille Meeussen <https://orcid.org/0000-0002-5869-4936>
Ann Milbau <https://orcid.org/0000-0003-3555-8883>
Fernando E. Moyano <https://orcid.org/0000-0002-4090-5838>
Isla Myers-Smith <https://orcid.org/0000-0002-8417-6112>
Pekka Niittynen <https://orcid.org/0000-0002-7290-029X>
Ivan Nijs <https://orcid.org/0000-0003-3111-680X>
Andrej Palaj <https://orcid.org/0000-0001-7054-4183>
Harald Pauli <https://orcid.org/0000-0002-9842-9934>
William D. Pearse <https://orcid.org/0000-0002-6241-3164>
Shyam S. Phartyal <https://orcid.org/0000-0003-3266-6619>
Mihai Pușcaș <https://orcid.org/0000-0002-2632-640X>
Lisa Rew <https://orcid.org/0000-0002-2818-3991>
Christian Rossi <https://orcid.org/0000-0001-9983-8898>
Olivier Roupsard <http://orcid.org/0000-0002-1319-142X>
Jhonatan Sallo-Bravo <https://orcid.org/0000-0001-9007-4959>
Thomas Scholten <https://orcid.org/0000-0002-4875-2602>
Rebecca A. Senior <https://orcid.org/0000-0002-8208-736X>
Zuzana Sitková <https://orcid.org/0000-0001-6354-6105>
Stuart W. Smith <https://orcid.org/0000-0001-9396-6610>
Ben Somers <https://orcid.org/0000-0002-7875-107X>
James D. M. Speed <http://orcid.org/0000-0002-0633-5595>
Klaus Steinbauer <https://orcid.org/0000-0002-3730-9920>
Jörg G. Stephan <http://orcid.org/0000-0001-6195-7867>
Martin Svátek <https://orcid.org/0000-0003-2328-4627>
Miroslav Svoboda <https://orcid.org/0000-0003-4050-3422>
Andrew Thomas <https://orcid.org/0000-0002-1360-1687>
Haydn Thomas <https://orcid.org/0000-0001-9099-6304>
Marcello Tomaselli <https://orcid.org/0000-0003-4208-3433>
Pavel Dan Turtureanu <https://orcid.org/0000-0002-7422-3106>
Masahito Ueyama <https://orcid.org/0000-0002-4000-4888>
Josef Urban <https://orcid.org/0000-0003-1730-947X>
Tudor-Mihai Ursu <https://orcid.org/0000-0002-4898-6345>
Liesbeth van den Brink <https://orcid.org/0000-0003-0313-8147>
Pieter Vangansbeke <https://orcid.org/0000-0002-6356-2858>
Andrej Varlagin <https://orcid.org/0000-0002-2549-5236>
Koenraad Van Meerbeek <https://orcid.org/0000-0002-9260-3815>
Lore T. Verryckt <https://orcid.org/0000-0002-9452-5216>
Pascal Vittoz <https://orcid.org/0000-0003-4218-4517>
Ronja E. M. Wedegärtner <https://orcid.org/0000-0003-4633-755X>
Jan Wild <https://orcid.org/0000-0003-3007-4070>
Martin Wilmking <https://orcid.org/0000-0003-4964-2402>
Manuela Winkler <http://orcid.org/0000-0002-8655-9555>
Sonja Wipf <http://orcid.org/0000-0002-3492-1399>
Florian Zellweger <https://orcid.org/0000-0003-1265-9147>
Jian Zhang <https://orcid.org/0000-0003-0589-6267>

1) Research Group PLECO (Plants and Ecosystems), University of Antwerp, 2610 Wilrijk, Belgium, 2) Finnish Meteorological Inst., P.O. Box 503, FI-00101 Helsinki, Finland, 3) Dept of Geosciences and Geography, Gustaf Hällströmin katu 2a, FIN-00014 Univ. of Helsinki, Finland, 4) Centre for Sustainable Ecosystem Solutions, School of Biological Sciences, University of Wollongong, Wollongong, Australia, 5) Australian Museum, Sydney, Australia, 6) Forest & Nature Lab, Department of Environment, Ghent University, Geraardsbergsesteenweg 267, 9090 Melle-Gontrode, Belgium, 7) Institute of Botany of the Czech Academy of Sciences, Zámek 1, CZ-25243, Průhonice, Czech Republic, 8) Faculty of Forestry and Wood Sciences, Czech University of Life Sciences Prague, Kamýcká 129, CZ-165 21, Prague 6 - Suchbát, Czech Republic, 9) UR 'Ecologie et Dynamique des Systèmes Anthropisés' (EDYSAN, UMR 7058 CNRS-UPJV), Univ. de Picardie Jules Verne, Amiens, France, 10) Environment and Sustainability Institute, University of Exeter, Penryn Campus, Penryn, UK, TR10 9FE, 11) CIRAD, UMR Eco&Sols, B.P. 1386, CP 18524, Dakar, Senegal, 12) Eco&Sols, Univ Montpellier, CIRAD, INRAE, IRD, Institut Agro, Montpellier, France, 13) Laboratorio de Invasiones Biológicas (LIB), Facultad de Ciencias Forestales, Universidad de Concepción, Concepción, Chile, 14) Instituto de Ecología y Biodiversidad (IEB), Santiago, Chile, 15) School of Education and Social Sciences, Adventist University of Chile, Chile, 16) Landscape Ecology, Institute of Terrestrial Ecosystems, Department of Environmental Systems Science, ETH Zürich, 8092 Zürich, Switzerland, 17) Unit of Land Change Science, Swiss Federal Research Institute WSL, 8903 Birmensdorf, Switzerland, 18) Department of Biological and Environmental Sciences, Qatar University, Doha, Qatar, 19) Environmental Science Center, Qatar University, Doha, Qatar, 20) Department of Biology, Norwegian University of Science and Technology, 7491 Trondheim, Norway, 21) Asian School of Environment, Nanyang Technological University, 42 Nanyang Ave, Singapore 639815, Singapore, 22) Department of Earth Sciences, University of Gothenburg, P.O. Box 460, SE-40530 Gothenburg, Sweden, 23) Gothenburg Global Biodiversity Centre, P.O. Box 461, SE-405 30 Gothenburg, Sweden, 24) Experimental Plant Ecology, Institute of Botany and Landscape Ecology, University of Greifswald, D-17487 Greifswald, Germany, 25) Plant Ecology, Albrecht-von-Haller-Institute for Plant Sciences, University of Goettingen, 37073 Goettingen, Germany, 26) German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Leipzig, Germany, 27) Institute of Biology, Leipzig University, Leipzig, Germany, 28) Institute of Biology / Geobotany and Botanical Garden, Martin Luther University Halle-Wittenberg, Halle (Saale), Germany, 29) Isotope Bioscience Laboratory - ISOFYS, Ghent University, Coupure Links 653, 9000 Gent, Belgium, 30) Mountains of the Moon University, P.O Box 837, Fort Portal, Uganda, 31) Department of Forest Botany, Dendrology and Geobiocoenology, Mendel University in Brno, Czech Republic, 32) Siberian Federal University, Krasnoyarsk, Russia, 33) School of Ecology and Environment Studies, Nalanda University, Rajgir, India, 34) Department of Forestry and NR, H.N.B. Garhwal University, Srinagar-Garhwal, India, 35) WSL Institute for Snow and Avalanche Research SLF, 7260 Davos, Switzerland, 36) Swiss National Park, Chastè Planta-Wildenberg, 7530 Zernez, Switzerland, 37) A. Borza Botanical Garden and Department of Taxonomy and Ecology, Faculty of Biology and Geology, Babeş-Bolyai University, Cluj-Napoca, Romania, 38) A. Borza Botanical Garden, Babeş-Bolyai University, Cluj-Napoca, Romania, 39) A.N. Severtsov Institute of Ecology and Evolution, Russian Academy of Sciences, 119071, Leninsky pr.33, Moscow, Russia, 40) Grupo de Ecología de Poblaciones de Insectos, IFAB (INTA - CONICET), Isla Victoria 4450, Bariloche, Argentina, 41) Biological and Environmental Sciences, Faculty of Natural Sciences, University of Stirling, Scotland, FK9 4LA, 42) Centre for Sustainable Ecosystem Solutions, School of Earth, Atmospheric and Life Sciences, University of Wollongong, Wollongong, New South Wales, 2522, Australia, 43) Climate Impacts Research Centre, Department of Ecology and Environmental Sciences, Umeå University, Abisko, Sweden, 44) CNR - Institute for mediterranean Agricultural and Forest Systems, Via Patacca 85, ercolano (napoli), Italy, 45) Dartmouth College, Hanover, NH, USA, 46) Department of Biological Sciences and Bjerknes Centre for Climate Research, University of Bergen, N-5020 Bergen, Norway, 47) Department of Biological Sciences, University of Notre Dame, Notre Dame, IN 46556, USA, 48) Department of Biology and Ecology Center, Utah State University, 5305 Old Main Hill, Logan, UT 84322, USA, 49) Department of Life Sciences, Imperial College, Silwood Park Campus, Ascot, Berkshire SL5 7PY, UK, 50) Department of Bioscience and Arctic Research Centre, Grenåvej 14, 8410 Rønne, Denmark, 51) Department of Botany, University of Granada, 18071, Granada, Spain, 52) Department of Chemistry, Life Sciences and Environmental Sustainability, University of Parma, Parco Area delle Scienze 11/A, 43124 Parma, Italy, 53) Department of Earth and Environmental Sciences, Celestijnenlaan 200E, 3001 Leuven, Belgium, 54) Department of Ecology, Environment and Plant Sciences and Bolin Centre for Climate Research, Stockholm University, 106 91 Stockholm, Sweden, 55) Department of Ecology, Environment and Plant Sciences, Stockholm University, SE-106 91 Stockholm, Sweden, 56) Department of Environmental Systems Science, ETH Zurich, Universitaetstrasse 2, 8092 Zurich, Switzerland, 57) Faculty of Science, Department of Botany, University of South Bohemia, Na Zlaté Stoece 1, 37005 České Budějovice, Czech Republic, 58) Department of Geography and Earth Sciences, Aberystwyth University, Wales, UK, 59) Department of Geography, York St John University, Lord Mayor's Walk, York, YO31 7EX, United Kingdom, 60) Department of Geo-information in Environmental Management, Mediterranean Agronomic Institute of Chania, PO Box 85, 73100 Chania, Greece, 61) Department of Geology, Geography and Environment. University of Alcalá. 28805 Alcalá de Henares, Madrid, Spain., 62) Department of Land Resources and Environmental Sciences, Montana State University, Bozeman MT, USA, 59717, 63) Department of Natural History, NTNU University Museum, Norwegian University of Science and Technology, NO-7491 Trondheim Norway, 64) Department of Physical Geography and Ecosystem Science, Lund University, Sölvegatan 12 223 62 Lund Sweden, 65) Department of Science and High Technology, Insubria University, Via Valleggio 11, 22100 Como, Italy, 66) Department of Theoretical and Applied Sciences, Insubria University, Via Dunant 3, 21100 Varese, Italy, 67) Ecological Plant Geography, Faculty of Geography, University of Marburg, Deutschhausstr. 10, 35032, Marburg, Germany, 68) EnvixLab, Dipartimento di Bioscienze e Territorio, Università degli Studi del Molise, Via Duca degli Abruzzi s.n.c., 86039 Termoli, Italy, 69) Facultad de Ciencias Exactas y Naturales, Universidad Nacional de Cuyo, 70) Instituto Argentino de Nivología, Glaciología y Ciencias Ambientales (IANIGLA), CONICET, CCT-Mendoza, 71) Faculty of Ecology and Environmental Sciences, Technical University in Zvolen, T.G.Masaryka 24, 960 01

Zvolen, Slovakia, 72) Faculty of Forestry, Technical University in Zvolen, T.G.Masaryka 24, 960 01 Zvolen, Slovakia, 73) Georgian Institute of Public Affairs, Tbilisi, Georgia, 74) GLORIA Coordination, Institute for Interdisciplinary Mountain Research, Austrian Academy of Sciences (ÖAW) & Department of Integrative Biology and Biodiversity Research, University of Natural Resources and Life Sciences Vienna (BOKU), Silbergasse 30/3, 1190 Vienna, Austria, 75) Graduate School of Life and Environmental Sciences, Osaka Prefecture University, 599-8531, Japan, 76) Grupo de Ecología de Invasiones, INIBIOMA, CONICET/ Universidad Nacional del Comahue, Av. de los Pioneros 2350, Bariloche 8400, Argentina, 77) Institute of Biological Research Cluj-Napoca, National Institute of Research and Development for Biological Sciences, Bucharest, Romania, 78) Institute of Botany and Landscape Ecology, University Greifswald, D-17487 Greifswald, Germany, 79) Institute of Earth Surface Dynamics, Faculty of Geosciences and Environment, University of Lausanne, Géopolis, 1015 Lausanne, Switzerland, 80) Institute of Ecology and Earth Sciences, University of Tartu, Lai 40, Tartu 51005, Estonia, 81) Institute of Landscape Ecology Slovak Academy of Sciences, Štefánikova 3, 81499 Bratislava, Slovakia, 82) Jolube Consultor Botánico. C/Mariano R de Ledesma, 4. E-22700 Jaca, Huesca, SPAIN, 83) Key Laboratory of Geographical Processes and Ecological Security in Changbai Mountains, Ministry of Education, School of Geographical Sciences, Northeast Normal University, Changchun 130024, China, 84) Majella Seed Bank, Majella National Park, Colle Madonna, 66010 Lama dei Peligni, Italy, 85) National Forest Centre, Forest Research Institute Zvolen, T. G. Masaryka 22, 96001 Zvolen, Slovakia, 86) Plant Ecology Group, Department of Evolution and Ecology, University of Tübingen, Tübingen, Germany, 87) the County Administrative Board of Västra Götaland, SE-403 40 Gothenburg, Sweden, 88) Regional Centre for Integrated Environmental Monitoring, Odesa National I.I. Mechnikov University, 7 Mayakovskogo lane, 65082 Odesa, Ukraine, 89) Research Group Plants and Ecosystems (PLECO), University of Antwerp, 2610 Wilrijk, Belgium, 90) Research Institute for Nature and Forest (INBO), Havenlaan 88, bus 73, 1000 Brussel, Belgium, 91) School of GeoSciences, University of Edinburgh, King's Buildings, Edinburgh, EH9 3FF, United Kingdom, 92) School of Life Sciences, Arizona State University, Tempe, AZ, USA, 93) Department of Environmental Science, Policy, and Management, University of California, Berkeley, CA 94720 USA, 94) Swedish University of Agricultural Sciences, Swedish Species Information Centre, Almas allé 8 E, 75651 Uppsala, Sweden, 95) Swiss Federal Research Institute WSL, 8903 Birmensdorf, Switzerland, 96) TERRA Teaching and Research Center, Faculty of Gembloux Agro-Bio Tech, University of Liege, Passage des déportés, 2, 5030 Gembloux, Belgium, 97) UK Centre for Ecology & Hydrology, Bush Estate, Penicuik, Midlothian, EH26 0QB, United Kingdom, 98) University of Goettingen, Bioclimatology, Büsngenweg 2, 37077 Göttingen, Germany., 99) University of Notre Dame, Department of Biological Sciences and the Environmental Change Initiative, 100) Woodrow Wilson School of Public and International Affairs, Princeton University, Princeton, NJ 08540, USA, 101) ARAID Research and Development, Zaragoza, Spain, 102) Alfred Wegener Institute Helmholtz Center for Polar and Marine Research, Telegrafenberg A45, 14473 Potsdam, Germany, 103) Geography Department, Humboldt-Universität zu Berlin, Germany, 104) Chair of Soil Science and Geomorphology, Department of Geosciences, University of Tuebingen, 72070 Tuebingen, Germany, 105) Department of Wildlife Ecology and Conservation, University of Florida, Gainesville, FL 32611, USA, 106) School of Natural Resources and Environment, University of Florida, Gainesville, FL 32611, USA, 107) Zhejiang Tiantong Forest Ecosystem National Observation and Research Station, School of Ecological and Environmental Sciences, East China Normal University, Shanghai 200241, China, 108) Biodiversity and Landscape, TERRA research centre, Gembloux Agro-Bio Tech, University of Liège, Gembloux, 5032, Belgium ; Research Group PLECO (Plants and Ecosystems), University of Antwerp, 2610 Wilrijk, Belgium, 109) Faculty of Arts and Sciences, Department of Molecular Biology and Genetics, Ordu University, 52200, Ordu, Turkey, 110) Universidad Nacional de San Antonio Abad del Cusco, Cusco, Peru, 111) Department of Geography, Masaryk University, Brno, Czech Republic, 112) Climate Change Unit, Environmental Protection Agency of Aosta Valley, Sain Christophe, Aosta, Italy, 113) Senckenberg Research Institute and Natural History Museum Frankfurt, 63571 Gelnhausen, Germany, 114) Faculty of Biology, University of Duisburg-Essen, 45141 Essen, Germany, 115) Remote Sensing Laboratories, Dept. of Geography, University of Zurich, Winterthurerstrasse 190, 8057 Zurich, Switzerland, 116) Research Unit Community Ecology, Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Zürcherstrasse 111, 8903 Birmensdorf, Switzerland

1 **Abstract**

2 Current analyses and predictions of spatially-explicit patterns and processes in ecology most often rely
3 on climate data interpolated from standardized weather stations. This interpolated climate data
4 represents long-term average thermal conditions at coarse spatial resolutions only. Hence, many
5 climate-forcing factors that operate at fine spatiotemporal resolutions are overlooked. This is
6 particularly important in relation to effects of observation height (e.g. vegetation, snow and soil
7 characteristics) and in habitats varying in their exposure to radiation, moisture and wind (e.g.
8 topography, radiative forcing, or cold-air pooling). Since organisms living close to the ground relate
9 more strongly to these microclimatic conditions than to free-air temperatures, microclimatic ground
10 and near-surface data are needed to provide realistic forecasts of the fate of such organisms under
11 anthropogenic climate change, as well as of the functioning of the ecosystems they live in.

12 To fill this critical gap, we highlight a call for temperature time series submissions to SoilTemp, a
13 geospatial database initiative compiling soil and near-surface temperature data from all over the
14 world. Currently this database contains time series from 7538 temperature sensors from 51 countries
15 across all key biomes. The database will pave the way towards an improved global understanding of
16 microclimate and bridge the gap between the available climate data and the climate at fine
17 spatiotemporal resolutions relevant to most organisms and ecosystem processes.

18 **Keywords:** microclimate, soil climate, climate change, topoclimate, database, temperature, species
19 distributions, ecosystem processes

20 **Introduction**

21 Current ecological research increasingly deals with large-scale patterns and processes, with global
22 databases of species distributions and traits becoming increasingly available (Bruehlheide *et al.*, 2018,
23 Kissling *et al.*, 2018, Kattge *et al.*, 2019). Analyses of these patterns and processes – and their
24 predictions under anthropogenic climate change – often rely on global climatic grids at coarse spatial
25 resolutions interpolated from standardized weather stations that represent long-term average
26 atmospheric conditions (Lembrechts *et al.*, 2018). Moreover, sensors in these weather stations are
27 shielded from direct solar radiation and located at ~2 meters above a frequently mown lawn (free-air
28 temperature or 'macroclimate', Jarraud, 2008). These climatic grids thus ignore many climate-forcing
29 processes that operate near the ground surface, at fine spatiotemporal resolutions, and in
30 environments that vary in their exposure to winds, radiation and moisture ('microclimate', Daly, 2006,
31 Bramer *et al.*, 2018, Körner & Hiltbrunner, 2018). Importantly, while these microclimatic processes
32 often operate at fine spatiotemporal resolutions, they can affect ecological relations both at the local
33 and the global scale (De Frenne *et al.*, 2013, Ashcroft *et al.*, 2014, Lembrechts *et al.*, 2019). For
34 example, they can potentially protect ground-dwelling biota against long-term climate variability,
35 providing microrefugia for these species to survive in locations deemed unsuitable in models using
36 climate data at coarse spatial resolutions, or buffer organisms against short-term extreme events (De
37 Frenne *et al.*, 2013, Lenoir *et al.*, 2017, Bramer *et al.*, 2018, Suggitt *et al.*, 2018). Microclimates can
38 however also expose organisms to more extreme temperatures, in which case distribution models
39 that ignore such microclimates may erroneously predict species survival instead of extinction
40 (Pincebourde & Casas, 2019). In order to provide realistic forecasts of species distributions and
41 performance, as well as of the functioning of the ecosystems they operate in, climate data that
42 incorporates microclimatic processes, ideally measured *in-situ*, are thus urgently needed (Körner &
43 Hiltbrunner, 2018).

44 **Horizontal and vertical features driving microclimate**

45 The offset between micro- and macroclimate is particularly pronounced around the soil surface, as
46 temperatures measured at 2 m above the ground can differ substantially from those at ground level,
47 or in the layers just above and below it (Geiger, 1950, Lembrechts *et al.*, 2019). This offset can result
48 from both 'horizontal' and 'vertical' features (Fig. 1), and can exceed several degrees centigrade in
49 annual averages. For example, Kearney (2019) modelled coarse-scale soil temperatures at various
50 depths considering the vertical features affecting the radiation balance. These vertical features include
51 the effects of vegetation characteristics (e.g. structure and cover), snow cover and soil characteristics
52 (e.g. moisture content, geological types, texture and bulk density) (Li, 1926, Zhang *et al.*, 2008,

53 Lembrechts *et al.*, 2019). The result of these vertical features is not only an instantaneous temperature
54 offset between air and soil temperatures, but also a buffering effect, i.e. the temporal variability in
55 temperature changes is lower in the soil than in the air (Geiger, 1950, Ashcroft & Gollan, 2013).
56 Horizontal processes on the other hand relate more to the spatial resolution of the climatic data. They
57 can be broken up into those that require only fine-resolution environmental information for specific
58 sites (e.g. effects of slope and aspect on radiation balances; Bennie *et al.*, 2008), and those where
59 temperatures are also affected by neighboring locations (e.g. topographic shading, cold-air drainage
60 and atmospheric temperature inversions, which are landscape context dependent; Whiteman, 1982,
61 Ashcroft & Gollan, 2012).

62 How horizontal and vertical features interact to define differences between soil and air temperature
63 may differ with the biome, season and day time. For example, in grasslands during summer, incoming
64 short-wave solar radiation is usually the dominant factor determining daytime soil surface
65 temperatures, which in turn result in higher air temperatures through convective heating (Geiger,
66 1950). However, during winter, horizontal processes such as cold-air drainage and coastal buffering
67 can have larger effects, especially on overnight air temperatures, when air temperatures may be
68 driving soil temperatures rather than vice-versa (Vitasse *et al.*, 2017). In dense forests, the situation is
69 even more complex: upper canopies block the bulk of short wave solar radiation, such that sub-canopy
70 temperatures are determined by convective heat transfer between the air surrounding the canopy
71 and direct conductance through physical contact of different parts of the canopy layer, in addition to
72 the limited radiation that does permeate the canopy (Körner & Paulsen, 2004, Lenoir *et al.*, 2017,
73 Zellweger *et al.*, 2019). As a result, horizontal processes such as passing fronts, and winds blowing in
74 hotter or colder air from outside the forest, will in large part define the – dampened – temperature
75 patterns under forest canopies (Ashcroft *et al.*, 2008).

76 ***The need for microclimate data across the field of ecology***

77 Many organisms living in the soil and close to the soil surface (e.g. soil micro-organisms like fungi,
78 ground arthropods, herbs, mosses, tree seedlings and small vertebrates) only experience fine-scale
79 soil and/or near-surface temperatures, and thus likely relate less strongly to free-air temperatures
80 (Randin *et al.*, 2009, Niittynen & Luoto, 2017, Lembrechts *et al.*, 2019). This may be reflected in a
81 species' distribution, but also their morphology, physiology and behavior (Körner & Paulsen, 2004,
82 Kearney *et al.*, 2009, Opedal *et al.*, 2015, de Boeck *et al.*, 2016). Many species indeed survive, live and
83 reproduce where average background climate appears unsuitable, and equally may be gone from sites
84 within apparently suitable areas where microclimatic extremes exceed their limits (Suggitt *et al.*,
85 2011). Without microclimate data, we not only lack information on the potential thermal

86 heterogeneity that is available for species to thermoregulate in situ, but also on the true magnitude
87 of climate change that species will be exposed to (Pincebourde *et al.*, 2016, Maclean *et al.*, 2017).
88 Accurately predicting how species' ranges will shift under climate change requires a good
89 understanding of the variety of climate niches truly available to them (Maclean *et al.*, 2015, Lenoir *et*
90 *al.*, 2017). The latter requires both a good understanding of what defines current microclimates, as
91 well of how climate change will interact with the drivers of microclimatic conditions (Maclean, 2019).
92 Additionally, it is the soil temperature rather than the air temperature that defines many ecosystem
93 functions in and close to the soil, like evapotranspiration, decomposition, root growth,
94 biogeochemical cycling and soil respiration (Pleim & Gilliam, 2009, Portillo-Estrada *et al.*, 2016, Hursh
95 *et al.*, 2017, Gottschall *et al.*, 2019, Medinets *et al.*, 2019). Given the repeatedly proven sensitivity of
96 many of these processes to temperatures (Rosenberg *et al.*, 1990, Coûteaux *et al.*, 1995, Schimel *et*
97 *al.*, 1996), here again having accurate measurements will be of utmost importance. The carbon
98 balance in boreal forests, for example, is largely dependent on soil thaw and thus soil rather than air
99 temperatures (Goulden *et al.*, 1998).

100 These realizations highlight the urgency to start using soil and near-surface microclimate data when
101 modelling the ecology and biogeography of surface and soil-dwelling organisms, as well as the
102 functioning of soil ecosystems, instead of readily available coarse-scaled free-air climate data (from
103 e.g. CHELSA (Karger *et al.*, 2017), TerraClimate (Abatzoglou *et al.*, 2018) or WorldClim (Fick & Hijmans,
104 2017)). While a suit of models now exist that produce fine-scale climate data (Bramer *et al.*, 2018,
105 Lembrechts *et al.*, 2018), we do not yet fully understand whether models using data that represent
106 average conditions over large areas provide adequate “mean field approximations” of (i.e. are
107 representative for) more complex spatiotemporal effects driven by the climatic conditions that
108 organisms experience (Bennie *et al.*, 2014). To accomplish the latter, global in-situ data is needed for
109 large-scale fine-resolution calibration and validation of these models. However, while the quality and
110 resolution of free-air temperature data and models at the global scale is rapidly improving (Bramer *et*
111 *al.*, 2018), soil temperature datasets used in biogeography and biogeochemistry are still largely
112 restricted to the landscape or regional scale, at best, and from intensively studied regions only
113 (Ashcroft *et al.*, 2008, Ashcroft *et al.*, 2009, Carter *et al.*, 2015, Aalto *et al.*, 2018), or they are derived
114 from models lacking fine-grained ground-truthing data (e.g. Copernicus Climate Change Service (C3S),
115 2019). Land surface temperatures as obtained from satellite data, on the other hand, are hampered
116 by their inability to measure below the vegetation cover (Bramer *et al.*, 2018).

117 In order to accurately describe and predict the (future) distribution and/or traits of surface and soil-
118 dwelling species at larger scales, we need to improve our general knowledge of the offsets and

119 spatiotemporal changes in variability between soil-level and free-air temperatures (Aalto *et al.*, 2018,
120 Lembrechts *et al.*, 2019). There is an urgent need to work towards globally available soil and near-
121 surface temperature data based on in-situ measurements and at relevant spatiotemporal resolutions
122 (Ashcroft & Gollan, 2012, Pradervand *et al.*, 2014, Slavich *et al.*, 2014, Opedal *et al.*, 2015, Meineri &
123 Hylander, 2017).

124 ***Launch of the SoilTemp database***

125 To tackle these issues, we launch an ambitious database initiative, compiling soil and near-surface
126 temperature data from all over the world into a global geospatial database: SoilTemp. At the time of
127 writing, we brought together temperature data from 7538 sensors placed both below, at and above
128 (up to 2 m) the soil surface (Fig. 2a), which is an accumulation of over 180.000 months of temperature
129 data with measurement intervals between 1 and 240 minutes (>30% every 60 minutes). The database
130 hosts loggers from 51 different countries spread across all continents, with a broad distribution across
131 the world's climatic space (Fig. 2b). There is a dominance of time series from Europe and areas below
132 1500 m a.s.l. (Fig. 2c, d). More than 75% of sensor measurements occurred within the last decade, but
133 the database does contain several time series covering longer time periods as well, with a maximum
134 of 42 years (Fig. 2d).

135 When the remaining critical gaps in our spatial coverage will be filled (see below), this database will
136 allow global assessments of the long-established theories on boundary layer climatology in
137 heterogeneous environments (Geiger, 1950), which has so far been lacking. The growing database
138 provides a unique opportunity to disentangle the role of the different horizontal and vertical features
139 influencing soil and near-surface temperature across all biomes of the world, with high spatial and
140 temporal resolutions. It will allow relating patterns in soil temperature to processes in the lower air
141 layers and calibrate and validate global models of soil temperature and (micro)climate (Kearney *et al.*,
142 2014a, Kearney *et al.*, 2014b, Carter *et al.*, 2015, Maclean *et al.*, 2017). It will also allow us to create
143 global maps of a wide array of general and microclimate-specific bioclimatic variables (e.g. growing
144 degree days, growing season length) at relevant spatiotemporal resolutions (Körner & Hiltbrunner,
145 2018).

146 Ultimately, this joint global effort and the resulting global microclimatic products will enable us to
147 improve analyses of the relationships between species' macroecology and the microclimate they
148 experience, identify microrefugia and stepping stones and improve global models of ecosystem
149 functioning and element cycling. Indeed, replacing the coarse-scaled free-air temperature averages
150 used traditionally in models in all fields of ecology with these more relevant soil-specific data products

151 is likely to increase their descriptive and predictive power, as the countless above-mentioned regional
152 studies exemplify (Lembrechts *et al.*, 2019). Additionally, this first global effort to combine and collect
153 in-situ measurements will help solve long-standing issues regarding sensor comparability and data
154 collection variability (Bramer *et al.*, 2018), as well as the question at what spatial scale microclimate
155 data can prove most informative for ecological modelling (Jucker *et al.*, 2020). The temperature time
156 series in the database, many of which are covering increasingly long time periods of up to a decade or
157 more, will also allow fine-tuning forecasts of microclimate data into the future by deepening our
158 understanding of the link between microclimatic dynamics in the soil and the air (Lenoir *et al.*, 2017,
159 Wason *et al.*, 2017, Bramer *et al.*, 2018, Maclean, 2019), improving our predictions of biodiversity and
160 ecosystem functioning under climate change.

161 ***Dig out your loggers! A call for contributions***

162 To reach these goals, we encourage scientists owning in-situ measured temperature data to submit
163 these to the growing SoilTemp database. All time series spanning one month or more, with
164 temperature measurements a maximum of 4 hours apart, all soil depths, all heights above the ground
165 up till two meters, all biomes, and all sensor types and brands will be accepted. Note that both spatially
166 dense and sparse logger networks, as well as single loggers are accepted. The achieved spatial
167 resolution is dependent on the provision of spatially precise coordinates to achieve a good relationship
168 with potential explanatory variables (e.g. high resolution remotely sensed environmental data). If we
169 have these coordinates and thus the location and distance between loggers, we can effectively obtain
170 the extent and spacing for each logger network (Western *et al.*, 2002).

171 We include data from both observational and experimental plots, yet sensors have to be measuring
172 in-situ and not in pots, and experiments manipulating the local climate (e.g. open-top chambers, rain-
173 out shelters or vegetation-removal experiments) are excluded (Table 1). Given currently less well-
174 represented climate regions, we especially encourage submissions from extreme cold and hot
175 environments to fill the remaining gaps in our global coverage. More specifically, hot tropical climates
176 (both tropical rainforests and tropical seasonal forests and savannas) and cold and hot deserts are
177 currently still largely underrepresented (Fig. 2b), in particular from Africa, Asia, Antarctica and the
178 Americas (Fig. 2a). Data contributors will be invited as co-authors on the main global papers resulting
179 from this database (see Supplementary Materials for details on terms of use and data ownership).

180 By encouraging sampling and submissions from remote areas, we aim to help solve the global sampling
181 bias in soil ecological data (Cameron *et al.*, 2018, Guerra *et al.*, 2019), and we hope to build a truly
182 global network representing – and actively engaging - scientists from a wide diversity of cultural

183 backgrounds (Maestre & Eisenhauer, 2019). More information is available on the SoilTemp website,
184 accessible via Figshare (DOI [10.6084/m9.figshare.12126516](https://doi.org/10.6084/m9.figshare.12126516)).

185 When fully established, the SoilTemp database and its derivative products (e.g. bioclimatic variables)
186 will be made freely available to facilitate the analysis of global patterns in microclimates, increase the
187 comparability between regional studies and simplify the use of accurate microclimatic data in ecology
188 (Bramer *et al.*, 2018). At the moment, critical metadata is already freely accessible via Figshare (DOI
189 [10.6084/m9.figshare.12126516](https://doi.org/10.6084/m9.figshare.12126516)). Given the absence of and the need for globally available soil
190 microclimate data products at relevant spatial resolutions for use in ecological analyses, we believe
191 that SoilTemp has the potential to become a highly important resource that will enable a step change
192 in ecological modelling.

193

194 **Table**

195 *Table 1: Minimal data requirements and obligatory metadata for submission to the database. For*
196 *more details, see Supplementary Material.*

197

Minimum data requirements	Obligatory metadata
Minimum one consecutive month of in-situ measured temperature time series	Accurate (handheld GPS or finer) spatial coordinates of the loggers (+ estimated accuracy)
Maximum time interval between measurements: 4 hours	Height/depth of the sensor relative to the soil surface
No climate manipulation experiments (only control plots of those experiments, or observational studies)	Type or brand of temperature sensor used, and type of shelter (e.g. no shelter, home-made shelter, Stevenson screen...)
No modelling studies (only empirical data)	Temporal resolution of the sensor Habitat classification

198

199

200 **Figures**

201 **Figure 1: the horizontal and vertical drivers of the offset between in-situ soil and free-air**
202 **temperatures.** Conceptually, there are two different sets of features responsible for the offset between
203 coarse-scale free air temperatures (top left, e.g. WorldClim, Fick & Hijmans, 2017) and fine-scale soil
204 temperatures (bottom right, e.g. Ashcroft & Gollan, 2012, Lembrechts et al., 2019),. Firstly, one can
205 incorporate fine-scale horizontal climate-forcing factors like topography and terrain-related features,
206 land cover types and distance to water bodies to go from coarse-scaled to finer resolutions (top right,
207 e.g. Aalto et al., 2017, Macek et al., 2019). Secondly, one can consider observation height, and the
208 effects of vegetation characteristics (like structure and cover), snow cover and soil characteristics (like
209 moisture, geological types, texture and bulk density) on the radiation balance to convert from free-air
210 to soil temperatures (e.g. Kearney, 2019). Both horizontal and vertical features can introduce positive
211 or negative differences (offset values) between soil and air temperatures through their effects on
212 processes related to the radiation balance, like wind, convective heat transfer and surface albedo. The
213 complexities of these horizontal and vertical processes can vary with biome, season and time of day.
214 Temperatures are represented here using an unspecified temperature range from cold (blue) to warm
215 (red).

216 **Figure 2: Overview of the status of the SoilTemp-database as of March 2020.** Spatial (a), climatic (b),
217 elevational (c) and temporal (d) distribution of sensors in the SoilTemp-database as of March 2020. (a)
218 Background world map in WGS1984, hexagons with a resolution of approximately 70.000 km² using
219 the *dggridR*-package in R. (b) Colors of hexagons indicate the number of sensors at each climatic
220 location, with a 40 × 40 bin resolution. Small dots in the background represent the global variation in
221 climatic space (obtained by sampling 1.000.000 random locations from the CHLSA world maps at a
222 spatial resolution of 2.5 arc minutes. Overlay with dotted lines and numbers (from 1 to 9) depict a
223 delineation of Whittaker biomes (adapted from Whittaker, 1970): (1) tundra and ice, (2) boreal forest,
224 (3) temperate seasonal forest, (4) temperate rainforest, (5) tropical rainforest, (6) tropical seasonal
225 forest/savanna, (7) subtropical desert, (8) temperate grassland/desert, (9) woodland/shrubland. (c)
226 Number of sensors in each elevation class. (d) Time span covered by each sensor in the database,
227 ranked by starting date. Data showed from 1992 onwards, note that the time period covered by 4
228 loggers with starting dates in 1976 is truncated.

229 Acknowledgments

230 We thank Sylvain Pincebourde and an anonymous reviewer for their critical evaluation of our manuscript. This work was
231 supported by the Research Foundation Flanders (FWO) through a postdoctoral fellowship to Jonas J. Lembrechts (12P1819N)
232 and a Research Network Grant (WOG W001919N). We gratefully acknowledge all data contributors, all staff of the author
233 institutions engaged in field measurements and equipment maintenance (namely Erik Herberg, Iris Hamersveld, Ida Westman,
234 Fredrik Brounes, Pernille Eidsen, Eleanor Walker and the teachers participating in the *Tepåseförsöket 2015*) and the ILTER-
235 network, and thank local peoples for permission to collect data on their lands. Temperature data collection on European GLORIA
236 summits was funded by European Union FP-5 project GLORIA-Europe (EVK2-CT-2000-0006) and the Swiss MAVA Foundation
237 project 'Climate change impacts in protected areas of the Alps and high mountains of Eastern Europe and the Mediterranean
238 region', on the Eastern Swiss GLORIA summits by the Swiss Federal Office for the Environment (FOEn), the Research
239 Commission and staff of the Swiss National Park, and the Foundation Dr. Joachim de Giacomo, on Tenerife in the framework of
240 the Flexible Pool project (W47014118) of Sylvia Haider funded by the German Centre for Integrative Biodiversity Research (iDiv)
241 Halle-Jena-Leipzig, on Livingston Island, Antarctica by different research projects of the Govern of Spain (PERMAPLANET
242 CTM2009-10165-E; ANTARPERMA CTM2011-15565-E; PERMASNOW CTM2014-52021-R), and the PERMATHERMAL
243 arrangement between the University of Alcalá and the Spanish Polar Committee and on the Western Swiss GLORIA summits by
244 Département de la culture et des sports du Valais, Fondation Mariétan, Société académique de Genève, Swiss Federal Office of
245 Education and Science and Swiss Federal Office for the Environment. Jan Wild, Martin Macek, Martin Kopecký, Lucia Hederová,
246 Matěj Man and Josef Brůna were supported by the Czech Science Foundation (project 17-13998S) and the Czech Academy of
247 Sciences (project RVO 67985939), Meelis Pärtel by an Estonian Research Council grant (PRG609) and by the European
248 Regional Development Fund (Centre of Excellence EcolChange), Lena Muffler, Juergen Kreyling, Robert Weigel, Mario Trouillier,
249 Martin Wilmking and Jonas Schmeddes by DFG GraKo 2010 Response, Juha M. Alatalo by Qatar Petroleum (QUEx-CAS-QP-
250 RD-18/19), the authors from Odesa National I. I. Mechnikov University (Sergiy Medinets and Volodymyr Medinets) by EU FP6
251 The nitrogen cycle and its influence on the European greenhouse gas balance (NitroEurope), EU FP7 Effects of Climate Change
252 on Air Pollution Impacts and Response Strategies for European Ecosystems (ÉCLAIRE), Ukrainian national research projects
253 (No. 505, 550, 574) funded by Ministry of Education and Science of Ukraine and GEF-UNEP funded 'Towards INMS' project, see
254 www.inms.international for more details. Florian Zellweger was supported by the Swiss National Science Foundation (grant no.
255 172198), Peter Barančok, Róbert Kanka, Jozef Kollár and Andrej Palaj by the Slovak Scientific Grant Agency (project VEGA
256 2/0132/18), Jonas Ardö by a infrastructure grant from faculty of Science, Lund University, Julia Kempinen by the Doctoral
257 Programme in Geosciences at the University of Helsinki, Jan Altman by the Czech Science Foundation (projects 17-07378S and
258 20-05840Y), the Czech Academy of Sciences (project RVO 67985939) and Ministry of Education, Youth and Sport of the Czech
259 Republic, program Inter-Excellence, subprogram Inter-Action (project LTAUSA19137), Toke Thomas Høye by the Carlsberg
260 Foundation (grant no. CF16-0896) and the Villum Foundation (grant no. 17523), Jiri Dolezal by the Czech Science Foundation
261 (projects 17-19376S), and Ministry of Education, Youth and Sport of the Czech Republic, program Inter-Excellence, subprogram
262 Inter-Action (project LTAUSA18007), Nico Eisenhauer, Felix Gottschall and Simone Cesarz by the German Centre for Integrative
263 Biodiversity Research (iDiv) Halle-Jena-Leipzig, funded by the German Research Foundation (FZT 118), Stuart W. Smith by
264 AfricanBioServices project funded by the EU Horizon 2020 grant number 641918, Haydn Thomas by a K Natural Environmental
265 Research Council doctoral training partnership grant NE/L002558/1, Isla H. Myers-Smith by the UK Natural Environmental
266 Research Council ShrubTundra Project NE/M016323/1, Anibal Pauchard, Rafael Garcia and Eduardo Fuentes-Lillo by the
267 projects Fondecyt 1180205, Fondecyt 11170516 and CONICYT PIA APOYO CTE AFB170008, Rafaela Canessa, Maaiké Y.
268 Bader, Liesbeth van den Brink, and Katja Tielbörger by the DFG Priority Programme 1803 EarthShape (projects 1 (BA 3843/6-
269 1) and 11 (TI 338/14-1&2)), Martin Svátek by a grant from the Ministry of Education, Youth and Sports of the Czech Republic
270 (grant number: INTER-TRANSFER LTT17017), Mihai Pușcaș by ODYSSEE project (ANR-13-ISV7-0004 France, PN-III-ID-JRP-
271 RO-FR-2012 UEFISCDI Romania), Pavel Dan Turtureanu by UEFISCDI in Romania, MEMOIRE grant no. PN-III-P1-1.1-PD2016-
272 0925, Jonathan Lenoir by the Agence Nationale de la Recherche (ANR) within the framework of the IMPRINT project "IMpacts

273 des PProcessus microclimatiques sur la redistribution de la biodiversité forestière en contexte de réchauffement du macroclimat"
274 (grant number: ANR-19-CE32-0005-01), Radim Matula and Roman Plichta by a grant Inter-Excellence (project: INTER-
275 TRANSFER LTT17033) from the Ministry of Education, Youth and Sports of the Czech Republic, Lisa Rew by the National
276 Institute of Food and Agriculture, U.S. Department of Agriculture Hatch MONB00363, Tim Seipel and Christian Larson by a grant
277 from the United States National Institute of Food and Agriculture grant 2017-70006-27272, Nina Buchmann by the SNF (projects
278 M4P 40FA40_154245, ICOS-CH 20FI21_148992, 20FI20_173691, InnoFarm 407340_172433) and the EU (SUPER-G contract
279 no. 774124) for the Swiss FluxNet, Mana Gharun by the SNF project ICOS-CH Phase 2 20FI20_173691, Sanne Govaert by the
280 Research Foundation Flanders (FWO) (project G0H1517N Pieter De Frenne, Camille Meeussen. and Pieter Van Gansbeke by
281 the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (ERC
282 Starting Grant FORMICA 757833), Olivier Roupsard by EU-LEAP-Agri (RAMSES II), Agropolis and Total Foundation (DSCATT),
283 CGIAR (GLDC) and EU-DESIRA (CASSECS), Zuzana Sitková by the Slovak Research and Development Agency under the
284 project No. APVV-16-0325 and project ITMS 26220220066 co-funded by the ERDF, Brett Ryan Scheffers by National Geographic
285 Society (grant no. 9480-14 and WW-240R-17), James D. M. Speed by the Research Council of Norway (262064), William D.
286 Pearse and the Pearse Lab by National Science Foundation ABI-1759965, NSF EF-1802605 and United States Department of
287 Agriculture Forest Service agreement 18-CS-11046000-041, Isla H. Myers-Smith by the UK Natural Environmental Research
288 Council ShrubTundra Project NE/M016323/1, Andrew D Thomas by a Leverhulme Trust Research Fellowship under Government
289 of Botswana permit EWT8/36/4 VIII(4), Shengwei Zong by National Natural Science Foundation of China (No. 41971124), Roman
290 Plichta by the post-doc project 7.3 of Institutional plan of Mendel University in Brno 2019–2020, František Máliš by the Slovak
291 Research and Development Agency project APVV-15-0270, Filip Hrbacek and Kamil Laska by the projects LM2015078 and
292 CZ.02.01/0.0/0.0/16_013/0001708 of Ministry of Youth and Sports of the Czech Republic, T-M. Ursu was supported by the
293 Ministry of Research and Innovation through Projects for Excellence Financing in RDI: Contract no. 22 PFE/2018 and PN2019-
294 2022/19270201 – Ctr. 25N BIODIVERS 3-BIOSERV and Andrej Varlagin by RFBR project number 19-04-01234-a. Lore T.
295 Verryckt is funded by a PhD fellowship from the Research Foundation Flanders (FWO) and acknowledges support from the
296 European Research Council Synergy Grant; ERC-2562013-SyG-610028 IMBALANCE-P and Pallieter De Smedt holds a
297 postdoctoral fellowship of the Research Foundation-Flanders (FWO) and The Kreinitz Experiment is a cooperative research
298 project initiated by the Helmholtz Centre for Environmental Research - UFZ. We also acknowledge project 18-74-10048 from the
299 Russian Science Foundation, the Dirección General de Cambio Climático del Gobierno de Aragón, the Ordesa y Monte Perdido
300 National Park and the Servicio de Medio Ambiente de Soria de la Junta de Castilla y León, the National Swiss Fund for research
301 (SNSF, project "Lif3web", n°162604).

302 **Conflict of Interest:** The authors declare that they have no conflict of interest.

303 **References**

- 304 Aalto J, Riihimäki H, Meineri E, Hylander K, Luoto M (2017) Revealing topoclimatic heterogeneity
305 using meteorological station data. *International Journal of Climatology*, **37**, 544-556.
- 306 Aalto J, Scherrer D, Lenoir J, Guisan A, Luoto M (2018) Biogeophysical controls on soil-atmosphere
307 thermal differences: implications on warming Arctic ecosystems. *Environmental Research*
308 *Letters*, **13**, 074003.
- 309 Abatzoglou JT, Dobrowski SZ, Parks SA, Hegewisch KC (2018) TerraClimate, a high-resolution global
310 dataset of monthly climate and climatic water balance from 1958–2015. *Scientific data*, **5**,
311 170191.
- 312 Ashcroft MB, Cavanagh M, Eldridge MDB, Gollan JR (2014) Testing the ability of topoclimatic grids of
313 extreme temperatures to explain the distribution of the endangered brush-tailed rock-
314 wallaby (*Petrogale penicillata*). *Journal of biogeography*, **41**, 1402-1413.
- 315 Ashcroft MB, Chisholm LA, French KO (2008) The effect of exposure on landscape scale soil surface
316 temperatures and species distribution models. *Landscape Ecology*, **23**, 211-225.
- 317 Ashcroft MB, Chisholm LA, French KO (2009) Climate change at the landscape scale: predicting fine-
318 grained spatial heterogeneity in warming and potential refugia for vegetation. *Global change*
319 *biology*, **15**, 656-667.
- 320 Ashcroft MB, Gollan JR (2012) Fine-resolution (25 m) topoclimatic grids of near-surface (5 cm)
321 extreme temperatures and humidities across various habitats in a large (200 x 300 km) and
322 diverse region. *International Journal of Climatology*, **32**, 2134-2148.
- 323 Ashcroft MB, Gollan JR (2013) Moisture, thermal inertia, and the spatial distributions of near-surface
324 soil and air temperatures: Understanding factors that promote microrefugia. *Agricultural*
325 *and Forest Meteorology*, **176**, 77-89.
- 326 Bennie J, Huntley B, Wiltshire A, Hill MO, Baxter R (2008) Slope, aspect and climate: Spatially explicit
327 and implicit models of topographic microclimate in chalk grassland. *Ecological Modelling*,
328 **216**, 47-59.
- 329 Bennie J, Wilson RJ, Maclean IMD, Suggitt AJ (2014) Seeing the woods for the trees - when is
330 microclimate important in species distribution models? *Global change biology*, **20**, 2699-
331 2700.
- 332 Bramer I, Anderson B, Bennie J, Bladon A, De Frenne P, Hemming D, Hill RA, Kearney MR, Körner C,
333 Korstjens AH, Lenoir J, Maclean IMD, Marsh CD, Morecroft MD, Ohlemüller R, Slater HD,
334 Suggitt AJ, Zellweger F, Gillingham PK (2018) Advances in monitoring and modelling climate
335 at ecologically relevant scales. *Advances in Ecological Research*, **58**, 101-161.
- 336 Bruelheide H, Dengler J, Purschke O, Lenoir J, Jiménez-Alfaro B, Hennekens SM, Botta-Dukát Z,
337 Chytrý M, Field R, Jansen F (2018) Global trait–environment relationships of plant
338 communities. *Nature ecology & evolution*, **2**, 1906.
- 339 Cameron EK, Martins IS, Lavelle P, Mathieu J, Tedersoo L, Gottschall F, Guerra CA, Hines J, Patoine G,
340 Siebert J (2018) Global gaps in soil biodiversity data. *Nature ecology & evolution*, **2**, 1042.
- 341 Carter A, Kearney M, Mitchell N, Hartley S, Porter W, Nelson N (2015) Modelling the soil
342 microclimate: does the spatial or temporal resolution of input parameters matter? *Frontiers*
343 *of Biogeography*, **7**, 138-154.
- 344 Copernicus Climate Change Service (C3s) (2019) C3S ERA5-Land reanalysis. (ed Copernicus Climate
345 Change Service).
- 346 Coûteaux M-M, Bottner P, Berg B (1995) Litter decomposition, climate and litter quality. *Trends in*
347 *ecology & evolution*, **10**, 63-66.
- 348 Daly C (2006) Guidelines for assessing the suitability of spatial climate data sets. *International journal*
349 *of climatology*, **26**, 707-721.
- 350 De Boeck HJ, Van De Velde H, De Groote T, Nijs I (2016) Ideas and perspectives: Heat stress: more
351 than hot air. *Biogeosciences*, **13**, 5821-5825.

352 De Frenne P, Rodríguez-Sánchez F, Coomes DA, Baeten L, Verstraeten G, Vellend M, Bernhardt-
353 Römermann M, Brown CD, Brunet J, Cornelis J (2013) Microclimate moderates plant
354 responses to macroclimate warming. *Proceedings of the National Academy of Sciences*, **110**,
355 18561-18565.

356 Fick SE, Hijmans RJ (2017) WorldClim 2: new 1-km spatial resolution climate surfaces for global land
357 areas. *International Journal of Climatology*, **37**, 4302-4315.

358 Geiger R (1950) *The climate near the ground*, Cambridge, Massachusetts, USA, Harvard University
359 Press.

360 Gottschall F, Davids S, Newiger-Dous TE, Auge H, Cesarz S, Eisenhauer N (2019) Tree species identity
361 determines wood decomposition via microclimatic effects. *Ecology and evolution*, **9**, 12113-
362 12127.

363 Goulden M, Wofsy S, Harden J, Trumbore SE, Crill P, Gower S, Fries T, Daube B, Fan S-M, Sutton D
364 (1998) Sensitivity of boreal forest carbon balance to soil thaw. *Science*, **279**, 214-217.

365 Guerra CA, Heintz-Buschart A, Sikorski J, Chatzinotas A, Guerrero-Ramírez N, Cesarz S, Beaumelle L,
366 Rillig MC, Maestre FT, Delgado-Baquerizo M (2019) Blind spots in global soil biodiversity and
367 ecosystem function research. *bioRxiv*, 774356.

368 Hursh A, Ballantyne A, Cooper L, Maneta M, Kimball J, Watts J (2017) The sensitivity of soil
369 respiration to soil temperature, moisture, and carbon supply at the global scale. *Global
370 change biology*, **23**, 2090-2103.

371 Jarraud M (2008) Guide to meteorological instruments and methods of observation (WMO-No. 8).
372 World Meteorological Organisation: Geneva, Switzerland.

373 Jucker T, Jackson T, Zellweger F, Swinfield T, Gregory N, Williamson J, Slade E, Phillips J, Bittencourt
374 P, Blonder B, Boyle M, Ellwood M, Hemprich-Bennett D, Lewis O, Matula R, Senior RA,
375 Shenkin A, Svatek M, Coomes D (2020) A research agenda for microclimate ecology in
376 human-modified tropical forests. *Frontiers in Forests and Global Change*, **2**.

377 Karger DN, Conrad O, Böhner J, Kawohl T, Kreft H, Soria-Auza RW, Zimmermann NE, Linder HP,
378 Kessler M (2017) Climatologies at high resolution for the earth's land surface areas. *Scientific
379 data*, **4**, 170122.

380 Kattge J, Bönsch G, Diaz S, Lavorel S, Prentice IC, Leadley P, Tautenhahn S, Werner G, Günther A
381 (2019) TRY plant trait database-enhanced coverage and open access. *Global change biology*,
382 **26**, 119-188.

383 Kearney M, Shine R, Porter WP (2009) The potential for behavioral thermoregulation to buffer “cold-
384 blooded” animals against climate warming. *Proceedings of the National Academy of
385 Sciences*, **106**, 3835-3840.

386 Kearney MR (2019) MicroclimOz—A microclimate data set for Australia, with example applications.
387 *Austral Ecology*, **44**, 534-544.

388 Kearney MR, Isaac AP, Porter WP (2014a) microclim: Global estimates of hourly microclimate based
389 on long-term monthly climate averages. *Scientific data*, **1**, 140006.

390 Kearney MR, Shamakhy A, Tingley R, Karoly DJ, Hoffmann AA, Briggs PR, Porter WP (2014b)
391 Microclimate modelling at macro scales: a test of a general microclimate model integrated
392 with gridded continental-scale soil and weather data. *Methods in Ecology and Evolution*, **5**,
393 273-286.

394 Kissling WD, Walls R, Bowser A, Jones MO, Kattge J, Agosti D, Amengual J, Basset A, Van Bodegom
395 PM, Cornelissen JH (2018) Towards global data products of Essential Biodiversity Variables
396 on species traits. *Nature ecology & evolution*, **2**, 1531-1540.

397 Körner C, Hiltbrunner E (2018) The 90 ways to describe plant temperature. *Perspectives in plant
398 ecology, evolution and systematics*, **30**, 16-21.

399 Körner C, Paulsen J (2004) A world-wide study of high altitude treeline temperatures. *Journal of
400 biogeography*, **31**, 713-732.

401 Lembrechts J, Nijs I, Lenoir J (2018) Incorporating microclimate into species distribution models.
402 *Ecography*, **42**, 1267-1279.

403 Lembrechts JJ, Lenoir J, Roth N, Hattab T, Milbau A, Haider S, Pellissier L, Pauchard A, Ratier Backes
404 A, Dimarco RD (2019) Comparing temperature data sources for use in species distribution
405 models: From in-situ logging to remote sensing. *Global Ecology and Biogeography*, **28**, 1578-
406 1596.

407 Lenoir J, Hattab T, Pierre G (2017) Climatic microrefugia under anthropogenic climate change:
408 implications for species redistribution. *Ecography*, **40**, 253-266.

409 Li T-T (1926) *Soil temperature as influenced by forest cover*.

410 Macek M, Kopecký M, Wild J (2019) Maximum air temperature controlled by landscape topography
411 affects plant species composition in temperate forests. *Landscape Ecology*, **34**, 2541-2556.

412 Maclean IM (2019) Predicting future climate at high spatial and temporal resolution. *Global change
413 biology*, **26**, 1003-1011.

414 Maclean IMD, Hopkins JJ, Bennie J, Lawson CR, Wilson RJ (2015) Microclimates buffer the responses
415 of plant communities to climate change. *Global Ecology and Biogeography*, **24**, 1340-1350.

416 Maclean IMD, Suggitt AJ, Wilson RJ, Duffy JP, Bennie JJ (2017) Fine-scale climate change: modelling
417 spatial variation in biologically meaningful rates of warming. *Global change biology*, **23**, 256-
418 268.

419 Maestre FT, Eisenhauer N (2019) Recommendations for establishing global collaborative networks in
420 soil ecology. *Soil organisms*, **91**, 73.

421 Medinets S, Gasche R, Kiese R, Rennenberg H, Butterbach-Bahl K (2019) Seasonal dynamics and
422 profiles of soil NO concentrations in a temperate forest. *Plant and Soil*, **445**, 335-348.

423 Meineri E, Hylander K (2017) Fine-grain, large-domain climate models based on climate station and
424 comprehensive topographic information improve microrefugia detection. *Ecography*, **40**,
425 1003-1013.

426 Niittynen P, Luoto M (2017) The importance of snow in species distribution models of arctic
427 vegetation. *Ecography*, **41**, 1024-1037.

428 Opedal OH, Armbruster WS, Graae BJ (2015) Linking small-scale topography with microclimate, plant
429 species diversity and intra-specific trait variation in an alpine landscape. *Plant Ecology &
430 Diversity*, **8**, 305-315.

431 Pincebourde S, Casas J (2019) Narrow safety margin in the phyllosphere during thermal extremes.
432 *Proceedings of the National Academy of Sciences*, **116**, 5588-5596.

433 Pincebourde S, Murdock CC, Vickers M, Sears MW (2016) Fine-scale microclimatic variation can
434 shape the responses of organisms to global change in both natural and urban environments.
435 *Integrative and Comparative Biology*, **56**, 45-61.

436 Pleim JE, Gilliam R (2009) An indirect data assimilation scheme for deep soil temperature in the
437 Pleim–Xiu land surface model. *Journal of Applied Meteorology and Climatology*, **48**, 1362-
438 1376.

439 Portillo-Estrada M, Pihlatie M, Korhonen JFJ, Levula J, Frumau AKF, Ibrom A, Lembrechts JJ, Morillas
440 L, Horvath L, Jones SK, Niinemets U (2016) Climatic controls on leaf litter decomposition
441 across European forests and grasslands revealed by reciprocal litter transplantation
442 experiments. *Biogeosciences*, **13**, 1621-1633.

443 Pradervand J-N, Dubuis A, Pellissier L, Guisan A, Randin C (2014) Very high resolution environmental
444 predictors in species distribution models: Moving beyond topography? *Progress in Physical
445 Geography*, **38**, 79-96.

446 Randin CF, Vuissoz G, Liston GE, Vittoz P, Guisan A (2009) Introduction of snow and geomorphic
447 disturbance variables into predictive models of alpine plant distribution in the Western Swiss
448 Alps. *Arctic, Antarctic, and Alpine Research*, **41**, 347-361.

449 Rosenberg NJ, Kimball B, Martin P, Cooper C (1990) From climate and CO₂ enrichment to
450 evapotranspiration. *Climate change and US water resources.*, 151-175.

451 Schimel DS, Braswell B, Mckeown R, Ojima DS, Parton W, Pulliam W (1996) Climate and nitrogen
452 controls on the geography and timescales of terrestrial biogeochemical cycling. *Global
453 Biogeochemical Cycles*, **10**, 677-692.

454 Slavich E, Warton DI, Ashcroft MB, Gollan JR, Ramp D (2014) Topoclimate versus macroclimate: how
455 does climate mapping methodology affect species distribution models and climate change
456 projections? *Diversity and Distributions*, **20**, 952-963.

457 Suggitt AJ, Gillingham PK, Hill JK, Huntley B, Kunin WE, Roy DB, Thomas CD (2011) Habitat
458 microclimates drive fine-scale variation in extreme temperatures. *Oikos*, **120**, 1-8.

459 Suggitt AJ, Wilson RJ, Isaac NJ, Beale CM, Auffret AG, August T, Bennie JJ, Crick HQ, Duffield S, Fox R
460 (2018) Extinction risk from climate change is reduced by microclimatic buffering. *Nature*
461 *Climate Change*, **8**, 713.

462 Vitasse Y, Klein G, Kirchner JW, Rebetez M (2017) Intensity, frequency and spatial configuration of
463 winter temperature inversions in the closed La Brevine valley, Switzerland. *Theoretical and*
464 *applied climatology*, **130**, 1073-1083.

465 Wason JW, Bevilacqua E, Dovciak M (2017) Climates on the move: Implications of climate warming
466 for species distributions in mountains of the northeastern United States. *Agricultural and*
467 *Forest Meteorology*, **246**, 272-280.

468 Western AW, Grayson RB, Blöschl G (2002) Scaling of soil moisture: A hydrologic perspective. *Annual*
469 *review of earth and planetary sciences*, **30**, 149-180.

470 Whiteman CD (1982) Breakup of temperature inversions in deep mountain valleys: Part I.
471 Observations. *Journal of Applied Meteorology*, **21**, 270-289.

472 Whittaker RH (1970) Communities and ecosystems. *Communities and ecosystems*.

473 Zellweger F, De Frenne P, Lenoir J, Rocchini D, Coomes D (2019) Advances in microclimate ecology
474 arising from remote sensing. *Trends in ecology & evolution*, **34**, 327-341.

475 Zhang Y, Wang S, Barr AG, Black T (2008) Impact of snow cover on soil temperature and its
476 simulation in a boreal aspen forest. *Cold Regions Science and Technology*, **52**, 355-370.

477