

OAK RIDGE NATIONAL LABORATORY

operated by

UNION CARBIDE CORPORATION NUCLEAR DIVISION

for the

U.S. ATOMIC ENERGY COMMISSION



ORNL - TM - 2194

DATE -

Neutron Physics Division

SOLAR NEUTRON TRANSPORT IN THE EARTH'S ATMOSPHERE

R. G. Alsmiller, Jr., and R. T. Boughner*

Abstract

The neutron flux per unit energy induced in the earth's atmosphere by solar neutrons has been calculated as a function of atmospheric depth for depths of less than 300 g/cm² (~30,000 ft). The calculated flux is compared with estimates of the cosmic-ray neutron flux per unit energy, and it is found that for many flares the solar neutron flux per unit energy is sufficiently above the cosmic-ray neutron flux per unit energy to produce a measurable effect at both balloon and aircraft altitudes.

N 68-27208

(THRU)

(COPY)

(CATEGORY)

(ACCESSION NUMBER)

(PAGES)

(NASA CR OR TMX OR AD NUMBER)

FACILITY FORM 602

GPO PRICE \$

CFSTI PRICE(S) \$

Hard copy (HC) 3.00

Microfiche (MF) .65

ff 653 July 65

NOTE:

This Work Partially Supported by
NATIONAL AERONAUTICS AND SPACE ADMINISTRATION
Under Order R-104(1)

*Mathematics Division.

NOTICE This document contains information of a preliminary nature and was prepared primarily for internal use at the Oak Ridge National Laboratory. It is subject to revision or correction and therefore does not represent a final report.



LEGAL NOTICE

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

1. *Introduction.* It has often been suggested that the high-energy protons emitted from the sun during solar flares will interact with the solar photosphere and produce high-energy neutrons. In a series of papers, hereinafter referred to as 1, *Lingenfelter et al.* [1965] have estimated the intensity and energy spectra of solar neutrons which might exist in the vicinity of the earth following a solar flare. In 1, it is estimated that following a major flare high-energy neutrons should be observable in the vicinity of the earth for times of the order of one hour after the flare. It is furthermore estimated that these neutrons produce no observable effect on the surface of the earth, and it is suggested that the search for solar neutrons be "conducted with a satellite in equatorial orbit or a balloon at high altitude and low latitude carrying a neutron counter sensitive above 10 Mev."

In this paper, the alternate possibility of observing solar neutrons by neutron spectral measurements at aircraft altitudes is considered. When high-energy neutrons enter the atmosphere, they interact with the atmospheric nuclei to produce a nuclear cascade in much the same manner as do the incident cosmic-ray protons and alpha particles. The question considered here is the magnitude of the neutron flux per unit energy at various depths in the atmosphere due to incident solar-flare neutrons compared to that due to incident cosmic rays. Throughout the paper, consideration is restricted to latitudes where the earth's magnetic field shields the atmosphere from solar-flare protons, and thus neutrons induced in the atmosphere by such protons need not be considered.

Using the estimate of the solar-neutron flux given in 1, high-energy neutron transport calculations have been carried out, and the neutron flux per unit energy at various depths in the atmosphere has been obtained. Since

in 1 it is estimated that the solar neutrons reaching the earth's atmosphere are essentially monoenergetic with the energy varying as a function of time, the calculations have been carried out for incident energies of 50, 100, 200, 300, 400, and 500 Mev. Calculated results are given at balloon altitudes as well as at greater atmospheric depths. The calculated neutron flux is compared with estimates of the cosmic-ray neutron-flux background and is found to be sufficiently above this background, at least under some circumstances, to make measurements of the solar neutrons at balloon and airplane altitudes quite feasible.

In section 2 the details of the calculations are given. In section 3 the results are presented and discussed.

2. *Calculational Details.* The geometry considered throughout this paper is that of a normally incident neutron flux on a half-space of air. The air is assumed to be composed of 79% nitrogen and 21% oxygen. An explicit assumption about the density variation in the upper atmosphere is not required since it can be shown from the transport equations that if the depth is measured in g/cm^2 the calculated results are independent of the density variation.

The transport calculations were carried out using the nucleon transport code NTC [Kinney, 1964]. Since the operation of this code has been described in detail by Kinney, only a brief description of the physical processes will be given here. The interactions of the incident neutrons with the nuclei of the atmosphere initiate a complex avalanche of secondary neutrons and protons which proceed through the atmosphere increasing in number and decreasing in energy. In general, the first nuclear interactions of the incident high-energy neutrons produce several nucleons having

energies ranging from a few Mev to the source energy and quite complex angular distributions. The higher energy secondary neutrons and protons collide with other atmospheric nuclei thus producing additional particles and expanding the cascade. The protons continually lose energy between their nuclear collisions. The low-energy neutrons undergo elastic nuclear collisions which degrade their energy and change their direction of travel. In the calculations presented here, Monte Carlo methods are employed, and all of these various processes are taken into account. The nonelastic cross section for nucleon-nucleus collisions and the energy and angular distributions of the secondary nucleons from such collisions when the incident particle energy is greater than 25 Mev are taken from the calculations of *Bertini* [1966, 1967]. Elastic nucleon-nucleus collisions at energies of greater than 25 Mev are neglected. Protons of energy less than 25 Mev are assumed to slow down and stop without undergoing nuclear collision. Elastic neutron collisions below 25 Mev are treated using primarily experimental data² and nonelastic neutron collisions below 25 Mev are treated using the evaporation calculations of *Dresner* [1961].

²The master cross-section tape compiled by *D. C. Irving* for use in the neutron transport code, together with references to all the data used, is available on request from the Radiation Shielding Information Center of the Oak Ridge National Laboratory.

3. *Results and Conclusions.* Calculations have been carried out for monoenergetic incident neutrons of 50, 100, 200, 300, 400, and 500 Mev. At each of these energies, the normally incident flux (neutrons/cm²/sec) was taken to correspond to the estimate of the incident solar-neutron flux given in 1 for a proton flare spectrum with a characteristic rigidity of 125 Mv and a time-integrated proton flux in the vicinity of the earth of 1.4×10^9 protons/cm² with energy greater than 30 Mev. The incident flux and the approximate time after the flare when neutrons of this energy will be incident on the atmosphere are shown in Table 1. A time-integrated proton flux of 1.4×10^9 protons/cm² with energy greater than 30 Mev corresponds to a very large flare which can not be expected to occur very often. The values in Table 1 and consequently all of the calculated results are, however, directly proportional to the assumed proton flux, so the results may easily be scaled to correspond to a smaller flare. The values in Table 1 are also dependent on the characteristic rigidity of the flare used. The scaling factors that must be applied to the results to make them correspond to flare spectra of other characteristic rigidities may be obtained from 1 (page 4090).

The omnidirectional neutron flux per unit energy, that is, the angular neutron flux integrated over all angles, for each of the incident energies considered is shown at atmospheric depths of 5, 10, 25, 50, 100, 200, and 300 g/cm² in Figures 1-7, respectively. The neutron flux at energies of less than 0.1 Mev is not shown, but values are given in Table 2. The proton flux, integrated over rather large energy intervals because of the relatively large statistical error associated with this small flux, is given in Table 3. There is no calculated flux of protons with energy less than

TABLE 1. Incident Neutron Flux at Each Energy
Used in the Calculations

Energy (Mev)	Time (sec)	Incident Neutron Flux (Neutrons/cm ² /sec)
50	1590	23
100	1160	42
200	880	34
300	763	25.4
400	698	15.9
500	657	7.5

TABLE 2. Neutron Flux Per Unit Energy as a
Function of Depth in Atmosphere
(Averaged over the energy interval 0 to 0.1 Mev)

Incident Neutron Energy (Mev)	Neutrons/Mev/cm ² /sec						
	5 g/cm ²	10 g/cm ²	25 g/cm ²	50 g/cm ²	100 g/cm ²	200 g/cm ²	300 g/cm ²
50	5.43	8.95	19.53	24.51	19.55	5.68	1.40
100	5.92	11.29	20.27	28.92	28.00	14.01	5.25
200	7.07	10.90	22.31	31.88	35.22	21.40	10.19
300	8.43	11.69	25.75	40.51	40.23	29.80	16.07
400	7.12	13.07	26.50	40.52	49.89	39.52	22.01
500	9.02	17.10	27.20	47.42	56.66	45.10	37.7

TABLE 3. Proton Flux as a Function of Atmospheric Depth

Incident Neutron Energy (Mev)	10^{-2} Protons/cm ² /sec						
	5 g/cm ²	10 g/cm ²	25 g/cm ²	50 g/cm ²	100 g/cm ²	200 g/cm ²	300 g/cm ²
(Integrated Over Energy Interval 25 to 100 Mev)							
50	0.19	0.27	0.08	0.13	0.10	0.0	0.0
100	1.74	1.04	1.13	0.76	1.12	0.26	0.05
200	3.34	2.28	3.63	4.91	3.04	2.02	0.29
300	2.47	3.29	2.95	6.76	2.34	3.17	2.04
400	1.84	1.45	4.42	4.89	10.2	2.92	3.06
500	3.65	7.99	8.25	8.69	7.69	5.81	3.72
(Integrated Over Energy Interval 100 Mev to Initial Energy)							
50	0.0	0.0	0.0	0.0	0.0	0.0	0.0
100	0.0	0.0	0.0	0.0	0.0	0.0	0.0
200	1.44	1.77	2.09	1.79	1.08	0.52	0.14
300	1.96	2.98	5.85	7.15	2.79	1.35	1.44
400	1.83	4.95	6.97	8.77	9.05	4.15	1.39
500	4.50	5.47	11.4	13.7	14.1	5.58	3.27




Fig. 1. Neutron Flux Per Unit Energy vs Energy at an Atmospheric Depth of 5 g/cm².




Fig. 2. Neutron Flux Per Unit Energy vs Energy at an Atmospheric Depth of 10 g/cm².




Fig. 3. Neutron Flux Per Unit Energy vs Energy at an Atmospheric Depth of 25 g/cm².

ORNL-DWG 68-3085




Fig. 4. Neutron Flux Per Unit Energy vs Energy at an Atmospheric Depth of 50 g/cm².

ORNL-DWG 68-3086




Fig. 5. Neutron Flux Per Unit Energy vs Energy at an Atmospheric Depth of 100 g/cm².




Fig. 6. Neutron Flux Per Unit Energy vs Energy at an Atmospheric Depth of 200 g/cm².

ORNL-DWG 68-3088




Fig. 7. Neutron Flux Per Unit Energy vs Energy at an Atmospheric Depth of 300 g/cm².

25 Mev because these particles are assumed in the calculation to stop where they originate. The large 'spikes' at the right of each curve in the figures represent the uncollided flux. For purposes of plotting, the monoenergetic uncollided particles have arbitrarily been spread over a 10-Mev interval. While detailed results are not given, it is to be understood that the calculated flux at the lower energies has essentially an isotropic angular distribution while the calculated uncollided and high-energy fluxes are highly directional.

The spectral shape of the cosmic-ray neutron spectrum at the top of the atmosphere is not well established. *Boella et al.*³ [1965] have recently measured the flux of cosmic-ray neutrons with energy less than 20 MeV as a function of atmospheric depth at 42° latitude, and it is possible to compare the calculated results with these measurements. This comparison is shown in Figure 8. For the flare considered and all incident energies considered, the low-energy neutron flux induced in the atmosphere by the solar neutrons is well above the low-energy cosmic-ray neutron spectrum at all depths. The flare proton flux would have to be reduced by one to two orders of magnitude, that is, would have to be of the order of 10^7 to 10^8 protons/cm² with energy greater than 30 Mev, before the solar-neutron-induced low-energy neutron flux would become comparable with the low-energy cosmic-ray neutron flux. The comparison in Figure 8 is for 42° latitude where the measurements were made. At equatorial latitudes where the geomagnetic cut-off is higher, the cosmic-ray neutron flux will be smaller. The magnitude

³A comprehensive list of references to the earlier measurements of the cosmic-ray neutron flux in the atmosphere and a discussion of the degree of agreement of these measurements are given by *Boella* [1965] and *Haymes* [1965].




Fig. 8. Flux of Neutrons with Energy of $< 20 \text{ Mev}$ vs Atmospheric Depth.

of this effect is depth dependent and is not well determined, but the existing measurements [Soberman, 1956] indicate that the cosmic-ray flux in Figure 8 is a factor of three to five larger than the corresponding flux at the equator.

In Figures 4, 5, 6, and 7, that is, at depths of 50, 100, 200, and 300 g/cm², the cosmic-ray neutron flux per unit energy given by Hess *et al.* [1959, 1961] is shown for comparison purposes. At 200 g/cm², this flux is the result of measurements while at the other depths it is due to calculations. At the lower energies, there are other measurements and calculations [Boella *et al.*, 1965; Haymes, 1965]. The results of Hess *et al.* [1959, 1961] are used here not because they are thought to be more accurate but because in general they are higher than those obtained by other workers, and thus give a more pessimistic comparison for our purposes.

In Figures 4, 5, 6, and 7, the solar-neutron flux per unit energy is in general well above the cosmic-ray neutron flux per unit energy at all secondary energies and at all depths. The major exception to this is in Figure 7 where the flux from 50-Mev incident neutrons is comparable to the cosmic-ray flux. The uncollided solar-neutron flux is well above the cosmic-ray flux at all depths. At the lower energies the reduction in flare magnitude that makes the solar-neutron flux comparable to the cosmic-ray flux is smaller than that determined from Figure 8 because the flux of Hess *et al.* is larger than that of Boella [1965]. The comparisons in Figures 4 to 7 are for 44° latitude where the results of Hess *et al.* apply. At equatorial latitudes a reduction in the low-energy cosmic-ray flux due to the higher geomagnetic cutoff, such as that discussed previously, is still to be expected. At higher energies some reduction may also be expected, but the magnitude of this reduction is not known.

In conclusion, it seems that the neutron flux induced in the atmosphere by solar neutrons for a variety of solar flares will be sufficiently above the cosmic-ray flux to produce measurable effects at both balloon and aircraft altitudes. The search for solar neutrons in the atmosphere can, in principle, be carried out by making time-dependent neutron-flux measurements using a counter that is sensitive to either the low-energy secondary neutrons or to the high-energy uncollided incident solar neutrons.

Acknowledgments. It is a pleasure to thank Dr. W. N. Hess for pointing out the need for the calculations reported here and for several helpful discussions.

This work was partially supported by the National Aeronautics and Space Administration [Order R-104(1)], under Union Carbide Corporation's contract with the U. S. Atomic Energy Commission.

REFERENCES

- Bertini, H. W., Results of low-energy, intranuclear-cascade calculation, *Nucl. Phys.* 87, 138-140, 1966.
- Bertini, H. W., Secondary particle spectra from the interaction of 30- to 340-MeV protons on complex nuclei: experimental data and comparison with theory, *Phys. Rev.* 162, 976-982, 1967.
- Boella, G., *et al.*, Measurement of the cosmic ray neutron flux at 4.6 billion volts geomagnetic cutoff rigidity, *J. Geophys. Res.* 70, 1019-1030, 1965.
- Dresner, L., EVAP - a Fortran program for calculating the evaporation of various particles from excited nuclei, ORNL-TM-196, Oak Ridge National Laboratory, 1961.
- Haymes, R. C., Terrestrial and solar neutrons, *Rev. Geophys.* 3, 345-364, 1965.
- Hess, W. N., *et al.*, Cosmic-ray neutron energy spectrum, *Phys. Rev.* 116, 445-457, 1959.
- Hess, W. N., E. H. Canfield, and R. E. Lingenfelter, Cosmic-ray neutron demography, *J. Geophys. Res.* 66, 665-677, 1961.
- Kinney, W. E., The nucleon transport code, NTC, ORNL-3610, Oak Ridge National Laboratory, 1964.
- Lingenfelter, R. E., *et al.*, High-energy solar neutrons. 1. Production in flares, *J. Geophys. Res.* 70, 4077-4086, 1965. 2. Flux at the earth, *J. Geophys. Res.* 70, 4087-4095, 1965.
- Soberman, R. K., High altitude cosmic-ray neutron intensity variations, *Phys. Rev.* 102, 1399-1409, 1956.

NASA-High Energy Listing

- Tino Ahrens, Advanced Research Corporation, 715 Miami Circle, N.E., Atlanta,
Georgia 30324
- Louis Avrami, Explosives Laboratory, Bldg. 407, Pictinny Arsenal, Dover,
New Jersey 07301
- M. Awschalom, Princeton Penn. Accelerator, P. O. Box 682, Princeton,
New Jersey 08540
- M. Barbier, CERN, Geneva Switzerland
- N. Barr, Radiological Physics Branch, Division of Biology and Medicine,
U. S. Atomic Energy Commission, Washington, D. C. 20545 (5 copies)
- C. K. Bauer, Dept. 72-34, Z-26, Lockheed-Georgia Company, Marietta,
Georgia 30060
- P. R. Bell, TH, Chief of Lunar and Earth Sciences Division, Manned Space-
craft Center- NASA, Houston, Texas 77058
- S. Bresticker, Grumman Aircraft Engineering Corp., Space Sciences Group,
Plant 5, Bethpage, L. I., New York 11714
- Karan O. Brien, Health and Safety Laboratory, Radiation Physics Division,
376 Hudson Street, New York, New York 10014
- Brooks Air Force Base, Radiobiology Department, Chief, San Antonio,
Texas 78235
- M. O. Burrell, M-RP-NIP, National Aeronautics and Space Adm., Marshall
Space Flight Center, Huntsville, Alabama 35812
- B. W. Colston, U. S. Atomic Energy Commission, Sandia Area Office,
P. O. Box 5400, Albuquerque, New Mexico 87115
- R. G. Cochran, Department of Nuclear Engineering, A and M College of
Texas, College Station, Texas
- Ted Colvin, Bendix Systems Division, 3300 Plymouth Road, Ann Arbor,
Michigan 48105
- E. A. Cosbie, Argonne National Laboratory, Argonne, Illinois 60440
- Frederick P. Cowan, Brookhaven National Laboratory, Upton, L. I., New
York 11973
- Director, Defense Atomic Support Agency, Pentagon, Washington, D. C.
20301

Charles A. Dempsey, 6570 AMRL (MRT), Wright-Patterson AFB, Ohio 45433

H. DeStaebler, Jr., Stanford Linear Accelerator Center, Stanford University,
Stanford, California 94305

Herman J. Donnert, U. S. Army Nuclear Defense Laboratory, AMXND-C, Edge-
wood Arsenal, Maryland 21010

D. W. Drawbaugh, Westinghouse Electric Corp., Astronuclear Laboratory, P. O.
Box 10864, Pittsburgh, Pennsylvania 15236

John E. Duberg, National Aeronautics and Space Adm., Langley Research Center,
Langley Field, Virginia 23365

D. L. Dye, The Boeing Company, Mail Stop S3-72, Seattle, Washington 98124

Ronald F. Edge, Department of Physics, University of South Carolina,
Columbia, South Carolina

Nat Edmunson, Code R-RP-N, National Aeronautics and Space Adm., Marshall
Space Flight Center, Huntsville, Alabama 35812

Robley D. Evans, Professor of Physics, Room 6-315, Massachusetts Institute
of Technology, Cambridge, Massachusetts 02139

E. M. Finkelman, Grumman Aircraft Engineering Corp., LEM Project, Plant 25,
Bethpage, L.I., New York 11714

Trutz Foelsche, National Aeronautics and Space Adm., Langley Research
Center, Langley Field, Virginia 23365

R. E. Fortney, Northrop Space Laboratories, 3401 West Broadway, Hawthorne,
California 90250

Leo Fox, Code RBH, Biotechnology and Human Research Division, National
Aeronautics and Space Adm., Washington, D. C. 20546

Stan Freden, Aerospace Corp., Box 95085, Los Angeles, California 90045

J. Y. Freeman, Division MPS, CERN, Geneva 23, Switzerland

J. Geibel, CERN, Geneva, Switzerland

R. C. Good, Jr., General Electric Company, Room M7023H - VFSTC, P. O. Box
8555, Philadelphia, Pennsylvania 19101

F. Gordon, Code 716, National Aeronautics and Space Adm., Goddard Space
Flight Center, Greenbelt, Maryland 10027

Raymond M. Hansen, MS 235, National Aeronautics and Space Adm., Langley
Research Center, Mail Stop 235, Langley Field, Virginia 23365

Harry Harrison, Code RRE, National Aeronautics and Space Adm., Hdqts.,
Washington, D. C. 20546

Russell Heath, Phillips Petroleum Company, P. O. Box 2067, Idaho Falls,
Idaho 83401

Herbert D. Hendricks, National Aeronautics and Space Adm., Langley Research
Center, MS499, Langley Field, Virginia 23365

W. N. Hess, National Aeronautics and Space Adm., Goddard Space Flight
Center, Greenbelt, Maryland 20771

R. H. Hilberg, Bellcomm, Inc., 1100 17th Street, N.W., Washington, D. C.
20036

Charles W. Hill, Dept. 73-69, Zone 280, Lockheed-Georgia Company, Marietta,
Georgia 30060

L. Hoffman, CERN, Geneva, Switzerland

J. T. Holloway, Grants and Research Contracts, Office of Space Sciences,
National Aeronautics and Space Adm., Washington, D. C. 20546 (5 copies)

William C. Hulten, National Aeronautics and Space Adm., Mail Stop 235, Langley
Research Center, Langley Field, Virginia 23365

T. Inada, National Institute Radiological Sciences, 250 Kurosuna-Cho, Chiba-
Shi, JAPAN

Lt. Joseph F. Janni, WLRB-1, Air Force Weapons Laboratory, Kirtland Air Force
Base, New Mexico 87117

Dale W. Jenkins, Chief, Environmental Biology, Office of Space Sciences,
NASA, Washington, D. C. 20546

Philippe Tardy-Joubert, Serv. De Protection Contre Les Radiation, Centre D.
Etudes Nucleaires De Saclay, B. P. No. 2, Gif-Sur-Yvette (Seine et Oise),
FRANCE

Clyde Jupiter, General Atomic, P. O. Box 608, San Diego, California 92112

Irving Karp, NASA/ Lewis Research Center, Cleveland, Ohio 44135

Ludwig Katz, Air Force Cambridge Research Center, L. G. Hanscom Field,
Massachusetts

Glenn Keister, Boeing Airplane Company, Aerospace Division, P. O. Box 3707,
Seattle, Washington 98124

J. Warren Keller, Code RV-1, National Aeronautics and Space Adm., Washington,
D. C. 20546 (3 copies)

- James F. Kenny, Boeing Scientific Research Laboratory, P. O. Box 3981,
Seattle, Washington 98124
- E. C. Kidd, Zone S71, Dept. 61-2, General Dynamics/Fort Worth, P. O. Box 748,
Fort Worth, Texas 76101
- David King, Department of Physics, University of Tennessee, Knoxville,
Tennessee 37916
- Robert L. Kloster, McDonnell Aircraft Corp., P. O. Box 516, St. Louis,
Missouri 63166
- George A. Kolstad, Assistant Director Physics-Math Programs, Division of
Research, U. S. Atomic Energy Commission, Washington, D. C. 20545
- Eugene B. Konecni, Department of Management BEOB-200, University of Texas,
Austin, Texas 78712
- W. Kreger, Code 940, U. S. Naval Radiological Defense Laboratory, San
Francisco, California 94135
- Ed Kuhn, Republic Aviation Corporation, Power Conversion Systems Division,
Farmingdale, L. I., New York 11735
- Wright H. Langham, Los Alamos Scientific Laboratory, Los Alamos, New
Mexico 87544
- Borje Larsson, University of Uppsala, The Gustaf Werner Institute, Uppsala,
Sweden
- J. M. Lavie, Centre D. Etudes Nucleaires, De Saclay, S. E. C. R. - B. P.
No. 2, 91-Gif-Sur-Yvette, FRANCE
- Martin Leimdorfer, Industri-Matematik AB, De Geersgatan 8, Stockholm No.
SWEDEN
- S. H. Levine, Northrop Space Laboratories, 3401 W. Broadway, 2452/61,
Hawthorne, California 90250
- Lynn R. Lewis, Dept. 250, Bendix Systems Division, 3300 Plymouth Road, Ann
Arbor, Michigan 48105
- John R. Lilley, A-830-BBFO-78, Missile Space Systems Division, Douglas
Aircraft Co., Inc., Santa Monica, California
- S. J. Lindenbaum, Brookhaven National Laboratory, Upton, L. I., New
York 11973
- Major Russell E. Linkous, Air Force Systems Command (SCTR), Andrews AFB,
Maryland 20331

- M. Stanley Livingston, Cambridge Electron Accelerator, 42 Oxford Street
Cambridge, Massachusetts 02139
- Robert Macklin, Jet Propulsion Laboratory, Pasadena, California 91103
- Brian Mar, Boeing Airplane Co., MS 23-82, Aerospace Division, P. O. Box
3707, Seattle, Washington 98124
- Thomas J. McGuire, Systems Engineering Group (SESSV), Wright-Patterson AFB,
Ohio 45433
- E. J. McLaughlin, Space Medicine, NASA - Code MM, Washington, D. C. 20546
- R. V. Meghreblian, Jet Propulsion Laboratory, 4800 Oak Grove Drive, Pasadena,
California 91103
- Albert E. Metzger, Jet Propulsion Laboratory, 354-401F, 4800 Oak Grove,
Pasadena, California
- J. M. Miller, Chemistry Department, Columbia University, New York, New
York 10027
- R. A. Miller, Zone S71, Dept. 61-2, General Dynamics/Fort Worth, P. O. Box
748, Fort Worth, Texas 76101
- Jerry L. Modisette, National Aeronautics and Space Adm., Manned Spacecraft
Center, Houston, Texas 77001 (5 copies)
- Winnie M. Morgan, Technical Reports, Grants and Research Contracts, Office of
Space Sciences, NASA, Washington, D. C. 20546 (25 copies)
- B. J. Moyer, University of California, Lawrence Radiation Laboratory,
6141 Building 50A, Berkeley, California 94720
- R. F. Mozley, SLAC, Stanford University, Stanford, California 94305
- Capt. J. D. Munson, Space Systems Division (SSTDS), Los Angeles Air Force
Station, Los Angeles, California 90045
- Sam V. Nablo, ION Physics Corporation, Burlington, Massachusetts 01803
- R. R. Nash, Code RRM, National Aeronautics and Space Adm., Washington,
D. C. 20546
- John P. Neissel, MC-506, 175 Courtner Avenue, General Electric Company,
San Jose, California 95125
- W. R. Nelson, Stanford Linear Accelerator Center, Stanford, California
94305

- W. K. H. Panofsky, Stanford Linear Accelerator Center, Stanford University,
Stanford, California 94305
- Wade Patterson, University of California, Lawrence Radiation Laboratory,
Berkeley, California 94720
- Maynard Pearson, Boeing Airplane Company, Aerospace Division, P. O. Box 3707,
Seattle, Washington 98124
- Col. John E. Pickering, USAF, National Aeronautics and Space Adm., (Code MM),
Washington, D. C. 20546
- G. F. Pieper, Code 600, National Aeronautics and Space Adm., Goddard Space
Flight Center, Greenbelt, Maryland 10027
- Robert Pruett, P. O. Box 95085, Los Angeles, California 90045
- Arthur Reetz, Code RV-1, National Aeronautics and Space Adm., Washington,
D. C. 20546 (3 copies)
- O. Reynolds, Director, Bio-Science Programs, Office of Space Sciences,
National Aeronautics and Space Adm., Washington, D. C. 20546 (5 copies)
- Robert G. Riedesel, Douglas Aircraft Company, Missile and Space Division,
3000 Ocean Park Blvd., Santa Monica, California 90405
- Don Robbins, ET32, National Aeronautics and Space Adm., Manned Space-
craft Center, Houston, Texas 77058
- H. J. Schaefer, U. S. Naval School of Aviation Medicine, U. S. Naval
Aviation Medical Center-54, Pensacola, Florida 32512
- W. Wayne Scott, Chattanooga State Technical Institute, 4501 Amnic Highway,
Chattanooga, Tennessee 37401
- Robert L. Seale, University of Arizona, Tuscon, Arizona 85721
- Pierre Lafore Sepp, Commissariat A L. Energie Atomique, Centre D. Etudes
Nucleaires, De Fontenay-Aux-Roses (Seine), Boite Postale No. 6, 92
Fontenay Aux Roses, FRANCE
- Jérôme L. Shapiro, Division Engineering and Applied Science, Pasadena,
California 91109
- R. D. Shelton, Code R-RP-N, National Aeronautics and Space Adm., Marshall
Space Flight Center, Huntsville, Alabama
- Robert T. Siegel, Department of Physics, College of William and Mary,
Williamsburg, Virginia 23185
- J. J. Singh, M.S. 234, NASA. Langley Research Center, Langley Station,
Hampton, Virginia 23365

- G. D. Smith, Ames Research Center, Moffett Field, California 94035
- Jerry Speakman, 6570 AMRL (MRBBR), Wright-Patterson AFB, Ohio
- Dwain F. Spencer, Jet Propulsion Laboratory, Pasadena, California 31103
- Stanford Linear Accelerator Center, ATTN: Library, P. O. Box 4349, Stanford, California 94305
- William Steigelmann, Kuljian Corp., 1200 North Broad Street, Philadelphia, Pennsylvania 19121
- Henry Stern, R-RP-N, NASA, Marshall Space Flight Center, Huntsville, Alabama 35812
- T. R. Strayhorn, S-71, General Dynamics, Fort Worth, Texas 76101
- S. Tom Taketa, Mail Stop N 236-5, NASA, Ames Research Center, Moffett Field, California 94035
- Eizo Tajima, Rikkyo University, Ikebukuro, Toshimaku, Tokyo, JAPAN
- Ralph H. Thomas, Rutherford High Energy Laboratory, Chilton, Didcot, Berks, ENGLAND
- O. Lyle Tiffany, Chief Scientist, Bendix Systems Division, 3300 Plymouth Road, Ann Arbor, Michigan 48103
- Cornelius Tobias, University of California, Lawrence Radiation Laboratory, Berkeley, California 94720
- Jacob I. Trombka, Code SM, National Aeronautics and Space Adm., Washington, D. C. 20546
- W. Turchinetz, Massachusetts Institute of Technology, 155 Massachusetts Avenue, Cambridge, Massachusetts
- Werner Von Braun, Director, George C. Marshall Space Flight Center, NASA, Huntsville, Alabama 35812
- G. P. Wachtell, Franklin Institute, 20th and Parkway, Philadelphia, Pennsylvania 19103
- Roger Wallace, University of California, Bldg. 72, Lawrence Radiation Laboratory, Berkeley, California 94720
- G. T. Western, Y-71, General Dynamics, Fort Worth, Texas 76101
- Glenn A. Whan, Associate Professor, Nuclear Engineering Laboratory, The University of New Mexico, Albuquerque, New Mexico 87106

Ralph Wiley, Mail Zone Y-42, P. O. Box 748, Fort Worth, Texas 76101

Maurice Wilkinson, The Boeing Company, M. S. 23-82, Seattle, Washington
98124

W. R. Yucker, A-2-833, Douglas Aircraft Co., Nuclear Department, 3000 Ocean
Park Boulevard, Santa Monica, California 90405

Marcello Zocchi, Reactor and Radiation, National Bureau of Standards,
Washington, D. C. 20234

K. Ziock, Department of Physics, University of Virginia, Charlottesville,
Virginia 32901

Internal Distribution

- | | | | |
|--------|----------------------|--------|----------------------------|
| 1-3. | L. S. Abbott | 48. | R. T. Santoro |
| 4. | F. S. Alsmiller | 49. | D. Sundberg |
| 5-29. | R. G. Alsmiller, Jr. | 50. | D. K. Trubey |
| 30. | T. W. Armstrong | 51. | J. W. Wachter |
| 31. | H. W. Bertini | 52. | W. Zobel |
| 32-36. | R. T. Boughner | 53. | E. R. Cohn (consultant) |
| 37. | C. E. Clifford | 54. | B. C. Diven (consultant) |
| 38. | T. A. Gabriel | 55. | W. N. Hess (consultant) |
| 39. | M. P. Guthrie | 56. | M. H. Kalos (consultant) |
| 40. | D. C. Irving | 57. | L. V. Spencer (consultant) |
| 41. | D. F. King | 58-59. | Central Research Library |
| 42. | W. E. Kinney | 60. | Document Reference Section |
| 43. | T. A. Love | 61-74. | Laboratory Records Dept. |
| 44-45. | F. C. Maienschein | 75. | Laboratory Records ORNL RC |
| 46. | H. S. Moran | 76. | ORNL Patent Office |
| 47. | R. W. Peelle | | |

External Distribution

- 77. P. B. Hemmig, Division of Reactor Development and Technology, U. S. Atomic Energy Commission, Washington, D. C.
- 78. I. F. Zartman, Reactor and Development, U. S. Atomic Energy Commission, Washington, D. C.
- 79. Herbert Goldstein, 242 Mudd Building, Columbia University, New York
- 80-255. Given NASA-High Energy Distribution as shown on preceding pages
- 256-270. Division of Technical Information Extension (DTIE)
- 271. Division of Reactor Development (ORO)